

# Complex Analysis

LIN150117

TSINGHUA UNIVERSITY.

linzj23@mails.tsinghua.edu.cn

November 4, 2024

## Contents

<b>2</b>	<b>Complex Functions</b>	<b>4</b>
2.1	Analytic functions and rational functions . . . . .	4
2.1.1	Harmonic function . . . . .	4
2.1.2	Polynomials and rational function . . . . .	4
2.2	Power Series . . . . .	7
2.2.1	Power series . . . . .	7
2.3	Exponential, Trigonometric and logarithmic functions . . . . .	7
2.3.1	Exponential and Trigonometric function . . . . .	7
2.3.2	Logarithmic Functions . . . . .	7
<b>3</b>	<b>Conformal Mappings</b>	<b>8</b>
3.1	Basic topology . . . . .	8
3.1.1	Connectedness . . . . .	8
3.1.2	Compactness . . . . .	8

3.1.3	Continuous Functions . . . . .	9
3.2	Conformality, geometric consequences of the existence of a derivative . . . . .	9
3.2.1	Arcs and closed curves . . . . .	9
3.2.2	Analytic Functions in Regions . . . . .	10
3.2.3	Conformal Mappings . . . . .	10
3.2.4	Length and Area . . . . .	11
3.3	Möbius Transformation . . . . .	11
3.3.1	Cross ratio . . . . .	12
3.3.2	Symmetry . . . . .	13
3.3.3	Steiner Circles, circular net . . . . .	15
<b>4</b>	<b>Elementary Conformal mapping</b>	<b>16</b>
4.1	Elementary Riemann surfaces . . . . .	17
<b>5</b>	<b>Complex Integration</b>	<b>18</b>
5.1	Fundamental Theorems . . . . .	18
5.1.1	Line integral and rectifiable arcs . . . . .	18
5.1.2	The fundamental theorem of Calculus for integrals in $\mathbb{C}$ . . . . .	19
5.1.3	Cauchy's theorem for a rectangle . . . . .	22
5.1.4	Cauchy's Theorem for a disk . . . . .	23
5.2	Cauchy's integral formula . . . . .	24
5.2.1	Index of a point with respect to a closed curve . . . . .	24
5.2.2	Cauchy's integral formula . . . . .	26
5.2.3	Higher derivatives . . . . .	27
5.2.4	Consequences of Cauchy . . . . .	28
5.3	Local properties of analytic functions . . . . .	31
5.3.1	Removable Singularities and Taylor's Theorem . . . . .	31
5.3.2	Zeros and poles . . . . .	33

5.3.3	The Local Mappings . . . . .	38
5.3.4	The Maximum Principle . . . . .	41
5.4	The General Form of Cauchy's Theorem . . . . .	42
5.4.1	Chains and Cycles . . . . .	42
<b>Index</b>		<b>43</b>
<b>List of Theorems</b>		<b>45</b>

## 2 Complex Functions

### 2.1 Analytic functions and rational functions

#### 2.1.1 Harmonic function

**Definition 2.1** (Cauchy-Riemann equation).

$$\frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

**Definition 2.2** (Harmonic function). A function  $u$  is **harmonic** if it satisfied **Laplace equation**  $\triangle u = 0$ .

If two harmonic function  $u$  and  $v$  satisfies Cauchy-Riemann equations, then we say that  $v$  is **conjugate harmonic function of  $u$**   $\Rightarrow u$  is conjugate harmonic of  $-v$ .

#### 2.1.2 Polynomials and rational function

The polynomial  $P(z) = \sum_{j=0}^n a_j z^j$  is analytic in  $\mathbb{C}$ .

We will prove the fundamental theorem of algebra

**Theorem 2.3** (Fundamental Theorem of Algebra). *Every polynomial with degree  $n > 0$  has at least one point.*

**Theorem 2.4** (Gauss-Lucus theorem). *The smallest convex polygon that contain the zeros of  $P$  also contains the zeros of  $P'$ .*

*Proof.* Only need to check.

We can get this equation.

$$\frac{P'(\alpha)}{P(\alpha)} = \sum_{j=1}^n \frac{1}{\alpha - \alpha_j} = 0 \Rightarrow \sum_{j=1}^n \frac{\overline{\alpha - \alpha_j}}{|\alpha - \alpha_j|^2}$$

Hence  $\alpha$  is linearly represented by  $\alpha_j$ . □

**Proposition 2.5.** *Let  $P$  and  $Q$  be two polynomial with no common zeros. Then the rational function  $R(z) = \frac{P(z)}{Q(z)}$  is analytic away from the zeros of  $Q$ .*

*The zeros of  $Q$  are called **poles** of  $R$ , and the **order of a pole** is equal to the order of the corresponding zero of  $Q$ .*

We often view  $R$  as a function from  $\hat{\mathbb{C}}$  to  $\hat{\mathbb{C}}$ .  $R_1(z) := R(\frac{1}{z})$ .

If  $R_1(0) = 0$ , the order of the zero at  $\infty$  (of  $R$ ) is the order of the zero of  $R_1(z)$  at  $z = 0$ .

If  $R_1(0) = \infty$ , the order of the pole at  $\infty$  (of  $R$ ) is the order of the pole of  $R_1(z)$  at  $z = 0$ .

Suppose

$$R(z) = \frac{a_n z^n + \cdots + a_1 z + a_0}{b_m z^m + \cdots + b_1 z + b_0}, a_n \neq 0, b_m \neq 0$$

Then

$$R_1(z) = z^{m-n} \frac{a_0 z^n + \cdots + a_n}{b_0 z^m + \cdots + b_m}$$

By discussing  $m$  and  $n$ , we can infer the situation of  $R(z)$  at  $\infty$ .

By adding the order of poles and zeros at  $\infty$ , we can get the following theorem.

**Theorem 2.6.** *The total number of zeros and poles of a rational function are the same.*

**Remark 2.7.** This common number is called the **order of the rational function**.

**Corollary 2.8.** *Suppose a rational function  $R$  has order  $p$ . Then every equation  $R(z) = a$  has exactly  $p$  roots.*

*Proof.*  $\hat{R}(z) = R(z) - a$  has the same poles as  $R$ . □

A rational function of order 1 is a **linear fraction**  $R(z) = \frac{az+b}{cz+d}, ad - bc \neq 0$

Such fraction is often called **Möbius transformation**

Every rational function has a representation by **partial fractions**.

- If  $R$  has a pole at  $\infty$ . Then we can write

$$R(z) = G(z) + H(z) \quad (*)$$

where  $G$  is a polynomial without constant term, and  $H$  is finite at  $\infty$ .

The degree of  $G$  is the order of the pole of  $R$  at  $\infty$ .  $G$  is called the **singular part** of  $R$  at  $\infty$ .

- Let the distinct finite poles of  $R$  be  $\beta_1, \dots, \beta_k$ . Let  $R_j(\psi) = R(\beta_j + \frac{1}{\psi})$ . Then  $R_j$  is a rational function with a pole at  $\infty$ . As in  $(*)$ , we can write

$$R_j = G_j + H_j$$

with  $H_j$  finite at  $\infty$ . Then

$$R(z) = G_j\left(\frac{1}{z - \beta_j}\right) + H\left(\frac{1}{z - \beta_j}\right)$$

with  $G_j$  is a polynomial in  $\frac{1}{z - \beta_j}$  without constant term called the **singular point** of  $R$  at  $\beta_j$ .

- Let  $F(z) = R(z) - G(z) - \sum_{j=1}^k G_j\left(\frac{1}{z - \beta_j}\right)$ .

Then  $F$  is a rational function which can only have poles among  $\beta_j, \infty$

Since by our construction,  $F$  is finite at every  $\beta_j, 1 \leq j \leq k$  and  $\infty$ .

So  $F$  is a constant.

In particular,  $R(z) = G(z) + \sum_{j=1}^k G_j\left(\frac{1}{z - \beta_j}\right) + c$ .

## 2.2 Power Series

### 2.2.1 Power series

**Theorem 2.9** (Abel's theorem). *If  $\sum a_n$  converges, then  $f(z) = \sum a_n z^n \rightarrow f(1)$  as  $z \rightarrow 1$  in such a way that  $\frac{|1-z|}{1-|z|}$  remains bounded.*

## 2.3 Exponential, Trigonometric and logarithmic functions

### 2.3.1 Exponential and Trigonometric function

The **exponential function** is defined as the solution of the differential equation

$$\begin{cases} f'(z) = f(z) \\ f(0) = 1 \end{cases}$$

We denote  $e^z = \exp z = \sum_{n=0}^{\infty} \frac{z^n}{n!}$ .

The **trigonometric function** are defined by

$$\cos z = \frac{e^{iz} + e^{-iz}}{2} \quad \sin z = \frac{e^{iz} - e^{-iz}}{2i}$$

### 2.3.2 Logarithmic Functions

The **logarithmic function**  $\ln$  is defined by  $z = \ln w$  is a root of the equation  $e^z = w$ .

For  $w \neq 0$ , we write  $z = x + iy$ , then

$$e^{x+iy} = w \Leftrightarrow \begin{cases} e^x = |w| \\ e^{iy} = \frac{w}{|w|} \end{cases}$$

The first equation has a unique solution  $x = \ln |w|$ .

The second equation  $e^{iy} = \frac{w}{|w|}$  has a unique solution  $y_0 \in [0, 2\pi)$ .

If we write  $w = re^{i\theta}$ , then  $x = \ln w$ ,  $y = \theta = \arg w$ .

Thus, for  $w \neq 0$ , we have

$$\ln w = \ln |w| + i \arg w$$

The function  $\ln$  is actually not single-valued. But we can define a single-valued function  $Ln$

We define

$$a^b = \exp(b \ln a)$$

We will prove  $Ln$  is analytic in  $\mathbb{C} - (-\infty, 0]$  but not continuous in  $(-\infty, 0]$ .

$Ln$  is the principal branch of the logarithm.

## 3 Conformal Mappings

### 3.1 Basic topology

#### 3.1.1 Connectedness

**Theorem 3.1.** *A nonempty open set in  $\mathbb{C}$  is connected iff any two of its points can be joined by a polygon which lies in the set, i.e. Connectedness is equivalent to Path Connectedness*

An nonempty connected subset is called a **region**

#### 3.1.2 Compactness

**Definition 3.2.** A set  $X$  is **totally bounded** if  $\forall \varepsilon > 0$ ,  $X$  can be covered by finitely many balls of radius  $\varepsilon$



**Theorem 3.3.** *A set is compact iff it is complete and totally bounded.*

**Theorem 3.4.** *A subset  $X \subset \mathbb{C}$  is compact iff every infinite sequence of  $X$  has a limit point in  $X$ .*

### 3.1.3 Continuous Functions

**Theorem 3.5.** *Continuous function maps connected space to connected space.*

**Theorem 3.6.** *Continuous function maps compact space to compact space.*

## 3.2 Conformality, geometric consequences of the existence of a derivative

### 3.2.1 Arcs and closed curves

The equation of an **arc**  $r$  in  $\mathbb{C}$  can be represented by one of the terms

- $x = x(t), y = y(t), \alpha \leq t \leq \beta, x, y$  are continuous at  $t$
- $z(t) = x(t) + iy(t), \alpha \leq t \leq \beta.$
- The continuous mapping  $\gamma : [\alpha, \beta] \rightarrow \mathbb{C}.$

For a non-decreasing function  $\varphi : [\alpha, \beta] \rightarrow [\alpha', \beta'], z = z(\varphi(t)), \alpha' \leq \tau \leq \beta'$  is **change of parameter** of  $z(t).$

The change is **reversible** iff  $\varphi$  is strictly increasing.

If  $\gamma$  is differentiable, then call  $\gamma$  a **curve**.

$\gamma$  is **simple**, or a **Jordan curve**, if  $\gamma$  is injective.

$\gamma$  is **closed curve** if  $\gamma(0) = \gamma(1).$

### 3.2.2 Analytic Functions in Regions

A function  $f$  is analytic on an arbitrary set  $A$  if it is the restriction to  $A$  of a function which is analytic in some open set containing  $A$ .

**Theorem 3.7.** *An analytic function in a region (i.e. open and connected)  $\Omega$  whose derivative is 0 must reduce to a constant. The same hold if the real part, the imaginary part, the modulus, or the argument is constant.*

### 3.2.3 Conformal Mappings

Suppose  $f : \Omega \rightarrow \mathbb{C}$  is analytic in  $\Omega$ .  $r_1 = z_1(t), r_2 = z_2(t), \alpha \leq t \leq \beta$ .

$$z_0 = z_1(t_0) = z_2(t'_0), z'_1(t_0) \neq 0, z'_2(\hat{t}_0) \neq 0, \alpha < t_0, \hat{t}_0 < \beta.$$

$$f'(z_0) \neq 0, w_1(t) = f(z_1(t)), w_2 = f(z_2(\hat{t}_0))$$

$$\Gamma_1 = \{w_1(t) | \alpha \leq t \leq \beta\}, \Gamma_2 = \{w_2(t) | \alpha \leq t \leq \beta\}$$

Then

$$w'_1(t) = f'(z_1(t))z'_1(t)$$

$$w'_2(t) = f'(z_2(\hat{t}))z'_2(\hat{t})$$

$\Rightarrow$

$$w'_1(t_0) \neq 0, w'_2(t_0) \neq 0$$

$$\arg w'_1(t_0) = \arg f'(z_1(t_0))z'_1(t_0)$$

$$\arg w'_2(t_0) = \arg f'(z_2(\hat{t}_0))z'_2(\hat{t}_0)$$

So the "angle"  $\arg w'_1(t_0) - \arg w'_2(t_0) = \arg z'_1(t_0) - \arg z'_2(\hat{t}_0)$  remains the same.

Now we give the definition.

**Definition 3.8.**  $w = f(z)$  is said to be **conformal** in  $\Omega$  if  $f$  is analytic in  $\Omega$  and  $f'(z) \neq 0$  for  $\forall z \in \Omega$ .

Easy to prove that linear change of scale at  $z_0$  is independent of the direction.

$$i.e. |f'(z_0)| = \lim_{z \rightarrow z_0} \frac{\delta\sigma}{\delta s}$$

### 3.2.4 Length and Area

The **length** of a differentiable arc  $\gamma$  with the equation  $z(t) = x(t) + iy(t)$ ,  $a \leq t \leq b$

$$L(\gamma) = \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt = \int_a^b |z'(t)| dt$$

For  $\Gamma = f(\gamma)$  where  $f$  conformal mapping.

Then

$$L(\Gamma) = \int_a^b |f'(z(t))| \cdot |z'(t)| dt$$

The **area** of  $E \subset \mathbb{R}$  is  $A(E) = \iint_E dx dy$

Then by the differentiable functional transformation, the area  $\hat{E} = f(E)$  is

$$A(\hat{E}) = \iint_E |u_x v_y - u_y v_x| dx dy$$

If  $f$  is the conformal mapping of an open set containing  $E$ , then by Cauchy-Riemann equation

$$A(\hat{E}) = \iint_E |f'(z)|^2 dx dy$$

## 3.3 Möbius Transformation

Recall that a **Möbius transformation** is a function of the form

$$w = s(z) = \frac{az + b}{cz + d}, \quad ad - bc \neq 0$$

Then it has an inverse  $z = S^{-1}(w) = \frac{dw - b}{-cw + a}$ .

We may define  $S(\infty) = \lim_{z \rightarrow \infty} S(z) = \frac{a}{c}$ ,  $S(\frac{-d}{c}) = \infty$

With these definition,  $S : \hat{\mathbb{C}} \rightarrow \hat{\mathbb{C}}$  is a topological mapping. Here one may use the chordal metric to define the topology.

$$S'(z) = \frac{ad - bc}{(cz + d)^2}$$

Then  $S$  is conformal in  $\hat{\mathbb{C}} - \{-\frac{d}{c}, \infty\}$ .

$w = z + \alpha$  is called a **parallel translation**.

$w = kz$  with  $|k| = 1$  is a **rotation**.

$w = kz$  with  $k > 0$  is a **homothetic transformation**.

$x = \frac{1}{z}$  is called an **inversion**.

**Proposition 3.9.** *Every Möbius transformation is a composition of the above four operations.*

### 3.3.1 Cross ratio

For three distinct points  $z_2, z_3, z_4 \in \hat{\mathbb{C}}$ , we can find a Möbius transformation  $S$  such that  $S(z_2) = 0$ ,  $S(z_3) = 1$ ,  $S(z_4) = \infty$ .

**Lemma 3.10.** *The Möbius transformation satisfying the above conditions is unique.*

The **cross ratio**  $(z_1, z_2, z_3, z_4)$  is the image  $z_1$  under the Möbius transformation which maps  $z_2$  to 1,  $z_3$  to 0 and  $z_4$  to  $\infty$ .

**Theorem 3.11.** *If  $z_1, z_2, z_3, z_4 \in \hat{\mathbb{C}}$  are distinct, and  $T$  is any Möbius transformation, then  $(Tz_1, Tz_2, Tz_3, Tz_4) = (z_1, z_2, z_3, z_4)$ .*

**Lemma 3.12.** *Let  $T$  be a Möbius transformation,  $T(\mathbb{R})$  is either a circle or a straight line.*

**Theorem 3.13.** *The cross ratio  $(z_1, z_2, z_3, z_4)$  is real iff the four points lie on a circle or a straight line.*

**Remark 3.14.** One may prove the theorem by elementary geometry

**Theorem 3.15.** *A Möbius transformation maps circles into circles.*

### 3.3.2 Symmetry

Suppose  $T$  is a Möbius transformation which maps  $\hat{\mathbb{R}}$  onto a circle  $C$ .

We say that  $w = Tz$  and  $w^* = T\bar{z}$  are **symmetric w.r.t.  $C$** .

**Remark 3.16.** This definition is independent of  $T$ . Suppose  $S$  is another Möbius transformation which maps  $\hat{\mathbb{R}}$  onto  $C$ , then  $S^{-1}T$  maps  $\hat{\mathbb{R}}$  to  $\hat{\mathbb{R}}$ , and this  $S^{-1}w = S^{-1}Tz$  and  $S^{-1}w^* = S^{-1}T\bar{z}$  are conjugate.

The points  $z$  and  $z^*$  are **symmetric w.r.t  $C$  through  $z_1, z_2, z_3$**  iff  $(z^*, z_1, z_2, z_3) = \overline{(z, z_1, z_2, z_3)}$ .

This can be another definition.

Note that only the points on  $C$  are symmetric to themselves.

The mapping  $z \mapsto z^*$  is 1-1 and is called **reflection w.r.t.  $C$** .

### Geometric Meaning of Symmetry

Case1:  $C$  is a straight line. We may assume  $z_3 = \infty$ .

$z, z^*$  are symmetric w.r.t.  $C$  if and only if

$$\frac{z^* - z_2}{z_1 - z_2} = \frac{\bar{z} - \bar{z}_2}{\bar{z}_1 - \bar{z}_2}$$

Then

$$|z^* - z_2| = |z - z_2|, \quad \forall z_2 \in C \text{ and } z_2 \neq \infty$$

$$\operatorname{Im} \frac{z^* - z_2}{z_1 - z_2} = \operatorname{Im} \frac{\bar{z} - \bar{z}_2}{\bar{z}_1 - \bar{z}_2}$$

So  $C$  is the bisecting normal of the segment between  $z$  and  $z^*$ .

Case2:  $C$  is the circle  $|z - a| = R$ .

$$\begin{aligned} \text{Then for } \forall \text{ distinct } z_1, z_2, z_3 \in \mathbb{C}, \overline{(z, z_1, z_2, z_3)} &= \overline{(z - a, z_1 - a, z_2 - a, z_3 - a)} \\ &= (\bar{z} - \bar{a}, \bar{z}_1 - \bar{a}, \bar{z}_2 - \bar{a}, \bar{z}_3 - \bar{a}) = (\bar{z} - \bar{a}, \frac{R^2}{z_1 - a}, \frac{R^2}{z_2 - a}, \frac{R^2}{z_3 - a}) = (\frac{R^2}{\bar{z} - \bar{a}}, z_1 - a, z_2 - a, z_3 - a) \\ &= (\frac{R^2}{\bar{z} - \bar{a}}, z_1, z_2, z_3). \end{aligned}$$

Then the symmetric point of  $z$  w.r.t.  $C$  is

$$z^* = \frac{R^2}{\bar{z} - \bar{a}} + a$$

or

$$(z^* - a)(\bar{z} - \bar{a}) = R^2$$

$\Rightarrow$

$$\begin{cases} |z^* - a| \cdot |z - a| = R^2 \\ \frac{z^* - a}{z - a} = \frac{(z^* - a)(\bar{z} - \bar{a})}{|z - a|^2} > 0 \end{cases}$$

**Theorem 3.17** (The Symmetric principle). *If a Möbius transformation maps a circle  $C_1$  onto a circle  $C_2$ , then it transforms any pair of symmetric points w.r.t.  $C_1$  into a pair of symmetric points w.r.t.  $C_2$ .*

*Proof.* Case1:  $C_1 = \hat{\mathbb{R}}$ . Let  $T$  be the Möbius transformation which maps  $\hat{\mathbb{R}}$  onto  $C_2$ .

$\forall z \in \mathbb{C}$ , by definition,  $w = Tz$  and  $w^* = T\bar{z}$  are symmetric w.r.t.  $C_2$ .

Case2:  $C_1$  is a general circle. Let  $T : C_1 \rightarrow C_2$  and  $S : \mathbb{R} \rightarrow C_2$  be Möbius transformation.

Suppose  $w, w^*$  are symmetric w.r.t.  $C_1$ . Then there exists  $z$  s.t.  $w = Sz, w^* =$

$S\bar{z}$ .

Then we can find  $Tw = TSz, Tw^* = TS\bar{z}$  are symmetric w.r.t.  $C_2$  since  $TS : \hat{\mathbb{R}} \rightarrow C_2$  □

**Remark 3.18.** (1). The Möbius transformation from  $C_1$  to  $C_2$  satisfies  $z_1 \mapsto w, z_2 \mapsto w_2, z_3 \mapsto w_3$  where  $z_1, z_2, z_3 \in C_1, w_1, w_2, w_3 \in C_2$  is given by

$$(w, w_1, w_2, w_3) = (z, z_1, z_2, z_3)$$

(2). The Möbius transformation from  $C_1$  to  $C_2$  satisfies  $z_1 \mapsto w_1, z_2 \mapsto w_2$  where  $z_1 \in C_1, z_2 \notin C_1, w_1 \in C_2, w_2 \notin C_2$  is given by

$$(w, w_1, w_2, w_2^*) = (z, z_1, z_2, z_2^*)$$

### 3.3.3 Steiner Circles, circular net

For  $S(z) = \frac{az + b}{cz + d}, S'(z) = \frac{ad - bc}{(cz + d)^2}$ .

A point  $z \notin$  a circle  $C$  is said to on the **right(left, resp.)** of  $C$  if  $\text{Im}(z, z_1, z_2, z_3) > 0(\text{Im}(z, z_1, z_2, z_3) < 0)$

**Remark 3.19.**

(1). This agrees with everyday use since  $(i, 1, 0, \infty) = i$

(2). This distinct between left and right is the same for all triples, while the meaning may be reversed.

(If  $C = \hat{\mathbb{R}}$ , then  $(z, z_1, z_2, z_3) = \frac{az + b}{cz + d}$  with  $a, b, c, d \in \mathbb{R} \Rightarrow \text{Im}(z, z_1, z_2, z_3) = \frac{ad - bc}{|cz + d|^2} \text{Im}(z)$ )

(3). We can define an absolute positive orientation of all finite circles by requiring that  $\infty$  should be lie to the right of the oriented circles.

Consider a Möbius transformation of the form

$$w = k \cdot \frac{z - a}{z - b}$$

Here,  $z = a \mapsto w = 0, z = b \mapsto w = \infty$ .

Then circles through  $a, b$  maps to straight line through  $0, \infty$ .

The concentric circle about the origin,  $|w| = \rho$ , correspond to circles with the equation

$$\left| \frac{z - a}{z - b} \right| = \frac{\rho}{|k|}$$

These are the circles of **Apollonius** with limit points  $a$  and  $b$ .

Denote by  $C_1$  the circles through  $a, b$  and  $C_2$  the circles of Apollonius with these limit points. The configuration formed by all the circles  $C_1$  and  $C_2$  is called the **Steiner circles**(or **circular net**)

**Theorem 3.20.**

- (a) *There is exactly one  $C_1$  and one  $C_2$  through each point in  $\hat{\mathbb{C}} \setminus \{a, b\}$*
- (b) *Every  $C_1$  meets every  $C_2$  under right angle.*
- (c) *Reflection in a  $C_1$  transforms every  $C_2$  into itself and every  $C_1$  into another  $C_1$ .*
- (d) *The limit points  $a, b$  are symmetric w.r.t. each  $C_2$ , but not w.r.t. other circles.*

*Proof.* If the limit points are  $0, \infty$ , those properties are trivial in the  $w$ -plane. The general case follows since all properties are invariant under Möbius transformations. □

## 4 Elementary Conformal mapping

**Example 4.1.**  $w = z^\alpha$  where  $\alpha > 0$ .



Let  $S(u_1, u_2)$  with  $0 < \varphi_2 - \varphi_1 \leq 2\pi$  be  $\{z \in \mathbb{C} : z \neq 0, \varphi_1 < \arg(z) < \varphi_2\}$  where  $\arg(z)$  can be chosen as any value of it.

Then  $S(\varphi_1, \varphi_2)$  is a region.

In this region, a unique value of  $w = z^\alpha$  is defined by  $\arg w = \alpha \arg z$ .

This function is analytic with  $\frac{dw}{dz} = \alpha \frac{w}{z}$ .

This function is 1-1 only if  $\alpha(\varphi_2 - \varphi_1) \leq 2\pi$ .

**Example 4.2.**  $w = e^z$  maps  $\{z \in \mathbb{C} : -\frac{\pi}{2} < \text{Im}(z) < \frac{\pi}{2}\}$  onto  $\{w \in \mathbb{C} : \text{Re}(w) > 0\}$

**Example 4.3.**  $w = \frac{z-1}{z+1}$  maps  $\{z \in \mathbb{C} : \text{Re}(z) > 0\}$  onto  $\{w \in \mathbb{C} : |w| < 1\}$

**Example 4.4.**

$$\mathbb{C} \setminus [-1, 1] \xrightarrow{z_1 = \frac{z+1}{z-1}} \mathbb{C} \setminus (-\infty, 0] \xrightarrow{z_2 = \sqrt{z_1}} \{\text{Re}(z_2) > 0\} \xrightarrow{w = \frac{z_2-1}{z_2+1}} \{w \in \mathbb{C} : |w| < 1\} \quad (4.1)$$

## 4.1 Elementary Riemann surfaces

**Example 4.5.**  $w = z^n, n \in \mathbb{Z}_+$  and  $n > 1$ .

There is a 1-1 correspondence between each angle  $\frac{(k-1)2\pi}{n} < \arg z < \frac{k \cdot 2\pi}{n}, k = 1, 2, \dots, n$  and while  $w$ -plane except for the positive real axis.

**Example 4.6.**  $w = e^z$ . This function maps each parallel strip  $(k-1)2\pi < \text{Im } z < k \cdot 2\pi, k \in \mathbb{Z}$  onto a sheet with a cut along the positive axis.

## 5 Complex Integration

### 5.1 Fundamental Theorems

#### 5.1.1 Line integral and rectifiable arcs

Let  $f(t) = u(t) + iv(t)$  be a complex-valued defined on  $t \in [a, b] \subset \mathbb{R}$  where  $u, v$  are real-valued functions. If  $f$  is continuous on  $[a, b]$ , we may define the **integral**

$$\int_a^b f(t)dt := \int_a^b u(t)dt + i \int_a^b v(t)dt$$

Let  $\gamma$  be a piecewise differential arc in  $\mathbb{C}$  with the equation  $z = z(t), a \leq t \leq b$ . If  $f$  is continuous on  $\gamma$ , then  $f(z(t))$  is continuous on  $[a, b]$ , and we define

$$\int_{\gamma} f(z)dz = \int_a^b f(z(t))z'(t)dt \quad (5.1)$$

The integral defined in 5.1 is independent of the parametrization of  $\gamma$ . Suppose that another parametrization of  $\gamma$  is  $\gamma : (\alpha, \beta) \rightarrow \mathbb{C}, \tau \mapsto z(t(\tau))$ , where  $t : (\alpha, \beta) \rightarrow (a, b), \tau \mapsto t(\tau)$  is piecewise differentiable. Then we have

$$\int_a^b f(z(t))z'(t)dt = \int_{\alpha}^{\beta} f(z(t(\tau)))z'(t(\tau))t'(\tau)d\tau = \int_{\alpha}^{\beta} f(z(t(\tau)))\frac{dz(t(\tau))}{d\tau}d\tau \quad (5.2)$$

For an arc  $\gamma$  with equation  $z = z(t), a \leq t \leq b$ , we define  $-\gamma$  by  $z = z(-t), -b \leq t \leq a$ .

Then we have

$$\int_{-\gamma} f(z)dz = \int_{-b}^{-a} f(z(-t))\frac{dz(-t)}{dt}dt$$

$$\begin{aligned}
&= - \int_{-a}^{-b} f(z(-t))z'(-t)dt \\
&= - \int_a^b f(z(\tau))z'(\tau)d\tau \\
&= - \int_{\gamma} f(z)dz
\end{aligned}$$

So we have those properties:

**Proposition 5.1.**

(a)  $\int_{-\gamma} f(z)dz = - \int_{\gamma} f(z)dz$

(b) Let  $f$  and  $g$  be two continuous functions on the piecewise differentiable arc  $\gamma$ , then

$$\int_{\gamma} (\lambda_1 f + \lambda_2 g)dz = \lambda_1 \int_{\gamma} f dz + \lambda_2 \int_{\gamma} g dz, \forall \lambda_1, \lambda_2 \in \mathbb{C}$$

(c) If  $\gamma$  can be subdivided into two pieces differentiable arcs  $\gamma_1$  and  $\gamma_2$ , and  $f$  is continuous on  $\gamma_1$ , then

$$\int_{\gamma} f dz = \int_{\gamma_1} f dz + \int_{\gamma_2} f dz$$

(d) (c) implies that the integral of a closed curve doesn't depend on the starting point on the curve

**Example 5.2.** Evaluate  $\int_{\gamma} \frac{1}{z-a} dz$  where  $\gamma$  is the circle centered at  $a \in \mathbb{C}$  with radius  $R$ .

Let  $z = z(t) = a + Re^{it}$ . Then the integral is  $2\pi i$

### 5.1.2 The fundamental theorem of Calculus for integrals in $\mathbb{C}$

The line integral w.r.t.  $\bar{z}$  is defined by

$$\int_{\gamma} f(z) d\bar{z} = \int_{\gamma} \overline{f(\bar{z})} dz$$

With this notation, line integrals w.r.t.  $x = \operatorname{Re}(z)$  and  $y = \operatorname{Im}(z)$  can be defined by

$$\int_{\gamma} f(z) dx = \frac{1}{2} \left[ \int_{\gamma} f(z) dz + \int_{\gamma} f(z) \overline{dz} \right]$$

$$\int_{\gamma} f(z) dy = \frac{1}{2i} \left[ \int_{\gamma} f(z) dz - \int_{\gamma} f(z) \overline{dz} \right]$$

if we write  $f(z) = \mu + i\nu$ , we have

$$\int_{\gamma} f(z) dz = \int_{\gamma} f(z) dx + i \int_{\gamma} f(z) dy = \int_{\gamma} (\mu dx - \nu dy) + i \int_{\gamma} (\nu dx + \mu dy)$$

**Remark 5.3.** It is followed by the intuition. We can view the integration as the multiplication between  $f$  and  $dz$ .

The integral w.r.t. **arc length** is defined by

$$\int_{\gamma} f(z) |dz| = \int_a^b f(z(t)) |z'(t)| dt$$

This integral is again independent of the parametrization. It is easy to check

$$\int_{-\gamma} f(z) |dz| = \int_{\gamma} f(z) |dz|$$

Now we define **length** of a curve  $\gamma$ :  $L(\gamma) = \int_{\gamma} |dz|$

We have the inequality:

$$\left| \int_{\gamma} f dz \right| \leq \int_{\gamma} |f| \cdot |dz| \leq L(\gamma) \cdot \sup_{z \in \gamma} |f(z)|$$

The length of an arc  $\gamma$  ( $z = z(t)$ ) can also be defined as the least upper bound of

all sums

$$\sum_{i=1}^n |z(t_i) - z(t_{i-1})|$$

where  $a = t_0 < t_1 < \dots < t_n = b$ . If this least upper bound is finite, we say that the arc is **rectifiable**.

It is easy to show that piecewise differentiable arcs are rectifiable.

The integral of a continuous function  $f$  on a rectifiable arc may be defined as

$$\int_{\gamma} f(z) dz = \lim \sum_{k=1}^n f(z(\psi_k)) [z(t_k) - z(t_{k-1})]$$

**Theorem 5.4.** *Let  $\Omega \subset \mathbb{C}$  be a region, and  $P, Q$  two (possibly complex-valued) functions that are continuous on  $\Omega$ ,  $\gamma$  closed curve. The integral  $\int_{\gamma} p(x, y) dx + Q(x, y) dy$  depends only on the end point of  $\gamma$  iff there exists a function  $U(x, y)$  on  $\Omega$  with  $\frac{\partial U}{\partial x} = P, \frac{\partial U}{\partial y} = Q$ .*

*Proof.* " $\Leftarrow$ ": If such a  $U$  exists, then

$$\int_{\gamma} P dx + Q dy = \int_{\gamma} \frac{\partial U}{\partial x} dx + \frac{\partial U}{\partial y} dy = \int_{\gamma} \frac{dU}{dt} dt = U(\gamma(b)) - U(\gamma(a))$$

" $\Rightarrow$ ": Fix a point  $(x_0, y_0) \in \Omega$ . We define  $U(x, y) = \int_{\gamma} P dx + Q dy$  where  $\gamma$  is any curve between  $(x_0, y_0)$  and  $(x, y)$ . Easy to check that it is true.  $\square$

**Theorem 5.5** (Fundamental theorem of Calculus for integrals on  $\mathbb{C}$ ). *Let  $f$  be continuous on a region  $\Omega$  containing  $\gamma$ .  $\int_{\gamma} f dz$  depends on the endpoints iff  $f$  is the derivative of an analytic function  $F$  in  $\Omega$ .*

**Remark 5.6.** We will prove  $\int_{\gamma} f dz = F(\omega_2) - F(\omega_1)$  where  $\gamma$  begins at  $\omega_1$  and ends at  $\omega_2$ .

*Proof.* Transform the line integration into the composition of two real integration.  $\square$

**Corollary 5.7.** *If  $F$  is analytic on  $\Omega$  with  $F' = f$ , and  $\gamma$  is a closed curve in  $\Omega$ , then  $\int_{\gamma} f dz = 0$ . Conversely if  $f$  is continuous on  $\Omega$  and  $\int_{\gamma} f dz = 0$  for any closed curve in  $\Omega$ , then  $f$  is the derivative of an analytic function  $F$  in  $\Omega$ .*

### 5.1.3 Cauchy's theorem for a rectangle

There is some notes in this section:

$R$  is the rectangle in  $\mathbb{C}$ ,  $R = \{x + iy \in \mathbb{C} : a \leq x \leq b, c \leq y \leq d\}$ . And  $\partial R$  is boundary curve oriented in the counterclockwise direction.

**Theorem 5.8** (Cauchy's theorem for a rectangle). *If  $f$  is analytic on an open set which contains  $R$ , then  $\int_{\partial R} f(z) dz = 0$*

*Proof.* For  $\forall$  rectangle  $\tilde{R}$  inside  $R$ , we define  $Z(\tilde{R}) = \int_{\partial \tilde{R}} f(z) dz$ . Then  $Z(R) = Z(R_1) + Z(R_2)$  if  $R$  is divided into  $Z_1, Z_2$ .

Since we can divide  $R$  into four equal rectangles, and find a rectangle with  $|Z(R^{(1)})| \geq \frac{1}{4}|Z(R)|$ . Then repeat the above steps and we obtain a sequence of nested rectangles  $R \supset R^{(1)} \supset \dots$  with the property

$$|Z(R^{(n)})| \geq \frac{1}{4}|Z(R^{(n-1)})| \geq \dots \geq \frac{1}{4^n}|Z(R)| \quad (5.3)$$

$\forall \delta > 0, \exists n \in \mathbb{N}$  s.t.  $R^{(n)} \subset \{z \in \mathbb{C} : |z - z_0| < \delta\}$ ,  $\forall n \geq N$ , where  $z_0$  is the limit of  $R^{(n)}$  as  $n \rightarrow \infty$ .

$f$  is analytic in  $R \Rightarrow \forall \varepsilon, \exists \delta > 0$  s.t.

$$\left| \frac{f(z) - f(z_0)}{z - z_0} - f'(z_0) \right| < \varepsilon, \forall z \text{ with } |z - z_0| < \delta \quad (5.4)$$

We assume that  $\delta$  satisfies both conditions. We have

$$Z(R^{(n)}) = \int_{\partial R^{(n)}} f(z) dz = \int_{\partial R^{(n)}} [f(z) - f(z_0) - (z - z_0)f'(z_0)] dz$$

$$\Rightarrow |Z(R^{(n)})| \leq \varepsilon \int_{\partial R^{(n)}} |z - z_0| dz \text{ by 5.4}$$

Let  $d_n$  be the length of diagonal of  $R^{(n)}$ ,  $L_n$  be the length of its perimeter. Then

$$|z - z_0| \leq d_n, \forall z \in \partial R^{(n)}.$$

$$\Rightarrow |Z(R^{(n)})| \leq \varepsilon d_n L_n = \varepsilon \frac{D}{2^n} \cdot \frac{L}{2^n} \text{ where } D, L \text{ are the diameter and perimeter of } R.$$

$$\Rightarrow |Z(R)| \stackrel{5.3}{\leq} 4^n |Z(R^{(n)})| \leq \varepsilon DL \Rightarrow Z(R) = 0 \text{ since } \varepsilon \text{ is arbitrary.} \quad \square$$

We will next prove the following stronger theorem:

**Theorem 5.9** (stronger version of Cauchy's theorem for a rectangle). *Let  $f$  be analytic on  $R' = R \setminus \{\psi_1, \dots, \psi_m\}$ ,  $m \in \mathbb{N}$ . If  $\lim_{z \rightarrow \psi_j} (z - \psi_j)f(z) = 0, \forall 1 \leq j \leq m$ , then*

$$\int_{\partial R} f(z) dz = 0.$$

*Proof.* WLOG, we may assume  $f$  is not analytic at only one point  $\psi \in R$ . If we put  $\psi$  into a small rectangle  $S_0$ , then the previous theorem tells us  $\int_{\partial R} f(z) dz = \int_{\partial S_0} f(z) dz$ .

$$\forall \varepsilon > 0, \text{ we may choose } S_0 \text{ small enough such that } |f(z)| \leq \frac{\varepsilon}{|z - \psi|}, \forall z \in \partial S_0$$

$$\Rightarrow \left| \int_{\partial R} f(z) dz \right| \leq \varepsilon \int_{\partial S_0} \frac{|dz|}{|z - \psi|} \leq \varepsilon \frac{1}{\frac{l}{2}} \cdot 4l = 8\varepsilon$$

$$\Rightarrow \int_{\partial R} f(z) dz = 0 \text{ since } \varepsilon \text{ is arbitrary.} \quad \square$$

### 5.1.4 Cauchy's Theorem for a disk

$$\Delta := \{z \in \mathbb{C} : |z - z_0| < R\} \text{ where } R > 0.$$

**Theorem 5.10** (Cauchy's Theorem for a disk). *If  $f$  is analytic in an open disk  $\Delta$ , then  $\int_{\gamma} f(z) dz = 0$  for closed curve  $\gamma$  in  $\Delta$ .*

*Proof.* Suppose the center of  $\Delta$  is  $z_0 = x_0 + iy_0$ ,  $z = x + iy$ . We define

$$F(z) = \int_{\gamma} f(z) dz$$

where  $\gamma$  is the horizontal line segment from  $z_0$  to  $(x, y_0)$  added with vertical line segment from  $(x, y_0)$  to  $z$ . We have

$$\frac{\partial F}{\partial y} = \lim_{\delta y \rightarrow 0} \frac{F(x, y + \delta y) - F(x, y)}{\delta y} = \lim_{\delta y \rightarrow 0} \frac{1}{\delta y} \int_{\delta \gamma} f(z) dz = i f(z) \quad (5.5)$$

By Cauchy's theorem on rectangles, one has  $F(z) = -\int_{\tilde{\gamma}} f(z) dz$ , where  $\tilde{\gamma}$  is the vertical line segment from  $z_0$  to  $(x_0, y)$  added with horizontal line segment from  $(x_0, y)$  to  $z$ .

Similarly,  $\frac{\partial F}{\partial x} = f(z)$ .

$\Rightarrow \frac{\partial F}{\partial x} = -i \frac{\partial F}{\partial y} \Rightarrow F$  is analytic in  $\Delta$  with derivative  $f$ . By Fundamental Theorem 5.5 of Calculus  $\Rightarrow \int_{\gamma} f(z) dz = 0$  for  $\forall$  closed curve in  $\Delta$ .  $\square$

Here is a stronger version.

**Theorem 5.11** (stronger version of Cauchy's Theorem for a disk). *Let  $f$  be analytic in a region  $\Delta' = \Delta \setminus \{\psi_1, \dots, \psi_m\}$  with  $m \in \mathbb{N}$ . If  $f$  satisfies  $\lim_{z \rightarrow \psi_j} (z - \psi_j) f(z) = 0, \forall 1 \leq j \leq m$ , then  $\int_{\gamma} f(z) dz = 0, \forall \gamma$  closed in  $\Delta'$*

*Proof.* It is similar to the above proof.

For the case no  $\psi_j$  lies on  $x = x_0$  and  $y = y_0$ , we can find a similar curve  $\gamma$  with last segment is a vertical one. Let  $F(z) = \int_{\gamma} f(z) dz$ . And continue the process of proof of the previous theorem.

For the case that  $\exists \psi_j$  lies on the lines  $x = x_0, y = y_0$ , we actually can move the center to another point s.t. no  $\psi_j$  lies on the lines  $x = x'_0, y = y'_0$ .  $\square$

## 5.2 Cauchy's integral formula

### 5.2.1 Index of a point with respect to a closed curve

**Lemma 5.12.** *If the piecewise differentiable closed curve  $\gamma$  does not pass through  $z \in \mathbb{C}$ , then the value of the integral  $\int_{\gamma} \frac{d\zeta}{\zeta - z}$  is a multiple of  $2\pi i$ .*



*Proof.*  $\gamma : \zeta = \zeta(t), \alpha \leq t \leq \beta$ .  $h(t) = \int_{\alpha}^t \frac{\zeta'(s)}{\zeta(s) - z} ds$ .

$z \in \gamma \Rightarrow h$  is defined and continuous on  $[\alpha, \beta]$ . For all  $t$  s.t.  $\zeta'(t)$  is continuous, we have

$$h'(t) = \frac{\zeta'(t)}{\zeta(t) - z} \Rightarrow \frac{d}{dt} [e^{-h(t)}(\zeta(t) - z)] = 0$$

So  $e^{-h(t)}(\zeta(t) - z)$  is constant on  $[\alpha, \beta]$ .

$$\text{Then } e^{h(t)} = \frac{\zeta(t) - z}{\zeta(\alpha) - z} \Rightarrow e^{h(\beta)} = 1 \Rightarrow h(\beta) \in \{2k\pi i : k \in \mathbb{Z}\}.$$

□

The **index of the point**  $z$  w.r.t. the closed curve  $\gamma$  is the number

$$n(\gamma, z) = \frac{1}{2\pi i} \int_{\gamma} \frac{d\zeta}{\zeta - z}$$

$n$  is also called the **winding number**.

**Theorem 5.13.** *Let  $\gamma$  be a piecewise differentiable closed curve. The function  $z \mapsto n(\gamma, z)$  is constant on each connected set of  $\mathbb{C} \setminus \gamma$ , and zero if this set is unbounded.*

*Proof.* Define  $f : \mathbb{C} \setminus \gamma \rightarrow \mathbb{C}, z \mapsto n(\gamma, z) = \frac{1}{2\pi i} \int_{\gamma} \frac{d\zeta}{\zeta - z}$ .

Then

$$|f(z) - f(z_0)| = \frac{1}{2\pi} \left| \int_{\gamma} \frac{z - z_0}{(\zeta - z)(\zeta - z_0)} d\zeta \right| \leq \frac{|z - z_0|}{2\pi} \int_{\gamma} \frac{1}{|\zeta - z| \cdot |\zeta - z_0|} |d\zeta|$$

$\Rightarrow f$  is continuous on each open connected set of  $\mathbb{C} \setminus \gamma$ . Let  $\Omega$  be any open connected set of  $\mathbb{C} \setminus \gamma$ . We have  $f(\Omega)$  is connected  $\xrightarrow{f(\Omega) \subset \mathbb{Z}} f(\Omega)$  contains at most one point  $\Rightarrow f$  is constant on  $\Omega$ .

If  $|z|$  is sufficient large,  $\exists$  a disk of radius  $R$ ,  $B(0, R)$ , s.t.  $\gamma \subset B(0, R)$  but  $z \notin B(0, R)$ . Cauchy's theorem for a disk 5.10 tells us that  $f(z) = n(\gamma, z) = 0$ . So it is zero if this set is unbounded. □

**Lemma 5.14.** *Let  $z_1, z_2$  be two points on a closed curve  $\gamma$  and  $0 \notin \gamma$ .*

Suppose  $z_1$  in the lower half space and  $z_2$  in upper half space. If  $\gamma_1 \cap \{(x, 0) : x \leq 0\} = \emptyset$ , and  $\gamma_2 \cap \{(x, 0) : x \geq 0\} = \emptyset$ , then  $n(\gamma, 0) = 1$ .

**Remark 5.15.** One method to prove this lemma is to create two segment from  $z_i$  to the point in the unit circle. By divide the curve into two parts, we can easily remove the part of previous curve by using the theorem 5.13, since 0 is in the unbounded set.

In this proof, we can find that Theorem 5.13 is such powerful that we can change any curve to a more simple curve easily!

### 5.2.2 Cauchy's integral formula

**Theorem 5.16** (Cauchy's integral formula). Suppose that  $f$  is analytic in an open disk  $\Delta$ , and let  $\gamma$  be a closed curve in  $\Delta$ . For  $\forall z \notin \gamma$ ,

$$n(\gamma, z)f(z) = \frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$$

where  $n(\gamma, z)$  is the index of  $z$  w.r.t.  $\gamma$ .

*Proof.* If  $z \notin \Delta$ , The both sides of the equation is 0.

So we may assume  $z \in \Delta$  and  $z \notin \gamma$ . Define  $F : \Delta \setminus \{z\} \rightarrow \mathbb{C}, \zeta \mapsto \frac{f(\zeta) - f(z)}{\zeta - z}$ .

Then  $F$  is analytic in  $\Delta \setminus \{z\}$ , and  $\lim_{\zeta \rightarrow z} (\zeta - z)F(\zeta) = f(z)$ .

By Cauchy's Theorem 5.9  $\Rightarrow \int_{\gamma} F(\zeta) d\zeta = 0 \Rightarrow \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta = f(z) \int_{\gamma} \frac{1}{\zeta - z} d\zeta = f(z) \cdot 2\pi i \cdot n(\gamma, z)$  □

**Remark 5.17.** This proof let us find that for a good-enough function, its integral over a closed curve is a constant.

The theorem still holds if  $f$  is analytic except at a finite number of  $\zeta_j$  s.t.

$$\lim_{\zeta \rightarrow \zeta_j} (\zeta - \zeta_j)f(\zeta) = 0$$

and  $z \neq \zeta_j$  for each  $j$ , since Cauchy's theorem is still applicable.

**Theorem 5.18** (The mean value property for analytic functions).  *$f$  is analytic in a region  $\Omega$  which contain  $\overline{B(z, R)}$ . Then*

$$f(z) = \frac{1}{2\pi} \int_0^{2\pi} f(z + Re^{i\theta}) dt$$

*Proof.* The previous theorem 5.16  $\Rightarrow$

$$f(z) = \frac{1}{2\pi i} \int_{|\zeta-z|=R} \frac{f(\zeta)}{\zeta - z} d\zeta \stackrel{\zeta=z+Re^{it}}{=} \frac{1}{2\pi} \int_0^{2\pi} f(z + Re^{it}) dt$$

□

If  $f$  is analytic in an open disk  $\Delta$ , and  $\gamma$  is a closed curve in  $\Delta$ . And  $n(\gamma, z) = 1$ .

Then

$$f(z) = \frac{1}{2\pi i} \int_{\gamma} \frac{f(\zeta)}{\zeta - z} d\zeta$$

This is usually referred to as **Cauchy's integral formula**

### 5.2.3 Higher derivatives

**Lemma 5.19.** *Let  $\Omega \subset \mathbb{C}$  be a region and  $\gamma$  be an arc in  $\Omega$ . If  $\varphi$  is continuous on  $\gamma$ , then the function*

$$F_n(z) := \int_{\gamma} \frac{\varphi(\zeta)}{(\zeta - z)^n} d\zeta$$

*is analytic in each of the regions  $\Omega \setminus \gamma$ , and its derivative is  $F'_n(z) = nF_{n+1}(z)$*

*Proof.* We prove it by induction.

The lemma is true if  $n = 0$ :  $F_0(z) = \int_{\gamma} \varphi(\zeta) d\zeta$  and  $F'_0(z) = 0 = 0 \cdot F_1(z)$ .

We suppose that the lemma holds for  $n - 1$  with  $n \in \mathbb{N}$ :  $\forall$  continuous  $\varphi$  on  $\gamma$ ,  $F_{n-1}$  is analytic in  $\Omega \setminus \gamma$  and  $F'_{n-1}(z) = (n - 1)F_n(z)$ ,  $\forall z \in \Omega \setminus \gamma$ .

Fix  $z_0 \in \Omega \setminus \gamma$ . For  $\forall z \in B(z_0, \frac{\delta}{2})$ , with  $B(z_0, \delta) \subset \Omega \setminus \gamma$ , we have  $|\zeta - z| > \frac{\delta}{2}$ ,  $\forall \zeta \in$

$\gamma$ .

For  $\forall$  continuous  $\varphi$  on  $\gamma$ ,

$$\begin{aligned} F_n(z) - F_n(z_0) &= \int_{\gamma} \frac{\varphi(\zeta)(\zeta - z + z - z_0)}{(\zeta - z)^n(\zeta - z_0)} d\zeta - \int_{\gamma} \frac{\varphi(\zeta)}{(\zeta - z_0)^n} d\zeta \\ &= \left[ \int_{\gamma} \frac{\varphi(\zeta)}{(\zeta - z)^{n-1}(\zeta - z_0)} d\zeta - \int_{\gamma} \frac{\varphi(\zeta)}{(\zeta - z_0)^{n-1}(\zeta - z)} d\zeta \right] \\ &\quad + (z - z_0) \int_{\gamma} \frac{\varphi(\zeta) d\zeta}{(\zeta - z)^n(\zeta - z_0)} \end{aligned}$$

Let  $\psi(\zeta) = \frac{\varphi(\zeta)}{\zeta - z_0}$ , which is continuous except  $\gamma$ .

Using the induction condition to  $\psi$ , we can finish the proof.  $\square$

**Theorem 5.20.** *An analytic function on a region  $\Omega$  has derivatives of all orders which are analytic in  $\Omega$ . More precisely,  $\forall z_0 \in \Omega$ , choose  $B(z, \delta) \subset \Omega$  and a circle  $C \subset B(z_0, \delta)$  with center  $z_0$ . For  $\forall z$  in the interior of  $C$ , Cauchy's integral formula gives*

$$f(z) = \frac{1}{2\pi i} \int_C \frac{f(\zeta)}{\zeta - z} d\zeta$$

Then the previous lemma implies  $f'(z) = \frac{1}{2\pi i} \int_C \frac{f(\zeta)}{(\zeta - z)^2} d\zeta$  is analytic in the interior of  $C$ . More generally, for  $\forall n \in \mathbb{N}$ ,

$$f^{(n)}(z) = \frac{n!}{2\pi i} \int_C \frac{f(\zeta)}{(\zeta - z)^{n+1}} d\zeta \quad (5.6)$$

## 5.2.4 Consequences of Cauchy

**Theorem 5.21 (Morera's Theorem).** *If  $f$  is continuous in a region  $\Omega$ , and if  $\int_{\gamma} f(z) dz = 0$  for  $\forall$  closed curve  $\gamma$  in  $\Omega$ . Then  $f$  is analytic in  $\Omega$ .*

*Proof.* We proved in Corollary 5.7 that under the hypothesis of theorem,  $f = F'$

where  $F$  is analytic in  $\Omega$ . The last theorem  $\Rightarrow f$  is analytic.  $\square$

Suppose  $f$  is analytic in a disk,  $\overline{B(z_0, R)}$ , and bounded on the circle  $\gamma$  given by  $|z - z_0| = R$ . Then  $\forall z \in \gamma, |f(z)| \leq M$  for some  $M \geq 0$ . By (5.6),

$$|f^{(n)}(z)| \leq \frac{n!}{2\pi} \int_C \frac{|f(\zeta)|}{|\zeta - z_0|^{n+1}} |d\zeta| \leq \frac{n!}{2\pi} \cdot \frac{M}{R^{n+1}} \cdot 2\pi R = MR^{-n}n! \quad (5.7)$$

This inequality is known as **Cauchy's estimate**.

**Theorem 5.22** (Liouville's Theorem). *A bounded entire function (i.e. analytic in  $\mathbb{C}$ ) is constant.*

*Proof.* Suppose  $|f(z)| \leq M, \forall z \in \mathbb{C}$ . Cauchy's estimate  $\Rightarrow$

$$|f'(z)| \leq \frac{M}{R}, \forall z \in \mathbb{C}, \forall R > 0 \quad (5.8)$$

$\square$

$$\xrightarrow{R \rightarrow \infty} f'(z) = 0 \text{ for } z \in \mathbb{C} \Rightarrow f = 0.$$

**Theorem 5.23** (Fundamental Theorem for Algebra). *Every polynomial of degree  $n \geq 1$  has  $n$  roots.*

*Proof.* It suffices to prove it has at least one root.

Suppose  $P(z) = a_n z^n + \cdots + a_1 z + a_0$  with  $a_0 \neq 0$  does not have a root.

Then  $f(z) := \frac{1}{P(z)}$  is an entire function. As  $z \rightarrow \infty, \lim_{|z| \rightarrow \infty} \frac{|P(z)|}{|z|^n} = |a_n| \Rightarrow$

$$\lim_{|z| \rightarrow \infty} \frac{1}{|P(z)|} = 0.$$

So  $f$  is bounded. By Liouville's Theorem,  $f$  is a constant. Where  $f = f(\infty) = 0$ .

That causes contradiction.  $\square$

**Theorem 5.24** (Power series). *If  $f$  is analytic in a region  $\Omega$  which contains a closed disk  $\overline{B(z_0, R)}$ , then  $f$  has a power series expansion at  $z_0$ ,*

$$f(z) = \sum_{n=0}^{\infty} \frac{f^{(n)}(z_0)}{n!} (z - z_0)^n, \quad \forall z \in B(z_0, R) \quad (5.9)$$

*Proof.*  $\forall z \in B(z_0, R), \forall \zeta$  with  $|\zeta - z_0| = R$ .

$$\begin{aligned} \frac{1}{\zeta - z} &= \frac{1}{(\zeta - z_0) - (z - z_0)} \\ &= \frac{1}{\zeta - z_0} \cdot \frac{1}{1 - \frac{z - z_0}{\zeta - z_0}} \\ &= \frac{1}{\zeta - z_0} \sum_{n=0}^{\infty} \left( \frac{z - z_0}{\zeta - z_0} \right)^n \\ &= \sum_{n=0}^{\infty} \frac{(z - z_0)^n}{(\zeta - z_0)^{n+1}} \end{aligned} \quad (5.10)$$

This series converges uniformly in  $\zeta$  with  $|\zeta - z_0| = R$ .

For  $\forall z \in B(z, R)$ ,

$$\begin{aligned} f(z) &= \frac{1}{2\pi i} \int_{|\zeta - z| = R} \frac{f(\zeta)}{\zeta - z} d\zeta \\ &= \frac{1}{2\pi i} \int_{|\zeta - z| = R} f(\zeta) \sum_{n=0}^{\infty} \frac{(z - z_0)^n}{(\zeta - z_0)^{n+1}} d\zeta \\ &\stackrel{\text{uniformly}}{=} \sum_{n=0}^{\infty} \frac{1}{2\pi i} \int_{|\zeta - z| = R} \frac{f(\zeta)}{(\zeta - z_0)^{n+1}} d\zeta \cdot (z - z_0)^n \\ &\stackrel{(5.6)}{=} \sum_{n=0}^{\infty} \frac{f^{(n)}(z_0)}{n!} (z - z_0)^n \end{aligned} \quad (5.11)$$

□

## 5.3 Local properties of analytic functions

### 5.3.1 Removable Singularities and Taylor's Theorem

We remarked that Cauchy's integral formula holds if  $f$  is analytic except at a finite number of point  $\zeta_j$  s.t.  $\lim_{\zeta \rightarrow \zeta_j} (\zeta - \zeta_j)f(\zeta) = 0$ . We will prove  $f$  can be extended to an analytic function in  $\Delta$ . In other word,  $\zeta_j$  are **removable singularities**.

**Theorem 5.25** (Riemann's Removable Singularities Theorem). *Suppose that  $f$  is analytic in the region  $\Omega' = \Omega \setminus \{\zeta_0\}$  where  $\Omega$  is also a region. Then there exists an analytic function in  $\Omega$  which coincides with  $f$  in  $\Omega'$  if and only if  $\lim_{z \rightarrow \zeta_0} (z - \zeta_0)f(z) = 0$ .*

*Proof.* The uniqueness and " $\Rightarrow$ " part is trivial since the extended function is continuous at  $\psi_0$ .

" $\Leftarrow$ ": Cauchy's integral formula  $\Rightarrow$

$$f(z) = \frac{1}{2\pi i} \int_C \frac{f(\zeta)}{\zeta - z} d\zeta, \quad \forall z \in \Delta \text{ and } z \neq \zeta_0 \quad (5.12)$$

Lemma 5.19  $\Rightarrow$  the RHS of the last equation 5.12 is analytic in  $z \in \Delta$ . Then

$$\hat{f}(z) = \begin{cases} f(z), & z \neq \zeta_0 \\ \frac{1}{2\pi i} \int_C \frac{f(\zeta)}{\zeta - \zeta_0} d\zeta, & z = \zeta_0 \end{cases} \quad (5.13)$$

is analytic in  $\Omega$ . □

We apply Theorem 5.25 to the function  $F(z) = \frac{f(z) - f(\zeta)}{z - \zeta}$ , where  $f$  is analytic in a region  $\Omega$ . Note that

$$\lim_{z \rightarrow \zeta_0} (z - \zeta)F(z) = 0, \quad \lim_{z \rightarrow \zeta} F(z) = f'(\zeta) \quad (5.14)$$

Theorem 5.25  $\Rightarrow \exists$  analytic function  $f_1$  on  $\Omega$  s.t.

$$f_1(z) = \begin{cases} F(z), & z \neq \zeta_0 \\ f'(\zeta), z = \zeta_0 \end{cases} \quad (5.15)$$

we may thus write  $f(z) = f(\zeta) + (z - \zeta)f_1(z)$ .

Repeating this process for  $f_1$ , we get an analytic function  $f_2$  on  $\Omega$  s.t.

$$f_1(z) = f_1(\zeta) + (z - \zeta)f_2(z) \quad (5.16)$$

where

$$f_2(z) = \begin{cases} \frac{f_1(z) - f_1(\zeta)}{z - \zeta}, & z \neq \zeta \\ f'_2(\zeta), & z = \zeta \end{cases} \quad (5.17)$$

Continuing the recursion, we have the general form

$$f_{n-1}(z) = f_{n-1}(\zeta) + (z - \zeta)f_n(z) \quad (5.18)$$

$\Rightarrow$

$$f(z) = f(\zeta) + (z - \zeta)f_1(\zeta) + \cdots + (z - \zeta)^{n-1}f_n(\zeta) + (z - \zeta)^n f_n(z) \quad (5.19)$$

Differentiating  $n$  times and setting  $z = \zeta \Rightarrow f^{(n)}(\zeta) = n!f_n(\zeta)$

We just prove **Taylor's Theorem**

**Theorem 5.26** (Taylor's Theorem). *If  $f$  is analytic in a region  $\Omega$ ,  $\zeta \in \Omega$ , then we have*

$$f(z) = f(\zeta) + (z - \zeta)f'(\zeta) + \cdots + \frac{f^{(n-1)}(\zeta)}{(n-1)!}(z - \zeta)^{n-1} + f_n(z)(z - \zeta)^n \quad (5.20)$$



where  $f_n$  is analytic in  $\Omega$ . Moreover,

$$f_n(z) = \frac{1}{2\pi i} \int_C \frac{f(\omega)}{(\omega - \zeta)^n(\omega - z)} d\omega \quad (5.21)$$

where  $C$  is a circle in  $\Omega$  s.t. its interior  $\Delta$  is also in  $\Omega$  and  $\zeta, z \in \Delta$

*Proof.* It suffices to prove the second part.

Cauchy's integral formula  $\Rightarrow f_n(z) = \frac{1}{2\pi i} \int_C \frac{f_n(\omega)}{\omega - z} d\omega, \forall z \in \Delta$ .

For  $f_n(z)$ , we substitute the expression from (5.20). The first term is

$$\frac{1}{2\pi i} \int_C \frac{f(\omega)}{(\omega - \zeta)^n(\omega - z)} d\omega \quad (5.22)$$

The remaining terms have the following form, except for constant factors:

$$g_k(\zeta) = \int_C \frac{1}{(\omega - \zeta)^n(\omega - z)} d\omega, \quad 1 \leq k \leq n \quad (5.23)$$

The lemma 5.19 applies to  $\varphi(\omega) = \frac{1}{\omega - z}, g'_k(\zeta) = k g_{k-1}(\zeta), k \in \mathbb{N}, \forall \zeta \in \Delta$ . So

$$\begin{aligned} g_1(\zeta) &= \int_C \frac{1}{(\omega - \zeta)(\omega - z)} d\omega \\ &= \frac{1}{\zeta - z} \left[ \int_C \frac{1}{\omega - \zeta} d\omega - \int_C \frac{1}{\omega - z} d\omega \right] \\ &= \frac{1}{\omega - z} [2\pi i - 2\pi i] = 0 \end{aligned} \quad (5.24)$$

So  $g_k(z) = 0, \forall k \in \mathbb{N}, \forall z \in \Delta$ . □

### 5.3.2 Zeros and poles

**Theorem 5.27.** If  $f$  is analytic in a region  $\Omega$  and  $\exists a \in \Omega$  s.t.  $f^{(n)}(a) = 0$  for  $\forall n \in \mathbb{N} \cup \{0\}$ , then  $f \equiv 0$  in  $\Omega$ .

*Proof.* Let  $B(a, R)$  be the disk s.t.  $\overline{B(a, R)} \subset \Omega$ . Let  $C = \partial B(0, R)$ .

Taylor's theorem  $\Rightarrow f(z) = (z - a)^n f_n(z)$  with

$$f_n(z) = \frac{1}{2\pi i} \int_C \frac{f(\omega)}{(\omega - a)^n (\omega - z)} d\omega, \forall n \in \mathbb{N} \cup \{0\}, \forall z \in B(a, R) \quad (5.25)$$

Let  $M = \max_{z \in C} |f(z)|$ .

$$\begin{aligned} \Rightarrow |f_n(z)| &\leq \frac{1}{2\pi} \cdot \frac{M}{R^n (R - |z - a|)} \cdot 2\pi R \\ \Rightarrow |f(z)| &\leq \frac{|z - a|^n}{R^n} \cdot \frac{MR}{R - |z - a|} \rightarrow 0 \text{ as } n \rightarrow \infty, \forall z \in B(0, R) \\ \Rightarrow f(z) &= 0, \forall z \in B(0, R) \end{aligned}$$

Now define

$$\begin{aligned} E_1 &= \{z \in \Omega \mid f^{(n)}(z) = 0, \forall n \in \mathbb{N} \cup \{0\}\} \\ E_2 &= \Omega \setminus E_1 = \{z \in \Omega \mid f^{(n)}(z) \neq 0, \text{ for some } n \in \mathbb{N} \cup \{0\}\} \end{aligned}$$

We just proved  $E_1$  is open.  $E_2$  is open because  $f^{(n)}$  is continuous in  $\Omega$  for  $\forall n \in \mathbb{N} \cup \{0\}$ .  $\Omega$  is a region  $\Rightarrow$  either  $R_1 = \emptyset$  or  $R_2 = \emptyset$ .

The assumption of the theorem  $\Rightarrow E_1 \neq \emptyset \Rightarrow E_1 = \Omega$ .  $\square$

Let  $f$  be analytic in  $\Omega$  which is not identically zero,  $f(a) = 0$  for some  $a \in \Omega$ . The previous theorem implies  $\exists$  first  $N \in \mathbb{N}$  s.t.  $f^{(N)}(a) \neq 0$ . Taylor's theorem implies that  $f(z) = (z - a)^N f_N(z)$  where  $f_N$  is analytic and  $f_N(a) \neq 0$ . We say that  $a$  is a **zero of order  $N$**  of  $f$ .

$f_N$  is continuous  $\Rightarrow \exists \delta > 0$  s.t.  $f(z) \neq 0$  for  $\forall z \in B(a, \delta) \setminus \{0\}$ .

So we have just proved an important result: Zeros of analytic functions are isolated, or equivalently, we have a famous theorem:

**Theorem 5.28 (Identity Theorem).** *If  $f$  and  $g$  are analytic in a region  $\omega$ , and  $f = g$  on a set which has an accumulation point in  $\Omega$ , then  $f(z) = g(z)$ .*

**Corollary 5.29.**

- (1) If  $f \equiv 0$  in a subregion of  $\Omega$  and  $f$  is analytic in  $\Omega$ , then  $f \equiv 0$  in  $\Omega$ .
- (2) If  $f$  is analytic in  $\Omega$  and vanishes on an arc in  $\Omega$  which doesn't reduce to a point, then  $f \equiv 0$  in  $\Omega$ .

If  $f$  is analytic in a neighborhood of  $a$ , but perhaps not at  $a$  itself, then  $a$  is called an **isolated singularity** of  $f$ .

If  $\lim_{z \rightarrow a} f(z) = \infty$ , then  $a$  is said to be a **pole** of  $f$ , and we set  $f(a) = \infty$ . Continuity implies  $\exists \delta > 0$  s.t.  $f(z) \neq 0$  for  $\forall z \in B(0, \delta) \setminus \{a\}$ . Thus,  $g(z) = \frac{1}{f(z)}$  is analytic in  $B(a, \delta) \setminus \{a\}$ .  $\lim_{z \rightarrow a} (z - a)g(z) = 0 \Rightarrow a$  is a removable singularity of  $g$ , and  $g$  has an analytic extension with  $g(a) = 0$ .  $g \not\equiv 0 \Rightarrow a$  is a zero of  $g$  with finite order. The **order of the pole** of  $f$  at  $a$  is the order  $N$  of the zero of  $g$  at  $a$ .

We can write

$$f(z) = (z - a)^{-N} f_N(z), \quad \forall z \in B(a, \delta) \setminus \{a\} \quad (5.26)$$

where  $f_N$  is analytic and nonzero in a neighborhood of  $a$ .

**Definition 5.30.** A function which is analytic in a region  $\Omega$  except for (isolated) poles is called a **meromorphic function**.

**Example 5.31.** If  $f$  and  $g$  are analytic in  $\Omega$  and  $g \not\equiv 0$ , then  $\frac{f}{g}$  is a meromorphic function in  $\Omega$ . (See the Identity Theorem 5.28)

**Remark 5.32.** The sum, the product and quotient (if denominator is not always zero) of two meromorphic functions are meromorphic.

If  $f$  has a pole of order  $N$  at  $a$ , then  $(z - a)^N f(z)$  is analytic at  $a$ , and Taylor's theorem 5.26 implies

$$(z - a)^N f(z) = b_N + b_{N-1}(z - a) + \cdots + b_1(z - a)^{N-1} + \varphi(z) \cdot (z - a)^N \quad (5.27)$$

where  $\varphi$  is analytic at  $a$ .

$$\Rightarrow f(z) = b_N(z-a)^{-N} + b_{N-1}(z-a)^{-(N-1)} + \cdots + b_1(z-a)^{-1} + \varphi(z), \forall z \neq a. \quad (5.28)$$

**Theorem 5.33.** *If  $f$  is analytic in a neighborhood of  $a$ , but perhaps not at  $a$  itself, then exactly one of the following 3 cases occurs:*

(i)  $f \equiv 0$  in this neighborhood.

$$(ii) \exists \text{ integer } N \in \mathbb{Z} \text{ s.t. } \lim_{z \rightarrow a} |z-a|^\alpha \cdot |f(z)| = \begin{cases} 0, & \alpha > N \\ \infty, & \alpha < N \end{cases}$$

(iii) neither  $\lim_{z \rightarrow a} |z-a|^\alpha \cdot |f(z)| = 0$  for any  $\alpha \in \mathbb{R}$  nor  $\lim_{z \rightarrow a} |z-a|^\alpha \cdot |f(z)| = \infty$  for any  $\alpha \in \mathbb{R}$

*Proof.*

① If  $\lim_{z \rightarrow a} |z-a|^\alpha \cdot |f(z)| = 0$  for  $\forall \alpha \in \mathbb{R}$ , then  $\lim_{z \rightarrow a} |z-a|^m \cdot |f(z)| = 0$  for  $\forall$  integer  $m > \alpha$ .

$\Rightarrow (z-a)^m f(z)$  has a removable singularity at  $a$  and vanishes at  $z = a$

$\Rightarrow$  Either  $f \equiv 0$  in  $B(a, \delta) \setminus \{a\}$ , which is case (i), or  $(z-a)^m f(z)$  has a zero of

finite order  $k$  at  $a \Rightarrow \lim_{z \rightarrow a} |z-a|^\alpha \cdot |f(z)| = \begin{cases} 0, & \alpha > m-k \\ \infty, & \alpha < m-k \end{cases}$

② If  $\lim_{z \rightarrow a} |z-a|^\alpha |f(z)| = \infty$  for some  $\alpha \in \mathbb{R}$ , then  $\lim_{z \rightarrow a} |z-a|^n \cdot |f(z)| = \infty$  for  $\forall$  integer  $n < \alpha$ .

$\Rightarrow (z-a)^n f(z)$  has a pole of finite order  $l$  at  $a$

$$\Rightarrow \lim_{z \rightarrow a} |z-a|^\alpha \cdot |f(z)| = \begin{cases} 0, & \alpha > n+l \\ \infty, & \alpha < n+l \end{cases}$$

□

**Remark 5.34.** In case (ii),  $N$  may be called the **algebraic order** of  $f$  at  $a$ .  $N > 0$  if  $a$  is a pole,  $N < 0$  if  $a$  is a zero, and  $N = 0$  if  $f$  is analytic at  $a$  and  $f(a) \neq 0$ . The order is always an integer, there is no analytic function which tends to 0 or  $\infty$ , like a fractional power of  $|z - a|$ .

In some sense, three cases depends on whether  $\lim_{z \rightarrow a} (z - a)^N f(z)$  converges for some  $N$ .

In case (iii), the point  $a$  is an **essential isolated singularity**.

**Example 5.35.**  $f(z) = \exp(\frac{1}{z})$  has an essential isolated singularity  $z = 0$ .

**Theorem 5.36** (Weierstrass). *An analytic function comes arbitrarily close to any complex value in every neighborhood of an essential singularity. Or equivalently, the codomain of  $f$  on every neighborhood of an essential singularity is dense in  $\mathbb{C}$ .*

*Proof.* Suppose the statement is false.

$\exists A \in \mathbb{C}, \delta > 0$  and  $\varepsilon > 0$  s.t.

$$|f(z) - A| > \delta, \forall z \text{ with } 0 < |z - a| < \varepsilon \quad (5.29)$$

$\Rightarrow \lim_{z \rightarrow a} |z - a|^\alpha \cdot |f(z) - A| = \infty$  for  $\forall \alpha < 0$ .  $\Rightarrow a$  is not an essential singularity of  $f(z) - A$ .

The previous theorem  $\Rightarrow \exists \beta \in \mathbb{R}$  s.t.  $\lim_{z \rightarrow a} |z - a|^\beta \cdot |f(z) - A| = 0$ , and we may choose  $\beta > 0$ .

Then  $\lim_{z \rightarrow a} |z - a|^\beta \cdot |A| = 0 \Rightarrow \lim_{z \rightarrow a} |z - a|^\beta \cdot |f(z)| = 0$  by the triangular inequality.

So  $a$  is not an essential singularity of  $f$ , which causes contradiction!

So the statement has to be true. □

**Remark 5.37.** If  $f$  is analytic in  $|z| > R$ . We treat  $\infty$  as an isolated singularity. Removable singularity, pole or essential singularity of  $f$  at  $\infty$  is defined according to  $g(z) = f(\frac{1}{z})$  at  $z = 0$ .

### 5.3.3 The Local Mappings

**Theorem 5.38** (The Argument Principle). *Let  $f$  be analytic in a disk  $\Delta$  s.t.  $f$  does not vanish identically. Let  $z_j$  be the zeros of  $f$ , each zero being counted as many times as its order indicates. For every closed curve  $\gamma$  in  $\Delta$  which does not pass through a zero, we have*

$$\sum_j n(\gamma, z_j) = \frac{1}{2\pi i} \int_{\gamma} \frac{f'(z)}{f(z)} dz \quad (5.30)$$

where the sum has only a finite number of terms with nonzero value.

*Proof.*

Case I:  $f$  has exactly  $n$  zeros  $z_1, \dots, z_n$ .

By repeated application of Taylor' Theorem 5.26, we can write

$$f(z) = (z - z_1)(z - z_2) \cdots (z - z_n)g(z), \quad z \in \Delta \quad (5.31)$$

where  $g$  is analytic in  $\Delta$  and  $g(z) \neq 0$  for  $\forall z \in \Delta$ .  $\Rightarrow$

$$\frac{f'(z)}{f(z)} = \frac{1}{z - z_1} + \frac{1}{z - z_2} + \cdots + \frac{1}{z - z_n} + \frac{g'(z)}{g(z)}, \quad \forall z \in \Delta \text{ and } z \neq z_j \quad (5.32)$$

Cauchy' Theorem 5.10  $\Rightarrow$

$$\int_{\gamma} \frac{g'(z)}{g(z)} dz = 0 \Rightarrow \frac{1}{2\pi i} \int_{\gamma} \frac{f'(z)}{f(z)} dz = \sum_{j=1}^n n(\gamma, z_j) \quad (5.33)$$

Case II:  $f$  has infinitely many zeros in  $\Delta$ . Then  $\gamma$  is inside a concentric disk  $\Delta'$  smaller than  $\Delta$ .

$f \not\equiv 0 \Rightarrow$  There is only a finite number of zeros in  $\Delta'$ .

So we can apply (5.33) to the disk  $\Delta' \Rightarrow$  (5.30) holds since  $n(\gamma, z_j) = 0$  if  $z \notin \Delta'$ . □

**Remark 5.39.**

- The function  $\omega = f(z)$  maps  $\gamma$  onto a closed curve  $\Gamma$  in the  $\omega$ -plane, and we have

$$\int_{\Gamma} \frac{d\omega}{\omega} = \int_{\gamma} \frac{f'(z)}{f(z)} dz \quad (5.34)$$

Then (5.30) can be interpreted as  $n(\Gamma, 0) = \sum_j n(\gamma, z_j)$ .

- The most useful application of the theorem is to the case when  $\gamma$  is a circle (or more generally a simple closed curve). So that

$$n(\gamma, z) = \begin{cases} 1, & z \text{ is inside } \gamma \\ 0, & z \text{ is outside } \gamma \end{cases} \quad \text{Then (5.30) yields a formula for the total number of zeros enclosed by } \gamma.$$

Let  $a \in \mathbb{C}$ . Apply the previous theorem to  $f(z) - a$

$$\sum_j n(\gamma, z_j(a)) = \frac{1}{2\pi i} \int_{\gamma} \frac{f'(z)}{f(z) - a} dz$$

where  $z_j(a)$  are zeros of  $f - a$  (or roots of  $f(z) = a$ ), and  $\gamma$  is a closed curve in  $\Delta$  which doesn't pass  $z_j(a) \Rightarrow$

$$n(\Gamma, a) = \sum_j n(\gamma, z_j(a))$$

If  $a$  and  $b$  are in the same region determined by  $\Gamma$ , then  $n(\Gamma, a) = n(\Gamma, b) \Rightarrow$

$$\sum_j n(\gamma, z_j(a)) = \sum_j n(\gamma, z_j(b)) \quad (5.35)$$

If  $\gamma$  is a circle, then  $f$  takes the values  $a$  and  $b$  equally many times inside  $\gamma$ , counted as many times as their orders indicate.

We have the equation that

$$\begin{aligned}
\frac{1}{2\pi i} \int_{\gamma} \frac{f'(z)}{f(z) - a} dz &= n(\Gamma, a) = n(\Gamma, b) \\
&= \frac{1}{2\pi i} \int_{\Gamma} \frac{d\omega}{\omega - b} = \frac{1}{2\pi i} \frac{f'(z) dz}{f(z) - b} \\
&= \text{card}\{z \text{ inside } \gamma : f(z) = b\}
\end{aligned} \tag{5.36}$$

**Theorem 5.40.** Suppose  $f$  is analytic at  $z_0$ , and  $f(z) - \omega_0$  has a zero of order  $N \in \mathbb{N}$  at  $z_0$ . Then for  $\forall \varepsilon > 0$  sufficiently small,  $\exists \delta > 0$  s.t. for  $\forall a$  with  $|a - \omega_0| < \delta$ , the equation  $f(z) = a$  has exactly  $N$  roots in the disk  $|z - z_0| < \varepsilon$

*Proof.* We choose  $\varepsilon > 0$  s.t.

- (1)  $f$  is analytic in  $|z - z_0| \leq \varepsilon$
- (2)  $z_0$  is the only zero of  $f(z) - \omega_0$  in this disk.
- (3)  $f'(z) \neq 0$  for  $\forall z$  with  $0 < |z - z_0| < \varepsilon$

Let  $\gamma$  be the circle  $|z - z_0| < \varepsilon$  and  $\Gamma = f \circ \gamma$ .

$$\omega_0 \notin \Gamma \Rightarrow \exists \delta > 0 \text{ s.t. } B(\omega_0, \delta) \cap \Gamma = \emptyset.$$

The consequence of the argument principle 5.38, i.e. (5.36)  $\Rightarrow f$  takes all values  $a \in B(\omega_0, \delta)$  the same number of times  $N$  inside  $\gamma$ , since  $f(z) = \omega_0$  has exactly  $N$  coinciding roots inside  $\gamma$ .

$$(3) \Rightarrow \text{all roots } f(z) = a \text{ with } a \in B(\omega_0, \delta) \setminus \{\omega_0\} \text{ are simple} \quad \square$$

**Corollary 5.41** (open mapping theorem). A nonconstant analytic function maps open sets onto open sets.

*Proof.* The previous theorem  $\Rightarrow \forall \varepsilon > 0, f(B(z_0, \varepsilon)) \supset B(\omega_0, \delta)$   $\square$



**Corollary 5.42.** *If  $f$  is analytic at  $z_0$  with  $f'(z_0) \neq 0$ . It maps a neighborhood of  $z_0$  conformally and topologically onto a region.*

*Proof.* This is the case  $N = 0$ . The previous theorem  $\Rightarrow$  There is 1-1 corresponding between the disk  $|\omega - \omega_0| < \delta$  and an open subset of  $|z - z_0| < \varepsilon$ . The open mapping theorem 5.41  $\Rightarrow f^{-1}$  is continuous  $\Rightarrow f$  is a topological map. And  $f$  is conformal on  $|z - z_0| < \varepsilon$   $\square$

**Remark 5.43.** Under the assumption of Corollary 5.42,  $f^{-1}$  is continuous  $\Rightarrow f^{-1}$  is analytic  $\Rightarrow f^{-1}$  is conformal map.

If  $f : \Omega \rightarrow \mathbb{C}$  is 1-1 and analytic, Theorem 5.40 can hold only with  $N = 1 \Rightarrow f'(z) \neq 0$  for  $\forall z \in \mathbb{C}$ . So this condition is stronger than the conformal condition.

### 5.3.4 The Maximum Principle

**Theorem 5.44** (The maximum principle). *If  $f$  is analytic and nonconstant in a region  $\Omega$ , then its modulus  $|f|$  has no maximum in  $\Omega$ .*

*Proof.*  $\forall z_0 \in \Omega$ , the open mapping theorem 5.41  $\Rightarrow \exists$  an open disk  $|\omega - f(z_0)| < \delta$  contained in  $F(\Omega)$ . In this disk,  $\exists \omega$  s.t.  $|\omega| > |f(z_0)| \Rightarrow |f(z_0)|$  is not the maximum of  $|f|$ .  $\square$

**Theorem 5.45** (The maximum principle). *If  $f$  is defined and continuous on a closed bounded set  $E$  and analytic in the interior of  $E$ , then the maximum of  $|f|$  on  $E$  is assumed on the boundary of  $E$ .*

**Remark 5.46.** The maximum principle can also be proved by the mean value theorem 5.18 for analytic functions.

**Theorem 5.47** (Schwarz Lemma). *If  $f$  is analytic in the disk  $|z| < 1$  and satisfies  $f(0) = 0$ ,  $|f(z)| \leq 1$ ,  $\forall z \in B(0, 1)$ , then  $|f(z)| \leq |z|$  and  $|f'(0)| \leq 1$ . Furthermore, if  $|f(z)| = |z|$  for some  $z \neq 0$ , or if  $|f'(0)| = 1$ , then  $f(z) = cz$  where  $c \in \mathbb{C}$  with  $|c| = 1$ .*

*Proof.* We define  $g(z) \begin{cases} \frac{f(z)}{z}, & z \neq 0, z \in B(0, 1) \\ f'(0), & z = 0 \end{cases}$ .

Then  $g$  is analytic with  $g'(0) = \frac{f'(0)}{2}$  using Taylor series (5.20).

The maximum principle implies that  $|g'(z)| \leq \frac{1}{r}, \forall z \in \overline{B(0, r)}$  where  $0 < r < 1$ .

Setting  $r \rightarrow 1$ , we get  $|g(z)| \leq 1, \forall |z| < 1$ .

If  $|f(z)| = |z|$  for some  $z \neq 0$ , or  $|f'(0)| = 1$ , then  $|g| = 1$  attains its maximum at some interior points. By maximum principle,  $g$  has to be a constant.  $\square$

**Remark 5.48.** For a general analytic function  $f : B(0, R) \rightarrow B(0, M), z_0 \mapsto w_0$ .

$$\text{Let } T(z) = \frac{\frac{z}{R} - \frac{z_0}{R}}{1 - \frac{\bar{z}_0}{R} \cdot \frac{z}{R}}$$

$$S(\omega) = \frac{\frac{\omega}{M} - \frac{\omega_0}{M}}{1 - \frac{\bar{\omega}_0}{M} \cdot \frac{\omega}{M}}.$$

Then  $S \circ f \circ T^{-1}$  satisfies  $S \circ f \circ T^{-1}(0) = 0$  and  $|S \circ f \circ T^{-1}(z)| \leq 1 \xrightarrow{\text{Schwarz lemma}}$

$$|S \circ f \circ T^{-1}(\zeta)| \leq |\zeta|.$$

$$\Rightarrow |S \circ f(z)| \leq |T(z)| \Rightarrow$$

$$\left| \frac{M(f(z) - \omega_0)}{M^2 - \bar{\omega}_0 f(z)} \right| \leq \left| \frac{R(z - z_0)}{R^2 - \bar{z}_0 z} \right|, \forall z \in B(0, R)$$

## 5.4 The General Form of Cauchy's Theorem

### 5.4.1 Chains and Cycles

Let  $\Omega \subset \mathbb{C}$  be open. Let  $\gamma_j : [\alpha_j, \beta_j] \rightarrow \Omega$  be piecewise continuously differentiable curves in  $\Omega$ . The sum  $\gamma_1 + \gamma_2 + \cdots + \gamma_N$ , which need not be a curve is called a **chain**. The **integral** of a continuous  $f$  in  $\Omega$  along this chain is defined by

$$\int_{\gamma_1 + \gamma_2 + \cdots + \gamma_N} f = \sum_{j=1}^N \int_{\gamma_j} f. \quad (5.37)$$

Two chains are **identical** if they yield the same line integrals for all function  $f$ .

# Index

- algebraic order, 37
- Apollonius, 16
- arc, 9
- area, 11
  
- Cauchy's estimate, 29
- Cauchy's integral formula, 27
- chain, 42
  - identical, 42
  - integral, 42
- change of parameter, 9
- circular net, 16
- conformal, 11
- conjugate harmonic function of  $u$  , 4
- cross ratio, 12
- curve, 9
  - closed curve, 9
  - Jordan curve, 9
  - simple , 9
  
- essential isolated singularity, 37
- exponential function , 7
  
- harmonic, 4
- homothetic transformation, 12
  
- index of the point  $z$ , 25
- integral, 18
  - $\bar{z}$ , 19
  - arc length, 20
  - inversion, 12
  - isolated singularity, 35
  
- Laplace equation, 4
- left, 15
- length, 11, 20
- linear fraction, 5
- logarithmic function, 7
  
- meromorphic function, 35
- Möbius transformation, 5, 11
  
- order of a pole, 5
- order of the pole, 35
- order of the rational function, 5
  
- parallel translation, 12
- partial fractions, 5
- pole, 35
- poles, 5
  
- rectifiable, 21
- reflection, 13
- region, 8
- removable singularities, 31
- reversible , 9
- right, 15

rotation, [12](#)

singular part, [6](#)

singular point, [6](#)

Steiner circles, [16](#)

symmetric, [13](#)

symmetric w.r.t  $C$  through  $z_1, z_2, z_3$ , [13](#)

Taylor's Theorem, [32](#)

totally bounded , [8](#)

trigonometric function, [7](#)

winding number, [25](#)

zero of order  $N$ , [34](#)

# List of Theorems

2.3	Theorem (Fundamental Theorem of Algebra)	4
2.4	Theorem (Gauss-Lucus theorem)	4
2.5	Proposition	5
2.6	Theorem	5
2.9	Theorem (Abel’s theorem)	7
3.1	Theorem	8
3.3	Theorem	9
3.4	Theorem	9
3.5	Theorem	9
3.6	Theorem	9
3.7	Theorem	10
3.9	Proposition	12
3.11	Theorem	12
3.13	Theorem	13
3.15	Theorem	13
3.17	Theorem (The Symmetric principle)	14
3.20	Theorem	16
5.1	Proposition	19
5.4	Theorem	21
5.5	Theorem (Fundamental theorem of Calculus for integrals on $\mathbb{C}$ )	21
5.8	Theorem (Cauch’s theorem for a rectangle)	22
5.9	Theorem (stronger version of Cauchy’s theorem for a rectangle)	23
5.10	Theorem (Cauchy’s Theorem for a disk)	23
5.11	Theorem (stronger version of Cauchy’s Theorem for a disk)	24
5.13	Theorem	25

5.16 Theorem (Cauch's integral formula) . . . . .	26
5.18 Theorem (The mean value property for analytic functions) . . . . .	27
5.20 Theorem . . . . .	28
5.21 Theorem (Morera's Theorem) . . . . .	28
5.22 Theorem (Liouville's Theorem) . . . . .	29
5.23 Theorem (Fundamental Theorem for Algebra) . . . . .	29
5.24 Theorem (Power series) . . . . .	29
5.25 Theorem (Riemann's Removable Singularities Theorem) . . . . .	31
5.26 Theorem (Taylor's Theorem) . . . . .	32
5.27 Theorem . . . . .	33
5.28 Theorem (Identity Theorem) . . . . .	34
5.33 Theorem . . . . .	36
5.36 Theorem (Weierstrass) . . . . .	37
5.38 Theorem (The Argument Principle) . . . . .	38
5.40 Theorem . . . . .	40
5.44 Theorem (The maximum principle) . . . . .	41
5.45 Theorem (The maximum principle) . . . . .	41
5.47 Theorem (Schwarz Lemma) . . . . .	41