

Homework 1

Lin Zejin

May 10, 2025

-
- **Collaborators:** I finish this homework by myself.
-

Problem 1. (a) When $\text{OPT} \geq c$, assume with $\frac{1}{T}$ algorithm A outputs a solution of value at least s . $T \in O(\text{poly}(n))$. Run algorithm A for $T \cdot n$ iterations. Then with $(1 - \frac{1}{T})^{Tn} < e^{-n}$ probability, the algorithm A outputs a solution of value less than s .

So with at least $1 - e^{-n}$ probability, the algorithm A outputs a solution of value at least s .

(b)

$$s = \mathbb{E}[\text{outputs}] \leq \Pr[\text{outputs} \geq s - \frac{1}{n^a}] \cdot \text{poly}(n) + (1 - \Pr[\text{outputs} \geq s - \frac{1}{n^a}]) \cdot (s - \frac{1}{n^a})$$

Then

$$\Pr[\text{outputs} \geq s - \frac{1}{n^a}] \geq \frac{\frac{1}{n^a}}{\text{poly}(n) - s + \frac{1}{n^a}} = \frac{1}{n^a(\text{poly}(n) - s) + 1}$$

Here we end the proof.

Problem 2.

Problem 3. (a) Let $k = c$, $U = \{1, 2, \dots, c\}^q$ where q large enough. Introduce

$$S_{i,b} = \{e \in U : e_i = b\}, i \in [q], b \in [c]$$

Choose $S_{i,t}$, $1 \leq t \leq c$ and the coverage is 1.

$x_{i,b}^* = \frac{1}{q}$ will also achieves coverage 1.

Then we cover each $j \in U$ with probability

$$1 - (1 - \frac{1}{c})^c$$

So the expected coverage of rounding is

$$1 - (1 - \frac{1}{c})^c$$

AS c large enough, the expected coverage of rounding is $1 - \frac{1}{e}$.

(b) With instance $k = c$, $U = \{0, 1\}^q$, $n = 2^q$ and

$$S_{i,b} = \{e \in U : e_i = b\}, i \in [q], b = 0, 1$$

The LP solution $x_{i,b}^* = \frac{1}{q}$.

$$\alpha x_{i,b}^* = \frac{(1-\epsilon) \ln n}{q} = (1-\epsilon) \ln 2 < \ln 2.$$

Then

$$\Pr[j \text{ is covered}] = 1 - (1 - \alpha x_{i,b}^*)^q < 1 - (1 - \ln 2)^q < 1 - (2^{-1.5})^{\log_2 n} = 1 - n^{-3/2}$$

So

$$\Pr[U \text{ is all covered}] < (1 - n^{-3/2})^n < 1 - n^{-\frac{1}{2}+\epsilon}$$

as n large enough. So the randomized rounding algorithm may not be able to find a feasible solution with probability at least $n^{-\frac{1}{2}+\epsilon}$.

Problem 4. (a)

(b) No, since the rounding algorithm gives a solution with expected value large than $(1 - \frac{1}{e})\text{LP}$. So

$$\text{OPT} \geq (1 - \frac{1}{e})\text{LP}$$

always holds.

Problem 5. (a) Suffices to prove the decision problem that if there exists a clique of size k is whether or not NP-complete.

For an 3-SAT instance with clauses c_1, \dots, c_m and variables x_1, \dots, x_n , we can construct a graph G with vertices $c_1, \dots, c_m, x_1, \bar{x}_1, x_2, \bar{x}_2, \dots, x_n, \bar{x}_n$ and additional clauses $x_{j_1} \wedge x_{j_2} \wedge x_{j_3}$ if there is some c_j is composed by $x_{j_1}, x_{j_2}, x_{j_3}$ or some of its negation. Let $k = 2m + n$

First construct a complete graph.

Remove the edges between x_i and \bar{x}_i

Remove those edges that connects $c_j = a \vee b \vee c$ and $\bar{a} \wedge \bar{b} \wedge \bar{c}$.

For a clause $a \wedge b \wedge c$, which means a, b, c holds in the same time, we remove the edges between $a \wedge b \wedge c$ and those additional clauses that contains one of $\bar{a}, \bar{b}, \bar{c}$

Remove the edges between $a \wedge b \wedge c$ and $\bar{a}, \bar{b}, \bar{c}$

Then, that a graph is clique is equivalent to this conditions:

- (1) If $a \wedge b \wedge c$ belongs to it, then additional clauses that contains $\bar{a}, \bar{b}, \bar{c}$ cannot belong to the same clique, and $c_j = \bar{a} \vee \bar{b} \vee \bar{c}$ cannot belong if it exists.
- (2) If x_i belongs to it, then \bar{x}_i cannot belong to the same clique and those additional clauses contains \bar{x}_i cannot belong to the same clique.

Certainly, the size of the clique in this graph cannot be larger than $2m + n$ since each clauses c_j corresponds to 8 additional clauses but at most one of them is contained.

Therefore, if 3-SAT is satisfiable, then we choose all true variables and all clauses that is true in the solution. Then c_j will be chosen all and exactly one of eight additional clauses that corresponds to c_j will be chosen, so $k = 2m + n$ is reachable.

If there exists a clique of size $k = 2m + n$, which means choosing n of independent variables, c_j and exactly one of eight additional clauses that corresponds to c_j . Those clauses will be TRUE when the variables that is chosen is assumed TRUE.

So it is a feasible instance for max-clique problem.

Therefore, It is NP-Hard.

(b) We have proved in (a) that if the instance has optimum 1, then the induced instance has optimum 1.

Now we suffice to prove soundness.