

KÄCHELE, H. (2013). Manualization as tool in psychodynamic psychotherapy research and clinical practice - Commentary on six studies. *Psychoanalytic Inquiry*, 33:626-630.

Manualization as Tool in Psychodynamic Psychotherapy Research and Clinical Practice - Commentary on Six Studies

Horst Kächele

It is an honour to comment on six fine papers on these recently developed treatments that share the distinctive features of using treatment manuals. The first generation of shorter psychodynamic treatments like Malan (1976), Mann (1973) Sifneos (1979) and Davanloo (1980) were presented in a traditional textbook format. The term 'manual' had not yet become popular. What is a manual? The New Oxford American Dictionary describes it as "a book of instructions, esp. for operating a machine or learning a subject; a handbook: *a computer manual* | *a training manual*".

In the seventies of the last century the term 'manual' began to replace the textbook notion; Lester Luborsky (1984) pointed out that "less than a decade ago a small revolution in psychotherapy research style demanded that an official 'manual' be devised for each psychotherapy so that methods and theories could effectively be implemented and compared" (p. XIX). He proudly pointed out that his concise primer on the principles of supportive-expressive psychotherapy was one of the first such manuals of which a first version, entitled "The Task of the Psychotherapist" was copyrighted on April 5, 1976. The result was a carefully designed guide for the practice of psychoanalytically oriented psychotherapy for practitioners, researchers, and supervisors.

According to Luborsky (1984, p. 4) therapeutic manuals need to fulfill the following three criteria:

- a) The specification of technique needs to be as complete as the type of treatment permits.
- b) The manual should make clear the treatment principles and the approaches which the therapist is supposed to perform.

- c) The manual should have an accompanying set of scales to measure the degree to which the therapists have complied with its main technique.

Psychotherapy manuals should ensure that therapists do what they are supposed to do according to the specified theoretical and technical framework (i.e. ensuring adherence). The first step in the development of manual consists in defining and clarifying the elements that are characteristic to a particular therapy method. For this reason, Gill recognized the importance of describing the elements of therapeutic approaches in his preface to Strupp and Binder's (1984) guide to "time-limited dynamic psychotherapy" (Gill 1984, p. VII).

For many clinicians, especially psychodynamically oriented ones, the introduction of such manuals into clinical practice induced the fear that the manualization of therapies would restrict the freedom, spontaneity, and variety of individual practice. However, if for a given therapy A, an individual action is recommended within the framework of defined rules, this instruction is only a guideline to evaluate therapeutic action. Guidelines are like sign posts: they do not tell us where to go (Wittgenstein). A rigid conformity to the rules will make therapy A less effective. For psychoanalytic therapies, this principle leads to fertile reflections about structuring a psychoanalytic treatment situation (Thomä & Kächele, 1994, p. 250).

All six papers in this issue utilize manualized treatments (Levy et al.; de Jonghe et al.; Busch & Milrod; Lemma et al.; Vinnars et al.; Bateman & Fonagy). For the purpose of establishing scientific credibility in the era of evidence-based medicine this is not surprising. The acceptance of such guidelines has advanced quite substantially. The agreement across the diverse 'systems of psychotherapy' (Ford, 1963) that structure-providing treatments can be more reliably assessed seems strong. The time when only cognitive-behavioral treatments were manualized and psychodynamic treatments were believed not to require these efforts is over. The impact of the Evidence-Based Medicine movement has dramatically changed psychodynamic therapists' view of manuals. We have recently compiled a (most likely incomplete) list of psychodynamic manuals that appeared in the period between 1980 and 2011 which demonstrates the exponential growth of manualized psychodynamic treatments (Seybert et al. 2012).

What is surprising in all of the papers, however, is the nearly complete absence of clinical concerns about manualized treatments. As noted above, it often was posited that manual-based specifications for psychodynamic treatments would be limiting or overly rigid. Although not addressed in the six papers in this issue, we may ask the following questions: Is there evidence that conducting therapy according to a manual does not interfere with positive outcome? What role does clinical experience play in the outcome of studies on treatment manuals? That is, is clinical experience as or possibly more important than manualized treatment in influencing outcome? There are a few studies on the issue that may help to answer these questions.

Luborsky et al. (1982), in a project with drug-addicted patients, studied three forms of manualized treatments: psychoanalytic psychotherapy, cognitive-behavioral therapy, and structured drug counseling. They demonstrated for the three modalities that therapists with the highest 'level of purity' in their technique – measured by evaluating the convergence of applied interventions with the theoretical frame - also had the best therapeutic results. The results of this first study on manual adherence suggested that one must reckon with the considerable variation of therapeutic quality within and between therapists.

In addition to therapist quality, the level of patient's psychopathology may also play a role in outcome. It has repeatedly been found that a large amount of variability in treatment outcome is due to the severity of patient pathology (Lambert & Ogles, 2004). Thus attention must be given to the technical problems of psychotherapeutic treatments with patients with more severe psychopathology. Within this problematic population, therapeutic competency could be decisive. In an illustration of this diversity Crits-Christoph et al. (1991) analyzed data from 15 comparative studies examining factors that could affect outcome, including the usage of a therapy manual, the degree of the therapist's expertise (competence), the treatment's duration, and the type of treatment (psychodynamic vs. cognitive-behavioral). They reported an average of 12% variance derived from therapist competence effects (with a dispersion of 50-0%), indicating a significant degree of impact.

The importance of therapist experience was also highlighted in the NIMH project, the Treatment of Depression Collaborative Research Program (TDCRP) (Elkin et al. 1985). The TDCRP was planned and executed by several senior members of the

NIMH extramural staff, who designed an extensive, elegant, and well-conducted randomized clinical trial that compared several forms of brief treatment for depression: 16 weeks of medication (imipramine, the antidepressant medication of choice at the time [IMI-CM]) and a double-blind placebo, PLA-CM, both with clinical management (CM; minimal support but no specific therapeutic intervention), with 16 sessions of manual- directed cognitive_behavioral therapy (CBT) or interpersonal psychotherapy (IPT). Eighteen experienced psychiatrists and 10 experienced PhD clinical psychologists (with a mean clinical experience of 11 years) saw patients at one of three treatment sites (medical schools at George Washington University, University of Oklahoma, and University of Pittsburgh). Surprisingly, Blatt et al. (1996) and also Kim et al. (2006) discovered a large effect for general therapeutic experience and its influence on patient's fear level, and only a low effect size from the use of specific therapeutic elements. The same tendency is shown in the results of a reanalysis of the "Multi-Center Collaborative Study for the Treatment of Panic Disorder" in Barlow's cognitive-behavioral group (Huppert et al. 2001). This study demonstrates even stronger results with regard to the impact of therapist experience on outcome than the re-analysis of the TDCRP.

In a study from Strupp's working group (Bein et al, 2000), the effect of a targeted training for experienced therapists in the Strupp-Binder manual (1984) was examined in terms of therapeutic process and outcome.. This study compared equally experienced therapists with and without explicit training in manualized therapy; the two groups did not show a significant difference in outcome. A Dutch cognitive-behaviorally oriented group also demonstrated that manualization did not have a greater impact compared to a nonmanualized treatment in the case of obsessive-compulsive disorders, either on the symptomatic level, or on the level of general functioning (Emmelkamp et al. 1994).

In the same vein Vinnars et al. (2005) reported that the comparison of a manualized supportive-expressive treatment versus non-manualized community-delivered psychodynamic therapy for patients with personality disorders – in contrast to their expectations – did not show any substantial differences. "The patients in the community therapy group received equally intensive psychotherapy, probably with a more supportive emphasis, delivered by experienced dynamic therapists receiving regular dynamic supervision" (p.1938).

Taken together a fair number of studies suggest that therapist experience prevails! In other words, general therapeutic knowledge is more important than expertise in a specific procedure. Phrased in a most convincing manner: "that which is relevant about experience may be general clinical contact rather than developing specific proficiencies" (Beutler et al. 2004, p. 240).

We are faced with a certain lack of clarity in the field. An extensive meta-analysis of 90 studies by Shadish et al. (2000) showed that the application of naturalistic psychotherapy and manualized treatment were on par. "First, psychological therapies are robustly effective across conditions that range from research-oriented to clinically representative" (p.522). Manuals are necessary for research purposes but there is no evidence that they are superior in treating disorders compared to the experienced clinician? Therefore the presentation of new manuals should include a specification of the degree of clinical expertise necessary for adequate implementation.

In addition to the description of treatments and research results in these informative papers that increase the diversity of psychodynamic interventions I would like to learn more about the impact of therapist competence and experience on outcome with the new treatments. Would the authors agree with findings that suggest that clinical expertise overrides the advantages of manualized treatments? I would also be interested in more information about how the authors think about implementation of the new tools into routine clinical practice. Are weekend trainings for already trained clinicians good enough or are the new products more suitable for basic teaching. Now that these treatments are increasingly accepted, questions about their implementation and role in clinical practice become paramount.

Bibliography

- Bein, E., Anderson, T., Strupp, H. H., Henry, W. P., Schacht, T. E., Binder, J. L., & Butler, S. F. (2000). The effects of training in time-limited dynamic psychotherapy: Changes in therapeutic outcome. *Psychotherapy Research*, 10, 119-32.
- Blatt, S. J., Sanislow, C., Zuroff, D. C., & Pilkonis, P. (1996). Characteristics of effective therapists: Further analyses of data from the National Institute of

- Mental Health Treatment of Depression Collaborative Research Program. *Journal of Consulting and Clinical Psychology*, 64, 76-84.
- Beutler, L. E., Malik, M., Alimohamed, S., Harwood, T. M., Talebi, H., Noble, S., & Wong, E. (2004). Therapist variables. In Lambert, M. J. (Ed.) *Bergin and Garfield's handbook of psychotherapy and behavior change* (5. Ed.). New York: Wiley, pp. 227-306.
- Crits-Christoph, P., Baranackie, K., Kurcias, J. S., & et al. (1991). Meta-analysis of therapist effects in psychotherapy outcome studies. *Psychotherapy Research*, 1, 81-91.
- Davanloo, H. (1980). *Short-term dynamic psychotherapy*. New York: Jason Aronson.
- Emmelkamp, P. M. G., Bouman, T. K., & Blaauw, F. (1994). Individualized versus standardized therapy: A comparative evaluation with obsessive-compulsive patients. *Clinical Psychology and Psychotherapy*, 1, 95-100.
- Elkin, I., Parloff, M.B., Hadley, S., Autry, J. (1985) NIMH Treatment of Depression Collaborative Research Program: Background and research plan. *Archives of General Psychiatry*, 42, 305-316
- Ford, D., & Urban, H. (1963). *Systems of psychotherapy. A comparative study*. New York: Wiley & Sons.
- Gill, M.M. (1984). Preface. In Strupp, H. H., & Binder, J. L. (1984) *Psychotherapy in a new key. A guide to time-limited dynamic psychotherapy*. New York: Basic Books, p. VII-X.
- Huppert, J., Bufka, L., Barlow, D., Gorman, J., Shear, M., & Woods, S. (2001). Therapists, therapists variables, and CBT outcome for panic disorders. *Journal of Consulting and Clinical Psychology*, 69, 747-755.
- Kim, D.M., Wampold, B.E., Bolt, D.M. (2006) Therapist effects in psychotherapy: A random-effects modeling of the National Institute of Mental Health Treatment of Depression Collaborative Research Program data. *Psychotherapy Research*, 16, 161-172.
- Lambert, M. J., & Ogles, B. (2004). The efficacy and effectiveness of psychotherapy. In Lambert, M. J. (Ed.) *Bergin and Garfield's handbook of psychotherapy and behavior change* (5. Ed.). New York Chichester Brisbane: Wiley, pp. 139-193.
- Luborsky, L. (1984). *Principles of psychoanalytic psychotherapy. A manual for supportive-expressive treatment*. New York: Basic Books.
- Luborsky, L., Woody, G. E., McLellan, A. T., O'Brien, C., & Rosenzweig, J. (1982). Can independent judges recognize different psychotherapies? An experience with manual-guided therapies. *Journal of Consulting and Clinical Psychology*, 50, 49-62.
- Malan, D. H. (1976). *Toward the validation of dynamic psychotherapy*. New York: Plenum Press.
- Mann, J. (1973). *Time-limited psychotherapy*. Cambridge, Mass.: Harvard University Press.
- Seybert, C., Erhardt, I., Levy, R.A., & Kächele, H. (2012). Manualized treatments for psychodynamic psychotherapy research. In Levy, RA., Ablon, JS., & Kächele, H. (Eds), *Psychodynamic Psychotherapy Research*. New York: Humana, pp. 650-7.
- Shadish, W., Matt, G., Navarro, A. M., & Phillips, G. (2000). The effects of psychological therapies under clinical representative conditions: A meta-analysis. *Psychol Bull*, 126, 512-29.
- Sifneos, P. (1979). *Short-term dynamic psychotherapy: Evaluation and technique*. New York: Plenum Medical Press.

- Strupp, H. H., & Binder, J. L. (1984). *Psychotherapy in a new key. A guide to time-limited dynamic psychotherapy*. New York: Basic Books.
- Thomä, H., & Kächele, H. (1994). *Psychoanalytic practice. vol. 2: Clinical studies*. New York: Jason Aronson.
- Vinnars B, Barber JP, Gallop R, Weinryb RM (2005) Manualized supportive-expressive psychotherapy versus nonmanualized community-delivered psychodynamic therapy for patients with personality disorders. *American Journal of Psychiatry*, 162, 1933-1944.

Prof. Dr.med. Dr. phil. Horst Kächele
International Psychoanalytic University.
Stromstr. 3. 10555 Berlin /FRG.
E-mail: horst.kaechele@ipu-berlin.de