

437. Brisch, K H, Munz D, Kächele H, Terinde R, Kreienberg R (2005) Effects of previous pregnancy loss on level of maternal anxiety after ultrasound screening for fetal abnormalities. *Journal of Loss and Trauma* 10: 131—153

Effect of previous loss of pregnancy on the level of maternal anxiety after prenatal ultrasound screening for fetal malformation

Karl Heinz Brisch

Department of Pediatric Psychosomatic Medicine and Psychotherapy, Dr. von Hauner Children's Hospital, Ludwig-Maximilians-University, Munich, Germany

Dorothee Munz and Horst Kächele

Department of Psychosomatic Medicine and Psychotherapy, University of Ulm, Ulm, Germany

Rainer Terinde and Rolf Kreienberg

Women's Hospital, Subdepartment of Prenatal Diagnosis and Sonography in Gynecology, University of Ulm, Ulm, Germany

Information about the authors:

Karl Heinz Brisch, M. D., head of the Department of Pediatric Psychosomatic Medicine and Psychotherapy at the Dr. von Hauner Children's Hospital, Ludwig-Maximilians-University, Munich / Germany, is a child- and adolescent psychiatrist, psychiatrist and neurologist, specialization in psychosomatic medicine, psychoanalyst. His main research areas include parent-infant-interaction, attachment, coping with high-tech medicine, diagnosis and treatment of attachment disorders

Doro Munz, Dr. rer. biol. hum., is a psychologist, researcher and psychotherapist in the Department of Psychosomatic Medicine and Psychotherapy, University of Ulm, Ulm / Germany. Her research areas include issues of coping with diagnostic procedures in medicine and psychotherapy.

Horst Kächele, M. D., director of the Department of Psychosomatic Medicine and Psychotherapy, University of Ulm, Ulm, Germany, is psychoanalyst and psychotherapist. His main research focuses on psychotherapy process and outcome research.###

Rainer Terinde, M. D., head of the, Subdepartment of Prenatal Diagnosis and Sonography in Gynecology, Women's Hospital, University of Ulm, Ulm, Germany, is a gynecologist and obstetrician, with specialization in prenatal diagnosis. His research areas are ###

Rainer Kreienberg, M. D., director of the Women's Hospital, University of Ulm, Ulm, Germany, is a gynecologist and obstetrician, with research specialization in oncology.###

Corresponding author:

Dr. med. Karl Heinz Brisch,
Abteilung Pädiatrische Psychosomatik und Psychotherapie,
Kinderklinik und Kinderpoliklinik,
Dr. von Haunersches Kinderspital,
Ludwig-Maximilians-Universität München
Pettenkoferstr. 10,
D-80336 München, Germany, email: Karl-Heinz.Brisch@med.uni-muenchen.de,
Phone: +49-89-5160-3709
Fax +49-89-5160-4730

Effect of previous loss of pregnancy on the level of maternal anxiety after prenatal ultrasound screening for fetal malformation

The impact of previous stillbirth, miscarriage, or preterm delivery on anxiety in pregnant women in various sub-groups at high risk for fetal abnormality in comparison with a non-risk control group of women with healthy uncomplicated pregnancies was studied longitudinally.

The level of anxiety in women (n=674) during early pregnancy was assessed by questionnaire. Data were collected at three points in time: immediately before the ultrasound scanning, at 5-6 weeks, and at 10-12 weeks after the prenatal examination for fetal malformation.

In general, all women with high-risk pregnancies (n=506) showed high levels of anxiety immediately before ultrasound scanning. There was a significant decrease in anxiety over the following 10-12 weeks. By contrast, the level of anxiety was not raised in the no-risk control group (n=168). However, pregnant women who had experienced miscarriage or stillbirth in previous pregnancies showed increasing or persistently high levels of anxiety over time.

We believe it is very important to screen women whose levels of anxiety are consistently high or escalating, and to offer psychotherapeutic counseling as a means for preventing pregnancy complications.

Introduction

Ultrasound scanning is a widely used diagnostic tool in prenatal care. In Germany, health insurance pays for two ultrasound screenings during pregnancy. One scan is done during the 18th-22nd week of gestation to detect fetal abnormalities. If a risk factor or an indication of fetal abnormality is detected, the pregnant woman is transferred to a center specializing in the detection of fetal malformation (prenatal diagnosis of Stage III). (Terinde, 1989). The fetus, placenta, and uterus can be screened in detail between the 18th and the 22nd week of gestation (Acton & Oats, 1984; see also Crowther et al., 1999). In a study by Götzmann et al. (2002), 96.4% of pregnant women were found to favor ultrasound scanning as one of the routine procedures for prenatal diagnosis, regardless of whether the suspected malformation was confirmed or not, or whether the pregnancy went to term or was terminated prematurely. With the widespread use and acceptance of ultrasound screening during pregnancy, more is known about the psychological impact of this technology (for a comprehensive review see Garcia et al., 2002). There is some indication that the visualization of a normal fetus during routine scanning is very attractive to women and their families (Garcia et al., 2002). However, results are contradictory with regard to the benefits of ultrasound visualization of the fetus (Kovacevic, 1993; Schönholzer, Götzmann, Zimmermann, & Buddeberg, 2000). Some studies have found positive effects in pregnant women who visualize the fetus in real time. In several studies, women reported a more positive attitude toward their pregnancy and more positive feelings of prenatal bonding (Black, 1992; Campbell et al., 1982; Kohn, Nelson, & Weiner, 1980; Milne & Rich, 1981). In other studies, however, these effects did not persist during the course of pregnancy and beyond birth (Hunter, Tsoi, Pearce, Chudleigh, & Campbell, 1987), or were not at all evident (Heidrich & Cranley, 1989; Kemp & Page, 1987). Baillie et al. (1999) showed in their review that most of these results are not very well documented by longitudinal research data based on empirical research. From our clinical experience with psychotherapeutic intervention—particularly with crisis intervention—at the Subdepartment of Prenatal Diagnosis and Sonography

in Gynecology (University of Ulm), we know that many pregnant women become very anxious while waiting for ultrasound scanning, especially if risk factors have been diagnosed previously (Brisch, 1998).

Prospective studies on the development of anxiety in the context of amniocentesis and chorion villus sampling are not reported on extensively here as no comparison group undergoing ultrasound scanning was involved. (Caccia, Johnson, Robinson, & Barna, 1991; Marteau et al., 1989; Möbus et al., 1992; Pauli, Blaser, & Herrman, 1990; Sjögren & Uddenberg, 1988; van Zuuren, 1993). The results of van Zuuren's study (1993) suggest that the invasive approach (needle insertion) to amniocentesis in particular may increase anxiety in pregnant women. Women are afraid that their fetus may be harmed by the technique, or that they may even lose their child, which is a known complication with this method. Van Zuuren (1993) was able to show that levels of anxiety in pregnant women were lower after amniocentesis than before, even though the women did not yet know the results of the amniocentesis. The results showed that the technique itself can induce anxiety (Caccia et al., 1991).

Lumley (1990) and Michelacci (1988) demonstrated that anxiety in pregnant women was high just before ultrasound scanning, but that levels were reduced if there was positive feedback indicating that the development of the fetus was normal. In contrast to these findings, Zlotogorski et al. (1996) could not explain the overall reduction in maternal anxiety after ultrasound scanning by different levels of feedback provided by the clinician (see also Eurenus, Axelsson, Gällstedt-Fransson, & Sjöden, 1997). With respect to urogenital abnormalities, Madarakan et al. (1990) found that levels of anxiety in pregnant women were high from the time the fetal abnormality was detected by

ultrasound and were not reduced until after birth, when the malformation was surgically corrected. In a study by Björkhem et al. (1997), anxiety decreased after normal fetal echocardiography both in women who had previously lost a child born to heart disease as well as in those whose child with a cardiac failure was still alive. Kowalcek et al. (2002) showed a significant reduction in anxiety after ultrasound scanning only in women with a nonsuspicious scan, but not in those with a suspected diagnosis of fetal malformation (see also Brisch et al., 2002). Detection of minor anomalies by prenatal scanning may generate disproportional maternal anxiety, as the findings of Thomas (2001) indicate. Levels of anxiety were significantly above the norm in both men and women before ultrasound scanning, but the men felt less stressed and experienced less anxiety than their pregnant partners (Götzmann, Kölble et al., 2002).

There was, however, no information on whether pregnant women had experienced complications in previous pregnancies, nor on how they coped with the results of ultrasound screening in a following pregnancy. It is well documented that a previous pregnancy loss due to congenital anomalies can influence maternal quality of life and emotional reactions toward later pregnancies (Hunfeld et al., 1993; 1996; Salvesen, Oyen, Schmidt, Malt, & Eik-Nes, 1997). In a longitudinal prospective study, Brisch et al. (2003) found that pregnant women fell into one of three broad categories in terms of coping strategies or patterns: cognitive-oriented coping, emotion-oriented coping, and action-oriented coping. A woman's specific coping strategy either decreased or increased anxiety, or had no influence on the level of anxiety longitudinally. Thus the pattern and quality of coping mechanisms might also explain the course of maternal anxiety after ultrasound screening for fetal malformation. Coping might be influenced by the information given by clinicians about the specific purposes of ultrasound scans and what they can and

what they cannot achieve (Garcia et al., 2002; Zlotogorski, Tadmor, Duniec, Rabinowitz, & Diamant, 1995), including information about the possibility of false positive and false negative results (Hutton & Spicer, 1994) and their psychological aftermath. Women might fear detection of a malformation as much as the possibility that it will be missed on examination (Kowalcek et al., 2002), and the ultrasound scanning as such could provoke an anxious maternal reaction (Teichmann, Rabinovitz, & Rabinowitz, 1991). Counseling might be helpful in this situation. Parents who received multidisciplinary counseling in the management of fetal surgical anomalies showed less anxiety than those counseled only by the obstetrician after birth (Alte et al., 2002).

Hypothesis

The objective of this prospective longitudinal study was to analyze the development of anxiety before and after ultrasound scanning over a period of 12 weeks with measurements taken at given intervals. We hypothesized that previous complications in pregnancies influence maternal anxiety, as discussed in the medical literature (Brisch, 1998; Feldman, 1990; Sjögren & Uddenberg, 1988; Sjögren & Uddenberg, 1990; Zuskar, 1987). The results should extend our knowledge about risk groups that potentially need psychotherapeutic intervention to cope with high levels of anxiety.

Methods

Design

The Ulm prospective longitudinal follow-up study on anxiety and coping processes after prenatal ultrasound scanning for fetal abnormalities was a joint project of the Department of Psychotherapy and Psychosomatic Medicine and the Subdepartment of Prenatal Diagnosis at the Women's Hospital, University of Ulm. The study was approved by the ethics committee of the University of Ulm.

All the women were seen during the waiting period immediately before ultrasound scanning (T0). Data were collected with questionnaires, and every 10th woman was asked to participate in a semi-structured interview. The questionnaires and the interview were repeated again after 4-5 weeks (T1) and after 8-10 weeks (T2). The purpose of the study was to collect longitudinal data on the development of anxiety. During the follow-up period some pregnant women had several ultrasound scans to confirm the diagnosis, or to get a clear picture of the further development of the fetus. Sometimes the diagnosis was not confirmed until at least 5-10 weeks after the first examination and the first data sampling. This suggests that some women had to cope with a longer period of uncertainty, as they did not have a precise diagnosis until 8-10 weeks post T0.

After informed consent, the pregnant women were asked to fill out a questionnaire on state and trait anxiety (STAI) (Laux, Glanzmann, Schaffner, & Spielberger, 1981; Spielberger, Gorsuch, & Lushene, 1970). State anxiety measures the situational anxiety that is provoked by the scanning for fetal malformation; trait anxiety is a characteristic feature of personality, with some women being more anxious than others regardless of situation. Further methods were applied for data collection:

a questionnaire on coping processes (Heim, Augustiny, Blaser, & Schaffner, 1991); a prenatal attachment scale (Brisch, Bemmerer-Mayer, & Nehb, 1994; Cranley, 1979; Cranley, 1981); a questionnaire on socio-demographic data, family and life situation, and on critical life-events (Brisch & Bemmerer-Mayer, 1994); a questionnaire on social support (Brisch, Bemmerer-Mayer, Buchheim, & Köhntop, 1994); and a questionnaire on personality (Fahrenberg, Selg, & Hampel, 1973). Diagnosis of fetal malformation by ultrasound scan was recorded in order to determine whether there was a difference in level of anxiety and coping styles as a result of positive findings or on the extent of the malformation. At T1 and at T2 the pregnant women were again asked to fill out the same questionnaires on anxiety, on coping, and on prenatal mother-fetus attachment. At T2 the latest most precise diagnosis was recorded.

Sample

The total sample size was n=674 pregnant women in the second trimester (16th-24th week of gestation): n=506 women were in the high-risk group, and n=168 women in the no-risk control group. The pregnant women in the high-risk group were contacted and informed about the aims of the study when they came to Women's Hospital for prenatal ultrasound scanning because of the risk of fetal malformation. The control group included pregnant women who were seen for routine ultrasound scanning at University Hospital or in the practice of gynecologists and obstetricians. No risk factors had been determined to that point. The women were included in the study after receiving a full explanation of the study and signing an informed consent. The acceptance of the study was very good: 88.7% of the pregnant women contacted consented to taking part in the study. The dropout rate was 21.6% as of the latest follow-up. The high-risk group was divided into five further sub-groups (Table 1):

Sub-group I: „Suspected fetal abnormality“

Pregnant women who had already been diagnosed by an obstetrician as showing signs of suspected malformation of the fetus (e.g., hydrocephalus, omphalocele, spina bifida, gastroschisis) were transferred to the University Center with its specialized diagnostic facilities, where the diagnosis was to be confirmed.

Sub-group II: „Complications in previous pregnancies“

Pregnant women who had experienced complications in previous pregnancies (e.g., abortus, fetal malformations, prematurity, miscarriage, stillbirth).

Sub-group III: „Maternal disease/medication“

Pregnant women who suffered from infections or chronic diseases and needed ongoing medication which might result in a risk of fetal malformation, miscarriage, or other complications in the current pregnancy (e.g., rubella infection, cytomegaly, borreliosis, epilepsy, diabetes mellitus).

Sub-group IV: „Advanced maternal age“

Pregnant women who were older than 35 and who were at risk for fetal abnormalities because of their age, and who did not accept amniocentesis or chorion villus sampling.

Sub-group V: „Endocrinological test“

Pregnant women who had undergone endocrinological diagnostic screening during the current pregnancy (e.g., alpha-fetoprotein-screening) and showed elevated blood levels resulting in a significantly increased risk of fetal malformation. High endocrine blood levels in particular proteins can be associated with a risk of Down Syndrome or spinal cord defects. Further diagnostic procedures such as ultrasound scanning were then recommended to come to at a more definitive diagnosis. Unfortunately, some pregnant women seemed to have been less than completely informed about the aims of blood sampling. Some did not know about the potential for false positive or false negative results.

Sub-group VI: „Multiple risks“

Pregnant women who had a combination of more than one risk factor.

here insert Table 1: Sub-Groups of Sample

The number of fetal malformations detected was quite different in these sub-groups (Table 2). In the control group, 2.4% of the pregnant women with normal pregnancies had pathological findings. The highest rate of fetal malformations (58.9%) was detected in the high-risk sub-group of women who had already been transferred to the center because of suspected fetal abnormalities.

here insert Table 2: Pathological Findings of Fetal Malformations in Subgroups

Statistical Analysis

The statistical analyses were done using SPSS 10.0 for Windows. Depending on the variable, we used the Chi-square test for discrete variables and, depending on the distribution of the data, we applied independent samples T-test and Mann-Whitney U-test, ANOVA, Kruskal-Wallis-H-test for several independent samples and Wilcoxon - two related samples tests to compare the results in the different groups. For comparison of results at different points in time we used the General Linear Model (GLM) with repeated measures.

Results

Description of sample

The high-risk group and the no-risk control group showed no significant differences in most of the socio-demographic variables (Table 3). There was a significant difference ($p=.000$) only for "maternal age," with a higher percentage of older women in the high-risk group. This was because the sub-group with the "advanced maternal age risk factor" was in the high risk sample. Furthermore, the level of education was significantly higher in the control than in the high-risk groups ($p=.012$). In addition, there was no statistical difference in the variables shown in Table 3 between the women in the control group who were recruited from the University Hospital and those who came from private practice.

here insert Table 3: Socio-demographic Data of the Sample

Levels of maternal anxiety

All pregnant women in the high-risk sub-groups showed significantly higher levels of anxiety immediately before ultrasound scanning (at T0) than did those in the control group. Two sub-groups had the highest levels of anxiety: a special sub-group that had been at risk because of very concrete signs of fetal malformation (Sub-Group I) and the sub-group that was at risk because of pathological "endocrinological tests " (Sub-Group V). These two groups had even higher levels of anxiety than the multiple-risks sub-group (Sub-Group VI). The lowest level of anxiety was seen in the sub-group with „advanced maternal age“ (Sub-Group IV). However, in this age-risk sub-group,

the anxiety level was still higher than in the control group, and lower than in the sub-group with „risks in previous pregnancies“ (Sub-Group II) (see Figure 1).

here insert Figure 1: Level of anxiety of sub-groups

Statistical analysis showed that the levels of anxiety within the sub-groups were significantly different at T0 ($p=.000$) and T1 ($p=.001$), but not any more at T2 ($p=.689$). With regard to the longitudinal development of anxiety, there was a significant decrease in level of anxiety in each sub-group from T0 to T1 and to T2 (Brisch et al., 2002).

Anxiety and pathological findings of fetal malformations

From the same sample, Brisch et al. (2002) reported that pregnant women who were confronted with a definitive diagnosis of fetal malformation had significantly higher levels of anxiety even before their ultrasound examination at T0, as well as at T1 and T2, than did women with no pathological findings. The difference in the reduction in anxiety from one point in time to the next was significantly greater in women with confirmed pathological findings in comparison to women with no diagnosis of fetal abnormality. In contrast to these results, the level of trait anxiety was not significantly elevated in women with confirmed fetal abnormalities.

Contrary to these findings, pregnant women in the control group with no risk of fetal abnormalities did not show raised levels of anxiety comparable to a female control group in Spielberger (1970). Control variables (e.g., socio-demographic data) did not show a significant correlation with level of anxiety at any of the three data sampling points.

Increase in levels of anxiety

Contrary to the overall decreases in the level of anxiety in the total sample, a small sub-sample of women (11.9%) showed increasingly high levels of anxiety from T0 to T2. The cut-off point for an „increase in anxiety“ was defined such that the mean difference in anxiety level had to increase more than one standard deviation from T0 to T2. Even in the control group of pregnant women with no risk factor, 13.3% (vs. 11.4% in the high risk sub-groups) showed a low level of anxiety at T0, but a high level at T1 and an even higher level at T2. These women were differentiated neither by sub-group category nor by the result of a confirmed diagnosis of fetal malformation, but were characterized by a significantly larger number of children and pregnancies and deliveries before this risk pregnancy. They had, in fact, experienced significantly more miscarriages and stillbirths in previous pregnancies. Other previous complications, such as preterm delivery and induced abortion, did not correspond significantly with an increase in anxiety (Table 4).

here insert Table 4: Increase in Levels of State Anxiety

Persistent high levels of anxiety

We also found another sub-group of pregnant women who had high levels of anxiety at T0, which stayed high at T1 and even at T2 (1.8% of the women in the control-group and 5.8% of the women in the high-risk sub-groups). A persistent high level of anxiety was defined by the highest value of anxiety at T0 that was reached by the women with a suspected fetal malformation (mean=54). The anxiety at T2 (mean=60.72, sd=5.14) remained at a level that was in the upper range (the possible maximum level of anxiety is 80) (Table 5). These pregnant women were neither differentiated by socio-demographic factors nor by their type

of high-risk sub-group, but they had significantly more confirmed fetal abnormalities after ultrasound scanning (Chi-Square=7.867, df=1, Fisher's Exact Test $p=.001$). Nevertheless, not all pregnant women with confirmed pathological findings showed persistent high levels of anxiety from T1 to T2. These two sub-groups of pregnant women with fetal malformations (one sub-group with a consistently high level of anxiety, and one with a decrease in anxiety from T0 to T2) were differentiated only by one additional variable from their previous pregnancies: the women with persistent high levels of anxiety had experienced stillbirths significantly more often in the past.

Table 5: Trait anxiety and persistent high levels of state anxiety at different points in time

In summary, the pregnant women who experienced extremely high anxiety levels at all three points in time also had a high trait level of anxiety at T0 (mean=47.89, sd= 8.48 vs mean=36.44, sd=8.13, in the group with decreasing anxiety) (Table 5). Furthermore, they showed significantly ($p=.000 - .029$) different results in some variables in the personality test: „less content with life situation,“ „higher level of arousal,“ „more somatic complaints,“ „more openness,“ „more emotionality.“

Discussion

Pregnant women who had risk factors for fetal malformation showed higher levels of anxiety before ultrasound scanning than those with no-risk pregnancies. In general, however, level of anxiety decreased significantly in the follow-up period after the ultrasound scanning in all high risk groups. Thus, for many women, anxiety can increase just before ultrasound examination and screening for fetal abnormalities, but they are then better able to cope with the results after the tests, as anxiety levels decrease during the following 12 weeks (Brisch et al., 2002).

Despite overall decreases in levels of anxiety in pregnant women in the high-risk sub-group, there was a small sub-sample that showed increasing levels of anxiety from T0 to T1 and even to T2. The results from this sub-sample that was recruited from *all* the high risk sub-groups, were opposite those of the mainstream development of level of anxiety in the longitudinal follow-up. In line with our hypothesis, this sub-group was distinguished from the other high-risk sub-groups by their correlation with the variables of previous stillbirth and miscarriage. This correlation was also found in the control group, which was characterized by a normal course of pregnancy without risks and complications. Nevertheless, the women in the control group who had previously experienced stillbirth and miscarriage had increasing level of anxiety despite the obstetrician's confirmation that the fetus was healthy and that the development of the present pregnancy was uncomplicated. This unusual increase in levels of anxiety might be influenced by the experience of previous—perhaps traumatic—complications. Women may be afraid to lose their baby again, as Hunfeld et al. (1993; 1996) found elevated stress reactions in these women to following pregnancies. If a previous stillbirth or miscarriage has not been adequately mourned (Beutel, Deckhardt, Rad, & Weiner, 1995), then the emotional background of this experience may be reactivated by the new pregnancy and increase anxiety (Hughes, Turton, Hopper, & Evans, 2002). This would explain why women who remained at a persistent high level of anxiety had a „higher level of arousal“ and „more

somatic complaints,“ which may be a sign of a posttraumatic stress disorder, as Seng et al. found in their study of post-pregnancy complications (2001). The actual normal development of the fetus does not seem to be an adequate source of reassurance (Bourne & Lewis, 1984; Condon, 1986; Radestad, Steineck, Nordin, & Sjögren, 1996).

In the high risk group, many of these pregnant women, especially those in the sub-group with complications in previous pregnancies, had experienced stillbirth or miscarriage, so that an increase in anxiety in these women was not surprising. But women with increasing levels of anxiety over time came from all the high-risk sub-groups. This result is very important, as 13.3% in the control and 11.4 % in the high-risk group showed this course of increasing anxiety.

A second extraordinary sub-sample of pregnant women was distinguished by an extremely high, persistent level of anxiety at both T0 and T2. These women were not comforted by the results of the ultrasound scanning, and therefore did not show any decrease in anxiety over time. These women also differed from the other high-risk samples in having experienced more than one stillbirth in the past. These past experiences, which had demonstrated that things can indeed go wrong during pregnancy, might have made them even more anxious in their following pregnancy regardless the course of this pregnancy. Their hopes for an uncomplicated pregnancy as well as their anxiety at being confronted again with the potentially traumatic experience of stillbirth may be very intense. In this highly ambivalent emotional state they may be „on alert“ during the entire pregnancy. Anxiety cannot decrease before the child is born because a healthy birth is the only outcome that removes the fear of another stillbirth and the anxiety that this fear causes.

The persistent high levels of anxiety in these women is warranted by their experience that fetal malformation is more often confirmed by ultrasound. These women had elevated trait levels as

well. This could indicate that a personality feature may also conflict with the results of consistently high levels of anxiety. Furthermore, one could argue that the experience of stillbirth, in particular, could have had a long-term effect on personality in the form of emotional alertness, which may be reflected in higher trait anxiety and in increasing or persistently high level of state anxiety.

These two special groups of pregnant women could form the target groups for psychological intervention as one might imagine that increasing or persistently high levels of anxiety during the course of pregnancy may interfere with the somatic development of the fetus and the integrity of the pregnancy as a result of anxiety-induced elevation in vascular resistance in the placenta and a decrease in alimentary and oxygen supply to the fetus (Teixeira, Fisk, & Glover, 1999).

Limitations

Johnson & Slade (2003) discuss in their review whether the State-Trait Anxiety Inventory (Spielberger et al., 1970) is valid as a measurement of anxiety in a pregnant sample. Although the stability of this instrument is explored as a viable anxiety measure in pregnant women, the test-retest reliability of the trait scale was low around the time of delivery. In our study we did not measure around the time of delivery. Nevertheless, Johnson & Slade (2003) question the use of this tool, especially during the third trimester of pregnancy. Newer instruments that focus on and evaluate specific anxieties in pregnancy in greater detail are tools that have not yet been well evaluated, and must still prove their validity and reliability (Johnson & Slade, 2003). This underscores our decision to evaluate anxiety in our study using Spielberger's State-Trait Anxiety Inventory.

Variations in periods of time that women had to wait to learn their diagnoses introduces extraneous variables that were not accounted for in any of the analyses. Some women had additional

ultrasounds and some had delayed diagnoses. Timing of the reception of the results could have affected anxiety levels.

Different levels of feedback given during or after the ultrasound examination were not registered in this study. However, findings from previous studies suggest that level of anxiety after ultrasound scanning for detection of malformation was unaffected by the amount of feedback given (Eurenius et al., 1997; Zlotogorski et al., 1996; Zlotogorski et al., 1995).

We did not study the amount and type of information women got from their obstetricians *before* the came to be scanned for fetal malformations. Especially in the group with endocrine testing, knowing that the endocrinological test was not definitive may have influenced their level of anxiety.

By design there should have been no women in the control group with previous histories of stillbirth or miscarriage. Although these women presented to the ultrasound examiner with a „normal, uncomplicated pregnancy and no previous history of pregnancy complications,“ the data from their questionnaires revealed that there had been previous pregnancy complications in 2.4%, which is about the basic rate of 3% defects for any pregnancy (Terinde, 1989). This discrepancy between oral communication to the examiner and written information in the questionnaire remains a factor of uncertainty.

Conclusions

Early prenatal ultrasound screening for fetal malformation can be a useful tool for detecting fetal anomalies during early pregnancy. Many women with high-risk factors for fetal malformation

showed a decrease in their level of anxiety, and in the following course of their pregnancy their anxiety was comparable to that of women with a healthy pregnancy.

The results revealed several sub-groups that should be addressed for further psychological intervention studies. Women with a confirmed diagnosis of malformation may need some form of longitudinal guidance and support (Leon, 1990; Stray-Pedersen & Stray-Pedersen, 1988). This could consist of clear information about risks, teaching of coping skills, anxiety reduction techniques, cognitive behavior therapy or more psychodynamically oriented psychotherapy to work through previous pathological or traumatic grieving (Condon, 1986; Ladker & Toedter, 1991; Lin & Lasker, 1996; Theut, Pedersen, Zaslow, & Rabinovich, 1988), or group counseling (Kaiser et al., 2002). Although their level of anxiety decreased over time, it still stayed higher than in the other sub-groups. This is not surprising, as more than 50% of these women received a confirmed diagnosis that the expected child would have a malformation. Obstetricians should not be misled by the decreasing anxiety overall, as these women still needed psychosocial support to cope with the first phase of emotional shock and for the long-term coping process (Franche & Culow, 1999).

Two groups that could potentially benefit from intervention were found during this study: pregnant women with increasing anxiety and those with extremely persistent high levels of anxiety. These women came from all sub-groups and even from the control group. As the percentage is not low (11.9% of the total sample), it seems prudent to offer them some form of focused, short-term psychotherapy (Leon, 1990). The background common to these two groups of women seems to be

past experience of miscarriage and stillbirth (cf Salvesen et al., 1997). The elevated level of anxiety could indicate that these previous events were not worked through, and were then reactivated by the new pregnancy—regardless of whether the new pregnancy was complicated or not (Beutel et al., 1995; Franche & Culow, 1999). These issues could perhaps be the focus of a more trauma-oriented psychotherapy (Seng et al., 2001). The results of this study underline the importance of psychological intervention for these women after a miscarriage, a stillbirth, or pregnancy complications of any kind. Results of a new study indicate that a high level of anxiety during pregnancy is not only a subjective emotional problem for these women, but can also decrease uterine blood circulation, negatively influencing the growth and well-being of the fetus (Teixeira et al., 1999). For these reasons, early psychological support could be used as a preventive measure for intervening in potentially problematic pregnancies.

Contributors

K.H. Brisch coordinated this study which was designed by K. H. Brisch, K. Bemmerer-Mayer and H. Kächele. K.H. Brisch, K. Bemmerer-Mayer and D. Munz were responsible for the collection of data and for overall data management. K.H. Brisch, K. Bemmerer-Mayer and D. Munz did statistical data analyses. R. Terinde and R. Kreienberg were responsible for the care of the patients. K. H. Brisch wrote the paper, with critical comments from the co-authors.

Acknowledgments

The research in this article was made possible by a grant from the German Research Council (DFG) to H. Kächele, R. Kreienberg, R. Terinde and K.H. Brisch (Grant KA 483/1-1, 1-2). Further grants from the University of Ulm and from the „Kässbohrer-Stiftung, Ulm“ (Germany) to K.H. Brisch supported this study. We are most grateful for this support. We are also indebted to the pregnant women for their generous longitudinal cooperation with the study, and to R. Terinde's team of co-workers who, with great interest and commitment, supported the recruitment of the sample.

References

- Acton, C. M., & Oats, J. N. (1984). Should an ultrasound scan be a routine in the obstetric patient? *Australasian Radiology*, 28(3), 257-261.
- Alte, L., Trucchi, A., Nahom, A., Spina, V., Bilancioni, E., & Bagolan, P. (2002). Multidisciplinary management of fetal surgical anomalies: The impact on maternal anxiety. *European Journal of Pediatric Surgery*, 12, Part 2, 90-94.
- Baillie, C., & Hewison, J. (1999). Should ultrasound scanning in pregnancy be routine? *Journal of Reproductive and Infant Psychology*, 17(2), 149-157.
- Beutel, M., Deckhardt, R., Rad, M. v., & Weiner, H. (1995). Grief and depression after miscarriage: Their separation, antecedents, and course. *Psychosomatic Medicine*, 57(6), 517-526.
- Björkhem, G., Jörgensen, C., & Hanséus, K. (1997). Parental reactions to fetal echocardiography. *The Journal of Maternal-Fetal Medicine*, 6, 87-92.
- Black, R. B. (1992). Seeing the Baby: The impact of ultrasound technology. *Journal of Genetic Counseling*, 1(1), 45-54.
- Bourne, S., & Lewis, E. (1984). Pregnancy after stillbirth or neonatal death. Psychological risks and management. *Lancet*, 2(8393), 31-33.
- Brisch, K. H. (1998). Die Bedeutung der Psychodynamik im Rahmen der pränatalen Fehlbildungsdiagnostik. *Speculum*, 16(1), 8-16.
- Brisch, K. H., & Bemmerer-Mayer, K. (1994). *Ulmer Fragebogen zur Lebens- und Familiensituation von Schwangeren (ULF-PRÄ)*. Ulm: Abteilung Psychotherapie und Psychosomatische Medizin und Psychosomatische Medizin.
- Brisch, K. H., Bemmerer-Mayer, K., Buchheim, A., & Köhntop, B. (1994). *Ulmer Skala zur sozialen Unterstützung-USU* (Unveröffentlichtes Manual.). Ulm: Abteilung Psychotherapie und Psychosomatische Medizin.
- Brisch, K. H., Bemmerer-Mayer, K., & Nehb, D. (1994). *Skala zur Entwicklung der pränatalen Mutter-Kind-Beziehung*. Ulm: Ambulanz für Kinder- und Jugendpsychiatrie und Psychotherapie, Abteilung Psychotherapie und Psychosomatische Medizin, Universität Ulm.
- Brisch, K. H., Munz, D., Bemmerer-Mayer, K., Terinde, R., Kreienberg, R., & Kächele, H. (2002). Ultrasound scanning for diagnosis of fetal abnormality and maternal anxieties in a longitudinal perspective. *Journal of Reproductive and Infant Psychology*, 20(4), 223-235.
- Brisch, K. H., Munz, D., Bemmerer-Mayer, K., Terinde, R., Kreienberg, R., & Kächele, H. (2003). Coping styles of pregnant women after prenatal ultrasound screening for fetal malformation. *Journal of Psychosomatic Research*, 55(2), 91-97.
- Caccia, N., Johnson, J. M., Robinson, G. E., & Barna, T. (1991). Impact of prenatal testing on maternal-fetal bonding: chorionic villus sampling versus amniocentesis. *American Journal of Obstetrics and Gynecology*, 165(4), 1122-1125.
- Campbell, S., Reading, A. E., Cox, D. N., Slemere, C. M., Mooney, R., Chudleigh, P., Beedle, J., & Ruddick, H. (1982). Ultrasound scanning in pregnancy: the short term psychological effects of real-time scans. *Journal of Psychosomatic Obstetrics and Gynaecology*, 1, 57-60.
- Condon, J. (1986). Management of established pathological grief reaction after stillbirth. *American Journal of Psychiatry*, 143(8), 987-992.
- Cranley, M. (1979). Maternal-Fetal Attachment Scale. State University at Buffalo.
- Cranley, M. S. (1981). Roots of attachment: the relationship of parents with their unborn. *Birth Defects*, 17(6), 59-83.
- Crowther, C., Kornmann, L., O'Callaghan, S., George, K., Furness, M., & Willson, K. (1999). Is an ultrasound assessment of gestational age at the first antenatal visit of value? A randomised clinical trial. *British Journal of Obstetrics and Gynaecology*, 106, 1273-1279.
- Eurenius, K., Axelsson, O., Gällstedt-Fransson, I., & Sjöden, P.-O. (1997). Perception of information, expectations and experiences among women and their partners attending a second-trimester routine ultrasound scan. *Ultrasound Obstet. Gynecol.*, 9, 86-90.

- Fahrenberg, J., Selg, H., & Hampel, R. (1973). *Freiburger Persönlichkeitsinventar (FPI)*. Göttingen: Hogrefe.
- Feldman, R., L. (1990). An analysis of coping processes in high-risk pregnant women. *Dissertation Abstracts International*, 50(8-B), 3743-3744.
- Franché, R.-L., & Culow, C. (1999). The impact of a subsequent pregnancy on grief and emotional adjustment following a perinatal loss. *Infant Mental Health Journal*, 20(2), 175-187.
- Garcia, J., Bricker, L., Henderson, J., Martin, M. A., Mugford, M., Nielson, J., & Roberts, T. (2002). Women's views of pregnancy ultrasound: A systematic review. *Birth*, 29, 225-250.
- Götzmann, L., Kölbl, N., Schönholzer, S. M., Klaghofer, R., Scheuer, E., Zimmerman, R., Huch, R., & Buddeberg, C. (2002). Verdachtsdiagnose einer fetalen Entwicklungsstörung im Ultraschall: Gefühle und Einstellungen zur Ultraschall-Technologie aus Sicht der Partner. *Ultraschall in der Medizin*, 23, 333-337.
- Götzmann, L., Schönholzer, S. M., Kölbl, N., Klaghofer, R., Scheuer, E., Huch, R., Buddeberg, C., & Zimmermann, R. (2002). Die Ultraschall-Untersuchung bei Verdacht auf fetale Entwicklungsstörungen: Zufriedenheit und Bewertungen aus Sicht der betroffenen Frauen. *Ultraschall in der Medizin*, 23, 27-32.
- Heidrich, S., & Cranley, M. (1989). Effect of fetal movement, ultrasound scans and amniocentesis on maternal-fetal attachment. *Nursing Research*, 38(2), 81-84.
- Heim, E., Augustiny, K. F., Blaser, A., & Schaffner, L. (1991). *Berner Bewältigungsformen (BEFO) Handbuch*. Bern, Göttingen, Toronto: Huber.
- Hughes, P., Turton, P., Hopper, E., & Evans, C. D. H. (2002). Assessment of guidelines for good practice in psychosocial care of mothers after stillbirth: a cohort study. *Lancet*, 360, 114-118.
- Hunfeld, J. A., Wladimiroff, J. W., Passchier, J., Venema-Van Uden, M. U., Frets, P. G., & Verhage, F. (1993). Emotional reactions in women in late pregnancy (24 weeks or longer) following the ultrasound diagnosis of a severe or lethal fetal malformation. *Prenat. Diagn.*, 13, 603-612.
- Hunfeld, J. A. M. e. a. (1996). Quality of life and anxiety in pregnancies after late pregnancy loss: A case-control study. *Prenatal Diagnosis*, 16, 783-790.
- Hunter, M. S., Tsoi, M. M., Pearce, M., Chudleigh, P., & Campbell, S. (1987). Ultrasound scanning in women with raised alpha fetoprotein: Long-term psychological effects. *Journal of Psychosomatic Obstetrics and Gynaecology*, 6, 25-31.
- Hutton, K. A. R., & Spicer, R. D. (1994). Benefits and adverse effects of prenatal diagnosis in non-urological surgical disorders. *European Journal of Pediatric Surgery*, 4, 161-164.
- Johnson, R. C., & Slade, P. (2003). Obstetric complications and anxiety during pregnancy: Is there a relationship. *Journal of Psychosomatics in Obstetrics and Gynecology*, 24, 1-14.
- Kaiser, A., Ferris, L. E., Pastuszak, A. L., Llewellyn-Thomas, H., Johnson, J.-A., Conacher, S., & Shaw, B. F. (2002). The effects of prenatal group genetic counselling on knowledge, anxiety and decisional conflict: Issues for nuchal translucency screening. *Journal of Obstetrics and Gynaecology*, 22(No. 3), 246 - 255.
- Kemp, V. H., & Page, C. K. (1987). Maternal prenatal attachment in normal and high risk pregnancies. *Journal of Obstetric, Gynecologic, and Neonatal Nursing*, 16, 179-184.
- Kohn, C. L., Nelson, A., & Weiner, S. (1980). Gravida's responses to realtime ultrasound fetal image. *Journal of Obstetric, Gynecologic, and Neonatal Nursing*, 9, 77-80.
- Kovacevic, M. (1993). The impact of fetus visualization on parents' psychological reactions. *Perinatal Psychology*, 8, 83-93.
- Kowalcek, I., Lammers, C., Brunk, J., Bieniakiewicz, I., & Gembruch, U. (2002). Angst der Schwangeren vor und nach der pränatalen Untersuchung bei unauffälligen und bei auffälligen Befunden. *Zentralblatt für Gynäkologie*, 124, 170-175.
- Ladker, J. N., & Toedter, L. J. (1991). Acute versus chronic grief: The case of pregnancy loss. *American Journal of Orthopsychiatry*(61), 510-522.
- Laux, L., Glanzmann, P., Schaffner, P., & Spielberger, C. D. (1981). Das State-Trait-Angstinventar. Weinheim: Beltz.

- Leon, I. G. (1990). *When a baby dies: Psychotherapy for pregnancy and newborn loss*. New Haven London: Yale University Press.
- Lin, S. X., & Lasker, J. N. (1996). Patterns of grief reaction after pregnancy loss. *American Journal of Orthopsychiatry*(66), 262-271.
- Lumley, J. (1990). Through a glass darkly: Ultrasound and prenatal bonding. *Birth*, 17(4), 214-217.
- Madarikan, B. A., Tew, B., & Lari, J. (1990). Maternal response to anomalies detected by antenatal ultrasonography. *British Journal of Clinical Practice*, 44(12), 587-589.
- Marteau, T. M., Johnston, M., Shaw, R. W., Michie, S., Kidd, J., & New, M. (1989). The impact of prenatal screening and diagnostic testing upon the cognitions, emotions and behaviour of pregnant women. *Journal of Psychosomatic Research*, 33(1), 7-16.
- Michelacci, L., Fava, G. A., Grandi, S., Bovicelli, L., Orlandi, C., & Trombini, G. (1988). Psychological reactions to ultrasound. Examination during pregnancy. *Psychotherapy and Psychosomatics*, 50(1), 1-4.
- Milne, L. S., & Rich, O. J. (1981). Cognitive and affective aspects of the responses of pregnant women to ultrasonography. *Maternal-Child Nursing Journal*, 10, 115-139.
- Möbus, V., Casper, F., Heintz, B., Kreienberg, R., Brockerhoff, P., & Knapstein, P. G. (1992). Amniozentese zur pränatalen Diagnostik aus psychischer Indikation. *Geburtshilfe und Frauenheilkunde*, 52(4), 225-229.
- Pauli, C., Blaser, A., & Herrman, U. (1990). Amniozentese: Psychische Belastung und deren Bewältigung bei der schwangeren Frau. *Geburtshilfe und Frauenheilkunde*, 50(4), 291-294.
- Radestad, I., Steineck, G., Nordin, C., & Sjögren, B. (1996). Psychological complications after stillbirth - Influence of memories and immediate management: Population based study. *British Medical Journal*, 312(7045), 1505-1508.
- Salvesen, K. A., Oyen, L., Schmidt, N., Malt, U. F., & Eik-Nes, S. H. (1997). Comparison of long-term psychological responses of women after pregnancy termination due to fetal anomalies and after perinatal loss. *Ultrasound Obstet. Gynecol.*, 9, 80-85.
- Schönholzer, S. M., Götzmann, L., Zimmermann, R., & Buddeberg, C. (2000). Psychologische Aspekte der Ultraschalluntersuchung während der Schwangeschaft. *Gynäkologisch-geburtshilfliche Rundschau*(40), 119-124.
- Seng, J. S., Oakley, D. J., Sampsel, C. M., Killian, C., Graham-Bermann, S., & Liberzon, I. (2001). Posttraumatic stress disorder and pregnancy complications. *Obstetrics and Gynecology*, 97(No. 1), 17 - 22.
- Sjögren, B., & Uddenberg, N. (1988). Prenatal diagnosis and maternal attachment to the child-to-be: A prospective study of 211 women undergoing prenatal diagnosis with amniocentesis or chorionic villi biopsy. *Journal of Psychosomatic Obstetrics and Gynaecology*, 9(2), 73-87.
- Sjögren, B., & Uddenberg, N. (1990). Prenatal diagnosis for psychological reasons: a comparison with other indications, advanced maternal age and known genetic risk. *Prenatal diagnosis*, 10(2), 111-120.
- Spielberger, C. D., Gorsuch, R. L., & Lushene, R. E. (1970). *Manual for the State-Trait Anxiety Inventory*. Palo Alto, California: Consulting Psychologists Press, Inc.
- Stray-Pedersen, B., & Stray-Pedersen, S. (1988). Recurrent abortion: The role of psychotherapy. In R. W. Bred & F. Sharp (Eds.), *Early pregnancy loss* (pp. 430-440). Berlin New York: Springer.
- Teichmann, Y., Rabinovitz, d., & Rabinowitz, Y. (1991). Emotional reactions of pregnant women to ultrasound scanning and postpartum. In C. D. Spielberger & I. G. Saranson & J. Strelau & J. M. T. Brebner (Eds.), *Stress and anxiety* (pp. 103-116). New York: Hemisphere Publishing.
- Teixeira, J. M. A., Fisk, N. M., & Glover, V. (1999). Association between maternal anxiety in pregnancy and increased uterine artery resistance index: Cohort based study. *British Medical Journal*(318), 153-157.

- Terinde, R. (1989). Normale und pathologische fetale Entwicklung im Ultraschallbild. In W. Dmoch & M. Stauber & L. Beck (Eds.), *Psychosomatische Gynäkologie und Geburtshilfe* (pp. 74-82). Berlin, Heidelberg, New York, Tokyo: Springer.
- Theut, S. K., Pedersen, F. A., Zaslow, M. J., & Rabinovich, B. A. (1988). Pregnancy subsequent to perinatal loss. Parental anxiety and depression. *Journal of American Academy of Child and Adolescent Psychiatry.*, 27(3), 289-292.
- Thomas, D. F. M. (2001). Prenatal diagnosis: Does it alter outcome? *Prenatal Diagnosis*, 21, 1004-1011.
- van Zuuren, F. J. (1993). Coping style and anxiety during prenatal diagnosis. Special issue: Psychological aspects of fetal screening and the new genetics. *Journal of Reproductive and Infant Psychology*, 11(1), 57-59.
- Zlotogorski, Z., Tadmor, O. P., Duniec, E., Rabinowitz, R., & Diamant, Y. Z. (1996). The effect of the amount of feedback on anxiety levels during ultrasound scanning. *Journal of Clinical Ultrasound*, 24(1), 21-24.
- Zlotogorski, Z., Tadmor, O., Duniec, E., Rabinowitz, R., & Diamant, Y. (1995). Anxiety levels of pregnant women during ultrasound examination: Coping styles, amount of feedback and learned resourcefulness. *Ultrasound Obstetrics and Gynecology*, 6, 425-429.
- Zuskar, D. M. (1987). The psychological impact of prenatal diagnosis of fetal abnormality: Strategies for investigation and intervention. *Women and Health*, 12(1), 91-103.

Table 3: Socio-demographic Data

Variable	Control Group n = 168	High Risk Groups n = 506	Test
Maternal Age	mean = 30.7 (sd=3.2; range=20-37)	mean = 32.3 (sd=4.7; range=17- 45)	t=-4.266 (df=668), p=.000***
School Education			
Senior High School	41.5%	30.3%	Chi- Square=9.734 (df=3), p=.021*
Junior High School	40.3%	42.0%	
Ext. Elementary School	18.2%	26.7%	
Special School Education	0%	1.0%	
Professional Education			ns
Professional Status			ns
Marital Status:			ns
married	81.4%	84.3%	
not married	14.9%	12.5%	
divorced/separated	3.7%	3.2%	
Living with a Partner			ns
yes	96.2%	98.4%	
no	3.8%	1.6%	

Note: Level of significance: *p<0.05, **p<0.01, p***<0.001.

t= Independent Samples T-test; Chi-Square = Pearson Chi-Square Test

Table 5: Trait Anxiety and Persistent High Levels of State Anxiety at Different Points in Time