

My Paper

Julian Reif*

January 3, 2020

Abstract

This paper provides an example of a document with tables and figures that were automated using Stata.

*University of Illinois and NBER.

1 Summary

This example paper includes tables and figures that were created using Stata and outputted into the `/analysis/results` project folder. Copy the contents of that folder to `/paper` to update the tables and figures in this document.

Table 1 reports summary statistics for Stata’s **auto.dta** dataset.¹ The average price of automobiles in this dataset is \$6,165. The price distribution, illustrated in Figure 1, is skewed right.

I estimate the association between automobile prices and fuel efficiency using the following linear model:

$$PRICE_i = \alpha + \beta X_i + \varepsilon \tag{1}$$

The outcome variable, $PRICE_i$, is the price of automobile i . The parameter of interest is β , a vector of coefficients. In my first specification (“spec 1”), the vector X_i includes miles per gallon. The second specification (“spec 2”) also includes the car’s weight. I estimate this model using ordinary least squares and report standard errors that are robust to heteroskedasticity. The analysis is performed first using Stata, and then repeated using R.


Table 2 reports my Stata estimates, separately for domestic and foreign cars. Column (1) reports that an increase in fuel efficiency of 1 mile per gallon is associated with a \$329 reduction in the price of domestic automobiles. Column (2) shows that this association becomes positive and insignificant when I also include weight as a regressor. Columns (3) and (4) show that these associations are similar for foreign automobiles.

Table 3 compares these Stata estimates to estimates from R. Panel A reproduces the Stata estimates that were presented in Table 2. Panel B of Table 3 reports estimates when I repeat this analysis in R using the **lm_robust** command from the *estimatr* package. The point estimates and the standard errors are identical across both software packages.

¹Type **sysuse auto, clear** at the Stata prompt to load this dataset.

2 Figures and Tables

Figure 1: Automobile prices



Notes: Data were obtained from Stata's built-in auto dataset.

Table 1: Summary statistics

	Mean	Stdev.	Min	Max	Count
Weight (pounds)	3,019	777	1,760	4,840	0
Miles per gallon	21.3	5.79	12	41	0
Price (1978 dollars)	6,165	2,949	3,291	15,906	0

Notes: Count reports the number of non-missing values for the variable.

Table 2: Association between automobile price and fuel efficiency

	(1)	(2)	(3)	(4)
	Domestic		Foreign	
	Spec 1	Spec 2	Spec 1	Spec 2
Miles per gallon	-329*** (81.2)	238 (203)	-250** (88.2)	-19.8 (51.7)
Weight (pounds)		4.42*** (1.34)		5.16*** (0.770)
N	52	52	22	22
R^2	0.254	0.483	0.399	0.785

Notes: Outcome variable is price (1978 dollars). Columns (1) and (2) report estimates of β from equation (1) for domestic automobiles. Columns (3) and (4) report estimates for foreign automobiles. Robust standard errors are reported in parentheses. A */**/** indicates significance at the 10/5/1% levels.

Table 3: Association between automobile price and fuel efficiency, Stata and R

	(1)	(2)	(3)	(4)
	Domestic		Foreign	
	Spec 1	Spec 2	Spec 1	Spec 2
A. Stata output (regress)				
Miles per gallon	-329*** (81.2)	238 (203)	-250** (88.2)	-19.8 (51.7)
Weight (pounds)		4.42*** (1.34)		5.16*** (0.770)
B. R output (lm_robust)				
Miles per gallon	-329*** (81.2)	238 (203)	-250** (88.2)	-19.8 (51.7)
Weight (pounds)		4.42*** (1.34)		5.16*** (0.770)

Notes: Outcome variable is price (1978 dollars). Columns (1) and (2) report estimates of β from equation (1) for domestic automobiles. Columns (3) and (4) report estimates for foreign automobiles. Robust standard errors are reported in parentheses. A */**/** indicates significance at the 10/5/1% levels.