

SUMMER

2026

July *Skies*

NORTHERN HEMISPHERE



clearskies.starstucklab.com

"The most beautiful thing we can experience is the mysterious."

— Albert Einstein

Sky Planner — July 2026

 Full: Jul 29 ·  Last Qtr: Jul 7 ·  New: Jul 14 ·  First Qtr: Jul 21

DATES	BEST TARGETS	BEST TIME	NOTE
Jul 1–6	M13 overhead, M57, Albireo, M27 rising E	9:30pm–1am	Waning Moon rises late — dark early evening
Jul 9	★ Venus 1° from Regulus — low WNW at dusk	8–8:45pm	Brightest planet beside Leo's heart
Jul 10–19	★ All deep sky — best dark skies of month	9:30pm–3am	New Moon Jul 14 — prime Milky Way window
Jul 20–24	M22, M11, M8 Sagittarius field, Moon (first quarter)	10pm–1am	First quarter Jul 21 — lunar terminator detail
Jul 25–28	M27, Albireo, Saturn rising late, Moon	10pm–midnight	Brightening Moon limits faint deep sky
Jul 29–31	Full Moon, Saturn, δ-Aquariid meteors (washed out)	10pm–midnight	Full Moon Jul 29 — skip faint targets

M13 → high south & overhead after 9:30pm · M57/Albireo → near zenith by midnight · M8/M22 Sagittarius → due south, low, 11pm–12:30am · M11 → south by 11pm · M27 → high east, best after midnight · Venus → very low WNW at dusk (1° from Regulus Jul 9) · Saturn → rises ESE late evening, best after midnight

The Moon — July 2026



First Qtr — Jul 21

Mare Crisium

The "Sea of Crises" — an isolated dark oval near the Moon's eastern limb, 555km across and ringed by mountains. Alone amid bright highlands, it is the first feature to catch the eye in the early crescent. At 36× the smooth floor and sharp rim stand out, foreshortened into a slim oval near the limb.

Best: Jul 16–18 · 25mm (36×)

Theophilus–Cyrillus–Catharina

A chain of three great craters on the western edge of Mare Nectaris. Theophilus (100km) is youngest, with terraced walls and a central peak; it overlaps the older Cyrillus, which joins the battered Catharina. With the terminator across them the shadows are dramatic at 100× — a fine crater-trio.

Best: Jul 18–20 · 9mm (100×)

Montes Apenninus

The Moon's grandest mountain range — a 600km arc along the southeastern shore of Mare Imbrium, with peaks over 5km. Near first quarter, sunrise lights the summits while the lowlands stay black and the range leaps into relief. Glorious at 36×; at 100× the peaks throw long shadows across the mare.

Best: Jul 20–22 · 25mm (36×)

Plato crater

A 101km crater with a remarkably smooth, dark lava-flooded floor, ringed by the peaks of the lunar Alps on Mare Imbrium's north shore. Its flatness and dark tone make it unmistakable. Just after first quarter the low sun throws the rim's shadow across the floor; hunt the tiny craterlets within at 100×.

Best: Jul 21–24 · 9mm (100×)

Planets & Events — July 2026

Venus



Constellation	Leo
Visible	Dusk, very low WNW
Sets after sunset	~1.5–2 hrs
Brightness	mag –4.5
At 36× (25mm)	Large thin crescent, ~45–55"
At 100× (9mm)	~25–30% lit, thinning crescent

Venus blazes in the WNW twilight as the brightest "star" in the sky, but it is dropping fast toward inferior conjunction in mid-August — sitting lower and thinner each evening. In the scope it is a large, dramatically thin crescent, obvious even at 36×. From Jul 4–14 it slides past Regulus, the heart of Leo, closest on Jul 9 at just 1.0° apart — a beautiful pairing of the brightest planet with a 1st-magnitude star. Catch it early in twilight from a clear western horizon before it sets.

Saturn



Constellation	Pisces
Visible	Late evening → morning, rises ESE
Rises after dark	~11pm (Jul 1) → ~9:30pm (Jul 31)
Brightness	mag +0.8
At 36× (25mm)	Pale disc, Titan visible, ring as a line
At 100× (9mm)	Rings only ~6–7° open — a slim bright ellipse

Saturn is climbing back into the late-evening sky, brightening week by week toward its October opposition. But its rings — edge-on through 2025 — have reopened only to about 6–7° in 2026, so instead of the wide-open showpiece you see a slim bright ellipse hugging the disc, almost a line drawn through the planet. It is still unmistakable, and Titan is easy at 36×. The narrow tilt puts the Cassini Division out of reach this year. Best after midnight, once Saturn climbs clear of the eastern haze.

★ Venus meets Regulus — July 9

The brightest planet passes just 1.0° from Regulus, the blue-white heart of Leo, low in the WNW about 45 minutes after sunset. Both fit easily in the same 36× field — Venus a large, brilliant crescent, Regulus a steady point of light beside it. Naked eye they make a striking close pair in the fading twilight. You will need a clear, low western horizon...

Milky Way core at its best

Through July the galactic core in Sagittarius and Scorpius rides due south in the late evening — the richest, brightest stretch of our galaxy. From a dark site the Milky Way band is an obvious naked-eye river of light, and the region is packed with deep-sky targets: the Lagoon (M8), the great globular M22, and the dense star-swarm of M1...

Delta Aquariid & Capricornid meteors — peak Jul 30

Two overlapping showers — the Southern Delta Aquariids and Alpha Capricornids — peak together on the night of Jul 30–31, radiating from low in the south. Together they can deliver 15–25 meteors an hour from a dark sky, including the occasional slow, bright Capricornid fireball. The catch this year: the Full Moon of Jul 29 floods the sky...

M13 — Great Hercules Cluster

Globular cluster · Hercules · mag 5.8 · 22,000 ly

★★★★★ OBSERVING EASE

BEST DATES Jul 1–28	BEST TIME 9:30pm–1am	DIRECTION Near the zenith, overhead
START WITH 25mm (36×)	THEN TRY 9mm (100×)	NEEDS Dark sky helps but not essential

DESCRIPTION

At 36× a large, obvious cotton-ball glow — big, bright, and unmistakable. At 100× the halo is rich and granular, the core blazes, and on a steady night the very edge of the halo breaks into individual star-points. In July M13 rides almost exactly overhead in the early evening — the best possible position, looking through the thinnest, steadiest column of air. The finest globular cluster in the northern sky and one of the best objects for any telescope. Show this first on any observing session.



9mm · 100×
rich halo, resolved edge stars

FINDER

Find the Keystone of Hercules — four stars forming a slightly squashed rectangle, nearly overhead at nightfall in July. M13 sits on the western edge of the Keystone, one-third of the way down from the top star (Eta) toward the bottom (Zeta). On a dark night it is faintly naked-eye as a fuzzy "star". Centre it in the 25mm, then switch to 9mm for the texture.

M8 — Lagoon Nebula

Emission nebula + open cluster · Sagittarius · mag 6.0 · ~4,100 ly

★ ★ ★ ★ ★ OBSERVING EASE

△ Low in the south — wait until it transits and the haze settles

BEST DATES Jul 10–24	BEST TIME 10:30pm–12:30am (transit ~11:30pm)	DIRECTION Due south, ~33° altitude
START WITH 25mm (36×)	THEN TRY 9mm (100×)	NEEDS Dark sky, clear southern horizon, no Moon

DESCRIPTION

One of the showpieces of the summer sky. At 36× a large, irregularly glowing cloud divided by a prominent dark lane — the "lagoon" that gives it its name — with the young star cluster NGC 6530 sprinkled across the eastern half. The overall extent is ~1.5° × 0.5°, so the 25mm field frames it comfortably. At 100× zoom into the embedded cluster and look for the bright knot of nebulosity near centre — the Hourglass region. Sitting at only ~33° altitude from 32°N, wait for it to transit due south for the steadiest, brightest view.



25mm · 36×
large glow, dark lane, star cluster

FINDER

Find the "Teapot" asterism in Sagittarius, low in the south. The lid star is Kaus Borealis (Lambda Sgr); M8 lies ~5° west-northwest of it, just above the spout. At 36× it is immediately obvious as a large, softly glowing cloud — far bigger than any globular, with visible extent at a glance. The dark lane is subtle but real on a transparent, moonless night.

M22 — Sagittarius Globular

Globular cluster · Sagittarius · mag 5.1 · ~10,400 ly

★★★★★ OBSERVING EASE

△ Low in the south — needs a clear horizon and transit timing

BEST DATES Jul 10–26	BEST TIME 11pm–1am (transit ~12am)	DIRECTION Due south, ~34° altitude
START WITH 25mm (36×)	THEN TRY 9mm (100×)	NEEDS Clear southern horizon, dark sky

DESCRIPTION

The brightest globular cluster easily visible from northern mid-latitudes, and one of the closest at ~10,400 light-years — only the far-southern Omega Centauri and 47 Tucanae outshine it. At mag 5.1 it is larger and looser than M13, and because it is so near, a 114mm resolves its outer halo into a swarm of star-points more readily than almost any other globular. At 36× it is a big, bright, slightly squashed glow; at 100× the granular edge breaks into individual suns on a steady night. The tragedy is its low altitude — only ~34° from 32°N — so wait for transit and a transparent sky to see it at its best.



9mm · 100×
large, partly resolved halo

FINDER

M22 sits just above the lid of the Sagittarius "Teapot". Find Kaus Borealis (Lambda Sgr), the top of the lid, then move ~2.5° to the northeast. M22 lies almost on the line between Kaus Borealis and the brighter Nunki, much closer to Kaus Borealis. At 36× sweep slowly for a large, soft round glow brighter than its low position would suggest — it is far easier than M8's nebulosity once you are on it.

M11 — Wild Duck Cluster


Open cluster · Scutum · mag 6.3 · ~6,200 ly

★★★★★ OBSERVING EASE

BEST DATES Jul 10–28	BEST TIME 10:30pm–1am (transit ~11pm)	DIRECTION South, ~52° altitude
START WITH 25mm (36×)	THEN TRY 9mm (100×)	NEEDS Dark sky brings out the richness

DESCRIPTION

One of the richest and most compact open clusters in the sky — so dense it almost looks like a loose globular. Some 2,900 stars are packed into a tight fan, set against one of the brightest star-clouds of the Milky Way (the Scutum Star Cloud). At 36× it is a bright, compact, slightly triangular haze with a single brighter star at its leading edge — the "head of the flock" that gives the Wild Duck its name. At 100× it explodes into a glittering wedge of dozens of resolved stars against a misty backdrop of fainter ones. A wonderful contrast to the smooth globulars — here every point is a sun you can pick out individually.



9mm · 100×
dense wedge, lead star at the head

FINDER

Follow the Milky Way south from Albireo down into the small constellation Scutum, just east of the Sagittarius Teapot. The cluster sits at the northern end of the bright Scutum Star Cloud. From the star Lambda Aquilae, hop ~3° southwest to a curving line of stars in Scutum; M11 lies at the end of that line. At 36× it is an obvious knot, denser and more concentrated than any other open cluster in the area.

M57 — Ring Nebula

Planetary nebula · Lyra · mag 8.8 · ~2,300 ly

★★★★★ OBSERVING EASE

BEST DATES Jul 1–28	BEST TIME 10pm–2am (near zenith by midnight)	DIRECTION Overhead, high east
USE 9mm (100×)	NEEDS Steady seeing, dark sky	CENTRAL STAR Not visible at 114mm

DESCRIPTION

The quintessential planetary nebula — the expelled shell of a dying star. At 100× it appears as a tiny, unmistakable grey-green smoke ring; a small oval distinctly darker at its centre. The central white dwarf is NOT visible at 114mm (it needs 200mm and up). The ring shape itself is clear and immediately satisfying. It is small — barely 1 arcminute across — so 100× is the minimum useful magnification. In July M57 rides high overhead through the late evening, giving the steadiest possible view; this is the best time of year to chase its delicate ring.



9mm · 100×
grey-green smoke ring

FINDER

Find Vega, the brilliant blue-white star almost overhead in the east. Lyra is the small parallelogram of stars hanging just south of Vega. M57 sits exactly halfway between the two southern stars of the parallelogram — Sheliak (Beta Lyrae) and Sulafat (Gamma Lyrae). At 100× look for the small out-of-focus smudge that will not sharpen, then look again for the darker centre — that is the ring.

Albireo — β Cygni

Double star · Cygnus · mags 3.1 + 5.1 · sep 34" · 430 ly

★★★★★ OBSERVING EASE

BEST DATES Jul 1–31	BEST TIME 10pm–3am (high overhead by midnight)	DIRECTION High east, climbing to the zenith
USE 25mm (36×) or 9mm (100×)	SEEING Any conditions — easy split at 34"	MOON CAUTION Any phase fine — bright pair

DESCRIPTION

The most celebrated colour-contrast double star in the sky. At 36× two stars split cleanly — one a warm golden-amber (the primary K-type giant), the other a piercing ice-blue (a hotter B-type star). The colour contrast is vivid and unlike anything else at the eyepiece. With 34 arcseconds of separation the pair splits with room to spare even at 36×. Cygnus climbs high overhead through July evenings, so Albireo is beautifully placed all night — and gets better as it rises and the sky darkens. The single best object to show anyone who has never looked through a telescope.



25mm · 36×
gold + blue, vivid contrast

FINDER

Find Vega blazing high in the east, then locate the large cross-shape of Cygnus — the Northern Cross — nearby, with bright Deneb at its top. Albireo is the star at the very foot of the cross, the "head of the swan". Naked eye it looks like a single faint star. Point the 25mm at it and prepare for the colour.

M27 — Dumbbell Nebula


Planetary nebula · Vulpecula · mag 7.4 · ~1,360 ly

★★★★★ OBSERVING EASE

BEST DATES Jul 10–31	BEST TIME 11pm–3am (high by midnight)	DIRECTION High east, near the zenith later
START WITH 25mm (36×)	THEN TRY 9mm (100×)	NEEDS Dark sky, averted vision

DESCRIPTION

The brightest and largest planetary nebula in the sky — and the easiest to see well in a small scope. Where M57 is a tiny ring, M27 is big and bold: at mag 7.4 and roughly 8 arcminutes across, it is obvious at 36× as a glowing grey patch. At 100× its distinctive shape emerges — a bright rectangular bar pinched in the middle like an apple core or a bow-tie, with fainter wings filling out the sides into the full "dumbbell". Averted vision swells it noticeably. No central star is visible at 114mm, but the structure itself is among the most rewarding deep-sky sights of summer. Climbing high in the east through July evenings, it is best after midnight near the zenith.



9mm · 100×
bright apple-core bar + faint wings

FINDER

Find Albireo at the foot of the Northern Cross, then drop ~8° south to the small arrow-shaped constellation Sagitta. M27 lies ~3° north of the arrow's tip (the star Gamma Sagittae), in neighbouring Vulpecula. At 36× sweep slowly north of Sagitta for a soft, sizeable grey glow — far larger and easier than M57. Once found, switch to 100× and use averted vision to tease out the pinched dumbbell waist.

Observing Conditions

Seeing (steadiness)

Poor seeing makes stars shimmer and prevents double-star splits. Good seeing gives steady, sharp images. Seeing varies nightly — try a different night if stars are boiling.

Transparency (clarity)

High transparency means dry, dust-free air. Low transparency hides faint galaxies. Haze and humidity are the enemy of faint objects.

Dark adaptation

Your eyes need ~20 minutes in full darkness to reach peak sensitivity. One bright light flash resets the process. Use red light only in the field.

Light pollution

Bortle 1–4 skies reveal galaxy detail and faint nebulosity. The Moon is the single biggest factor night-to-night.

Pre-session checklist

- ☐ Check seeing/transparency forecast (Clear Outside · Meteoblue)
- ☐ Dark sky window: **Jul 10–19**
- ☐ Cool telescope ≥ 20 min before observing
- ☐ Collimate if last session was ≥ 2 weeks ago
- ☐ Always start at 25mm (36 $\times$) — low-power first
- ☐ Switch to 9mm (100 $\times$) only after locating target
- ☐ 20 min dark adaptation before faint objects
- ☐ Use averted vision for galaxies and faint nebulae

Glossary

Magnitude (mag) Brightness scale — lower is brighter. Sun -26, naked-eye limit ~6, your scope reaches ~12.	Globular cluster A tight spherical swarm of 100,000–1,000,000 ancient stars gravitationally bound together. M13 and M22 are both prime July targets.	Open cluster A loose, young group of dozens to a few thousand stars born from the same gas cloud, scattered along the Milky Way. M11 (Wild Duck) is an unusually rich example.
Emission nebula A gas cloud ionised by nearby hot young stars, glowing in its own light. M8 (Lagoon) is a classic summer example.	Planetary nebula The expelled outer shell of a dying star, briefly glowing before it fades. M57 (a tiny ring) and M27 (a big dumbbell) are the two finest — nothing to do with planets.	Double star Two stars close together in the eyepiece. Albireo is the showpiece, splitting into a vivid gold-and-blue colour-contrast pair.
Averted vision Looking slightly off-centre to use the eye's more sensitive rod cells. Essential for globular halo texture and the faint wings of M27.	Seeing Atmospheric steadiness. Poor seeing blurs fine detail and prevents globular resolution even under a dark sky.	Transparency Atmospheric clarity. Poor transparency dims faint nebulae and the low Sagittarius objects even when stars look sharp.
Radiant The point in the sky a meteor shower appears to stream from. The Delta Aquariid radiant sits low in the south, rising after midnight.	Inferior conjunction When Venus passes between Earth and the Sun, briefly becoming invisible. Venus races toward this in mid-August 2026.	Milky Way core The dense galactic centre in Sagittarius, visible as a bright cloudy band low in the south on dark July nights from 32°N.

Your 114mm f/7.9 reflector's limits: limiting magnitude ~12 under dark skies · max useful magnification ~230× (100× is the practical ceiling most nights) · 265× more light than the naked eye · resolving power ~1 arcsecond on the steadiest nights · M22's nearness makes it the most resolvable globular this month, while M13 shows a full granular halo but needs 150mm+ for complete stellar resolution. Saturn's rings are nearly edge-on in 2026, so the Cassini Division is beyond reach until they open wider in coming years.

CLEAR SKIES — JULY 2026

Anish Mangal

With thanks to: Ishan Chungoo, Sohail Lalani

114mm f/7.9 Newtonian · 900mm focal length · 25mm (36×) & 9mm (100×)

"The most beautiful thing we can experience is the mysterious."

— Albert Einstein

