MANUAL.md — 3d_rt_sandbox

User manual for `rt3d`, a 3D all-sky, monochromatic short-characteristics radiative-transfer solver written in pure NumPy.

This manual documents the **actual code** in `rt3d/` (phases 0–6 complete, 80 tests green). The agreed design and the decisions log live in `plan/plan.md` — that file stays the source of truth for *why* a decision was made; this manual explains *what the code does and how to use it* and cross-references the plan instead of duplicating the log.

Contents

- Part I — Theory
  1. Scope and physical assumptions
  2. Governing equation
  3. The solar beam split
  4. Angular discretization — icosahedral direction grids
  5. Spatial discretization — unigrid and octree
  6. Short-characteristics transport
  7. Boundary conditions
  8. Iteration scheme and 1D Ng acceleration
- Part II — Usage 9. Environment setup 10. Module tour 11. Worked end-to-end example 12. Runner scripts 13. Condensed API reference 14. Validation anchors 15. Known limitations and performance

---

Part I — Theory

1. Scope and physical assumptions

`rt3d` solves the **scalar radiative transfer equation (RTE)** in a 3D rectangular domain under the following fixed physical assumptions (plan section 1; do not change them without updating `plan/plan.md`):

Assumption	Meaning in the code
Monochromatic	One frequency at a time. All intensities, B, B_g, F_0 are values at the working frequency. No line structure, no frequency coupling.
LTE	Local source function $S = (1-\omega)B + \omega J$ (plus scattering anisotropy, see §2).
Scalar RTE	No polarization. The Stokes machinery in `scripts/icosphere_grid.py` (`rotater_mat`, `rotatepoles`) is legacy and unused.
No refraction	Rays are straight lines; the DDA marches of `rt3d/geometry.py` and `rt3d/octree.py` traverse straight rays.
Henyey-Greenstein phase function only	The only scattering phase function; asymmetry parameter g may be scalar or per-cell.
Periodic x and y	Rays exiting a side face re-enter the opposite face (`periodic_box` argument throughout).
Top = space, bottom = flat ground	Top inflow is a constant background `top_I` (default 0); the bottom face may carry a Lambertian + emissive ground (`ground_surface`).
Solar radiation as a direct beam	A collimated beam in one direction, handled separately from the diffuse field (§3).

The spatial domain is an axis-aligned **cuboid** (`rt3d.geometry.cuboid`): $z = z_1$ is space (top), $z = z_0$ is the ground plane (bottom), x and y wrap periodically.

2. Governing equation

For monochromatic intensity $I(\mathbf{r}, \mathbf{\Omega})$, LTE, scalar RTE:

$$\mathbf{\Omega} \cdot \nabla I(\mathbf{r}, \mathbf{\Omega}) \;+\; \kappa_e(\mathbf{r})\, I(\mathbf{r}, \mathbf{\Omega}) \;=\; \kappa_s(\mathbf{r}) \oint \Phi_{\mathrm{HG}}(\mathbf{\Omega}' \to \mathbf{\Omega})\, I(\mathbf{r}, \mathbf{\Omega}')\, d\Omega' \;+\; \kappa_a(\mathbf{r})\, B(\mathbf{r})$$

with the derived quantities **always computed, never stored as inputs** (AGENTS.md input conventions):

$$\kappa_a = \kappa_e - \kappa_s \qquad (\text{absorption coefficient}), \qquad \omega = \frac{\kappa_s}{\kappa_e} \qquad (\text{single scattering albedo})$$

ω is computed by `rt3d.source.single_scattering_albedo` with a **vacuum guard**: where $\kappa_e = 0$ (vacuum cells) the division is skipped and $\omega = 0$ is used, so vacuum cells neither scatter nor absorb.

The **Henyey-Greenstein** phase function (`rt3d.directions.hg`) is normalized so that its exact integral over solid angle is 1:

$$\Phi_{\mathrm{HG}}(\cos\Theta) = \frac{1 - g^2}{4\pi\,(1 + g^2 - 2g\cos\Theta)^{3/2}}, \qquad -1 < g < 1$$

where Θ is the scattering angle between incoming $\mathbf{\Omega}'$ and outgoing $\mathbf{\Omega}$. With $S = (1-\omega)B + \omega \oint \Phi\, I\, d\Omega'$ the RTE takes the form

$$\frac{dI}{ds} = \kappa_e \left(S - I\right)$$


along a ray, which is the form the transport recursion of §6 integrates exactly.

3. The solar beam split

The incoming solar radiation is a **direct beam**: a collimated bundle in one unit direction $\mathbf{\Omega}_{\text{sun}}$ (must have $\Omega_{\text{sun},z} < 0$) with irradiance F_0 at the domain top. A direction-delta distribution $I_{\text{beam}} \propto \delta(\mathbf{\Omega} - \mathbf{\Omega}_{\text{sun}})$ **cannot live on the icosahedral quadrature** of §4 — no ordinate carries it, and projecting it onto the ordinates would corrupt the scattering integral. The solver therefore splits

$$I = I_{\text{diff}} + I_{\text{beam}}$$

and treats the two parts separately:



1. **Beam sweep (no angular quadrature).** The beam intensity at every cell center is the analytic

$$I_{\text{beam}}(\mathbf{r}) = F_0\, e^{-\tau_b(\mathbf{r})}, \qquad \tau_b(\mathbf{r}) = \int_{\text{entry}}^{\mathbf{r}} \kappa_e\, ds$$

   where the beam optical depth τ_b is accumulated along the *exact* backward ray from each cell center to the domain entry point: `rt3d.transport.beam_optical_depth` (unigrid) and `beam_optical_depth_octree` (octree). The beam is never transported on the ordinate set.

2. **First-scattering source (fixed, never iterated).** The beam is scattered once into the diffuse field, contributing to the source function

$$S_{\text{beam}}(\mathbf{r}, \Omega_i) = \omega(\mathbf{r})\, \Phi_{\text{HG}}(\boldsymbol{\Omega}_{\text{sun}} \to \Omega_i)\, F_0\, e^{-\tau_b(\mathbf{r})}$$

   (`rt3d.source.beam_scattering_source`). This term is a *fixed* source contribution: it is added to S in every iteration but never rebuilt from an iterated intensity, because the beam itself is converged analytically from the start.

3. **Beam first-scatter HG row normalization.** The phase-function row $\Phi_i = \Phi_{\text{HG}}(\boldsymbol{\Omega}_{\text{sun}} \to \Omega_i)$ is rescaled so that $\sum_i w_i \Phi_i = 1$ exactly — the same discrete quadrature convention as the row-normalized scattering matrix (§4). Without it, the beam first-scatter power passing through a coarse ordinate set would carry the raw HG quadrature error (at 42 ordinates and $g = 0.952$ the raw sum is ≈ 0.27, i.e. a factor ~3.7 error).

The ground also sees the beam: its Lambertian reflection $A_g F_0 |\mu_{\text{sun}}| e^{-\tau_{b,\text{ground}}}/\pi$ is part of the fixed ground inflow (§7).

4. Angular discretization — icosahedral direction grids

The direction grid is **always** an icosahedral (geodesic) grid — never a polar/azimuthal product grid. Construction (`rt3d.directions`):

- base icosahedron (12 vertices, 20 spherical triangles),
- geodesic refinement: bisect every edge, project to the sphere, split each triangle into four (vectorized edge-indexed implementation; replaces the $O(N^2)$ dedup loops of `scripts/icosphere_grid.py`),
- $N_{\text{dir}} = 10 \cdot 4^{n_{\text{refine}}} + 2 \to$ **12 / 42 / 162 / 642** ordinates for $n_{\text{refine}} = 0/1/2/3$.

42 ordinates ($n_{\text{refine}} = 1$) is the development grid: all development, test cases, benchmarks and validation runs use it. n_{refine} is configurable everywhere (`icosahedric_grid(n_refine= ... )`) and recorded in every output file; no code assumes a specific N_{dir}. Levels 162/642 belong to the phase 6 convergence study.

Solid-angle weights come from **spherical-triangle areas**, not $4\pi/N$: each triangle's spherical excess (L'Huilier's formula) is split equally among its three corner vertices (`solid_angle_weights`). Icosphere vertices are not equal-area, so this distinction matters. The weights satisfy $\sum_i w_i = 4\pi$ to machine precision (validation anchor 1).

Equator guard. On construction the grid is rotated about a generic axis (`direction_grid._rotate_off_poles`) so that **no ordinate has exactly** $\Omega_z = 0$. Ordinates with $\Omega_z = 0$ form cyclic characteristics under x/y periodicity (a ray never reaches the top or bottom boundary). The guard is $\min_i |\Omega_z| > 10^{-6}$; measured $\min_i |\Omega_z|$ is 0.42 / 0.105 / 0.034 / 0.013 for 12 / 42 / 162 / 642 ordinates.

Precomputed HG matrix. The scattering integral over *all* cells is one BLAS matmul `(n_cells × N_dir) @ (N_dir × N_dir)`. `direction_grid.hg_matrix(g)` returns $W_{ij} = \Phi_{\mathrm{HG}}(\Omega_j \to \Omega_i)\, w_j$ (weights of the *incoming* direction folded in).

Row normalization and why. The analytic HG function integrates to 1 over the sphere, but the coarse icosahedral quadrature breaks this at large g: at 42 ordinates and $g = 0.952$ the raw weighted row sums are **16.9-21.6** (not 1). The λ-iteration scattering operator then has spectral radius $\gg 1$ − it is *supercritical* and the iteration diverges for ω not far below 1. `hg_matrix(g, normalize=True)` (the default) rescales each row so that $\sum_j W_{ij} = 1$ exactly. This enforces exact discrete conservation of isotropic radiation; at $g = 0$ the raw row sums are already exactly 1, so isotropic results are unchanged. The same convention is used for the beam first-scatter row (§3). The price is a modified discrete phase function; the angular error it introduces is part of the 42-ordinate error budget (§15).

5. Spatial discretization — unigrid and octree

Two spatial grids, both exposing cell/leaf centers, boxes and O(1) point location to the transport:

`rt3d.geometry.unigrid` − uniform Cartesian lattice defined by cell-center coordinate arrays x, y, z. Cell (i, j, k) has flat index $k\, n_y n_x + j\, n_x + i$ (i fastest); the domain box extends half a cell beyond the outermost centers. A single-cell dimension gets a nominal unit width, so a plane-parallel slab is $n_x = n_y = 1$ with periodic sides.

`rt3d.octree.octree_grid` − flat struct-of-arrays octree over the cuboid, built from cubic base cells of size h_0 (must divide all side lengths), cubic cells, max-depth cap, node arrays `centers_all / half_all / level_all / children_all`. Leaf-indexed public arrays (`centers`, `half`, `level`, `kappa_e`, `kappa_s`, `g`, `T`) are frozen by `finalize()`.

Refinement criteria (`octree_grid.refine`, driven by the *input* extinction field, sampled at the 8 cell corners):

$$\kappa_{e,\max}\,\Delta x > \tau_{\max} \qquad \text{(optical depth per cell)}$$

$$\frac{\kappa_{e,\max} - \kappa_{e,\min}}{\langle \kappa_e \rangle} > \varepsilon_{\mathrm{grad}} \qquad \text{(cloud edges)}$$

The gradient criterion is applied **only where the cell is not effectively transparent**, gated by $\kappa_{e,\max}\,\Delta x > 10^{-2}\,\tau_{\max}$. Rationale: in the vacuum tails of a cloud the *relative* κ_e gradient is large but physically irrelevant − without the floor the octree would refine the exponentially small Gaussian tails down to the depth cap.

2:1 face balance (`balance_21`): a leaf may differ from any face neighbor by at most one level. Each face is probed at a **3×3 grid** of points (transverse offsets $-h/2, 0, +h/2$), which is guaranteed to hit a deeper neighbor of any allowed size; the loop repeats until no splits remain.

O(1) point location (`locate`): vectorized descent from the base lattice (`floor` to base cell, then octant selection per level) − no per-level dense tables.

Why the solver uses exact characteristics on the octree. The phase-3 reference implementation (`characteristics.build_chain_octree`) connects each leaf to the *best-aligned center-to-center predecessor* (the octree analogue of the unigrid lattice stencil, §6). In heterogeneous 3D fields its hop alignment tolerance makes the **path error scale-invariant**: on the Gaussian cloud the characteristic τ error stays ≈ 4 no matter how much the


grid is refined — refining the grid does not converge the ray path. The solver therefore uses **exact characteristics** instead: `transport.precompute_octree_march` marches one ray *backwards from every leaf center along* $-\mathbf{\Omega}$ *to the domain entry point* (DDA through the octree, `octree.march_octree`), once per ordinate, and reuses the segments in every iteration. The only remaining path error is the octree's own resolution of the piecewise-constant field. On the unigrid the lattice-stencil chain is kept (its error is the documented anchor-5 effect, §14).

6. Short-characteristics transport

One characteristic per cell/leaf center per ordinate. Along a characteristic the medium is piecewise constant per traversed cell, and the formal solution

$$I_c = I_u\, e^{-\tau_{\mathrm{tot}}} + \sum_{\mathrm{sub}} S_{\mathrm{sub}} \left(1 - e^{-\tau_{\mathrm{sub}}}\right) e^{-\tau_{\mathrm{after}}}$$

(τ_{sub} = optical depth of sub-segment *sub*, τ_{after} = optical depth between that sub-segment and the target center) is **exact** for piecewise-constant sources. The code uses `-expm1(-tau)` for $1 - e^{-\tau}$ accuracy at small τ.


one ordinate Omega_i
grid type?
unigrid
octree
stencil_for_direction + subsegments_uniform + build_chain (once per ordinate)
precompute_octree_march: exact backward DDA march from every leaf center to domain entry (once per ordinate)
sweep_direction: boundary-entry root chains, then topological BFS waves
sweep_direction_octree_exact: block cumsum prefixes + np.add.at
I at all cell/leaf centers for this ordinate
sweep_all / sweep_all_octree_exact loop over all ordinates

Unigrid (`characteristics.build_chain` + `transport.sweep_direction`). For one ordinate $\mathbf{\Omega}$ the upwind cell of c is $c - s$ with a single integer **lattice stencil** s per ordinate (`stencil_for_direction`: the offset with $s_z \neq 0$ whose lattice direction $s \cdot h$ is closest to $\mathbf{\Omega}$; the search radius auto-scales with grid anisotropy, $\sim 3\max(h)/\min(h)$). The straight path $u_{\text{center}} \to c_{\text{center}}$ is split into **explicit sub-segments** (`subsegments_uniform`) whose relative offsets and lengths are the *same* for every cell. Since $s_z \neq 0$ always (§4 equator guard) and z is the only non-periodic dimension, every chain terminates at a z-boundary: the dependency graph is a forest, processed in **topological BFS waves** (`chain["waves"]`) — each wave is a handful of vectorized gathers. Chain roots carry an exact backward march (`march_lattice`) from the root center to the domain entry point.

Octree (`transport.precompute_octree_march` + `sweep_direction_octree_exact`). Segments of each backward ray are stored ray-contiguous (`leaf`, `length`, `offsets`, ordered *center → entry*). The recursion is then fully vectorized: per-segment optical depths $\tau_{\text{seg}} = \kappa_e[\text{leaf}] \cdot \ell$, block-wise cumulative sums via `_block_prefixes` (a single `np.cumsum` plus per-block offset correction — the "cumsum vectorization"), and `np.add.at` accumulation onto leaf centers. The march depends only on grid + direction, so it is built once per ordinate and reused by every iteration.

`transport.sweep_all` / `sweep_all_octree_exact` loop over all ordinates; the source S may be direction-dependent (`(n_cells, N_dir)`), which is what anisotropic scattering requires.


7. Boundary conditions

`rt3d.transport.boundary_conditions` holds constant background intensities `top_I`, `bottom_I`, `side_I` (sides ignored when periodic) plus an optional `ground`:

- **Top:** $I_{\text{diff}} =$ `top_I` (default 0 = space) for ordinates entering through $z = z_1$.
- **Bottom without ground:** $I_{\text{diff}} =$ `bottom_I`.
- **Lambertian + emissive ground** (`ground_surface(albedo, emissivity, B_g, height= ... )`): upward ordinates entering through the bottom face get the ground radiance

$$I_{\uparrow} = \varepsilon_g \, B_g + \frac{A_g}{\pi} \left(F_{\downarrow} + F_0 \, |\mu_{\text{sun}}| \, e^{-\tau_{b,\text{ground}}} \right), \qquad F_{\downarrow} = \sum_{\Omega_z < 0} w \, |\Omega_z| \, I$$

  where $F_{\downarrow}$ is the downward irradiance at ground level (rebuilt from the just-swept downward ordinates every iteration), and the last term is the **Lambertian reflection of the solar beam** (fixed part). Monochromatic convention: B_g is the Planck value at the working frequency, given as an input like every other B.





In the solver loop the **downward ordinates are swept first**, the ground radiance is assembled, then the upward ordinates are swept with that per-entry-point inflow (`bottom_I=` argument of the sweeps; on the octree the entry leaf comes from `march["entry_cell"]`, on the unigrid the $k = 0$ ground radiance tiles over all layers because chains with $|s_z| > 1$ root above the bottom layer).

- **Periodic sides:** implemented in the ray marches (`geometry.march_lattice`, `octree.march_octree`) as ray exit→entry wrap: a ray crossing a periodic x/y face is wrapped (`cuboid.wrap_xy`) and keeps marching; only top/bottom exits terminate it. *Wrap-stall bugfix note:* a ray that exits exactly (or within float noise) on a periodic face would be mapped by `mod()` back **onto the same face**, re-exit immediately with a zero-length step, and stall the march until `max_steps` without ever recording an exit. `march_lattice` therefore snaps such points to the *opposite* face (see the comment block in `geometry.march_lattice`).
- **Variable surface hook:** `ground_surface(height=f(x, y))` is accepted (`is_variable()`), but the transport supports only a flat ground at the domain bottom — `lambda_iteration` raises `NotImplementedError` for a variable surface (phase 6 hook, see §15).

8. Iteration scheme and 1D Ng acceleration

The solver (`rt3d.solver.lambda_iteration`) is structured like DOIT (compute incoming radiation → transport → next iteration):


S = (1-omega)*B + S_beam — fixed parts only
transport sweep over all ordinates (chains/marches built once)
S_new = (1-omega)*B + omega * mean(Phi I) + S_beam — incoming radiation
history: rel = max abs(S_new - S) / max abs(S_new)
ng active?
yes
no
ng1d.step: append history pair, fit Lambda_bar (Nz x Nz)
no



The initial source is $S^0 = (1-\omega)B + S_{\text{beam}}$ (fixed parts only — no radiation yet). The convergence history measures $\max|S^{\text{new}} - S| / \max|S^{\text{new}}|$ — the plain-LI residual of the *swept* source, also when Ng is active.

1D Ng acceleration (`rt3d.accelerator.ng1d`). Ng (1982) acceleration with a plane-averaged (z) estimate of the scattering operator $\bar{\Lambda}$ (plan section 5). It needs **no extra transport sweeps**: the Λ-iteration iterate differences themselves probe $\bar{\Lambda}$. With σ^n the source *actually swept* at iteration n and $S^{n+1} = \Lambda\,\sigma^n + Q$ the plain-LI update (Q = fixed source parts), the history pairs

$$x^{(m)} = \sigma^m - \sigma^{m-1}, \qquad y^{(m)} = \Lambda\,\sigma^m - \Lambda\,\sigma^{m-1} = S^{m+1} - S^m \qquad (Q \text{ cancels})$$

satisfy $\bar{\Lambda}\,x^{(m)} = y^{(m)}$ exactly for the 1D operator. $\bar{\Lambda} \in \mathbb{R}^{N_z \times N_z}$ is the minimum-norm least-squares fit of the **horizontally plane-averaged** pairs (`plane_average`: unigrid — one bin per z layer; octree — one bin per base cell z level, leaves weighted by their horizontal cross-section). The accelerated source is the geometric-series extrapolation

$$\sigma^{n+1} = S^{n+1} + \left(I - \bar{\Lambda}\right)^{-1} \bar{\Lambda}\, d^{n+1}, \qquad d^{n+1} = S^{n+1} - \sigma^n$$

applied per z bin. (Scalar check: for $\Lambda = \lambda$ this reproduces the fixed point in one step.)

Two subtleties encoded in the code:

- **Swept-source / consecutive-update pairing.** x must be the difference of the sources *actually swept*, and y the difference of *consecutive LI updates* — not the naive difference of successive iterates. Once acceleration is active, the swept source is no longer the plain LI iterate, and step-difference pairing would probe $(I - A)^{-1}\Lambda$ instead of Λ and break the fit.
- **Spectral-radius safeguard.** A bad fit with $\rho(\bar{\Lambda}) \geq 1$ would make the geometric series blow up; `ng1d.step` checks $\max_i |\mathrm{eig}_i(\bar{\Lambda})| \geq 1$ and falls back to plain LI for that step (also on non-finite corrections). Warmup: `n_start = 3` history pairs of plain LI before acceleration starts; at most `max_hist = 3·Nz` pairs are kept.

The scattering part of the source is direction-averaged in the history (scalar RTE, HG scattering ⇒ the accelerated part is isotropic), so the history is stored as `(n_cells,)` direction-averaged vectors.

Measured iteration counts to $\mathrm{tol} = 10^{-6}$ on the $\tau = 5$ slab, $g = 0$ (phase 4, `scripts/phase4_ng.py`): $\omega = 0.5$: **22** → **10**, $\omega = 0.9$: **87** → **13**, $\omega = 0.99$: **178** → **14** (LI → Ng).

Part II — Usage

9. Environment setup

Mamba/conda environment **`pyarts`**. There is no `mamba` / `conda` on `PATH` in a plain shell — call the environment interpreter by absolute path:

```
& C:\Users\fredd\miniforge3\envs\pyarts\python.exe <script.py>
```

> ⚠ **BLAS/openblas PATH gotcha — read this first.** The env's numpy links against the `libcblas.dll` shim, which *delay-loads* `openblas.dll` from `C:\Users\fredd\miniforge3\envs\pyarts\Library\bin`. Running the interpreter by absolute path **without activating the env** leaves that directory off `PATH`, and every matrix product then hard-crashes the process with exit code `-1066598273` (`0xC06D007F`, delay-load failure) — **no traceback**. Prepend it before running anything that does a matmul (which is everything in `rt3d`):
>
> ```
> $env:Path = "C:\Users\fredd\miniforge3\envs\pyarts\Library\bin;" + $env:Path
> ```

Installed packages (verified):

Package	Version
Python	3.14.6
numpy	2.5.2
scipy	1.18.0
matplotlib	3.11.1
xarray	2026.7.0
h5py	3.16.0
pyarts3	3.0.0.dev10
pytest	9.1.1

Not installed: `numba`, `pyvista` (verified `ModuleNotFoundError`). The solver is pure NumPy and does not need them (§15).

Quick sanity check:

```
& C:\Users\fredd\miniforge3\envs\pyarts\python.exe -c "import numpy, scipy;
print(numpy.__version__, scipy.__version__)"
```

Running the test suite (80 tests, ≈ 2.5 min):

```
$env:Path = "C:\Users\fredd\miniforge3\envs\pyarts\Library\bin;" + $env:Path
& C:\Users\fredd\miniforge3\envs\pyarts\python.exe -m pytest tests -q
```

`tests/conftest.py` puts the repo root on `sys.path`, so `import rt3d` works from the repo root in scripts and tests alike. Plots must not block: use `matplotlib.use("Agg")` and `fig.savefig("plot/...")`; figure output goes to `plot/`.

10. Module tour

One section per `rt3d` module, with the real signatures.

`rt3d.directions` — direction grids and HG matrices

- `make_icosahedric_grid() → (vertices (12,3), triangles (20,3))` — base icosahedron (same construction as `scripts/icosphere_grid.py`).
- `refine_icosahedric_grid(vertices, triangles, n_refine=1) → (vertices, triangles)` — vectorized geodesic refinement (edge-indexed midpoint table; exact, fast enough for $n_{\text{refine}} = 3$).
- `solid_angle_weights(vertices, triangles) → weights (N,)` — per-vertex solid angles from spherical-triangle areas (L'Huilier); $\sum w = 4\pi$.
- `hg(cos_theta, g) → Phi` — HG phase function, $\oint \Phi \, d\Omega = 1$ analytically.
- `class direction_grid(n_refine=1)` — attributes `n_refine`, `directions (N_dir,3)`, `weights (N_dir,)`, `triangles`, `down_idx` / `up_idx` (indices with $\Omega_z < 0$ / > 0; a partition, since no ordinate has $\Omega_z = 0$), property `n_dir`. Methods: `cos_theta() → (N_dir,N_dir)`; `hg_matrix(g, with_weights=True, normalize=True) → W` with $W_{ij} = \Phi_{\text{HG}}(\Omega_j \to \Omega_i)\, w_j$ and unit row sums when `normalize=True` (§4).
- `icosahedric_grid(n_refine=1) → direction_grid` — the single place the default $n_{\text{refine}} = 1$ (42 ordinates) appears.

`rt3d.geometry` — domain, unigrid, ray marching, ground

- `class cuboid(x0, x1, y0, y1, z0, z1, periodic_x=True, periodic_y=True)` — properties `Lx, Ly, Lz`; `contains(points)`, `wrap_xy(points)`.
- `class unigrid(x, y, z)` — cell-center arrays; attributes `hx, hy, hz`, `nx, ny, nz, n_cells`, `centers (n_cells,3)` (flat-index order, i fastest), `box`. Methods `cell_box(idx)`, `unravel(idx)`, `ravel(i,j,k)`, `locate(points)` (O(1), clamped).
- `ray_box_exit(origin, direction, box) → (t_exit, exit_axis, exit_side)` — slab-method exit distance/face for points inside a box.
- `march_lattice(origin, direction, grid, max_steps=None, periodic_box=None) → (steps, exit_t, exit_axis, exit_side)` — vectorized Amanatides-Woo DDA of K rays sharing one direction through

the unigrid; `steps` is step-major `(ray_idx, cell_idx, length)`; with `periodic_box`, x/y exits wrap and only z exits terminate (§7 incl. wrap-stall fix).

- `class ground_surface(albedo=0.0, emissivity=1.0, B_g=0.0, height=0.0)` — flat Lambertian + emissive ground; `height(x, y)` / `is_variable()` are the variable-surface hook (transport: flat only, §7).

`rt3d.octree` — octree grid and DDA march

- `class octree_grid(cuboid, h0)` — cubic base cells, h_0 must divide all side lengths. Build: `refine(kappa_e_fn, tau_max=0.5, eps_grad=0.3, max_depth=6)`, `balance_21(max_depth=6)`, `finalize()` (freezes leaf arrays `centers, half, level, n_leaf` and allocates `kappa_e, kappa_s, g, T` slots); `sample_fields(kappa_e=…, kappa_s=…, g=…, T=…)` samples callables $(x, y, z) \mapsto$ array at leaf centers; `locate(points)` vectorized descent; `split(nodes)`, `leaves_all()`.
- `march_octree(origin, direction, oct, path_len, periodic_box=None) → (seg_ray, seg_leaf, seg_len)` — vectorized DDA, per-ray directions and per-ray path-length budgets, segments in traversal order per ray.
- `node_major_segments(seg_ray, seg_leaf, seg_len, n_nodes) → (seg_ray, seg_leaf, seg_len, offsets)` — regroup step-major segments into node-major contiguous blocks (stable sort preserves traversal order).

`rt3d.characteristics` — predecessor chains (unigrid + octree reference)

- `stencil_for_direction(omega, h, max_offset=None) → (s, vhat)` — integer lattice stencil per ordinate (§6); $s_z \neq 0$; automatic search radius $\min\big(24, \max(3, \lceil 3\max(h)/\min(h)\rceil)\big)$.
- `subsegments_uniform(s, h) → (rel (M,3), lengths (M,))` — exact sub-segments of the center-to-center path on the lattice.
- `build_chain(grid, omega, periodic_box=None, max_offset=None) → chain` — unigrid chain dict: `s`, `vhat`, `parent`, `roots`, `waves`, `sub_cells (M, n_cells)`, `sub_len`, `root_steps` (step-major backward march segments from root centers to entry), `root_exit_t`, `root_exit_axis`, `root_exit_side`. Used by the solver on the unigrid.
- `build_chain_octree(oct, omega, periodic_box=None, cos_min=0.9, max_hops=8) → chain` — **reference only, not used by the solver** (§5): best-aligned center-to-center connection with cycle peeling; kept for phase-3 validation (`scripts/phase3_octree.py` cell 1).

`rt3d.transport` — sweeps, boundary conditions, beam

- `class boundary_conditions(top_I=0.0, bottom_I=0.0, side_I=0.0, ground=None)` — face backgrounds + optional ground (§7); `inflow(exit_axis, exit_side)`.
- `bottom_mask(grid) → bool array` — cells/leaves touching the bottom face.
- `downward_irradiance(I, dg) → F_down` — $\sum_{\Omega_z<0} w\,|\Omega_z|\,I$ per cell (`I` restricted to `dg.down_idx` order).
- `ground_inflow(F_down, ground, beam_irr=0.0) → I_up` — $\varepsilon_g B_g + A_g(F_\downarrow + \mathrm{beam}_{\mathrm{irr}})/\pi$.

- `beam_irradiance_ground(grid, omega_sun, kappa_e, tau_b, F0, periodic_box=None)` — $F_0|\mu_{\rm sun}|e^{-\tau_b}$ extended through the bottom cell to the ground face.
- `sweep_direction(grid, chain, kappa_e, S, bc, bottom_I=None) → I (n_cells,)` — one unigrid ordinate sweep (recursion of §6).
- `sweep_all(grid, direction_grid, kappa_e, S, bc, periodic_box=None, chains=None) → (I (n_cells,N_dir), chains)` — all ordinates; pass prebuilt `chains` to reuse across iterations.
- `beam_optical_depth(grid, omega_sun, kappa_e, periodic_box=None) → tau_b` — exact backward-beam τ at cell centers (unigrid); raises unless $\Omega_{{\rm sun},z} < 0$.
- `precompute_octree_march(oct, omega, periodic_box=None) → march` — exact backward march per leaf center for one ordinate (§6); dict `leaf`, `length`, `offsets`, `ray_of`, `omega`, `entry_cell` (bottom-face leaf of each upward ray's ground entry point, for the ground BC).
- `sweep_direction_octree_exact(oct, march, kappa_e, S, bc, bottom_I=None) → I` — one octree ordinate sweep along exact characteristics.
- `sweep_all_octree_exact(oct, direction_grid, kappa_e, S, bc, periodic_box=None, marches=None) → (I, marches)` — all ordinates. **This is the octree transport used by the solver.**
- `sweep_direction_octree(oct, chain, kappa_e, S, bc)` and `sweep_all_octree( ... )` — sweeps along `build_chain_octree` chains (reference only).
- `beam_optical_depth_octree(oct, omega_sun, kappa_e, periodic_box=None) → tau_b`.

`rt3d.source` — source function assembly

- `single_scattering_albedo(kappa_e, kappa_s) → omega` — with the $\kappa_e \to 0$ guard (§2).
- `mean_intensity(I, dg) → J` — $J = \frac{1}{4\pi}\oint I\, d\Omega$.
- `flux_vector(I, dg) → F (n_cells,3)` — $F = \oint I\, \mathbf{\Omega}\, d\Omega$.
- `scattering_source(I, dg, omega, B, Phi_w=None, g=0.0, beam_scat=None, g_cells=None, g_decimals=2, Phi_cache=None) → S (n_cells,N_dir)` — $S_{ic} = (1-\omega_c)B_c + \omega_c \sum_j w_j \Phi(\Omega_j \to \Omega_i) I_{cj} + \mathrm{beam_scat}$. Spatially varying g (`g_cells`) is bucketed by `round(g, g_decimals)` with one BLAS matmul per bucket and an optional `{g: Phi_w}` `Phi_cache` (§2/§4).
- `beam_scattering_source(dg, omega_sun, F0, tau_b, omega, g, g_cells=None) → S_beam (n_cells,N_dir)` — fixed beam first-scatter term with normalized HG row (§3).

`rt3d.accelerator` — Ng acceleration

- `class ng1d(grid, max_hist=None, n_start=3)` — 1D Ng state (§8); methods `plane_average(v)`, `lambda_bar()` ($N_z \times N_z$ least-squares fit), `step(S_swept, S_new) → next source`.

`rt3d.solver` — the iteration loop

- `lambda_iteration(grid, dg, kappa_e, kappa_s, B, bc, g=0.0, periodic_box=None, beam=None, n_iter=100, tol=1e-8, ng=None) → result` — the entry point. `grid` is a `unigrid` or `octree_grid`; g scalar or `(n_cells,)`; `beam = {"omega_sun": (3,) unit, "F0": float}` or

`None`; `ng` is `None`/`False` (plain LI), `True` (build `ng1d(grid)`) or a prepared `ng1d`. Returns the result dict:

key	shape / type	meaning
`I`	`(n_cells, N_dir)`	converged center intensities (diffuse field)
`S`	`(n_cells, N_dir)`	converged source function
`J`	`(n_cells,)`	mean intensity
`F`	`(n_cells, 3)`	flux vector
`omega`	`(n_cells,)`	single scattering albedo
`history`	list of float	relative source change per iteration
`n_iter`	int	iterations performed
`converged`	bool	`history[-1] < tol`

Note there is **no** `SolverConfig` class (that name exists only in `plan/plan.md`); all configuration is via the keyword arguments above.

`rt3d.diagnostics` — diagnostics

- `heating_rate(kappa_a, J, B)` — net heating rate per volume, $4\pi\kappa_a(B - J)$ (positive = radiation gains energy from matter).
- `flux_divergence(grid, F) → per-cell div F` — central differences, periodic wrap in x/y, one-sided at top/bottom (uniform unigrid only).
- `energy_balance(grid, I, dg, kappa_a, B) → dict` — global closure on the unigrid: `out_top`, `out_bottom` (outward power through the faces, net F_z second-order extrapolated from the boundary layers), `heating` ($\int 4\pi\kappa_a(B - J)\,dV$), `residual = out_top + out_bottom − heating`. With a beam, subtract the first-scatter input $\int \kappa_s F_0 e^{-\tau_b}\,dV$ from the residual to close the diffuse-field balance (§11, §14 anchor 4).
- `allsky(grid, I, dg, face="top") → dict` — outgoing intensity images at a horizontal face for every leaving ordinate: `omega (n_out,3)`, `weights (n_out,)`, `images (n_out, ny, nx)`; `face="top"` uses upward ordinates at the top layer, `face="bottom"` downward ordinates at the bottom layer (uniform unigrid).

`rt3d.cases` — test cases


- `gauss(x, y, z, sigma=1.0, mu=(0,0,4))` — normalized 3D Gaussian (integral 1), as in `scripts/class_test.py`.
- `gaussian_cloud(sigma=1.0, mu=(0,0,4), tau_column_max=10.0, omega_cap=None) → case dict` — the `class_test.py` case as callables: `box` ($x, y \in [-5, 5]$, $z \in [0, 10]$) and `kappa_e`, `kappa_s`, `g`,

`T` fields $(x, y, z) \mapsto$ array. Peak column τ 10; local ω peaks at 1.0 (uncapped, faithful to `class_test.py`) or `omega_cap`; local g peaks ≈ 0.952. `class_test.py` stores `albedo`; this case converts to $\kappa_s = \omega_{\text{local}} \cdot \kappa_e$.

- Note: `slab_1d` / `clear_sky` from `plan/plan.md` **do not exist as case modules**; the plane-parallel slab lives inside `scripts/phase2_slab.py`.

11. Worked end-to-end example

Gaussian cloud + Lambertian ground + solar beam, on both grids, Ng accelerated. The script below was **executed with the pyarts interpreter (PATH fix applied) and its output verified**; the printed numbers are the actual measured values. Figures are written to `plot/`.



```
# gaussian_cloud_example.py — run from the repo root:
#   $env:Path = "C:\Users\fredd\miniforge3\envs\pyarts\Library\bin;" + $env:Path
#   & C:\Users\fredd\miniforge3\envs\pyarts\python.exe gaussian_cloud_example.py
import time

import numpy as np

import matplotlib
matplotlib.use("Agg")
import matplotlib.pyplot as plt

from rt3d.cases.gaussian_cloud import gaussian_cloud
from rt3d.directions import icosahedric_grid
from rt3d.geometry import unigrid, ground_surface
from rt3d.octree import octree_grid
from rt3d.transport import boundary_conditions, beam_optical_depth
from rt3d.solver import lambda_iteration
from rt3d.diagnostics import (allsky, energy_balance, flux_divergence,
                              heating_rate)

# ---------------------------------------------------------------- case setup
case = gaussian_cloud(omega_cap=0.9)          # capped core: Ng makes it tractable
box = case["box"]                              # x,y in [-5,5], z in [0,10]

dg = icosahedric_grid(n_refine=1)              # 42 ordinates (development grid)
print(f"N_dir = {dg.n_dir},  sum(w) - 4pi = {dg.weights.sum() - 4*np.pi:.3e}")

# solar beam: 30 deg from zenith, propagating in the x-z plane
mu_sun = np.cos(np.radians(30.0))
beam = {"omega_sun": np.array([np.sqrt(1.0 - mu_sun**2), 0.0, -mu_sun]),
        "F0": 1.0}

bc = boundary_conditions(top_I=1.0,
                         ground=ground_surface(albedo=0.3, emissivity=0.0, B_g=0.0))

# ---------------------------------------------------------------- unigrid run
n = 24
xs = np.linspace(box.x0, box.x1, n + 1)[:-1] + (box.Lx / n) / 2
grid = unigrid(xs, xs, np.linspace(box.z0, box.z1, n + 1)[:-1] + (box.Lz / n) / 2)
c = grid.centers
kappa_e = case["kappa_e"](*c.T)
kappa_s = case["kappa_s"](*c.T)
g_cells = case["g"](*c.T)                      # per-cell g -> bucketed matmul
```

```
B = np.zeros(grid.n_cells)

t0 = time.perf_counter()
res = lambda_iteration(grid, dg, kappa_e, kappa_s, B, bc,
                       g=g_cells, periodic_box=box, beam=beam,
                       n_iter=60, tol=1e-6, ng=True)
t_uni = time.perf_counter() - t0
print(f"unigrid:  n_iter={res['n_iter']}, converged={res['converged']}, "
      f"wall={t_uni:.2f} s")

# ──────────────────────────────────────────────────────────────── octree run
oct_ = octree_grid(box, h0=1.25)
oct_.refine(case["kappa_e"], tau_max=2.0, eps_grad=1.0, max_depth=3)
oct_.balance_21(max_depth=3)
oct_.finalize()
oct_.sample_fields(kappa_e=case["kappa_e"], kappa_s=case["kappa_s"],
                   g=case["g"], T=case["T"])
print(f"octree:   n_leaf={oct_.n_leaf}, max_level={oct_.max_level}")

t0 = time.perf_counter()
res_oct = lambda_iteration(oct_, dg, oct_.kappa_e, oct_.kappa_s,
                           np.zeros(oct_.n_leaf), bc,
                           g=oct_.g, periodic_box=box, beam=beam,
                           n_iter=60, tol=1e-6, ng=True)
t_oct = time.perf_counter() - t0
print(f"octree:   n_iter={res_oct['n_iter']}, converged={res_oct['converged']}, "
      f"wall={t_oct:.2f} s")

# ──────────────────────────────────────────────────────────── interpretation
J4 = res["J"].reshape(grid.nz, grid.ny, grid.nx)     # k-major → (k, j, i)

# mean-J profile on the 8 coarse base-cell bands (both grids)
nband = int(round(box.Lz / oct_.h0))
zband = np.linspace(box.z0, box.z1, nband + 1)
zb_uni = np.array([J4[(grid.z ≥ zband[b]) & (grid.z < zband[b + 1])].mean()
                   for b in range(nband)])
zi = np.clip(((oct_.centers[:, 2] - box.z0) / oct_.h0).astype(int), 0, nband - 1)
zb_oct = np.array([res_oct["J"][zi == b].mean() for b in range(nband)])

zc = 0.5 * (zband[:-1] + zband[1:])
print("z      J_unigrid  J_octree  rel.diff")
for b in (0, 2, 4, 5, 7):
    d = abs(zb_oct[b] - zb_uni[b]) / max(zb_uni[b], 1e-12)
    print(f"{zc[b]:5.2f}  {zb_uni[b]:9.4f}  {zb_oct[b]:8.4f}  {d*100:7.1f}%")
```

```
top_layer = oct_.centers[:, 2] + oct_.half ≥ box.z1 - 1e-9
print(f"top-of-domain mean F_z: unigrid "
      f"{res['F'][-grid.ny*grid.nx:, 2].mean():.4f}, "
      f"octree {res_oct['F'][top_layer, 2].mean():.4f}")

# diffuse-field energy closure (phase-5 recipe, §14 anchor 4)
kappa_a = kappa_e - kappa_s
bal = energy_balance(grid, res["I"], dg, kappa_a, B)
tau_b = beam_optical_depth(grid, beam["omega_sun"], kappa_e, periodic_box=box)
V = grid.hx * grid.hy * grid.hz
beam_scat = (kappa_s * beam["F0"] * np.exp(-tau_b)).sum() * V
res_dif = bal["residual"] - beam_scat
in_diff = np.sum(dg.weights[dg.down_idx]
                 * np.abs(dg.directions[dg.down_idx, 2])) * box.Lx * box.Ly
print(f"  diffuse closure residual {res_dif:9.2f}"
      f"  ({abs(res_dif) / (in_diff + beam_scat) * 100:.2f}% of diffuse input)")

# pointwise anchor 4: div F vs heating rate in the interior
divF = flux_divergence(grid, res["F"])
heat = heating_rate(kappa_a, res["J"], B)
kk = np.arange(grid.n_cells) // (grid.ny * grid.nx)
interior = (kk > 0) & (kk < grid.nz - 1)
err_rms = np.sqrt(np.mean((divF[interior] - heat[interior]) ** 2)) / np.abs(heat).max()
err_max = np.abs(divF[interior] - heat[interior]).max() / np.abs(heat).max()
print(f"pointwise div F vs heating (interior): rms {err_rms*100:.1f}%, "
      f"max {err_max*100:.1f}% of max|heating|")

# ──────────────────────────────────────────────────────────────── plots
fig, axes = plt.subplots(1, 2, figsize=(10, 4))
im = axes[0].imshow(J4[:, grid.ny // 2, :], origin="lower", aspect="auto",
                    extent=[box.x0, box.x1, box.z0, box.z1])
axes[0].set_xlabel("x"); axes[0].set_ylabel("z")
axes[0].set_title("J, y=0 slice (unigrid)")
plt.colorbar(im, ax=axes[0])
axes[1].plot(zb_uni, zc, label="unigrid 24$^3$")
axes[1].plot(zb_oct, zc, "--", label="octree")
axes[1].set_xlabel("band-mean J"); axes[1].set_ylabel("z")
axes[1].legend()
fig.savefig("plot/manual_example_j.png", dpi=120)

sky = allsky(grid, res["I"], dg, face="top")
n_out = sky["omega"].shape[0]
fig, axes = plt.subplots(6, 7, figsize=(12, 10))
mx = max(sky["images"].max(), 1e-12)
for a, img in zip(axes.ravel(), sky["images"]):
```


```
    a.imshow(img, origin="lower", vmin=0, vmax=mx, cmap="inferno")
    a.set_xticks([]); a.set_yticks([])
for a in axes.ravel()[n_out:]:
    a.axis("off")
fig.suptitle("all-sky outgoing intensity at top face (42 ordinates)")
fig.savefig("plot/manual_example_allsky.png", dpi=120)
```

Verified output (this machine; wall times vary a few %):

```
N_dir = 42,  sum(w) - 4pi = -1.776e-15
unigrid:  n_iter=32, converged=True, wall=4.23 s
octree:   n_leaf=25264, max_level=3
octree:   n_iter=30, converged=True, wall=107.00 s
z      J_unigrid  J_octree  rel.diff
 0.62     0.4372    0.3095     29.2%
 3.12     0.4529    0.3431     24.2%
 5.62     0.5589    0.4788     14.3%
 6.88     0.5870    0.5377      8.4%
 9.38     0.5893    0.5850      0.7%
top-of-domain mean F_z: unigrid -2.5215, octree -2.5312
  diffuse closure residual      9.49  (2.96% of diffuse input)
pointwise div F vs heating (interior): rms 1.6%, max 13.9% of max|heating|
```


Figures produced

Mean intensity J: left $-$ $y = 0$ slice through the cloud on the unigrid; right $-$ band-mean J profiles, unigrid 24^3 vs octree (the difference is the unigrid grazing-ordinate stencil error, anchor 5):



All-sky outgoing intensity at the top face — one image per upward ordinate (21 upward ordinates of the 42-ordinate development grid):

all-sky outgoing intensity at top face (42 ordinates)



Interpreting the result dict and the diagnostics:

- `n_iter` / `history`: Ng converges the near-conservative cloud ($\omega \leq 0.9$, $g \approx 0.95$ core) in ~30 iterations; plain LI would need several hundred (§8). `history[k]` is the relative source change at iteration k.
- **unigrid vs octree J**: the ~10–30% band-mean differences are the *unigrid* lattice-stencil path error for grazing ordinates in the strongly heterogeneous cloud (validation anchor 5, §14) — the octree exact-characteristic result is the better one; the top-of-domain flux, dominated by near-vertical paths, agrees to 0.4%.
- **energy closure**: `energy_balance` alone leaves the beam first-scatter input unaccounted; after subtracting $\int \kappa_s F_0 e^{-\tau_b}\, dV$ the global diffuse balance closes to **2.96%** of the incident diffuse power — the discretization error of the 42-ordinate / 24^3 solve, not a conservation bug.
- **pointwise div F vs heating**: rms 1.6% of max|heating| — anchor 4.
- Figures: `plot/manual_example_j.png` (J slice + profile), `plot/manual_example_allsky.png` (21 upward-ordinate all-sky images at the top face).

12. Runner scripts

All scripts in `scripts/` use `# %%` cell markers (VS Code / Spyder workflow), insert the repo root on `sys.path`, use `matplotlib.use("Agg")`, and write figures to `plot/`. Run any of them as

```
$env:Path = "C:\Users\fredd\miniforge3\envs\pyarts\Library\bin;" + $env:Path
& C:\Users\fredd\miniforge3\envs\pyarts\python.exe scripts/phase0_check.py
```

script	what it validates
`phase0_check.py`	direction grid rebuilds for n_{refine} 0-3 (12/42/162/642), $\sum w = 4\pi$, $\oint \Phi \, d\Omega = 1$, no $\Omega_z = 0$ ordinate, ray-box exit + DDA march consistency (anchors 1-2).
`phase1_beer.py`	uniform-grid short characteristics: beam $I = F_0 e^{-\tau_b}$, diffuse transmittance vs Beer-Lambert for every ordinate/cell (anchor 3), isothermal $I = B$ exactness.
`phase2_slab.py`	scattering + plain λ-iteration on a 1D slab ($\tau = 5$, $\omega = 0.5/0.9$) vs an independent 64-level Gauss-Legendre discrete-ordinates solver and the two-stream approximation. The slab case lives here (no `slab_1d` module).
`phase3_octree.py`	octree vs unigrid on the Gaussian cloud (anchor 5): chain-vs-exact-path sanity (`build_chain_octree` reference), octree self-convergence, column $\tau \to 10$, scattering run at $\omega \leq 0.7$.
`phase4_ng.py`	1D Ng acceleration: LI vs Ng iteration counts as $\omega \to 1$ (slab), cloud on unigrid ($\omega \leq 0.9$, per-cell g) and on the octree.
`phase5_ground.py`	periodic sides + Lambertian ground + all-sky output; analytic ground Beer-Lambert checks; pointwise div F vs heating and global energy balance incl. beam/ground terms (anchor 4).
`phase6_directions.py`	direction-grid convergence 42 → 162 → 642 on the cloud, $g = 0.952$ slab (known 42-ordinate limitation), performance pass (chain build + sweep time vs N_{dir}), npz record with `n_refine`.

`scripts/class_test.py` and `scripts/icosphere_grid.py` are **read-only legacy reference** (pre-existing research code); do not modify them.

13. Condensed API reference

Signatures + one-liners for all public functions/classes of `rt3d` (docstrings in the modules are authoritative for defaults).

`rt3d.directions`

signature	description
`make_icosahedric_grid()`	base icosahedron: 12 unit vertices, 20 triangles
`refine_icosahedric_grid(vertices, triangles, n_refine=1)`	vectorized geodesic refinement
`solid_angle_weights(vertices, triangles)`	per-vertex solid angles from spherical-triangle areas
`hg(cos_theta, g)`	HG phase function, $\oint \Phi \, d\Omega = 1$
`direction_grid(n_refine=1)`	icosahedral direction set with weights, up/down index split
`.cos_theta()`	(N, N) matrix of direction cosines
`.hg_matrix(g, with_weights=True, normalize=True)`	weighted, row-normalized HG matrix
`icosahedric_grid(n_refine=1)`	build a `direction_grid` (default 42 ordinates)

`rt3d.geometry`

signature	description
`cuboid(x0, x1, y0, y1, z0, z1, periodic_x=True, periodic_y=True)`	domain box with periodic x/y
`.contains(points)` / `.wrap_xy(points)`	inside test / periodic wrap
`unigrid(x, y, z)`	uniform Cartesian cell lattice from cell centers
`.cell_box(idx)` / `.unravel(idx)` / `.ravel(i,j,k)` / `.locate(points)`	cell boxes, index maps, O(1) point location
`ray_box_exit(origin, direction, box)`	exit distance + face of a box along a direction
`march_lattice(origin, direction, grid, max_steps=None, periodic_box=None)`	vectorized DDA march through the unigrid
`ground_surface(albedo=0.0, emissivity=1.0, B_g=0.0, height=0.0)`	flat Lambertian + emissive ground (variable surface = hook only)

`rt3d.octree`

signature	description
`octree_grid(cuboid, h0)`	flat struct-of-arrays octree, cubic base cells
`.refine(kappa_e_fn, tau_max=0.5, eps_grad=0.3, max_depth=6)`	top-down refinement on $\kappa_e \tau$ + gradient criteria
`.balance_21(max_depth=6)`	2:1 face balance via 3×3 face probing
`.finalize()` / `.sample_fields( ... )`	freeze leaf arrays / sample callable fields at leaf centers
`.locate(points)` / `.split(nodes)` / `.leaves_all()`	vectorized descent point location / node ops
`march_octree(origin, direction, oct, path_len, periodic_box=None)`	vectorized DDA march through the octree
`node_major_segments(seg_ray, seg_leaf, seg_len, n_nodes)`	regroup segments into node-major blocks

`rt3d.characteristics`

signature	description
`stencil_for_direction(omega, h, max_offset=None)`	best integer lattice stencil for one ordinate
`subsegments_uniform(s, h)`	exact sub-segments of the stencil center-to-center path
`build_chain(grid, omega, periodic_box=None, max_offset=None)`	unigrid predecessor chain (stencil + waves + root marches)
`build_chain_octree(oct, omega, periodic_box=None, cos_min=0.9, max_hops=8)`	octree center-to-center chain — **reference only, not used by the solver**

`rt3d.transport`

signature	description
`boundary_conditions(top_I=0.0, bottom_I=0.0, side_I=0.0, ground=None)`	face backgrounds + optional ground
`.inflow(exit_axis, exit_side)`	background inflow for a chain entry face
`bottom_mask(grid)`	cells/leaves touching the bottom face
`downward_irradiance(I, dg)`	$\sum_{\Omega_z<0} w\lvert\Omega_z\rvert I$ at ground level
`ground_inflow(F_down, ground, beam_irr=0.0)`	Lambertian ground radiance
`beam_irradiance_ground(grid, omega_sun, kappa_e, tau_b, F0, periodic_box=None)`	direct-beam irradiance at the ground face
`sweep_direction(grid, chain, kappa_e, S, bc, bottom_I=None)`	one unigrid ordinate sweep
`sweep_all(grid, direction_grid, kappa_e, S, bc, periodic_box=None, chains=None)`	all-ordinate unigrid transport
`precompute_octree_march(oct, omega, periodic_box=None)`	exact backward march per leaf center, one ordinate
`sweep_direction_octree_exact(oct, march, kappa_e, S, bc, bottom_I=None)`	one octree ordinate sweep, exact characteristics
`sweep_all_octree_exact(oct, direction_grid, kappa_e, S, bc, periodic_box=None, marches=None)`	all-ordinate octree transport (solver path)
`sweep_direction_octree(oct, chain, kappa_e, S, bc)` / `sweep_all_octree( ... )`	chain-based octree sweeps (reference only)
`beam_optical_depth(grid, omega_sun, kappa_e, periodic_box=None)`	beam τ at unigrid cell centers
`beam_optical_depth_octree(oct, omega_sun, kappa_e, periodic_box=None)`	beam τ at octree leaf centers

`rt3d.source`

signature	description
`single_scattering_albedo(kappa_e, kappa_s)`	$\omega = \kappa_s / \kappa_e$ with vacuum guard
`mean_intensity(I, dg)`	$J = \frac{1}{4\pi} \oint I \, d\Omega$
`flux_vector(I, dg)`	$F = \oint I \, \mathbf{\Omega} \, d\Omega$
`scattering_source(I, dg, omega, B, Phi_w=None, g=0.0, beam_scat=None, g_cells=None, g_decimals=2, Phi_cache=None)`	direction-dependent source $S = (1-\omega)B + \omega\langle \Phi I \rangle + \text{beam_scat}$
`beam_scattering_source(dg, omega_sun, F0, tau_b, omega, g, g_cells=None)`	fixed beam first-scatter source, normalized HG row

`rt3d.accelerator`

signature	description
`ng1d(grid, max_hist=None, n_start=3)`	1D plane-averaged Ng acceleration state
`.plane_average(v)` / `.lambda_bar()` / `.step(S_swept, S_new)`	z-bin averages / $\bar{\Lambda}$ fit / next source

`rt3d.solver`

signature	description
`lambda_iteration(grid, dg, kappa_e, kappa_s, B, bc, g=0.0, periodic_box=None, beam=None, n_iter=100, tol=1e-8, ng=None)`	the solver; returns `I, S, J, F, omega, history, n_iter, converged`

`rt3d.diagnostics`

signature	description
`heating_rate(kappa_a, J, B)`	$4\pi\kappa_a(B - J)$ per cell
`flux_divergence(grid, F)`	div F by central differences (unigrid)
`energy_balance(grid, I, dg, kappa_a, B)`	global face-flux vs heating closure
`allsky(grid, I, dg, face="top")`	outgoing intensity images per leaving ordinate at top/bottom face

`rt3d.cases`

signature	description
`gauss(x, y, z, sigma=1.0, mu=(0,0,4))`	normalized 3D Gaussian
`gaussian_cloud(sigma=1.0, mu=(0,0,4), tau_column_max=10.0, omega_cap=None)`	class_test.py cloud as box + callable fields

14. Validation anchors

The five physics anchors from AGENTS.md, with measured numbers (all re-verified for this manual unless marked "phase record"):

1. $\sum_i w_i = 4\pi$ **over the direction grid.** $|\sum_i w_i - 4\pi|$ = 3.6e-15 / 1.8e-15 / 7.1e-15 / 1.4e-14 for 12 / 42 / 162 / 642 ordinates — machine precision.

2. $\oint \Phi \, d\Omega = 1$ **for the HG phase function.** Analytically exact by construction. Discretely at 42 ordinates: raw weighted row sums are exactly 1 at $g = 0$, 1.000-1.002 at $g = 0.5$, and **16.9-21.6 at** $g = 0.952$ — the reason for row normalization (§4); after normalization the row sums are 1 to ≤ 3.3e-16.

3. $\kappa_s = 0$, $B = 0 \rightarrow$ **transmittance** $= e^{-\tau}$ **along every ray.** Octree exact march (uniform $\kappa_e = 0.7$, periodic box, all 42 ordinates): $\max|I - e^{-\tau}| = 9.7 \times 10^{-16}$, max τ error 7.1×10^{-14} — machine precision. (For contrast, the unigrid lattice stencil on the same test has path errors up to $\tau \approx 34$ for grazing ordinates — the documented stencil error, not a bug; see anchor 5.)

4. **div** F **equals the heating rate from the intensity field.** Cloud + ground + beam run (phase 5 setup; reproduced in §11): pointwise div F vs $4\pi\kappa_a(B - J)$ in the interior — **rms 1.6%, max 13.9%** of $\max|heating|$. Global closure (face fluxes

   - gas absorption − beam first-scatter input): **2.96%** of the incident diffuse power (phase 5 record: 3.2%). Both are discretization errors of the 42-ordinate/24^3 solve, not conservation failures.

5. **Octree converges to the uniform-grid result on the Gaussian cloud.** With identical piecewise-constant fields, the J difference between octree (exact characteristics) and unigrid (lattice stencil) is ≈ **18%** (phase 3 record; §11 band means show 0.7–29% depending on height). Interpretation: this is the **unigrid grazing-ordinate stencil error** — the best-aligned center-to-center characteristic of a 42-ordinate grazing ray deviates strongly from the true ray on a 24^3 lattice — *not* an octree error. The octree self-convergence series (phase 3, cell 2b) and the column-$\tau \to 10$ check confirm the octree side; the coarse uniform reference under-resolves the cloud (column τ 9.58).

15. Known limitations and performance

Angular resolution. 42 ordinates is the development grid but not a converged solution at large g: the HG lobe at $g \approx 0.95$ is under-resolved (≈ 6% J error in the cloud case, phase 6 record), and even **642 ordinates are not fully converged at** $g = 0.952$; angular convergence is **non-monotone at large** g (error cancellation between ordinates — do not extrapolate monotonically). The row normalization (§4) deliberately trades phase-function fidelity for discrete energy conservation.

Variable surface. `ground_surface(height=f(x,y))` is a hook only; the transport supports a flat ground at the domain bottom and `lambda_iteration` raises `NotImplementedError` otherwise (plan phase 6 hook).

Octree chain transport. `build_chain_octree` / `sweep_all_octree` are kept as phase-3 reference only: their center-to-center connection has a scale-invariant path error ($\tau \approx 4$ on the cloud) that refinement does not remove. The solver always uses exact characteristics (`precompute_octree_march`).

Performance (this machine, phase 6 record + re-measured):

- Transport sweep cost ≈ **2.1–2.3 ms per ordinate per** 24^3 **sweep** (re-measured for this manual: 2.6 ms/ordinate including loop overhead), **linear in** N_{dir}; chain/march precomputation is built once and reused.
- Full Ng cloud run (24^3, ground + beam) at **642 ordinates ≈ 84 s** (phase 6 record); the §11 example: unigrid 42 ordinates ≈ 4 s, octree (25 264 leaves) ≈ 100 s.
- The octree DDA march cost grows with the *finest* cell size (the step budget scales as $\mathrm{path}/h_{\min}$), so very deep octrees (max_depth ≥ 4 with small h_0) get disproportionately expensive — the documented cloud runs use $h_0 = 1.25$, max_depth = 3.
- **numba is not needed**: all heavy loops are vectorized NumPy (BFS waves, block cumsums, single BLAS matmul for the scattering integral); the test suite and all phase scripts run in pure NumPy.

Other caveats.

- `flux_divergence` and `energy_balance` are unigrid-only diagnostics (central differences on the uniform lattice).
- `energy_balance`'s `residual` does not include the beam first-scatter input; subtract $\int \kappa_s F_0 e^{-\tau_b}\, dV$ for the diffuse closure (§11).
- A single-cell unigrid dimension has nominal width 1.0 (slab convention); keep $n_x = n_y = 1$ only with periodic sides.
- The convergence history is the relative change of the *swept* source; with Ng active the swept source is the extrapolated one, so `history` measures the plain-LI residual of the accelerated iterate.