

Still water and other hard-to-model systems

Martin Büttner
Supervisor: Prof. Ted Johnson

16 March, 2015

The Shallow Water Equations

- ▶ Simplified fluid model:
 - ▶ Incompressible
 - ▶ Inviscid
 - ▶ *Shallow* (horizontal length scale \gg typical depth)
- ▶ Used to model oceans and atmosphere

The Shallow Water Equations

- ▶ Simplified fluid model:
 - ▶ Incompressible
 - ▶ Inviscid
 - ▶ *Shallow* (horizontal length scale \gg typical depth)
- ▶ Used to model oceans ($\sim 4\text{km}$ deep)
and atmosphere ($\sim 10\text{km}$ deep)

The Shallow Water Equations

- ▶ Simplified fluid model:
 - ▶ Incompressible
 - ▶ Inviscid
 - ▶ *Shallow* (horizontal length scale \gg typical depth)
- ▶ Used to model oceans ($\sim 4\text{km}$ deep)
and atmosphere ($\sim 10\text{km}$ deep)

$$h_t + (hu)_x = 0 \quad (1)$$

$$(hu)_t + \left(hu^2 + \frac{1}{2}gh^2 \right)_x = 0 \quad (2)$$


$$(hv)_t + (huv)_x = 0 \quad (3)$$

Finite Volume Methods

- Developed for *hyperbolic conservation laws*:

$$\mathbf{q}_t + \mathbf{f}(\mathbf{q})_x = 0 \quad (4)$$

- Discretisation:




Finite Volume Methods

- Developed for *hyperbolic conservation laws*:

$$\mathbf{q}_t + \mathbf{f}(\mathbf{q})_x = 0 \quad (4)$$

- Discretisation:




Finite Volume Methods

- Developed for *hyperbolic conservation laws*:

$$\mathbf{q}_t + \mathbf{f}(\mathbf{q})_x = 0 \quad (4)$$

- Discretisation:




Finite Volume Methods


- Developed for *hyperbolic conservation laws*:

$$\mathbf{q}_t + \mathbf{f}(\mathbf{q})_x = 0 \quad (4)$$

- Discretisation:




Riemann Problems




- Initial value problem: step function

Riemann Problems



- ▶ Initial value problem: step function
- ▶ Decompose into waves

Riemann Problems



- ▶ Initial value problem: step function
- ▶ Decompose into waves
- ▶ Finite speeds \Rightarrow only affect neighbouring cells

Godunov's Method

- ▶ Wave propagation form:

$$Q_i^{n+1} = Q_i^n - \frac{\Delta t}{\Delta x} (\mathcal{A}^+ \Delta Q_{i-1/2} + \mathcal{A}^- \Delta Q_{i+1/2}) \quad (5)$$

Godunov's Method

- Wave propagation form:

$$Q_i^{n+1} = Q_i^n - \frac{\Delta t}{\Delta x} (\mathcal{A}^+ \Delta Q_{i-1/2} + \mathcal{A}^- \Delta Q_{i+1/2}) \quad (5)$$

$$\mathcal{A}^- \Delta Q_{i-1/2} = \sum_{p=1}^{M_w} (s_{i-1/2}^p)^- \mathcal{W}_{i-1/2}^p \quad (6)$$

$$\mathcal{A}^+ \Delta Q_{i+1/2} = \sum_{p=1}^{M_w} (s_{i+1/2}^p)^+ \mathcal{W}_{i+1/2}^p \quad (7)$$

Implementation: Clawpack

- ▶ “**C**onservation **law package**”, by Randall J. LeVeque, 1994

Implementation: Clawpack

- ▶ “**C**onservation **law package**”, by Randall J. LeVeque, 1994
- ▶ Riemann solvers in Fortran 90
 - ▶ Take grid data, compute wave decomposition

Implementation: Clawpack

- ▶ “**C**onservation **law package**”, by Randall J. LeVeque, 1994
- ▶ Riemann solvers in Fortran 90
 - ▶ Take grid data, compute wave decomposition
- ▶ Configuration in Python 2
 - ▶ Grid setup
 - ▶ Initial conditions
 - ▶ Boundary conditions
 - ▶ Plotting instructions

Adding Source Terms

- Modelling additional effects:

$$h_t + (hu)_x = 0 \quad (8)$$

$$(hu)_t + \left(hu^2 + \frac{1}{2}gh^2 \right)_x = 0 \quad (9)$$

$$(hv)_t + (huv)_x = 0 \quad (10)$$

Adding Source Terms

- Modelling additional effects:

$$h_t + (hu)_x = 0 \quad (8)$$

$$(hu)_t + \left(hu^2 + \frac{1}{2}gh^2 \right)_x = -hB_x \quad (9)$$

$$(hv)_t + (huv)_x = 0 \quad (10)$$

Adding Source Terms

- Modelling additional effects:

$$h_t + (hu)_x = 0 \quad (8)$$

$$(hu)_t + \left(hu^2 + \frac{1}{2}gh^2 \right)_x = -hB_x + Khv \quad (9)$$

$$(hv)_t + (huv)_x = -Khu \quad (10)$$

Adding Source Terms

- Modelling additional effects:

$$h_t + (hu)_x = 0 \quad (8)$$

$$(hu)_t + \left(hu^2 + \frac{1}{2}gh^2 \right)_x = -hB_x + Khv \quad (9)$$

$$(hv)_t + (huv)_x = -Khu \quad (10)$$

- Full form of conservation law:

$$\mathbf{q}_t + \mathbf{f}(\mathbf{q})_x = \mathbf{s}(\mathbf{q}, x) \quad (11)$$

Adding Source Terms

- Split into two steps:

$$\mathbf{q}_t + \mathbf{f}(\mathbf{q})_x = 0 \quad (12)$$

$$\mathbf{q}_t = \mathbf{s}(\mathbf{q}, x) \quad (13)$$


Adding Source Terms

- Split into two steps:

$$\mathbf{q}_t + \mathbf{f}(\mathbf{q})_x = 0 \quad (12)$$

$$\mathbf{q}_t = \mathbf{s}(\mathbf{q}, x) \quad (13)$$

- The still water system:




Adding Source Terms

- Split into two steps:

$$\mathbf{q}_t + \mathbf{f}(\mathbf{q})_x = 0 \quad (12)$$

$$\mathbf{q}_t = \mathbf{s}(\mathbf{q}, x) \quad (13)$$


- The still water system:



Geostrophic Equilibria

- ▶ Coriolis force admits non-trivial equilibria when

$$u = 0, \quad v = \frac{h(h_s)x}{K} \quad (14)$$



- ▶ Geophysical flows close to geostrophic equilibrium at all times

Well-Balanced Methods


LeVeque

- ▶ LeVeque, 1998, “Balancing Source Terms and Flux Gradients in High-Resolution Godunov Methods: The Quasi-Steady Wave-Propagation Algorithm”

Well-Balanced Methods

LeVeque


- ▶ LeVeque, 1998, “Balancing Source Terms and Flux Gradients in High-Resolution Godunov Methods: The Quasi-Steady Wave-Propagation Algorithm”



Well-Balanced Methods

LeVeque


- ▶ LeVeque, 1998, “Balancing Source Terms and Flux Gradients in High-Resolution Godunov Methods: The Quasi-Steady Wave-Propagation Algorithm”



Well-Balanced Methods

LeVeque

- ▶ LeVeque, 1998, “Balancing Source Terms and Flux Gradients in High-Resolution Godunov Methods: The Quasi-Steady Wave-Propagation Algorithm”



$$\mathbf{f}(Q_i^+) - \mathbf{f}(Q_i^-) = \mathbf{s}(Q_i, x_i) \Delta x \quad (15)$$

Well-Balanced Methods

LeVeque

- ▶ Method originally developed for bathymetry term

Well-Balanced Methods

LeVeque

- ▶ Method originally developed for bathymetry term
- ▶ Derived analogous method including Coriolis terms

Well-Balanced Methods

LeVeque

- ▶ Method originally developed for bathymetry term
- ▶ Derived analogous method including Coriolis terms
- ▶ Pro:
 - ▶ Well-balanced for all equilibria

Well-Balanced Methods

LeVeque

- ▶ Method originally developed for bathymetry term
- ▶ Derived analogous method including Coriolis terms
- ▶ Pro:
 - ▶ Well-balanced for all equilibria
- ▶ Con:
 - ▶ Requires solving cubic for offset
 - ▶ Problematic away from equilibrium (transcritical flow)

Well-Balanced Methods

Rogers et al.

- ▶ Rogers, Borthwick, Taylor, 2003, “Mathematical balancing of flux gradient and source terms prior to using Roe’s approximate Riemann solver”

Well-Balanced Methods

Rogers et al.

- ▶ Rogers, Borthwick, Taylor, 2003, “Mathematical balancing of flux gradient and source terms prior to using Roe’s approximate Riemann solver”
- ▶ Change of variables: deviations from equilibrium

$$\mathbf{q}_t + \mathbf{f}(\mathbf{q})_x = \mathbf{s}(\mathbf{q}, x) \quad (16)$$

$$\rightarrow (\mathbf{q} - \mathbf{q}_{\text{eq}})_t + (\mathbf{f}(\mathbf{q}) - \mathbf{f}(\mathbf{q}_{\text{eq}}))_x = \mathbf{s}(\mathbf{q}, x) - \mathbf{s}(\mathbf{q}_{\text{eq}}, x) \quad (17)$$

$$\text{or } \mathbf{q}_t^* + \mathbf{f}_x^* = \mathbf{s}^* \quad (18)$$

Well-Balanced Methods

Rogers et al.

- ▶ Rogers, Borthwick, Taylor, 2003, “Mathematical balancing of flux gradient and source terms prior to using Roe’s approximate Riemann solver”
- ▶ Change of variables: deviations from equilibrium

$$\mathbf{q}_t + \mathbf{f}(\mathbf{q})_x = \mathbf{s}(\mathbf{q}, x) \quad (16)$$

$$\rightarrow (\mathbf{q} - \mathbf{q}_{\text{eq}})_t + (\mathbf{f}(\mathbf{q}) - \mathbf{f}(\mathbf{q}_{\text{eq}}))_x = \mathbf{s}(\mathbf{q}, x) - \mathbf{s}(\mathbf{q}_{\text{eq}}, x) \quad (17)$$

$$\text{or } \mathbf{q}_t^* + \mathbf{f}_x^* = \mathbf{s}^* \quad (18)$$

- ▶ At equilibrium, all terms vanish

Well-Balanced Methods

Rogers et al.

- ▶ Rogers, Borthwick, Taylor, 2003, “Mathematical balancing of flux gradient and source terms prior to using Roe’s approximate Riemann solver”
- ▶ Change of variables: deviations from equilibrium

$$\mathbf{q}_t + \mathbf{f}(\mathbf{q})_x = \mathbf{s}(\mathbf{q}, x) \quad (16)$$

$$\rightarrow (\mathbf{q} - \mathbf{q}_{\text{eq}})_t + (\mathbf{f}(\mathbf{q}) - \mathbf{f}(\mathbf{q}_{\text{eq}}))_x = \mathbf{s}(\mathbf{q}, x) - \mathbf{s}(\mathbf{q}_{\text{eq}}, x) \quad (17)$$

$$\text{or } \mathbf{q}_t^* + \mathbf{f}_x^* = \mathbf{s}^* \quad (18)$$

- ▶ At equilibrium, all terms vanish
- ▶ Jacobian of \mathbf{f} remains unchanged
 - ▶ easy to adapt existing solver

Well-Balanced Methods

Rogers et al.

- ▶ Derivation in original paper leads to

$$\mathbf{q}_t^* + \mathbf{f}'(\mathbf{q})\mathbf{q}_x^* = \mathbf{s}^* \quad (19)$$

- ▶ Implementation based on this failed away from equilibrium

Well-Balanced Methods

Rogers et al.

- Derivation in original paper leads to

$$\mathbf{q}_t^* + \mathbf{f}'(\mathbf{q})\mathbf{q}_x^* = \mathbf{s}^* \quad (19)$$

- Implementation based on this failed away from equilibrium
- Derived alternative form:

$$\mathbf{q}_t^* + \mathbf{f}'(\mathbf{q})\mathbf{q}_x^* = \mathbf{s}^* - \frac{\partial \mathbf{f}^*}{\partial \mathbf{q}_{\text{eq}}}(\mathbf{q}_{\text{eq}})_x \quad (20)$$

- Works, but less accurate than unbalanced method

Well-Balanced Methods

Rogers et al.

- ▶ Derivation in original paper leads to

$$\mathbf{q}_t^* + \mathbf{f}'(\mathbf{q})\mathbf{q}_x^* = \mathbf{s}^* \quad (19)$$

- ▶ Implementation based on this failed away from equilibrium
- ▶ Derived alternative form:

$$\mathbf{q}_t^* + \mathbf{f}'(\mathbf{q})\mathbf{q}_x^* = \mathbf{s}^* - \frac{\partial \mathbf{f}^*}{\partial \mathbf{q}_{\text{eq}}}(\mathbf{q}_{\text{eq}})_x \quad (20)$$

- ▶ Works, but less accurate than unbalanced method
- ▶ Again, extended original work to support Coriolis term

Well-Balanced Methods

Rogers et al.

- ▶ Derivation in original paper leads to

$$\mathbf{q}_t^* + \mathbf{f}'(\mathbf{q})\mathbf{q}_x^* = \mathbf{s}^* \quad (19)$$

- ▶ Implementation based on this failed away from equilibrium
- ▶ Derived alternative form:

$$\mathbf{q}_t^* + \mathbf{f}'(\mathbf{q})\mathbf{q}_x^* = \mathbf{s}^* - \frac{\partial \mathbf{f}^*}{\partial \mathbf{q}_{\text{eq}}}(\mathbf{q}_{\text{eq}})_x \quad (20)$$

- ▶ Works, but less accurate than unbalanced method
- ▶ Again, extended original work to support Coriolis term
- ▶ Also derived solver for geostrophic equilibria
 - ▶ Implementation does not work yet

Well-Balanced Methods



Rogers et al.

- ▶ Pro:
 - ▶ Works away from equilibrium
 - ▶ Fairly simple to derive and implement
- ▶ Con:
 - ▶ Tailored to a specific equilibrium
 - ▶ Need to know equilibrium a priori
 - ▶ Less accurate than unbalanced method away from equilibrium



Evaluation

- ▶ Implemented evaluation framework in Clawpack (Python)
- ▶ Supports 7 different bathymetries and 5 initial conditions
- ▶ Implemented 4 Riemann solvers in Fortran (unbalanced, LeVeque, Rogers for still water, Rogers for geostrophic equilibrium)



Results





Results





Results





Results





Results





Results





Results





Results



Results



Results



Results

