

Math 3000. Spring 2018. Information about the final exam.

The final exam will be given on Thursday, May 10, 9am-12pm in our regular classroom (Kerchof 317). It will contain 6-8 regular problems and possibly a bonus problem. The exam will be cumulative, but with more emphasis on the material since the cutoff point for the second midterm: the second part of Lecture 20 (inverse images) and Lectures 21-26 (the corresponding sections of the book are 6.2, 6.3 and 6.4, and the corresponding homework assignments are 10 and 11). You should expect 1 or 2 long problems on the final (like the practice problem 1 for the second midterm). Also, problems on the final may combine different topics.

Here are some basic suggestions for preparing for the final:

- (i) Go over your homework assignments; whenever you lost points on a problem, make sure to understand why. If you made a minor mistake, realizing what the mistake was is probably sufficient; if your solution was not close to being correct, redo the problem.
- (ii) Work on the practice problems that were assigned throughout the semester. Many of those problems are solved in the solution manual
- (iii) Go over the class notes and prepare questions about things you do not understand.
- (iv) Go over the sections of the book that you did not fully understand on the first reading.

Additional practice problems

Note: This list only includes problems on the material covered since the second midterm. Therefore this list should NOT be considered a model for the final.

1. Let $f : A \rightarrow B$ be a function. Prove that the following two conditions are equivalent:

- (a) f is injective
- (b) $f^{-1}(f(C)) = C$ for every subset C of A .

Hint: Use contrapositive for the proof in one of the two directions.

2. Give a short proof of the furthermore part of Theorem 6.2.8(d) from the book using Problem 1 above, the first assertion of Theorem 6.2.8(d) and Theorem 6.2.9(d)

3. Let A and B be sets. Prove that the following are equivalent:

- (i) there is an injection $f : A \rightarrow B$
- (ii) there is a surjection $g : B \rightarrow A$

Hint: If you are not sure how to start, take a look at the proof of Theorem 23.4. This should help you at least with the proof of the implication “(ii) \Rightarrow (i)”.

4. Construct an explicit bijection between $(-1, 1)$ and $[-1, 1]$. **Hint:** Imitate the construction of bijection between $[0, 1)$ and $[0, 1]$ from Lecture 24 or use the procedure from Problem 5 in HW#11.

5. Let $A = [-1, 1]$ and $B = (-1, 1)$. Define the functions $f : A \rightarrow B$ and $g : B \rightarrow A$ by $f(a) = \frac{a}{2}$ and $g(b) = b$ for all $a \in A$ and $b \in B$. It is straightforward to check that f and g are both injective, so one can apply the proof of the Schroeder-Bernstein theorem to those f and g to construct a bijection $\Phi : A \rightarrow B$. Find an explicit formula for such Φ (this is a good way to test if you understand the proof of the Schroeder-Bernstein theorem).

6. Let A be a set. Prove that $|A| \neq |\mathcal{P}(A)|$, that is, there is no bijection $\phi : A \rightarrow \mathcal{P}(A)$. **Hint:** Argue by contradiction: suppose that such ϕ exists. Consider the set $B = \{a \in A \mid a \notin \phi(a)\} \in \mathcal{P}(A)$, and show that B cannot lie in the image of ϕ . The contradiction you should get is very similar to the one in Russell's paradox.