

CORVINUS METROPOLIS TEMPLATE

EXAMPLES AND USAGE

John Doe

May 4, 2026

Example Department

Corvinus University of Budapest

Email: `john.doe@uni-corvinus.hu`

Download from: <https://github.com/m-marcell/corvinusmetropolis-updated>

OUTLINE

Definition of quadratic forms

Dyadic decomposition

- Dyad and completing the square

- High school method for a few variable only

- Professional method

Problems

Definiteness of a quadratic form

- Summary

DEFINITION OF QUADRATIC FORMS

DEFINITION OF QUADRATIC FORMS

A quadratic form is a special multi-variable function.

Definition

Let $A \in \mathbb{R}^{n \times n}$ be a symmetric matrix. Assume that $a_{k,j}$ denotes the appropriate element of that matrix. The function $Q : \mathbb{R}^n \rightarrow \mathbb{R}$

$$Q(x_1, x_2, \dots, x_n) = \sum_{k=1}^n \sum_{j=1}^n a_{k,j} x_k x_j$$

is called n variables quadratic function.

If $n = 1$ then $Q(x_1) = ax_1^2$.

If $n = 2$ then

$$Q(x_1, x_2) = a_{1,1}x_1^2 + a_{1,2}x_1x_2 + a_{2,1}x_2x_1 + a_{2,2}x_2^2.$$

Compute the case when $n = 3$.

DEFINITION OF QUADRATIC FORMS

Theorem

If $A \in \mathbb{R}^{n \times n}$ is a symmetric matrix then the quadratic form determined by A is

$$Q(x) = x \cdot Ax$$

for every $x \in \mathbb{R}^n$.

Example

Write the symmetric matrix of quadratic form given before. Write the quadratic form of a symmetric matrix given before.

Theorem

The quadratic form determined by $A + B$ is the same as the sum of quadratic forms determined by A and B respectively. Similarly, the quadratic form determined by αA is the same as the form determined by A multiplied by α .

DYADIC DECOMPOSITION

DYADIC DECOMPOSITION

DYAD AND COMPLETING THE SQUARE

The dyad is a special symmetric matrix representing the perfect squares.

Definition

A symmetric matrix $A \in \mathbb{R}^{n \times n}$ is called *dyad*, if there exists a vector $a \in \mathbb{R}^{n \times 1}$ for which $A = a \cdot a^T$

Theorem

A quadratic form is a complete square if and only if it is determined by a dyad. If

$A = a \cdot a^T$ where $a = (a_1, a_2, \dots, a_n)$, then

$Q(x_1, x_2, \dots, x_n) = (a_1x_1 + a_2x_2 + \dots + a_nx_n)^2$, here Q is the quadratic form determined by the dyad A .

DYADIC DECOMPOSITION

HIGH SCHOOL METHOD FOR A FEW VARIABLE
ONLY

Theorem

Every symmetric matrix is a linear combination of dyads. Thus every quadratic form is a linear combination of complete squares.

Example

Completing the square, rewrite the following quadratic forms as a linear combination of perfect squares.

$$Q(x_1, x_2) = x_1^2 + 4x_1x_2 - 5x_2^2,$$

$$Q(x_1, x_2) = 4x_1^2 + 6x_1x_2 + 9x_2^2,$$

$$Q(x_1, x_2) = 5x_1^2 - 6x_1x_2 + x_2^2,$$

$$Q(x_1, x_2, x_3) = 5x_1^2 - 6x_1x_2 + x_1x_3,$$

DYADIC DECOMPOSITION

PROFESSIONAL METHOD

DYADIC DECOMPOSITION USING GAUSS–JORDAN ELIMINATION

Every n -variable quadratic form can be decreased by a multiply of a dyad such that the remainder quadratic form has one variable less:

Theorem

Consider a symmetric matrix $A \in \mathbb{R}^{n \times n}$. Assume, that $a_{1,1} \neq 0$ and denote $d = \frac{1}{a_{1,1}}a_1$, where a_1 is the first column of matrix A . Then

$$R = A - a_{1,1}d \cdot d^T$$

is a symmetric matrix, where the first row (and column) is the zero vector.

Proof.

The sum of symmetric matrices is a symmetric matrix. If δ_k denotes the k -th coordinate of d , then $r_{k,j} = a_{k,j} - a_{1,1}\delta_k\delta_j$. If $k = 1$ we obtain

$$r_{1,j} = a_{1,j} - a_{1,1} \cdot 1 \cdot \frac{a_{1,j}}{a_{1,1}} = 0.$$



The k, j term of the matrix of the remainder quadratic form is

$$r_{k,j} = a_{k,j} - a_{1,1}\delta_k\delta_j = a_{k,j} - a_{1,1}\frac{a_{k,1}}{a_{1,1}}\delta_j = a_{k,j} - a_{k,1}\delta_j.$$

If $k > 1$ and $j > 1$, then the above formula just computes the new coordinates of the j -th column after the first column has entered to the base at the first position. Thus we proved the following theorem.

Theorem

Let $A \in \mathbb{R}^{n \times n}$ be a symmetric matrix and assume that $a_{1,1} \neq 0$. After separating the first complete square, as in the theorem above, the matrix of the remainder quadratic form is the bottom right $n - 1 \times n - 1$ matrix of the transformation table, when $a_{1,1}$ is the pivot term.

PROBLEMS

Write the following quadratic forms as a linear combination of complete squares.

$$1. Q(x_2, x_3) = 13x_2^2 - 4x_2x_3 + 8x_3^2,$$

$$2. Q(x_1, x_2, x_3) = 5x_1^2 + 6x_2^2 + 4x_3^2 - 4x_1x_2 - 4x_1x_3,$$

$$3. Q(z_1, z_2, z_3) = 2z_1^2 + \frac{3}{2}z_3^2 + 2z_1z_2 - 4z_1z_3 + 2z_2z_3,$$

$$4. Q(x_1, x_2, x_3, x_4) = 2x_1^2 + 2x_2^2 + 2x_3^2 + 2x_4^2 - 4x_1x_2 + 2x_1x_4 + 2x_2x_3 - 4x_3x_4$$

DEFINITENESS OF A QUADRATIC FORM

Classify a quadratic form with respect to the range!

Definition

Let $Q : \mathbb{R}^n \rightarrow \mathbb{R}$ be a nonzero quadratic form. It is called

positive definite, if $Q(x) > 0$ for every $x \in \mathbb{R}^n, x \neq 0$;

positive semidefinite, if $Q(x) \geq 0$ for every $x \in \mathbb{R}^n$, but there exists a $z \in \mathbb{R}^n, z \neq 0$ such that $Q(z) = 0$;

negative definite, if $Q(x) < 0$ for every $x \in \mathbb{R}^n, x \neq 0$;

negative semidefinite, if $Q(x) \leq 0$ for every $x \in \mathbb{R}^n$, but there exists a $z \in \mathbb{R}^n, z \neq 0$ such that $Q(z) = 0$;

indefinite, if there exist two vectors $x, y \in \mathbb{R}^n$ for which $Q(x) > 0$ but $Q(y) < 0$.

DEFINITENESS OF A QUADRATIC FORM

SUMMARY

DEFINITENESS AFTER COMPLETING THE SQUARE

It is not hard to decide the definiteness of a quadratic form if it is given with its dyad decomposition. Let Q be an n -variable none-zero quadratic form.

If the dyad decomposition

- includes positive and negative pivot terms then Q is an *indefinite* quadratic form.

Now assume the dyad decomposition includes exactly n complete squares. If

- all pivot terms are positive numbers then Q is *positive definite*;
- all pivot terms are negative numbers then Q is *negative definite*.

DEFINITENESS AFTER COMPLETING THE SQUARE ...

If the number of dyads is less than n , the same characterization holds true but the quadratic form is semidefinite only:

Thus if

- all pivot terms are positive numbers then Q is *positive semidefinite*;
- all pivot terms are negative numbers then Q is *negative semidefinite*.

THANK YOU FOR YOUR ATTENTION!



`john.doe@uni-corvinus.hu`

