NAGAR PATHOLOGY LABORATORY

Dr. G.K. Nagar
M.D. (Path. & Bact.)
Kalpavraksha Karyalaya, Tilak Marg
UJJAIN (M.P.) • Ph. : 2555119 (Lab.)

Dr. Vikas Nagar
M.B.B.S
5-A, Dhanwantari Marg, Freeganj
UJJAIN (M.P.) • Ph. : 4370150 (Lab.)

REPORT

Patient Name : MR.RADHESHYAM MITTAL **Date :** 15-Nov-2018

Reffered By : DR.

** HbA1c **

HbA1c : 8.10 %

In last 3 months the mean Blood Glucose Level was Apox. **186** mg./dl.

Reference Values:

Normal	:	<5.7 %
Pre Diabetes	:	5.7 - 6.4 %
Diabetes	:	>6.4 %

Test done on PocketChem "A1c-HbA1c analyzer" Arkray(Japan)
Test principle : Boronate Flourosence Quenching Technology (BFQT)

(Dr.G.K.Nagar) **(Dr.Vikas Nagar)**

What is HbA1c?
HbA1c is Glucose which is bound to Hemoglobin in the blood. It is also known as Glycated/Glycosylated/Glyco Hemoglobin. It gives average Blood Glucose for the last 3 months(90 days) and is a "True indicator" of Diabetes control. It is indicated as a % of total hemoglobin.
HbA1c can be performed at any time of the day(no need for fasting)

Why HbA1c?

As per UKPDS Study, a 1% reduction in HbA1c is associated with :
- 21% reduction in deaths due to diabetes.
- 37% reduction in risk of Microvascular complications.
- 14% reduction in risk of Myocardial Infarction.

Lowering A1C to approximately 7% or less has been shown to reduce microvascular complications of diabetes and if implemented soon after the diagnosis of diabetes, it is associated with long-term reduction in macrovascular disease.

When to test for HbA1c?
As per 2015 ADA Standards for Diabetes care,
- At least two times a year in patients who are meeting treatment goals(and who have stable glycemic control).
- Once in 3 months in patients whose therapy has changed or who are not meeting glycemic goals.

Use of point-of-care testing for A1C provides the opportunity for more timely treatment changes.

- This report is not for medicolegal purpose.
- Subject to Ujjain Jurisdiction only.
- In case of any doubt test can be repeated.