# Pooling Versus Ensembling for Ridge Regression Under Covariate Shift

**Anonymous Author(s)**
Affiliation
Address
email

## Abstract

Datasets in many settings naturally partition into clusters arising from sub-populations, batch effects, or aggregation across multiple sources. A common response to such heterogeneity is to ensemble learners trained on each cluster rather than fit a single model to the pooled data. Prior work motivating such approaches has typically considered settings in which both the covariate distribution and the conditional outcome model differ across clusters; the role of cluster-aware partitioning and ensembling based solely on the covariate distribution remains to be explored. We address this case for ridge-regularized least-squares regression under a linear outcome model and consider all ridge penalty values $\lambda \geq 0$, including the special case of the ridgeless predictor at $\lambda = 0$. By considering both fixed-effects and random-effects models, we argue that under random effects, an optimally tuned pooled ridge predictor always outperforms ensembles of individually optimally tuned predictors. For fixed effects, we derive a general formula for the pooled and ensembled predictors to characterize the role of both regression coefficients as well as the predictor distribution shifts. Together, these results generalize prior risk analyses of bagging and random-partition estimation using ridge and ridgeless regression predictors from the i.i.d. setting to encompass covariate shift and heterogeneity-aware partition structure.

## 1 Introduction

Heterogeneity in the distribution of covariates is a common artifact of modern datasets, where data can be composed of natural sub-populations, clusters, batch effects, and multiple sources [Chauhan et al., 2010, Goh et al., 2017]. A long-standing question in this setting is whether prediction models should explicitly acknowledge this structure or ignore it. This question has a substantial classical literature on generalization across multiple distributions [Ben-David et al., 2010, Crammer et al., 2008, Mansour et al., 2008] and on pooled estimation under heterogeneity [Li et al., 2022, Meinshausen and Bühlmann, 2015]. One natural response has been the development of ensembling frameworks, in which the training data is first separated into its component clusters or natural partitions, a learning algorithm is trained independently on each cluster, and the resulting single-cluster predictors are combined using weights that reward cross-cluster generalization on the training set [Deodhar and Ghosh, 2007, Guan et al., 2019, Patil and Parmigiani, 2018, Ramchandran and Mukherjee, 2021, Ramchandran et al., 2020, Trivedi et al., 2015]. Across a range of base learners, including neural networks, random forests, and regularized regression, such ensembles have been shown empirically to generalize more reliably than a single model trained on the pooled data.

Throughout the paper, we refer to this strategy simply as *ensembling*, and contrast it with *pooling*, which refers to training a single model on the entire training set with the same base learner. More specifically, we compare two procedures built from the same base learner. The *pooled* procedure trains a single predictor on the pooled training data. The *ensemble* procedure trains one predictor

Submitted to 40th Conference on Neural Information Processing Systems (NeurIPS 2026). Do not distribute.

separately on each known cluster and then averages, or more generally convexly combines, their predictions. Our goal is to determine whether the second strategy offers any advantage when clusters differ only in their covariate distributions and share the same conditional outcome model.

Although ensembling has been motivated by settings in which both the covariate distribution and the conditional outcome model vary across sources, it is far less clear what role cluster-aware partitioning plays when the heterogeneity is purely in the covariates. In this paper, we isolate this question by studying a setting with generalized covariate shift but no concept shift. Specifically, we analyze ridge-regularized least-squares regression with outcomes generated by a well-specified linear model whose coefficients are shared across clusters and test points; only the marginal distribution of the covariates varies both within the training set and between train and test sets.

### 1.1 Summary of Contributions

**Theoretical results.** We provide general formulas for the risk of the ensemble and pooled learners using ridge-regularized least-squares regression as the base learner under a well-specified linear outcome model and arbitrary cluster-specific covariate distributions. When the ridge penalty $\lambda > 0$, we prove that the pooled predictor has uniformly smaller (asymptotic) prediction risk than the ensemble built on the true cluster partitions. This holds independently of the angles, norms, and covariance structures of the cluster means. When the ridge penalty $\lambda = 0$ for the ridgeless predictor, we show that if both learners are in either the underparameterized or overparameterized regime, the pooled predictor similarly outperforms the ensemble; the only point at which this relationship switches is when the pooled predictor is at the interpolation threshold where the total number of samples equals the number of covariates, whereas the base learners constituting the ensemble are in the over-parameterized regime. In this case, the implicit regularization caused by overparameterization allows the ensemble to achieve lower variance than the pooled predictor.

**Simulation studies across regimes.** We confirm our theoretical results with simulations for both ridge and ridgeless predictors that span the underparameterized regime, the interpolation threshold, and the overparameterized regime. In every configuration we consider—varying cluster mean angles, cluster mean norms, and cluster covariance structures—the pooled predictor matches or outperforms the ensemble when $\lambda > 0$ (ridge regression). For ridgeless regression at $\lambda = 0$, the simulations confirm the theoretical result that the relationship between approaches depends on the regime, with the pooled learner largely dominating the ensemble when both are under- or over-parameterized. Furthermore, we find that the ensembling-versus-pooling gap is driven by the aspect ratio $p/n$, as opposed to the covariate distribution or partition strategy.

### 1.2 Related Work

Our work builds on two closely related lines of recent research. Hastie et al. [2022] provide the exact asymptotic risk formulas for ridge and ridgeless regression that underpin the qualitative behavior we observe, and Patil et al. [2023] analyze a class of bagging procedures (subagging and splagging) in a supervised setting similar to ours. We defer discussion of the related unsupervised stacked-SVD setting of Baharav et al. [2025] to Appendix A.1, since our focus is on supervised prediction.

**Asymptotic risk of ridge and ridgeless regression.** A line of recent work has produced precise asymptotic risk characterizations for high-dimensional regression in the proportional regime $p/n \to \gamma$, building on classical random matrix theory. Dobriban and Wager [2018] provide early asymptotic risk formulas for ridge regression and classification in this regime. Hastie et al. [2022] subsequently derive exact formulas for the prediction risk of the minimum-norm (ridgeless) least-squares estimator and of ridge regression under a well-specified linear model with arbitrary covariate covariance. Their formulas make explicit several phenomena central to our experimental story: the ridgeless risk diverges at the interpolation threshold $\gamma = 1$ and decreases beyond it (the "double descent" shape, identified empirically by Belkin et al. [2019] and analyzed as *benign overfitting* by Bartlett et al. [2020]), and even a small ridge penalty smooths the interpolation peak away. We use these results as a direct point of reference when interpreting the risk profile of our two learners.

**Bagging in overparameterized regression.** Patil et al. [2023] characterize the asymptotic prediction risk of two bagging variants for ridge and ridgeless least-squares predictors in the



Confidential reviewer copy. This manuscript is submitted to the 40th Conference on Neural Information Processing Systems (NeurIPS 2026). Unauthorized sharing, redistribution, or disclosure is strictly prohibited.

proportional-asymptotic regime: *subagging*, in which $K$ predictors are trained on possibly overlapping random subsamples, and *splagging* (split-aggregation), in which the data is partitioned via uniform random permutation into $K$ disjoint equal-size subsets and the per-partition predictors are averaged. Splagging is structurally similar to our ensemble learner with $K = 2$, with two key differences: their partitions are uniformly random, and they assume i.i.d. train and test data with no covariate shift or internal heterogeneity. Their main results are established under a well-specified linear model with arbitrary covariate covariance and arbitrary true coefficient vector. They show that properly tuned ridge regression is optimal among the methods considered, with no ensembling strategy surpassing it. One notable finding is that under isotropic covariates, the optimally-tuned subagged ridgeless predictor matches the asymptotic risk of optimally-tuned ridge; that is, bagging implicitly performs the regularization that ridge performs explicitly. This is further illustrated by comparing ensembles of ridgeless predictors trained on subsets to a single ridgeless predictor trained on the full dataset: the ensemble shows benefit largely at the full predictor's interpolation threshold, where each ensemble component has been pushed into the overparameterized regime and exhibits variance stabilization.

LeJeune et al. [2020] make this implicit-regularization view explicit, showing that in the proportional asymptotic regime, ensembles of ordinary least-squares predictors trained on random subsets behave like a single predictor with an effective ridge penalty determined by the subsample size. In a complementary fixed-dimension kernel-ridge setting, Zhang et al. [2015] analyze a divide-and-conquer estimator that partitions the data into $K$ disjoint subsets, fits kernel ridge regression on each, and averages the predictors; they show that for $K$ not too large, this matches the convergence rate of the predictor trained on the full dataset. Together, these results show that bagging and partitioning act as implicit regularization, with primary gains for linear predictors concentrated in regimes where interpolation would otherwise produce high variance; none, however, exceed optimally tuned ridge. Our work extends this picture by asking how covariate heterogeneity and heterogeneity-aware partitioning affect the benefit of ensembling versus pooling: rather than assuming i.i.d. data with uniformly random partitions, we study a mixture of clusters with heterogeneous covariate distributions and partition along the underlying cluster structure, investigating whether the conclusions of Patil et al. [2023] continue to hold.

# 2 Main Results

We organize the main results into subsections covering the mathematical setup, the analyses of the ensemble and pooled learners, and their comparison.

## 2.1 Setup

We theoretically compare the asymptotic risk of an ensemble predictor to that of a pooled predictor under a linear outcome model. The data are composed of $K$ clusters (or sources): $\mathcal{D}_\ell = (y_{i,\ell}, \boldsymbol{X}_{i,\ell})_{i=1}^{n_\ell}$, $\ell = 1, \dots, K$, with outcome $y \in \mathbb{R}$ and covariates $\boldsymbol{X} \in \mathbb{R}^p$. Let $\hat{f}_\ell$ be a predictor fit on each cluster $\mathcal{D}_\ell$, and let $\hat{f}_{\text{pool}}$ be a predictor fit on the pooled dataset $\mathcal{D} = \{\mathcal{D}_\ell : \ell = 1, \dots, K\}$. Our main results compare the prediction error weighted combinations of the $\hat{f}_\ell$'s to that of $\hat{f}_{\text{pool}}$, with all learners fit by ridge-regularized least squares.

To stage the analysis, we define

$$\mathbf{Y}_\ell = (Y_{1,\ell}, \dots, Y_{n_\ell,\ell})^\top \in \mathbb{R}^{n_\ell}, \qquad \mathbb{X}_\ell = (\boldsymbol{X}_{1,\ell}, \dots, \boldsymbol{X}_{n_\ell,\ell})^\top \in \mathbb{R}^{n_\ell \times p}$$

for the outcome vector and design matrix in cluster $\ell$, and let $\mathbf{Y}_{\text{pool}} = (\mathbf{Y}_1^\top, \dots, \mathbf{Y}_K^\top)^\top$ and $\mathbb{X}_{\text{pool}} = (\mathbb{X}_1^\top, \dots, \mathbb{X}_K^\top)^\top$ denote their pooled counterparts. The ridge predictors are then

$$\hat{f}_\ell(\mathbf{x}) = \hat{\boldsymbol{\beta}}_{\lambda_\ell}^\top \mathbf{x}, \qquad \hat{\boldsymbol{\beta}}_{\lambda_\ell} = (\mathbb{X}_\ell^\top \mathbb{X}_\ell + \lambda_\ell \mathbb{I})^{-1} \mathbb{X}_\ell^\top \mathbf{Y}_\ell, \qquad \lambda_\ell > 0;$$
$$\hat{f}_{\text{pool}}(\mathbf{x}) = \hat{\boldsymbol{\beta}}_{\lambda_0}^\top \mathbf{x}, \qquad \hat{\boldsymbol{\beta}}_{\lambda_0} = (\mathbb{X}_{\text{pool}}^\top \mathbb{X}_{\text{pool}} + \lambda_0 \mathbb{I})^{-1} \mathbb{X}_{\text{pool}}^\top \mathbf{Y}_{\text{pool}}, \qquad \lambda_0 > 0,$$

with ridgeless ($\lambda \downarrow 0$) counterparts defined via the Moore-Penrose pseudoinverse:

$$\hat{f}_{\ell,+}(\mathbf{x}) = \hat{\boldsymbol{\beta}}_{+,\ell}^\top \mathbf{x}, \qquad \hat{\boldsymbol{\beta}}_{+,\ell} = (\mathbb{X}_\ell^\top \mathbb{X}_\ell)^+ \mathbb{X}_\ell^\top \mathbf{Y}_\ell;$$
$$\hat{f}_{\text{pool},+}(\mathbf{x}) = \hat{\boldsymbol{\beta}}_{+}^\top \mathbf{x}, \qquad \hat{\boldsymbol{\beta}}_{+} = (\mathbb{X}_{\text{pool}}^\top \mathbb{X}_{\text{pool}})^+ \mathbb{X}_{\text{pool}}^\top \mathbf{Y}_{\text{pool}}.$$



Confidential reviewer copy. This manuscript is submitted to the 40th Conference on Neural Information Processing Systems (NeurIPS 2026). Unauthorized sharing, redistribution, or disclosure is strictly prohibited.

We work under the following assumptions.

**Assumption 2.1.** *The following hold for $\ell = 1, \dots, K$.*

*(i)* ***Covariate Distribution and Mean Shifts:*** *$\boldsymbol{X}_{i,\ell} = \boldsymbol{\mu}_\ell + \Sigma^{1/2}\mathbf{Z}_{i,\ell}$, where $\mathbf{Z}_{i,\ell} \in \mathbb{R}^p$ has i.i.d. mean-zero subgaussian coordinates and $\Sigma$ is positive definite with $\eta \le \lambda_1(\Sigma) \le \lambda_p(\Sigma) < \eta^{-1}$ for some fixed $\eta > 0$. Moreover, $K$ is fixed and the cluster means $\boldsymbol{\mu}_1, \dots, \boldsymbol{\mu}_K$ have uniformly bounded Euclidean norms.*

*(ii)* ***Outcome Regression:*** *$Y_{i,\ell} = \boldsymbol{\beta}^\top \boldsymbol{X}_{i,\ell} + \varepsilon_{i,\ell}$, with $\varepsilon_{i,\ell} \perp \boldsymbol{X}_{i,\ell}$ subgaussian, mean $0$, and variance $\sigma_\varepsilon^2$.*

*(iii)* ***Test Data:*** *$y^\star = \boldsymbol{\beta}^\top \mathbf{x}^\star + \varepsilon^\star$ with $\mathbb{E}(\mathbf{x}^\star) = 0$, $\mathrm{Var}(\mathbf{x}^\star) = \mathbb{I}$, $\varepsilon^\star \perp \mathbf{x}^\star$, $\mathbb{E}(\varepsilon^\star) = 0$, $\mathrm{Var}(\varepsilon^\star) = \sigma_\star^2$.*

*(iv)* ***Asymptotic Regime:*** *For $\ell \ge 0$, $p/n_\ell \to \gamma_\ell \in [0, \infty)$ and $\lambda_\ell = \rho_\ell n_\ell$ for fixed $\rho_\ell \in (0, \infty)$, where $n_0 = n = \sum_{\ell=1}^K n_\ell$.*

**Remark 2.2.** *Assumption 2.1 focuses on mean shifts as the source of covariate heterogeneity. This choice keeps the comparison between pooled and ensemble learners analytically transparent while still capturing a central form of between-cluster variation. The same random matrix arguments can be extended to more general forms of covariate shift, though at the cost of heavier notation. Our choice of a centered, isotropic test data point $\mathbf{x}^\star$ is also driven by the same goal of keeping our theoretical results relatively concise. Finally, the cluster labels are treated as known. This is natural when the data come from distinct studies or populations. When clusters are estimated from the data, additional uncertainty enters the analysis; we leave this extension for future work.*

Under these assumptions, we study the risk of ensemble versus pooled learners. For convex weights $\mathbf{w} = (w_1, \dots, w_K) \in \mathbb{R}_+^K$ with $\sum_{\ell=1}^K w_\ell = 1$ and $\boldsymbol{\rho} = (\rho_1, \dots, \rho_K)$ with $\lambda_\ell = \rho_\ell n_\ell$, define the ensemble estimators

$$\widehat{\boldsymbol{\beta}}_{\mathrm{ens},\boldsymbol{\rho}}(\mathbf{w}) = \sum_{\ell=1}^K w_\ell \widehat{\boldsymbol{\beta}}_{\lambda_\ell}, \qquad \widehat{\boldsymbol{\beta}}_{\mathrm{ens},+}(\mathbf{w}) = \sum_{\ell=1}^K w_\ell \widehat{\boldsymbol{\beta}}_{+,\ell}, \qquad w_\ell \ge 0,\ \sum_{\ell=1}^K w_\ell = 1.$$

Conditional on $\{\mathbb{X}_\ell\}_{\ell=1}^K$, their mean squared prediction errors are

$$R_{\mathrm{ens},\boldsymbol{\rho}}(\mathbf{w}) = \mathbb{E}\Big[(y^\star - \mathbf{x}^{\star\top}\widehat{\boldsymbol{\beta}}_{\mathrm{ens},\boldsymbol{\rho}}(\mathbf{w}))^2\Big], \qquad R_{\mathrm{ens},+}(\mathbf{w}) = \mathbb{E}\Big[(y^\star - \mathbf{x}^{\star\top}\widehat{\boldsymbol{\beta}}_{\mathrm{ens},+}(\mathbf{w}))^2\Big].$$

The pooled risks are defined analogously: for $\lambda_0 = \rho_0 n_0$,

$$R_{\mathrm{pool},\rho_0} = \mathbb{E}\Big[(y^\star - \mathbf{x}^{\star\top}\widehat{\boldsymbol{\beta}}_{\lambda_0})^2\Big], \qquad R_{\mathrm{pool},+} = \mathbb{E}\Big[(y^\star - \mathbf{x}^{\star\top}\widehat{\boldsymbol{\beta}}_{\mathrm{pool},+})^2\Big].$$

The next two subsections analyze the ensemble and pooled methods in turn; we compare them in Section 2.4. Stating the results requires further notation. Let $F_{\Sigma,p}$ denote the empirical spectral distribution of $\Sigma$, and for $\gamma, \rho > 0$ let $m_\gamma(\rho)$ be the unique solution [Knowles and Yin, 2017] of

$$\frac{1}{m_\gamma(\rho)} = \rho + \gamma \int \frac{x}{1 + x m_\gamma(\rho)}\, dF_{\Sigma,p}(x).$$

Define $Q_\gamma(\rho) = [\rho\{\mathbb{I} + m_\gamma(\rho)\Sigma\}]^{-1}$, $a_\gamma(\rho) = \frac{1}{p}\mathrm{Tr}Q_\gamma(\rho)$, $s_\gamma(\rho) = \frac{1}{p}\mathrm{Tr}\{\Sigma Q_\gamma(\rho)\}$, and the variance functional

$$V_\gamma(\rho) = \gamma\{a_\gamma(\rho) + \rho a_\gamma'(\rho)\},$$

with the derivative taken with respect to $\rho$. Cluster-specific values are written $Q_\ell = Q_{\gamma_\ell}(\rho_\ell)$, $a_\ell = a_{\gamma_\ell}(\rho_\ell)$, $V_\ell = V_{\gamma_\ell}(\rho_\ell)$. Finally, for deterministic $U \in \mathbb{R}^{p\times r}$ of fixed rank $r$ and uniformly bounded operator norm, set

$$\mathcal{K}_{\gamma,U}(\rho) = Q_\gamma(\rho) - Q_\gamma(\rho)U\big[\{1 + \gamma s_\gamma(\rho)\}\mathbb{I} + U^\top Q_\gamma(\rho)U\big]^{-1}U^\top Q_\gamma(\rho),$$

which when $U = \boldsymbol{\mu}$ is a single vector reduces to

$$\mathcal{K}_{\gamma,\boldsymbol{\mu}}(\rho) = Q_\gamma(\rho) - \frac{Q_\gamma(\rho)\boldsymbol{\mu}\boldsymbol{\mu}^\top Q_\gamma(\rho)}{1 + \gamma s_\gamma(\rho) + \boldsymbol{\mu}^\top Q_\gamma(\rho)\boldsymbol{\mu}}.$$

We are now ready to state the main results.



Confidential reviewer copy. This manuscript is submitted to the 40th Conference on Neural Information Processing Systems (NeurIPS 2026). Unauthorized sharing, redistribution, or disclosure is strictly prohibited.

## 2.2 Analysis of Ridge Regression Ensembles

We split by the assumption on $\boldsymbol{\beta}$, treating the random- and fixed-effects cases in turn. The next theorem collects both.

**Theorem 2.3.** *Assume 2.1.*

1. *Let $\boldsymbol{\beta}$ be random with $\mathbb{E}(\boldsymbol{\beta}) = 0$ and $\mathrm{Var}(\boldsymbol{\beta}) = \tau^2\mathbb{I}/p$. Then*

$$R_{\mathrm{ens},\boldsymbol{\rho}}(\mathbf{w}) = \sigma_\star^2 + \sum_{\ell=1}^{K} w_\ell^2 \left\{\tau^2\rho_\ell^2[-a'_{\gamma_\ell}(\rho_\ell)] + \sigma_\varepsilon^2 V_{\gamma_\ell}(\rho_\ell)\right\} + 2\tau^2 \sum_{1\le\ell<j\le K} w_\ell w_j \rho_\ell\rho_j \, \tfrac{1}{p}\mathrm{Tr}(Q_\ell Q_j) + o(1),$$
$$R_{\mathrm{ens},+}(\mathbf{w}) = \lim_{\substack{\rho_\ell\downarrow 0 \\ \ell=1,\dots,K}} R_{\mathrm{ens},\boldsymbol{\rho}}(\mathbf{w}) + o(1).$$

2. *If $\limsup \|\boldsymbol{\beta}\| < \infty$, set $\mathcal{K}_\ell(\rho) = \mathcal{K}_{\gamma_\ell,\boldsymbol{\mu}_\ell}(\rho)$ and $\psi_\ell(\rho) = \boldsymbol{\beta}^\top\mathcal{K}_\ell(\rho)\boldsymbol{\beta}$. Then*

$$R_{\mathrm{ens},\boldsymbol{\rho}}(\mathbf{w}) = \sigma_\star^2 + \sum_{\ell=1}^{K} w_\ell^2 \left\{\rho_\ell^2[-\psi'_\ell(\rho_\ell)] + \sigma_\varepsilon^2 V_{\gamma_\ell}(\rho_\ell)\right\} + 2\sum_{1\le\ell<j\le K} w_\ell w_j \rho_\ell\rho_j \, \boldsymbol{\beta}^\top\mathcal{K}_\ell(\rho_\ell)\mathcal{K}_j(\rho_j)\boldsymbol{\beta} + o(1),$$
$$R_{\mathrm{ens},+}(\mathbf{w}) = \lim_{\substack{\rho_\ell\downarrow 0 \\ \ell=1,\dots,K}} R_{\mathrm{ens},\boldsymbol{\rho}}(\mathbf{w}) + o(1).$$

In the proof, the first summand in each risk follows fairly directly from Hastie et al. [2019], with extra care for the mean shifts. The cross terms require more work, especially in the fixed-effects case: there we use a careful leave-two-out technique to control bilinear forms of resolvents of the sum of sample covariance matrices of non-mean-zero random vectors, where the vectors entering the bilinear form themselves depend on the resolvent's entries. Specializing to the optimally weighted ensemble requires the asymptotic behavior of the optimal weights, which we describe next.

**Theorem 2.4.** *Assume 2.1 and let $\boldsymbol{\beta}$ be random with $\mathbb{E}(\boldsymbol{\beta}) = 0$ and $\mathrm{Var}(\boldsymbol{\beta}) = \tau^2\mathbb{I}/p$. For each $\ell = 1,\dots,K$, let $\lambda_\ell^\star = n_\ell\rho_\ell^\star$ denote the optimal tuning for $\hat{f}_\ell(\mathbf{x}) = \mathbf{x}^\top\hat{\boldsymbol{\beta}}_{\lambda_\ell}$, and let $\mathbf{w}^\star$ denote the optimal convex weights for ensembling the optimally tuned $\hat{f}_\ell$. Define $A_{\mathrm{RE}}^\star \in \mathbb{R}^{K\times K}$ by*

$$A_{\mathrm{RE}}^\star(\ell,\ell) = \tau^2(\rho_\ell^\star)^2[-a'_{\gamma_\ell}(\rho_\ell^\star)] + \sigma_\varepsilon^2 V_{\gamma_\ell}(\rho_\ell^\star), \quad A_{\mathrm{RE}}^\star(\ell,\ell') = \tau^2\rho_\ell^\star\rho_{\ell'}^\star \, \tfrac{1}{p}\mathrm{Tr}(Q_\ell^\star Q_{\ell'}^\star),$$

*where $Q_\ell^\star = Q_{\gamma_\ell}(\rho_\ell^\star)$. Then:*

1. *The risk of $\sum_{\ell=1}^K w_\ell\hat{\boldsymbol{\beta}}_{\lambda_\ell^\star}$ equals $\sigma_\star^2 + \mathbf{w}^\top A_{\mathrm{RE}}^\star\mathbf{w} + o(1)$.*
2. *If some $\widetilde{\mathbf{w}} \in \arg\min_{\mathbf{w}\in\mathbb{R}^K:\mathbf{1}^\top\mathbf{w}=1} \mathbf{w}^\top A_{\mathrm{RE}}^\star\mathbf{w}$ satisfies $\widetilde{\mathbf{w}} \in \mathbb{R}_+^K$, the risk of $\sum_{\ell=1}^K w_\ell^\star\hat{\boldsymbol{\beta}}_{\lambda_\ell^\star}$ equals $\sigma_\star^2 + \left(\mathbf{1}^\top(A_{\mathrm{RE}}^\star)^{-1}\mathbf{1}\right)^{-1} + o(1)$.*

We state the optimally weighted, optimally tuned ensemble result only for the random-effects setting because this case yields a compact and interpretable expression. The same strategy can be applied in the fixed-effects setting, but the resulting optimal weights depend on the configuration of the mean shifts $\boldsymbol{\mu}_1,\dots,\boldsymbol{\mu}_K$ and on their relationship with the true coefficient vector $\boldsymbol{\beta}$. These relationships are challenging to estimate from data in the proportional asymptotic regime, which prevents a clean general comparison between pooled and ensemble learners under fixed effects. The same strategy as in Theorem 2.4 can nonetheless be applied to fixed-effects optimal ensembling, which we leave for future work.

## 2.3 Analysis of Pooled Ridge Regression

We split by the assumption on $\boldsymbol{\beta}$, treating the random- and fixed-effects cases in turn. The next theorem collects both.



Confidential reviewer copy. This manuscript is submitted to the 40th Conference on Neural Information Processing Systems (NeurIPS 2026). Unauthorized sharing, redistribution, or disclosure is strictly prohibited.

**Theorem 2.5.** *Assume Assumption 2.1 and $n_\ell/n \to \pi_\ell \in (0,1)$.*

1. *Let $\boldsymbol{\beta}$ be random with $\mathbb{E}(\boldsymbol{\beta}) = 0$ and $\mathrm{Var}(\boldsymbol{\beta}) = \tau^2\mathbb{I}/p$. Then*
$$R_{\mathrm{pool},\rho_0} = \sigma_\star^2 + \tau^2\rho_0^2[-a'_{\gamma_0}(\rho_0)] + \sigma_\varepsilon^2 V_{\gamma_0}(\rho_0) + o(1),$$
$$R_{\mathrm{pool},+} = \lim_{\rho_0 \downarrow 0} R_{\mathrm{pool},\rho_0} + o(1).$$

2. *If $\limsup \|\boldsymbol{\beta}\| < \infty$, set $U_0 = [\sqrt{\pi_1}\boldsymbol{\mu}_1, \ldots, \sqrt{\pi_K}\boldsymbol{\mu}_K]$, $\mathcal{K}_0(\rho) = \mathcal{K}_{\gamma_0,U_0}(\rho)$, and $\psi_0(\rho) = \boldsymbol{\beta}^\top\mathcal{K}_0(\rho)\boldsymbol{\beta}$. Then*
$$R_{\mathrm{pool},\rho_0} = \sigma_\star^2 + \tau^2\rho_0^2[-\psi'_0(\rho_0)] + \sigma_\varepsilon^2 V_{\gamma_0}(\rho_0) + o(1),$$
$$R_{\mathrm{pool},+} = \lim_{\rho_0 \downarrow 0} R_{\mathrm{pool},\rho_0} + o(1).$$

As in the ensemble analysis, most of the proof effort goes into the fixed-effects case, where we again must control bilinear forms of resolvents of the sum of sample covariance matrices of non-mean-zero random vectors, with the vectors entering the form depending on the resolvent's entries. With these results, we can now compare the two methods.

### 2.4 Comparison of Ensemble vs Pooled Learners

We compare the pooled and ensemble predictors via specific instances of Theorems 2.3 and 2.5. Setting $K = 2$, $\Sigma = \mathbb{I}$, and random effects on $\boldsymbol{\beta}$ admits a clean comparison and lets us quantify the precise benefit of the pooled learner over the optimally tuned, optimally weighted ensemble.

**Corollary 2.6.** *Assume 2.1 with $\Sigma = \mathbb{I}$ and let $\boldsymbol{\beta}$ be random with $\mathbb{E}(\boldsymbol{\beta}) = 0$ and $\mathrm{Var}(\boldsymbol{\beta}) = \tau^2\mathbb{I}/p$. Set $\gamma_0 = p/n$, $\gamma_\ell = p/n_\ell$ for $\ell = 1, \ldots, K$, $\alpha = \sigma_\varepsilon^2/\tau^2$, $\rho_\ell^\star = \gamma_\ell\alpha$ for $\ell \geq 0$, and*
$$\kappa_\ell := \kappa(\gamma_\ell) = \frac{-(\alpha - 1 + \gamma_\ell^{-1}) + \sqrt{(\alpha - 1 + \gamma_\ell^{-1})^2 + 4\alpha}}{2}, \qquad \ell \geq 0.$$
*With $\boldsymbol{\kappa} = (\kappa_1, \ldots, \kappa_K)^\top$, set $B = \mathrm{diag}(\kappa_1(1-\kappa_1), \ldots, \kappa_K(1-\kappa_K)) + \boldsymbol{\kappa}\boldsymbol{\kappa}^\top$. Then:*

1. *The optimally tuned pooled estimator is asymptotically no worse than the optimally tuned, optimally weighted ensemble if and only if*
$$\kappa_0 \leq \min_{\mathbf{w}=(w_1,\ldots,w_k)\in\mathbb{R}_+^K : \sum_{\ell=1}^K w_\ell = 1} \mathbf{w}^\top B\mathbf{w}.$$

2. *Suppose $K = 2$, and let $R_{\mathrm{ens}}^\star$ and $R_{\mathrm{pool}}^\star$ denote the mean squared prediction errors of the optimally weighted, optimally tuned ensemble and of the optimally tuned pooled regression. Then*
$$\frac{R_{\mathrm{ens}}^\star - \sigma_\star^2}{R_{\mathrm{pool}}^\star - \sigma_\star^2} = \frac{\kappa_1\kappa_2(1-\kappa_1\kappa_2)}{\kappa_0(\kappa_1 + \kappa_2 - 2\kappa_1\kappa_2)} + o_{\mathbb{P}}(1) > 1 + o_{\mathbb{P}}(1).$$

3. *Suppose $K = 2$ and $n_1 = n_2$, and let $\gamma = p/n$ where $n = n_1 + n_2 = 2n_1$. Let $R_{\mathrm{ens},+}$ and $R_{\mathrm{pool},+}$ denote the mean squared prediction errors of the optimally weighted ridgeless ensemble and of pooled ridgeless least squares. Then:*

   (a) *If $\gamma < \frac{1}{2}$, $\dfrac{R_{\mathrm{ens},+}}{R_{\mathrm{pool},+}} = \dfrac{1-\gamma}{1-2\gamma} + o(1) > 1 + o_{\mathbb{P}}(1)$.*

   (b) *If $\gamma > 1$, $\dfrac{R_{\mathrm{ens},+}}{R_{\mathrm{pool},+}} > 1 + o_{\mathbb{P}}(1)$ if and only if $\dfrac{\sigma_\varepsilon^2}{\tau^2} < \dfrac{(2\gamma+1)(2\gamma-1)(\gamma-1)}{4\gamma^2(3\gamma-1)}$.*

The Corollary above verifies that for random effects, the pooled learners perform better than ensembles—both for optimally tuned learners and the ridgeless case. Figure 1 provides an additional validation of the limit in Corollary 2.6 part (3a) in the ridgeless setting. We consider the case $K = 2$ with equal cluster sizes, as in the theoretical setup, and vary the aspect ratio $\gamma = p/n$ in the regime where both the pooled learner and the per-cluster base learners in the ensemble are underparameterized ($0 < \gamma < 0.5$). We find that the simulated MSE ratio closely follows the theoretical limit from Corollary 2.6 (3a).



Confidential reviewer copy. This manuscript is submitted to the 40th Conference on Neural Information Processing Systems (NeurIPS 2026). Unauthorized sharing, redistribution, or disclosure is strictly prohibited.





Figure 1: Ratio of *Ensemble* MSE to *Pooled* MSE for ridgeless learners with $K = 2$; total training sample size $n = 1000$, with $n_1 = n_2 = 500$, and $\gamma = p/n$ in the range $[0, p/n_1]$ so that both methods are underparameterized. For each cluster $\ell$, covariates are drawn from a $\mathbb{N}_p(\mu_\ell, \mathbb{I})$, with $\mu_1 = -\mu_2$. The theoretical limit from Corollary 2.6 (3a) is shown in red, matching our simulated results across all values of $\gamma$.

# 3 Simulations

We conduct a comprehensive simulation study to validate our theoretical results. We describe the common setup, then present results for ridge and ridgeless estimators with a particular focus on the effects of cluster heterogeneity. All experiments were run using 4 CPU cores on a local machine.

## 3.1 Simulation Setup

We simulate $n = 2000$ total training observations partitioned into $K = 2$ equal-sized clusters with $n_1 = n_2 = 1000$. Within each cluster $\ell \in \{1, 2\}$, the covariates are independent draws from a multivariate normal distribution with cluster-specific mean and identity covariance: $\mathbb{X}_\ell \sim \mathbb{N}_p(\mu_\ell, \mathbb{I})$ where $\mu_\ell \in \mathbb{R}^p$, so that all covariates are uncorrelated within and across clusters and have unit marginal variance. The first cluster's mean $\mu_1$ is a unit-norm random vector drawn from $\mathbb{N}_p(0, \mathbb{I})$ and rescaled so that $\|\mu_1\| = 1$; the second cluster's mean $\mu_2$ is constructed via Gram-Schmidt orthogonalization to have a prescribed angle with $\mu_1$ and a prescribed norm (the specific values used are described in Section 3.2). The means $\mu_1$ and $\mu_2$ are redrawn independently in every repetition.

The coefficient vector $\boldsymbol{\beta} \in \mathbb{R}^p$ used to generate the outcome is drawn from $\mathrm{N}_p(0, \mathbb{I})$ and normalized to unit norm ($\|\boldsymbol{\beta}\| = 1$); the same $\boldsymbol{\beta}$ is used to generate training and test outcomes within each repetition. For cluster $\ell \in \{1, 2\}$ and samples $i = 1, \ldots, n_\ell$, training data is generated as $\mathbb{X}_\ell \sim \mathrm{N}_p(\mu_\ell, \mathbb{I})$ with response $Y_\ell = \boldsymbol{\beta}^\top \boldsymbol{X}_\ell + \varepsilon_\ell$, where $\varepsilon_\ell \sim \mathrm{N}_{n_\ell}(0, \sigma^2 \mathbb{I})$ and $\sigma = 1/\sqrt{5}$, giving a signal-to-noise ratio $\mathrm{SNR} = \|\boldsymbol{\beta}\|^2/\sigma^2 = 5$ (matching the highest SNR considered by [Hastie et al., 2022]).

The test data $\boldsymbol{X}^\star \in \mathbb{R}^{n_{\text{test}} \times p}$ is drawn from a standard multivariate normal; $\boldsymbol{X}^\star \sim \mathrm{N}_p(0, \mathbb{I})$, with $n_{\text{test}} = 1000$. The outcome is simulated with no noise, $Y^\star = \boldsymbol{X}^\star \boldsymbol{\beta}$, isolating the prediction risk of each estimator on a fixed reference distribution and matching the theoretical setup of [Hastie et al., 2022]. The covariate dimension is varied across $p \in \{250, 500, 750, 1050, 1500, 2050, 3000, 4000, 5000, 6000\}$, spanning the underparameterized regime, the cluster interpolation threshold at $p = n/K = 1000$, the Pooled interpolation threshold at $p = n = 2000$, and the deep overparameterized regime. We avoid values of $p$ that exactly coincide with these thresholds to prevent numerical instability at the singular boundary. All results are averaged over 100 independent repetitions, with 95% confidence intervals computed from the $t$-distribution.

**Base Learners** We consider ridge and ridgeless base learners as defined in Section 2.1. We compute the ridgeless solution numerically using the SVD-based routine `numpy.linalg.lstsq` with the default rank-cutoff tolerance, which is mathematically equivalent to the pseudoinverse solution but more numerically stable. For the ridge solution, the regularization parameter $\lambda$ is selected by leave-one-out cross-validation over a logarithmic grid of 20 values spanning $[10^{-6}, 10^6]$ using the `RidgeCV` implementation in `scikit-learn` [Pedregosa et al., 2011]; $\lambda$ is tuned independently for every learner.



Confidential reviewer copy. This manuscript is submitted to the 40th Conference on Neural Information Processing Systems (NeurIPS 2026). Unauthorized sharing, redistribution, or disclosure is strictly prohibited.





Figure 2: MSE ratio (*Ensemble* / *Pooled*) in panels (a, b) and raw ensemble and pooled MSEs in panels (c, d); ridge regression in (a, c) and ridgeless regression in (b, d). Training data are drawn from two multivariate normal clusters with means $\mu_1, \mu_2$ and identity covariance under a linear outcome model, with $n_1 = n_2 = 1000$ ($n = 2000$); test points are $n_{\text{test}} = 1000$ draws from a standard multivariate normal. Mean results and 95% CIs are plotted over 100 replicates per configuration. Each panel overlays two scenarios: (i) angle between $\mu_1, \mu_2$ fixed at $60°$, $\|\mu_1\| = 1$, varying $\|\mu_2\|$; (ii) $\|\mu_1\| = \|\mu_2\| = 1$, varying angle. For ridge learners, *Pooled* uniformly outperforms *Ensemble*. For ridgeless learners, behavior is governed by two interpolation peaks (*Ensemble* at $p \approx n_1$, *Pooled* at $p \approx n$); away from these peaks, the two are nearly equivalent. Varying the angle or norms has a negligible effect throughout.

### 3.2 Parameter Configurations

To assess how cluster heterogeneity modulates the ensemble–pooled comparison, we vary two geometric parameters of the cluster means while holding the rest of the setup fixed:

- **Fixed angle, varying norm.** We fix the angle between $\mu_1$ and $\mu_2$ at $60°$ and set $\|\mu_1\| = 1$, $\|\mu_2\| \in \{1, 5, 10\}$. This sweeps the relative signal strength of the two clusters from balanced to highly asymmetric.

- **Fixed norm, varying angle.** We fix $\|\mu_1\| = \|\mu_2\| = 1$ and set the angle between them to $\{30°, 60°, 90°\}$. This sweeps the directional divergence between the two clusters from highly aligned to orthogonal.

We evaluate each parameter configuration for all 10 values of $p$ over 100 repetitions; we plot the MSE ratio of the ensemble to the pooled learner in Figure 2 (a) and (b), and the corresponding raw MSE values for each method in Figure 2 (c) and (d). We performed additional experiments varying the signal-to-noise ratio over the values $\{1, 10\}$, as well as considering fixed, non-zero $\boldsymbol{\beta}$ vectors for generating the outcome, and found that the results did not change; for simplicity, we show only the results corresponding to the specific parameter configurations discussed above.



Confidential reviewer copy. This manuscript is submitted to the 40th Conference on Neural Information Processing Systems (NeurIPS 2026). Unauthorized sharing, redistribution, or disclosure is strictly prohibited.

# 4 Discussion

This paper presents a precise asymptotic comparison between pooled and ensemble learners fit with ridge-regularized linear regression in a proportional asymptotic regime. Our results demonstrate the benefits of pooled methods in random-effects settings and quantify the precise efficiency gains over ensemble learners.

The simulations complement the theory and additionally capture behavior near the interpolation thresholds for ridgeless regression. For the ridgeless estimator, the MSE ratio shows distinct patterns across three regimes. When both the ensemble base learners and the pooled learner are underparameterized ($p < n_\ell$), the advantage of the pooled approach grows with $p$, vanishing only in the fixed-dimensional limit $p \ll n_\ell$ where the two methods perform comparably. At the ensemble's interpolation threshold ($p \approx n_\ell$), its MSE spikes and drives the ratio sharply up. As $p$ moves into the intermediate regime $n_\ell < p < n$, the ratio dips slightly below $1$ and continues to drop as $p$ approaches the pooled learner's interpolation threshold ($p \approx n$); at this point each ensemble base learner is already overparameterized, and the ensemble enjoys its largest advantage; this complements results from Patil et al. [2023]. Notably, the MSE spike of the ensemble is roughly a quarter the size of that for the pooled approach, indicating that the former's worst-case performance is considerably better than the latter's. Once both methods are overparameterized ($p > n$), the ratio approaches $1$ from below as both MSEs converge to the null risk. The picture differs sharply for ridge regression with $\lambda$ tuned optimally per learner: no regime favors the ensemble. The MSE ratio stays between $1.06$ and $1.62$ across all $p$, and the raw MSEs are smooth and monotone in $p$, with no spikes at either interpolation threshold. As in the ridgeless case, the ratio approaches $1$ as $p \gg n$, with both methods converging to the null risk.

Variations in cluster structure have essentially no effect on the comparison. Across all four panels of Figure 2, the curves for the different configurations overlap so tightly that the differences sit below the $95\%$ confidence intervals at almost every $p$. To probe whether this depends on partitioning along the true cluster structure, we ran an ablation in which data were generated under the two-cluster framework of Section 3.1 but randomly split into two halves to train the ensemble base learners. For both ridge and ridgeless regression, the results were identical to Figure 2, even though the random partitions bore no relation to the true clusters. We further considered a fully non-clustered regime aligned with Patil et al. [2023], in which both training and test data were drawn from a $p$-dimensional standard multivariate normal. Splitting the training data into two random equal halves for the ensemble, with the pooled learner trained on the full set, gave the same raw MSEs, and therefore the same MSE ratio, as the clustered experiments. The salient feature of the partition is not its alignment with any cluster structure but simply that it splits the data into two equal halves so that $2 \times p/n_\ell = p/n$. The dominant axis of variation is the aspect ratio; the geometry of the cluster means, and indeed the very presence of covariate heterogeneity, is by comparison irrelevant.

**Limitations and future work.** Our analysis focuses on well-specified linear models with shared coefficients and a shared noise variance across clusters. This is the appropriate setting for isolating pure covariate heterogeneity, but it does not cover model misspecification, nonlinear predictors, or clusters with different noise levels or signal-to-noise ratios. Several extensions are natural. The random matrix arguments developed here extend to joint shifts in mean and variance, beyond the mean shifts we consider. The case of estimated rather than known cluster memberships is open, as is the move beyond linear regression, which can follow the template of Hastie et al. [2019], where results were first established under linear models before being extended to general nonlinear settings. Theoretical analysis of the ridgeless estimator near the interpolation peaks under covariate shift is an additional future direction. Finally, when cluster effects are fixed rather than random, the optimal choice between pooled and ensemble strategies depends on unknown cluster parameters, making adaptive selection an interesting open question.



Confidential reviewer copy. This manuscript is submitted to the 40th Conference on Neural Information Processing Systems (NeurIPS 2026). Unauthorized sharing, redistribution, or disclosure is strictly prohibited.

# Bibliography


Tavor Z. Baharav, Phillip B. Nicol, Rafael A. Irizarry, and Rong Ma. Stacked SVD or SVD stacked? A random matrix theory perspective on data integration. *arXiv preprint arXiv:2507.22170*, 2025. URL https://arxiv.org/abs/2507.22170.

Zhidong Bai, Jack W Silverstein, et al. *Spectral analysis of large dimensional random matrices*, volume 20. Springer, 2010.

Peter L. Bartlett, Philip M. Long, Gábor Lugosi, and Alexander Tsigler. Benign overfitting in linear regression. *Proceedings of the National Academy of Sciences*, 117(48):30063–30070, 2020. doi: 10.1073/pnas.1907378117.

Mikhail Belkin, Daniel Hsu, Siyuan Ma, and Soumik Mandal. Reconciling modern machine-learning practice and the classical bias–variance trade-off. *Proceedings of the National Academy of Sciences*, 116(32):15849–15854, 2019. doi: 10.1073/pnas.1903070116.

Shai Ben-David, John Blitzer, Koby Crammer, Alex Kulesza, Fernando Pereira, and Jennifer Wortman Vaughan. A theory of learning from different domains. *Machine learning*, 79(1):151–175, 2010.

Ritu Chauhan, Harleen Kaur, and M Afshar Alam. Data clustering method for discovering clusters in spatial cancer databases. *International Journal of Computer Applications*, 10(6):9–14, 2010.

Koby Crammer, Michael Kearns, and Jennifer Wortman. Learning from multiple sources. *Journal of machine learning research*, 9(8), 2008.

Meghana Deodhar and Joydeep Ghosh. A framework for simultaneous co-clustering and learning from complex data. In *Proceedings of the 13th ACM SIGKDD International Conference on Knowledge Discovery and Data Mining*, KDD '07, page 250–259, New York, NY, USA, 2007. Association for Computing Machinery. ISBN 9781595936097. doi: 10.1145/1281192.1281222. URL https://doi.org/10.1145/1281192.1281222.

Edgar Dobriban and Stefan Wager. High-dimensional asymptotics of prediction: Ridge regression and classification. *The Annals of Statistics*, 46(1):247–279, 2018.

Wilson Wen Bin Goh, Wei Wang, and Limsoon Wong. Why batch effects matter in omics data, and how to avoid them. *Trends in Biotechnology*, 35(6):498–507, 2017.

Zoe Guan, Giovanni Parmigiani, and Prasad Patil. Merging versus ensembling in multi-study machine learning: Theoretical insight from random effects. *arXiv*, 2019.

Trevor Hastie, Andrea Montanari, Saharon Rosset, and Ryan J Tibshirani. Surprises in high-dimensional ridgeless least squares interpolation. *arXiv preprint arXiv:1903.08560*, 2019.

Trevor Hastie, Andrea Montanari, Saharon Rosset, and Ryan J. Tibshirani. Surprises in high-dimensional ridgeless least squares interpolation. *Annals of Statistics*, 50(2):949–986, 2022. doi: 10.1214/21-AOS2133.

Antti Knowles and Jun Yin. Anisotropic local laws for random matrices. *Probability Theory and Related Fields*, 169(1):257–352, 2017.

Daniel LeJeune, Hamid Javadi, and Richard Baraniuk. The implicit regularization of ordinary least squares ensembles. In *International Conference on Artificial Intelligence and Statistics*, pages 3525–3535. PMLR, 2020.

Sai Li, T Tony Cai, and Hongzhe Li. Transfer learning for high-dimensional linear regression: Prediction, estimation and minimax optimality. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 84(1):149–173, 2022.

Yishay Mansour, Mehryar Mohri, and Afshin Rostamizadeh. Domain adaptation with multiple sources. *Advances in neural information processing systems*, 21, 2008.




Confidential reviewer copy. This manuscript is submitted to the 40th Conference on Neural Information Processing Systems (NeurIPS 2026). Unauthorized sharing, redistribution, or disclosure is strictly prohibited.

Nicolai Meinshausen and Peter Bühlmann. Maximin effects in inhomogeneous large-scale data. *Annals of Statistics*, 43(4):1801–1830, 2015.

Prasad Patil and Giovanni Parmigiani. Training replicable predictors in multiple studies. *Proceedings of the National Academy of Sciences*, 115(11):2578–2583, 2018. ISSN 0027-8424. doi: 10.1073/pnas.1708283115. URL https://www.pnas.org/content/115/11/2578.

Pratik Patil, Jin-Hong Du, and Arun Kumar Kuchibhotla. Bagging in overparameterized learning: Risk characterization and risk monotonization. *Journal of Machine Learning Research*, 24(319): 1–113, 2023.

Fabian Pedregosa, Gaël Varoquaux, Alexandre Gramfort, Vincent Michel, Bertrand Thirion, Olivier Grisel, Mathieu Blondel, Peter Prettenhofer, Ron Weiss, Vincent Dubourg, et al. Scikit-learn: Machine learning in python. *the Journal of machine Learning research*, 12:2825–2830, 2011.

Maya Ramchandran and Rajarshi Mukherjee. On ensembling vs merging: Least squares and random forests under covariate shift. *arXiv preprint arXiv:2106.02589*, 2021.

Maya Ramchandran, Prasad Patil, and Giovanni Parmigiani. Tree-weighting for multi-study ensemble learners. *Pacific Symposium on Biocomputing*, 25:451–462, 2020. doi: https://doi.org/10.1101/698779.

Shubhendu Trivedi, Zachary A. Pardos, and Neil T. Heffernan. The utility of clustering in prediction tasks. *CoRR*, abs/1509.06163, 2015. URL http://arxiv.org/abs/1509.06163.

Yuchen Zhang, John Duchi, and Martin Wainwright. Divide and conquer kernel ridge regression: A distributed algorithm with minimax optimal rates. *The Journal of Machine Learning Research*, 16(1):3299–3340, 2015.

# A Appendix

## A.1 Further Related Work

**An unsupervised analog: stacking versus per-source SVD.** Baharav et al. [2025] study a closely related question in an unsupervised setting: given multiple datasets that share a common right singular subspace but differ in signal strength and noise level, they investigate whether it is better to stack all datasets and run a single SVD (*stacked SVD*) or to run SVD on each dataset separately and aggregate the per-source estimates (*stacked-SVDs*). Their main finding is that the *stacked SVD* approach dominates when dataset-specific weights are chosen optimally (with optimal weights downweighting noisier or lower-signal sources), but that without optimal weighting, neither approach dominates universally and per-source aggregation can occasionally win. Their setting differs from ours in the source of heterogeneity: all of their datasets share the same isotropic noise structure and signal direction but differ in signal-to-noise ratio, whereas in our supervised setting all clusters share the same signal and noise level but differ in their covariate distributions. Despite this difference, the two settings raise the same fundamental question: when multiple data sources differ in some structural way, is it optimal to combine them before fitting or to fit separately and aggregate?

## A.2 Background Results

Throughout we repeatedly use the following anisotropic local laws [Knowles and Yin, 2017].

**Proposition A.1** ([Knowles and Yin, 2017])**.** *Consider $n$ i.i.d. vectors $\boldsymbol{X}_i = \Sigma^{1/2}\mathbf{Z}_i$ with $\mathbf{Z}_i = (Z_{ij})_{j=1}^p$ i.i.d. from some distribution with mean $0$, variance $1$, and sub-gaussian tails. With $\hat{\Sigma} = \frac{1}{n}\sum_{i=1}^n \boldsymbol{X}_i\boldsymbol{X}_i^\top$ and constants $\alpha_0, \alpha_1 > 0$ define the event*

$$\Omega_n(\alpha_0, \alpha_1) = \left\{ \left| v^\top (\hat{\Sigma} + \lambda\mathbb{I})^{-1} v - v^\top \left(\mathbb{I} + \tilde{m}_n(-\lambda)\Sigma\right)^{-1} v \right| \leq \frac{1}{n^{\frac{1}{2}-\alpha_0}\lambda} \right\}$$



Confidential reviewer copy. This manuscript is submitted to the 40th Conference on Neural Information Processing Systems (NeurIPS 2026). Unauthorized sharing, redistribution, or disclosure is strictly prohibited.

*for all $\lambda \geq n^{-\frac{2}{3}+\alpha_1}$. Here $\tilde{m}_n(z)$ is the Stieltjes transform of the companion of the Marchenko-Pasteur law, defined as the solution of the following fixed-point equation:*

$$\frac{1}{\tilde{m}_n(z)} = -z + \gamma \frac{1}{p} \sum_{j=1}^{p} \frac{s_j(\Sigma)}{1 + s_j(\Sigma)\tilde{m}_n(z)} = -z + \gamma \int \frac{x}{1 + x\tilde{m}_n(z)} \, dF_{\Sigma,n}(x)\,, \tag{1}$$

*where $\{s_j(\Sigma)\}_{j\in[p]}$ are eigenvalues of $\Sigma$ and $\gamma = \lim \frac{p}{n}$. Then given any large $D > 0$ and small $\alpha_0, \alpha_1 > 0$, there exists $N_0 \geq 1$, such that $\mathbb{P}(\Omega_n^c(\alpha_0, \alpha_1)) \leq n^{-D}$ for $n \geq N_0$.*

### A.3 Proof of Main Results

Throughout we will assume $\tau^2 = \sigma_\varepsilon^2 = 1$. The general case follows verbatim through scaling.

#### A.3.1 Proof of Theorem 2.5

Recall the notation and definitions as follows.

$\hat{\Sigma}_\ell = \frac{1}{n_\ell}\mathbb{X}_\ell^\top \mathbb{X}_\ell$ and $\hat{\Sigma}_{\text{pool}} = \frac{1}{n}\mathbb{X}_{\text{pool}}^\top \mathbb{X}_{\text{pool}}$ where $n = \sum_{\ell=1}^{K} n_\ell$. Then define ridge regression as:

$$\hat{f}_\ell(\mathbf{x}) = \hat{\boldsymbol{\beta}}_{\lambda_\ell}^\top \mathbf{x}, \quad \hat{\boldsymbol{\beta}}_{\lambda_\ell} = (\mathbb{X}_\ell^\top \mathbb{X}_\ell + \lambda_\ell \mathbb{I})^{-1} \mathbb{X}_\ell^\top \mathbf{Y}_\ell, \quad \lambda_\ell > 0;$$
$$\hat{f}_{\text{pool}}(\mathbf{x}) = \hat{\boldsymbol{\beta}}_{\lambda}^\top \mathbf{x}, \quad \hat{\boldsymbol{\beta}}_{\lambda} = (\mathbb{X}_{\text{pool}}^\top \mathbb{X}_{\text{pool}} + \lambda \mathbb{I})^{-1} \mathbb{X}_{\text{pool}}^\top \mathbf{Y}_{\text{pool}}, \quad \lambda > 0.$$

Define ridge-less regression as:

$$\hat{f}_{\ell,+}(\mathbf{x}) = \hat{\boldsymbol{\beta}}_{+,\ell}^\top \mathbf{x}, \quad \hat{\boldsymbol{\beta}}_{+,\ell} = (\mathbb{X}_\ell^\top \mathbb{X}_\ell)^+ \mathbb{X}_\ell^\top \mathbf{Y}_\ell;$$
$$\hat{f}_{\text{pool},+}(\mathbf{x}) = \hat{\boldsymbol{\beta}}_{+}^\top \mathbf{x}, \quad \hat{\boldsymbol{\beta}}_{+} = (\mathbb{X}_{\text{pool}}^\top \mathbb{X}_{\text{pool}})^+ \mathbb{X}_{\text{pool}}^\top \mathbf{Y}_{\text{pool}}.$$

We first analyze $\hat{f}_{\text{pool}}(\mathbf{x})$. We divide the analysis into two cases and sub-cases. The first case pertains to the random effects analysis—further divided into fixed-$\rho$ analysis followed by the ridgeless. We subsequently follow the same template for the fixed effects analysis, i.e., fixed-$\rho$ analysis followed by the ridgeless. Each analysis follows from an initial template decomposition of the risk. Specifically, for any estimator $\hat{\boldsymbol{\beta}}$ of $\boldsymbol{\beta}$ the following holds under Assumption 2.1:

$$\mathbb{E}(y^\star - \hat{\boldsymbol{\beta}}^\top \mathbf{x}^\star)^2 = \sigma_\star^2 + \mathbb{E}\|\hat{\boldsymbol{\beta}} - \boldsymbol{\beta}\|^2.$$

Further, each $\hat{\boldsymbol{\beta}}$ in our analyses is a linear function of the outcome and can be written as $\mathbf{L}\mathbf{Y}_{\text{pool}}$ for some matrix $\mathbf{L}$ that is a measurable function of $\mathbb{X}_{\text{pool}}$. Therefore (using the standing convention $\sigma_\varepsilon^2 = 1$)

$$\mathbb{E}\|\hat{\boldsymbol{\beta}} - \boldsymbol{\beta}\|^2 = \mathbb{E}\|(L\mathbb{X}_{\text{pool}} - \mathbb{I})\boldsymbol{\beta}\|^2 + \text{Tr}(\mathbf{L}\mathbf{L}^\top).$$

The difference in analysis for fixed and random effects pertains to evaluating $\mathbb{E}\|(L\mathbb{X}_{\text{pool}} - \mathbb{I})\boldsymbol{\beta}\|^2$: for random effects this marginalizes over the distribution of $\boldsymbol{\beta}$, whereas in the fixed effects case it is evaluated at a given $\boldsymbol{\beta}$.

**Analysis of Pooled Ridge-Regularized Estimator – Random Effects:** Here $\mathbf{L} = (\mathbb{X}_{\text{pool}}^\top \mathbb{X}_{\text{pool}} + \lambda\mathbb{I})^{-1}\mathbb{X}_{\text{pool}}^\top$. Moreover, since $\mathbb{E}(\boldsymbol{\beta}) = 0$ and $\text{Var}(\boldsymbol{\beta}) = \mathbb{I}/p$ (using $\tau^2 = 1$), we have by direct calculations that

$$\begin{aligned}\mathbb{E}\|(L\mathbb{X}_{\text{pool}} - \mathbb{I})\boldsymbol{\beta}\|^2 &= \frac{1}{p}\text{Tr}((L\mathbb{X}_{\text{pool}} - \mathbb{I})^\top (L\mathbb{X}_{\text{pool}} - \mathbb{I})) \\ &= \rho^2 \frac{1}{p}\text{Tr}(G_{\text{pool}}^2(-\rho)),\end{aligned}$$

where $\rho = \lambda/n$ and $G_{\text{pool}}(z) = (\hat{\Sigma}_{\text{pool}} - z\mathbb{I})^{-1}$ for $z \notin \text{Spectrum}(\hat{\Sigma}_{\text{pool}})$. Similarly,

$$\text{Tr}(\mathbf{L}\mathbf{L}^\top) = \frac{\gamma_0}{p}\mathbb{E}\left(\text{Tr}(G_{\text{pool}}(-\rho) - \rho G_{\text{pool}}^2(-\rho))\right).$$



Confidential reviewer copy. This manuscript is submitted to the 40th Conference on Neural Information Processing Systems (NeurIPS 2026). Unauthorized sharing, redistribution, or disclosure is strictly prohibited.

Next we analyze we analyze $\frac{1}{p}\mathrm{Tr}(G^2_{\mathrm{pool}}(-\rho))$. To this end note that, we can write $\mathbb{X}_{\mathrm{pool}} = \mathbb{Z}_{\mathrm{pool}}\Sigma^{1/2} + M$ where $\mathrm{rank}(M) \leq K$. Specifically $M = DU$ where $D \in \mathbb{R}^{n\times K}$ with $\ell^{\mathrm{th}}$ column being a vector of 1's of length $n_\ell$ starting at $\sum_{j=1}^{\ell-1} n_j + 1$ and rest 0's, and $U = [\boldsymbol{\mu}_1, \ldots, \boldsymbol{\mu}_K]^\top \in \mathbb{R}^{K\times p}$. In this notation, we can write

$$\hat{\Sigma}_{\mathrm{pool}} = \hat{\Sigma}_{\mathrm{pool},0} + \frac{1}{n}\mathbb{Z}^T_{\mathrm{pool},0}M + \frac{1}{n}M^\top\mathbb{Z}_{\mathrm{pool},0} + \frac{1}{n}M^\top M$$

where $\mathbb{Z}_0 = \mathbb{Z}_{\mathrm{pool}}\Sigma^{1/2}$ and $\hat{\Sigma}_{\mathrm{pool},0} = \frac{1}{n}\mathbb{Z}_0^\top\mathbb{Z}_0$. Therefore $\mathrm{rank}(\hat{\Sigma}_{\mathrm{pool}} - \hat{\Sigma}_{\mathrm{pool},0}) \leq 3K$. Now let $\hat{F}_{\mathrm{pool}}$ and $\hat{F}_{\mathrm{pool},0}$ denote the empirical spectral distributions of $\hat{\Sigma}_{\mathrm{pool}}$, and $\hat{\Sigma}_{\mathrm{pool},0}$ respectively. Then [Bai et al., 2010, Theorem A.43] one has that

$$\sup_{x\in\mathbb{R}_+} |\hat{F}_{\mathrm{pool}}(x) - \hat{F}_{\mathrm{pool},0}(x)| \leq \frac{3K}{p} \to 0.$$

Next note that, for $t \in \mathbb{N}$, the function $f_{t,\rho}(x) = \frac{1}{(x+\rho)^t}$ is bounded and continuous on $\mathbb{R}_+$ for any fixed $\rho > 0$, with $\|f_{t,\rho}\|_\infty \leq \frac{1}{\rho^t}$. Therefore,

$$\frac{1}{p}\mathrm{Tr}(G^t_{\mathrm{pool},0}(-\rho)) \xrightarrow{\mathbb{P}} \int f_{t,\rho}(x)dF_{\gamma_0,H}$$

where $H$ is the limiting spectral distribution of $\Sigma$ and $F_{\gamma_0,H}$ is the generalized Marchenko-Pastur distribution [Bai et al., 2010]. However,

$$\left|\tfrac{1}{p}\mathrm{Tr}(G^t_{\mathrm{pool},0}(-\rho)) - \tfrac{1}{p}\mathrm{Tr}(G^t_{\mathrm{pool}}(-\rho))\right| = \left|\int f_{t,\rho}(x)d\hat{F}_{\mathrm{pool}}(x) - \int f_{t,\rho}(x)d\hat{F}_{\mathrm{pool},0}(x)\right| \leq \tfrac{1}{\rho^t}\cdot\tfrac{3K}{p} \to 0.$$

Hence,

$$\frac{1}{p}\mathrm{Tr}(G^t_{\mathrm{pool}}(-\rho)) \xrightarrow{\mathbb{P}} \int f_{t,\rho}(x)dF_{\gamma_0,H}(x).$$

Therefore

$$\mathbb{E}\|\hat{\boldsymbol{\beta}}_\lambda - \boldsymbol{\beta}\|^2 \xrightarrow{\mathbb{P}} \rho^2\int f_{2,\rho}(x)dF_{\gamma_0,H}(x) + \gamma_0\left[\int f_{1,\rho}(x)dF_{\gamma_0,H}(x) - \rho\int f_{2,\rho}(x)dF_{\gamma_0,H}(x)\right].$$

Now in terms of the notation of the theorem, it follows by direct calculations that

$$\int f_{1,\rho}(x)dF_{\gamma_0,H}(x) = a_\gamma(\rho);$$
$$\int f_{2,\rho}(x)dF_{\gamma_0,H}(x) = -a_\gamma^{'}(\rho).$$

This completes the proof for the pooled ridge regression when $\rho > 0$.

**Analysis of Pooled Ridgeless Estimator – Random Effects:** In this case, we start by noting that

$$\mathrm{Tr}(\mathbf{L}\mathbf{L}^\top) = \tfrac{1}{n}\mathrm{Tr}(\widehat{\Sigma}^+_{\mathrm{pool}});$$
$$\mathbb{E}\|(L\mathbb{X}_{\mathrm{pool}} - \mathbb{I})\boldsymbol{\beta}\|^2 = \frac{1}{p}\mathrm{Tr}(\mathbb{I} - P_{\mathcal{C}(\mathbb{X}^\top_{\mathrm{pool}})}),$$

where for any matrix $F$, $P_{\mathcal{C}(F)}$ denotes the orthogonal projection onto the column space $\mathcal{C}(F)$ of $F$ (so $P_{\mathcal{C}(\mathbb{X}^\top_{\mathrm{pool}})}$ is the projection onto the row space of $\mathbb{X}_{\mathrm{pool}}$). Subsequently we divide the analysis into two parts, depending on whether $\gamma_0$ is smaller or larger than 1, i.e., the under- versus overparameterized regimes.

**Underparameterized Regime - $\gamma_0 < 1$:** We borrow the notation of the subsection and note that in this regime $\hat{\Sigma}^+_{\mathrm{pool}} = \hat{\Sigma}^{-1}_{\mathrm{pool}}$ (henceforth we work on the event that $\hat{\Sigma}_{\mathrm{pool}}$ is invertible, which occurs with probability converging to 1) and $\hat{\Sigma}^+_{\mathrm{pool},0} = \hat{\Sigma}^{-1}_{\mathrm{pool},0}$. Subsequently, using the inverse identity that $A^{-1} - B^{-1} = -A^{-1}(B-A)B^{-1}$ we get that

$$\hat{\Sigma}^{-1}_{\mathrm{pool}} - \hat{\Sigma}^{-1}_{\mathrm{pool},0} = -\hat{\Sigma}^{-1}_{\mathrm{pool}}\left(\frac{1}{n}\mathbb{Z}^T_{\mathrm{pool},0}M + \frac{1}{n}M^\top\mathbb{Z}_{\mathrm{pool},0} + \frac{1}{n}M^\top M\right)\hat{\Sigma}^{-1}_{\mathrm{pool},0}.$$



Confidential reviewer copy. This manuscript is submitted to the 40th Conference on Neural Information Processing Systems (NeurIPS 2026). Unauthorized sharing, redistribution, or disclosure is strictly prohibited.

Therefore

$$\mathrm{rank}(\hat{\Sigma}_{\mathrm{pool}}^{-1} - \hat{\Sigma}_{\mathrm{pool},0}^{-1}) \leq 3K.$$

Moreover, there exists $c_{\gamma_0} < \infty$ such that with probability converging to 1 [Bai et al., 2010] one has that

$$\|\hat{\Sigma}_{\mathrm{pool}}^{-1} - \hat{\Sigma}_{\mathrm{pool},0}^{-1}\|_{\mathrm{op}} \leq c_{\gamma_0}.$$

Therefore with probability converging to 1

$$|\frac{1}{p}\mathrm{Tr}(\widehat{\Sigma}_{\mathrm{pool}}^{+}) - \frac{1}{p}\mathrm{Tr}(\widehat{\Sigma}_{\mathrm{pool},0}^{+})| \leq \frac{1}{p}\mathrm{rank}(\hat{\Sigma}_{\mathrm{pool}}^{-1} - \hat{\Sigma}_{\mathrm{pool},0}^{-1})\|\hat{\Sigma}_{\mathrm{pool}}^{-1} - \hat{\Sigma}_{\mathrm{pool},0}^{-1}\|_{\mathrm{op}} \leq \frac{3Kc_{\gamma_0}}{p} \to 0.$$

Moreover, in this regime $\frac{1}{p}\mathrm{Tr}(\mathbb{I} - P_{\mathcal{C}(\mathbb{X}_{\mathrm{pool}}^{\top})}) = 0$, so the bias contribution vanishes. Furthermore, in this regime [Hastie et al., 2019],

$$\frac{1}{n}\mathrm{Tr}(\widehat{\Sigma}_{\mathrm{pool},0}^{+}) \xrightarrow{\mathbb{P}} \gamma_0 \int_{x>0} \frac{1}{x} dF_{\gamma_0,H}(x).$$

This implies that

$$\mathbb{E}\|\hat{\boldsymbol{\beta}} - \boldsymbol{\beta}\|^2 \xrightarrow{\mathbb{P}} \gamma_0 \int_{x>0} \frac{1}{x} dF_{\gamma_0,H}(x) = \lim_{\rho_0 \downarrow 0} \left(\rho_0^2[-a'_{\gamma_0}(\rho_0)] + V_{\gamma_0}(\rho_0)\right),$$

as promised.

**Overparameterized Regime -** $\gamma_0 > 1$**:** Let $\Gamma = \frac{1}{n}\mathbb{X}_{\mathrm{pool}}\mathbb{X}_{\mathrm{pool}}^{\top}$ and $\Gamma_0 = \frac{1}{n}\mathbb{Z}_0\mathbb{Z}_0^{\top}$ where $\mathbb{Z}_0 = \mathbb{Z}_{\mathrm{pool}}\Sigma^{1/2}$. Now note that

$$\mathrm{Tr}(\hat{\Sigma}_{\mathrm{pool}}^{+}) = \mathrm{Tr}(\Gamma^{-1}), \quad \mathrm{Tr}(\hat{\Sigma}_{\mathrm{pool},0}^{+}) = \mathrm{Tr}(\Gamma_0^{-1}).$$

Arguing similarly to before,

$$\mathrm{rank}(\Gamma - \Gamma_0) \leq 3K,$$

and there exists $c_{\gamma_0} < \infty$ such that with probability converging to 1

$$\|\Gamma\|_{\mathrm{op}} \vee \|\Gamma_0\|_{\mathrm{op}} \leq c_{\gamma_0}.$$

Therefore

$$|\frac{1}{p}\mathrm{Tr}(\hat{\Sigma}_{\mathrm{pool}}^{+}) - \frac{1}{p}\mathrm{Tr}(\hat{\Sigma}_{\mathrm{pool},0}^{+})| \leq 6Kc_{\gamma_0}/p \to 0.$$

Therefore

$$\frac{1}{p}\mathrm{Tr}(\hat{\Sigma}_{\mathrm{pool}}^{+}) \xrightarrow{\mathbb{P}} \gamma_0 \int_{x>0} \frac{1}{x} dF_{\gamma_0,H}(x),$$

as before. Moreover it is easy to check by simple rank calculations that

$$\left|\tfrac{1}{p}\mathrm{Tr}(\mathbb{I} - P_{\mathcal{C}(\mathbb{X}_{\mathrm{pool}}^{\top})}) - \tfrac{1}{p}\mathrm{Tr}(\mathbb{I} - P_{\mathcal{C}(\mathbb{Z}_0^{\top})})\right| \leq K/p \to 0.$$

But $\frac{1}{p}\mathrm{Tr}(\mathbb{I} - P_{\mathcal{C}(\mathbb{Z}_0^{\top})}) = 1 - n/p \to 1 - 1/\gamma_0$. Therefore as before

$$\mathbb{E}\|\hat{\boldsymbol{\beta}} - \boldsymbol{\beta}\|^2 \xrightarrow{\mathbb{P}} \gamma_0 \int_{x>0} \frac{1}{x} dF_{\gamma_0,H}(x) = \lim_{\rho_0 \downarrow 0} \left(\rho_0^2[-a'_{\gamma_0}(\rho_0)] + V_{\gamma_0}(\rho_0)\right),$$

as promised.



Confidential reviewer copy. This manuscript is submitted to the 40th Conference on Neural Information Processing Systems (NeurIPS 2026). Unauthorized sharing, redistribution, or disclosure is strictly prohibited.

**Analysis of Pooled Ridge-Regularized Estimator – Fixed Effects:** First note that by direct calculations

$$\mathbb{E}\|\hat{\boldsymbol{\beta}}_\lambda - \boldsymbol{\beta}\|^2 = \rho_0^2\, \boldsymbol{\beta}^\top G_{\text{pool}}^2(-\rho_0)\boldsymbol{\beta} + \frac{1}{n}\text{Tr}\Big(\hat{\Sigma}_{\text{pool}}\, G_{\text{pool}}^2(-\rho_0)\Big) = B_n + V_n \quad \text{(say)}.$$

The analysis of $V_n \xrightarrow{\mathbb{P}} V_{\gamma_0}(\rho_0)$ does not depend on $\boldsymbol{\beta}$ and is identical to the random-effects case; we therefore focus on the bias $B_n$ below. Write as before

$$\mathbb{X}_{\text{pool}} = \mathbb{Z}_0 + M,$$

For each cluster define

$$\bar{z}_\ell = \frac{1}{n_\ell}\sum_{i=1}^{n_\ell} \boldsymbol{Z}_{i,\ell}, \qquad \widetilde{Z}_{i,\ell} = \boldsymbol{Z}_{i,\ell} - \bar{z}_\ell.$$

Let

$$S_z = \frac{1}{n}\sum_{\ell=1}^{K}\sum_{i=1}^{n_\ell} \Sigma^{1/2}\boldsymbol{Z}_{i,\ell}\boldsymbol{Z}_{i,\ell}^\top\Sigma^{1/2},$$

and

$$S_n^{(-)} = \frac{1}{n}\sum_{\ell=1}^{K}\sum_{i=1}^{n_\ell} \Sigma^{1/2}\widetilde{\boldsymbol{Z}}_{i,\ell}\widetilde{\boldsymbol{Z}}_{i,\ell}^\top\Sigma^{1/2}.$$

Recall

$$\pi_{\ell,n} = \frac{n_\ell}{n}, \qquad U_n = [\sqrt{\pi_{1,n}}\mu_1, \ldots, \sqrt{\pi_{K,n}}\mu_K],$$

and define

$$R_n = [\sqrt{\pi_{1,n}}\Sigma^{1/2}\bar{z}_1, \ldots, \sqrt{\pi_{K,n}}\Sigma^{1/2}\bar{z}_K].$$

Then

$$S_z = S_n^{(-)} + R_n R_n^\top.$$

Also, since inside cluster $\ell$,

$$\boldsymbol{X}_{i,\ell} = \boldsymbol{\mu}_\ell + \Sigma^{1/2}\bar{z}_\ell + \Sigma^{1/2}\widetilde{\boldsymbol{Z}}_{i,\ell},$$

and

$$\sum_{i=1}^{n_\ell}\widetilde{Z}_{i,\ell} = 0,$$

we get the exact decomposition

$$\widehat{\Sigma}_{\text{pool}} = S_n^{(-)} + (U_n + R_n)(U_n + R_n)^\top.$$

Now for $\rho > 0$, define the two resolvents

$$H_n(\rho) = (S_z + \rho\mathbb{I})^{-1}, \qquad G_n^{(-)}(\rho) = (S_n^{(-)} + \rho\mathbb{I})^{-1}.$$

**Lemma A.2.** *Fix $\rho$ in a compact subset of $(0,\infty)$. Then, locally uniformly in $\rho$,*

$$R_n^\top H_n(\rho) R_n \xrightarrow{\mathbb{P}} \frac{\gamma_0 s_{\gamma_0}(\rho)}{1+\gamma_0 s_{\gamma_0}(\rho)}\mathbb{I}_K,$$

*where $s_\gamma(\rho) = \frac{1}{p}\text{Tr}\{\Sigma Q_\gamma(\rho)\}$, and $\mathbb{I}_K$ is the identity matrix of order $K$. Moreover, for every deterministic $v \in \mathbb{R}^p$ with uniformly bounded norm,*

$$v^\top H_n(\rho) R_n \xrightarrow{\mathbb{P}} 0, \qquad R_n^\top H_n(\rho) v \xrightarrow{\mathbb{P}} 0.$$

*Proof.* Fix a cluster $\ell$, write $m = n_\ell$, and suppress the cluster index and dependence on resolvents on $\rho$ whenever clear from context. Let

$$\bar{z} = \frac{1}{m}\sum_{i=1}^{m} z_i, \qquad r_{\ell,n} = \sqrt{\pi_{\ell,n}}\Sigma^{1/2}\bar{z}.$$

Then

$$r_{\ell,n}^\top H_n r_{\ell,n} = \pi_{\ell,n}\bar{z}^\top\Sigma^{1/2}H_n\Sigma^{1/2}\bar{z}.$$



Confidential reviewer copy. This manuscript is submitted to the 40th Conference on Neural Information Processing Systems (NeurIPS 2026). Unauthorized sharing, redistribution, or disclosure is strictly prohibited.

Therefore

$$\bar{z}^\top \Sigma^{1/2} H_n \Sigma^{1/2} \bar{z} = \frac{1}{m^2} \sum_{i,j=1}^{m} \boldsymbol{Z}_i^\top \Sigma^{1/2} H_n \Sigma^{1/2} \boldsymbol{Z}_j.$$

We first treat the diagonal terms i.e. $i = j$. Let $H_n^{(-i)}$ be the resolvent with the row $\boldsymbol{Z}_i$ removed from $S_z$. That is,

$$H_n^{(-i)} = \left( S_z - \frac{1}{n} \Sigma^{1/2} \boldsymbol{Z}_i \boldsymbol{Z}_i^\top \Sigma^{1/2} + \rho \mathbb{I} \right)^{-1}.$$

Now let

$$B_i = \Sigma^{1/2} H_n^{(-i)} \Sigma^{1/2}, \qquad q_i = \boldsymbol{Z}_i^\top B_i \boldsymbol{Z}_i.$$

Then by Sherman-Morrison formula,

$$H_n = H_n^{(-i)} - \frac{\frac{1}{n} H_n^{(-i)} \Sigma^{1/2} \boldsymbol{Z}_i \boldsymbol{Z}_i^\top \Sigma^{1/2} H_n^{(-i)}}{1 + \frac{1}{n} \boldsymbol{Z}_i^\top B_i \boldsymbol{Z}_i}.$$

Therefore

$$\boldsymbol{Z}_i^\top \Sigma^{1/2} H_n \Sigma^{1/2} \boldsymbol{Z}_i = \frac{q_i}{1 + q_i/n}.$$

now note that $H_n^{(-i)}$ is independent of $z_i$ and there exists a constant $C_\rho < \infty$ such that with probability converging to 1 [Bai et al., 2010]

$$\|H_n^{(-i)}\|_{\mathrm{op}} \le \rho^{-1}, \qquad \|B_i\|_{\mathrm{op}} \le C_\rho,$$

Therefore by Hanson–Wright inequality we have by union bound that uniformly over $i$

$$q_i = \mathrm{Tr} B_i + O_{\mathbb{P}}(\sqrt{p \log n}).$$

Also, removing one row changes normalized traces by $o_{\mathbb{P}}(1)$ uniformly over rows [Bai et al., 2010, Theorem A.43]. Therefore uniformly over $i$

$$\frac{1}{p} \mathrm{Tr} B_i = \frac{1}{p} \mathrm{Tr}\{\Sigma^{1/2} H_n^{(-i)} \Sigma^{1/2}\} \xrightarrow{\mathbb{P}} s_{\gamma_0}(\rho).$$

Consequently,

$$\frac{1}{m^2} \sum_{i=1}^{m} \boldsymbol{Z}_i^\top \Sigma^{1/2} H_n \Sigma^{1/2} \boldsymbol{Z}_i = \frac{1}{m^2} \sum_{i=1}^{m} \frac{q_i}{1 + q_i/n}$$

satisfies

$$\frac{1}{m^2} \sum_{i=1}^{m} \frac{q_i}{1 + q_i/n} - \frac{1}{m} \cdot \frac{p\, s_{\gamma_0}(\rho)}{1 + \gamma_0 s_{\gamma_0}(\rho)} \xrightarrow{\mathbb{P}} 0.$$

Since

$$\frac{p}{m} = \frac{p/n}{n_\ell/n} \to \frac{\gamma_0}{\pi_\ell},$$

we obtain

$$\frac{1}{m^2} \sum_{i=1}^{m} \boldsymbol{Z}_i^\top \Sigma^{1/2} H_n \Sigma^{1/2} \boldsymbol{Z}_i \xrightarrow{\mathbb{P}} \frac{\gamma_0}{\pi_\ell} \frac{s_{\gamma_0}(\rho)}{1 + \gamma_0 s_{\gamma_0}(\rho)}.$$

Next, we consider the off-diagonal terms $i \neq j$. Here we employ a leave-two-out technique as follows. To streamline notation, set $\xi_i := \Sigma^{1/2} \boldsymbol{Z}_i$ and write $R_{ij} := H_n^{(-ij)}$ for the leave-two-out resolvent introduced below. We start by leaving out the index $i$ and write, using the Sherman–Morrison formula,

$$\boldsymbol{Z}_i^\top \Sigma^{1/2} H_n \Sigma^{1/2} \boldsymbol{Z}_j = \frac{\boldsymbol{Z}_i^\top \Sigma^{1/2} H_n^{(-i)} \Sigma^{1/2} \boldsymbol{Z}_j}{1 + \frac{1}{n} \boldsymbol{Z}_i^\top \Sigma^{1/2} H_n^{(-i)} \Sigma^{1/2} \boldsymbol{Z}_i}.$$

Applying the Sherman–Morrison formula once more to leave out the index $j$ gives

$$\boldsymbol{Z}_i^\top \Sigma^{1/2} H_n^{(-i)} \Sigma^{1/2} \boldsymbol{Z}_j = \frac{\boldsymbol{Z}_i^\top \Sigma^{1/2} H_n^{(-ij)} \Sigma^{1/2} \boldsymbol{Z}_j}{1 + \frac{1}{n} \boldsymbol{Z}_j^\top \Sigma^{1/2} H_n^{(-ij)} \Sigma^{1/2} \boldsymbol{Z}_j},$$



Confidential reviewer copy. This manuscript is submitted to the 40th Conference on Neural Information Processing Systems (NeurIPS 2026). Unauthorized sharing, redistribution, or disclosure is strictly prohibited.

where

$$H_n^{(-ij)} = \left( S_z - \frac{1}{n}\Sigma^{1/2}\boldsymbol{Z}_i\boldsymbol{Z}_i^\top\Sigma^{1/2} - \frac{1}{n}\Sigma^{1/2}\boldsymbol{Z}_j\boldsymbol{Z}_j^\top\Sigma^{1/2} + \rho\mathbb{I} \right)^{-1}.$$

Therefore

$$\boldsymbol{Z}_i^\top\Sigma^{1/2}H_n\Sigma^{1/2}\boldsymbol{Z}_j = \frac{\boldsymbol{Z}_i^\top\Sigma^{1/2}H_n^{(-ij)}\Sigma^{1/2}\boldsymbol{Z}_j}{(1+\frac{1}{n}\boldsymbol{Z}_i^\top\Sigma^{1/2}H_n^{(-i)}\Sigma^{1/2}\boldsymbol{Z}_i)(1+\frac{1}{n}\boldsymbol{Z}_j^\top\Sigma^{1/2}H_n^{(-ij)}\Sigma^{1/2}\boldsymbol{Z}_j)}.$$

Therefore, we essentially need to control

$$\frac{1}{m^2}\sum_{i\neq j=1}^{m}\frac{\boldsymbol{Z}_i^\top\Sigma^{1/2}H_n^{(-ij)}\Sigma^{1/2}\boldsymbol{Z}_j}{(1+\frac{1}{n}\boldsymbol{Z}_i^\top\Sigma^{1/2}H_n^{(-i)}\Sigma^{1/2}\boldsymbol{Z}_i)(1+\frac{1}{n}\boldsymbol{Z}_j^\top\Sigma^{1/2}H_n^{(-ij)}\Sigma^{1/2}\boldsymbol{Z}_j)}.$$

We now show that

$$\frac{1}{m^2}\sum_{i\neq j}\frac{\boldsymbol{Z}_i^\top\Sigma^{1/2}H_n^{(-ij)}\Sigma^{1/2}\boldsymbol{Z}_j}{\left(1+n^{-1}\boldsymbol{Z}_i^\top\Sigma^{1/2}H_n^{(-i)}\Sigma^{1/2}\boldsymbol{Z}_i\right)\left(1+n^{-1}\boldsymbol{Z}_j^\top\Sigma^{1/2}H_n^{(-ij)}\Sigma^{1/2}\boldsymbol{Z}_j\right)} = o_{\mathbb{P}}(1).$$

Letting

$$B_{ij} = \Sigma^{1/2}H_n^{(-ij)}\Sigma^{1/2},$$

the summands are not mutually independent, because $B_{ij}$ depends on the observations other than $i, j$. We handle this by decoupling. Set

$$U_m = \sum_{i\neq j}\boldsymbol{Z}_i^\top B_{ij}\boldsymbol{Z}_j.$$

Since $H_n^{(-ij)}$ removes rows $i$ and $j$, the matrix $B_{ij}$ is measurable with respect to $\sigma(\boldsymbol{Z}_k : k \neq i, j)$, and is therefore independent of $(\boldsymbol{Z}_i, \boldsymbol{Z}_j)$. Moreover, because $\rho$ is bounded away from zero and $\|\Sigma\|_{\mathrm{op}}$ is bounded,

$$\|B_{ij}\|_{\mathrm{op}} \leq C_\rho, \qquad \mathrm{tr}(B_{ij}^2) \leq C_\rho p$$

uniformly in $i, j$, with probability tending to one. We prove the required bound by a leave-four covariance calculation. Writing $\xi_a := \Sigma^{1/2}\boldsymbol{Z}_a$ for brevity (so each summand of $U_m$ equals $\xi_i^\top H_n^{(-ij)}\xi_j$), expanding the second moment gives

$$\mathbb{E}U_m^2 = \sum_{i\neq j}\sum_{k\neq\ell}\mathbb{E}\left[\xi_i^\top H_n^{(-ij)}\xi_j\,\xi_k^\top H_n^{(-k\ell)}\xi_\ell\right],$$

where we use the shorthand $R_{ij} := H_n^{(-ij)}$, $R_{ij\ell} := H_n^{(-i,-j,-\ell)}$, $R_{ijk\ell} := H_n^{(-i,-j,-k,-\ell)}$ throughout the rest of this argument. We classify the terms according to the overlap between the ordered pairs $(i, j)$ and $(k, \ell)$.

First, consider the diagonal and reversed-pair cases. If $(i, j) = (k, \ell)$, then, conditional on $R_{ij}$,

$$\mathbb{E}\left[(\xi_i^\top R_{ij}\xi_j)^2 \mid R_{ij}\right] \leq C\ \mathrm{tr}\{(\Sigma^{1/2}R_{ij}\Sigma^{1/2})^2\} \leq Cp.$$

The same bound holds when $(i, j) = (\ell, k)$. Since there are $O(m^2)$ such terms, their total contribution is $O(m^2p)$.

Next suppose that the two ordered pairs share exactly one index. For concreteness, take $i = k$, with $j \neq \ell$; the other cases are identical. By the Sherman–Morrison identity,

$$R_{ij} = R_{ij\ell} - \frac{n^{-1}R_{ij\ell}\xi_\ell\xi_\ell^\top R_{ij\ell}}{1+n^{-1}\xi_\ell^\top R_{ij\ell}\xi_\ell}.$$

Hence

$$\xi_i^\top R_{ij}\xi_j = \xi_i^\top R_{ij\ell}\xi_j - \frac{n^{-1}(\xi_i^\top R_{ij\ell}\xi_\ell)(\xi_\ell^\top R_{ij\ell}\xi_j)}{1+n^{-1}\xi_\ell^\top R_{ij\ell}\xi_\ell}.$$

Similarly,

$$R_{i\ell} = R_{ij\ell} - \frac{n^{-1}R_{ij\ell}\xi_j\xi_j^\top R_{ij\ell}}{1+n^{-1}\xi_j^\top R_{ij\ell}\xi_j}.$$



Confidential reviewer copy. This manuscript is submitted to the 40th Conference on Neural Information Processing Systems (NeurIPS 2026). Unauthorized sharing, redistribution, or disclosure is strictly prohibited.

Therefore

$$\xi_i^\top R_{i\ell}\xi_\ell = \xi_i^\top R_{ij\ell}\xi_\ell - \frac{n^{-1}(\xi_i^\top R_{ij\ell}\xi_j)(\xi_j^\top R_{ij\ell}\xi_\ell)}{1+n^{-1}\xi_j^\top R_{ij\ell}\xi_j}.$$

Conditional on $R_{ij\ell}$, the vectors $\xi_i, \xi_j, \xi_\ell$ are independent and centered. Thus the product of the two leading terms has conditional expectation zero:

$$\mathbb{E}\left[(\xi_i^\top R_{ij\ell}\xi_j)(\xi_i^\top R_{ij\ell}\xi_\ell) \mid R_{ij\ell}\right] = 0.$$

The remaining terms each contain at least one factor $n^{-1}$. Using $\|R_{ij\ell}\|_{\mathrm{op}} \le \rho^{-1}$, the boundedness of $\|\Sigma\|_{\mathrm{op}}$, and the standard quadratic-form bounds

$$\mathbb{E}\left[(\xi_a^\top R_{ij\ell}\xi_b)^2 \mid R_{ij\ell}\right] \le Cp \quad (a \neq b), \qquad \mathbb{E}\left[\xi_a^\top R_{ij\ell}\xi_a \mid R_{ij\ell}\right] \le Cp,$$

we obtain

$$\left|\mathbb{E}\left[\xi_i^\top R_{ij}\xi_j\, \xi_i^\top R_{i\ell}\xi_\ell\right]\right| \le C\,\frac{p}{n}.$$

Since there are $O(m^3)$ ordered quadruples with exactly one shared index, the total contribution of all one-overlap terms is

$$O\left(m^3\,\frac{p}{n}\right).$$

Finally suppose $i, j, k, \ell$ are all distinct. Apply Sherman–Morrison twice to express both $R_{ij}$ and $R_{k\ell}$ as rank-one updates of the common leave-four resolvent $R_{ijk\ell}$. The leading product is

$$(\xi_i^\top R_{ijk\ell}\xi_j)(\xi_k^\top R_{ijk\ell}\xi_\ell),$$

whose conditional expectation given $R_{ijk\ell}$ is zero, because $\xi_i, \xi_j, \xi_k, \xi_\ell$ are independent centered vectors. Every nonzero contribution must therefore come from resolvent-update terms. In the disjoint case, to obtain a nonzero conditional expectation, both factors must be updated: one update is needed to introduce $\xi_k$ or $\xi_\ell$ into the first factor, and another update is needed to introduce $\xi_i$ or $\xi_j$ into the second factor. Thus every nonzero term carries at least two factors of $n^{-1}$. Using again the boundedness of the leave-four resolvent and the quadratic-form moment bounds, each such term is bounded by

$$C\,\frac{p}{n^2}.$$

Hence, for four distinct indices,

$$\left|\mathbb{E}\left[\xi_i^\top R_{ij}\xi_j\, \xi_k^\top R_{k\ell}\xi_\ell\right]\right| \le C\,\frac{p}{n^2}.$$

There are $O(m^4)$ such quadruples, so their total contribution is

$$O\left(m^4\,\frac{p}{n^2}\right).$$

Combining the three cases,

$$\mathbb{E}U_m^2 \le C\left[m^2 p + m^3\,\frac{p}{n} + m^4\,\frac{p}{n^2}\right].$$

Since $m/n \to \pi_\ell \in (0,1)$, this simplifies to

$$EU_m^2 \le Cm^2 p.$$

Therefore

$$U_m = O_{\mathbb{P}}(m\sqrt{p}),$$

and hence

$$\frac{1}{m^2}U_m = O_P\left(\frac{\sqrt{p}}{m}\right) = o_P(1),$$

because $p/m \to \gamma_0/\pi_\ell < \infty$. This proves that the off-diagonal contribution is negligible.

Finally, for deterministic bounded $v$,

$$v^\top H_n(\rho) r_{\ell,n} = \sqrt{\pi_{\ell,n}}\, v^\top H_n(\rho)\Sigma^{1/2}\bar{z}_\ell.$$



Confidential reviewer copy. This manuscript is submitted to the 40th Conference on Neural Information Processing Systems (NeurIPS 2026). Unauthorized sharing, redistribution, or disclosure is strictly prohibited.

Expanding the empirical mean and applying leave-one-out gives

$$v^\top H_n \Sigma^{1/2} \bar{z}_\ell = \frac{1}{m} \sum_{i=1}^{m} v^\top H_n^{(-i)} \Sigma^{1/2} \boldsymbol{Z}_{i,\ell} + o_{\mathbb{P}}(1).$$

Conditional on $H_n^{(-i)}$, the summands have mean zero and uniformly bounded variance. Hence the average is $O_{\mathbb{P}}(m^{-1/2}) = o_{\mathbb{P}}(1)$. Therefore

$$v^\top H_n(\rho) r_{\ell,n} \xrightarrow{\mathbb{P}} 0.$$

This proves the lemma. □

The next lemma connects $H_n$ with $G_n^{(-)}$.

**Lemma A.3.** *Locally uniformly for $\rho$ in compact subsets of $(0, \infty)$,*

$$R_n^\top G_n^{(-)}(\rho) R_n \xrightarrow{\mathbb{P}} \gamma_0 s_{\gamma_0}(\rho) I_K,$$

*and for deterministic bounded $v$,*

$$v^\top G_n^{(-)}(\rho) R_n \xrightarrow{\mathbb{P}} 0, \qquad R_n^\top G_n^{(-)}(\rho) v \xrightarrow{\mathbb{P}} 0.$$

*Proof.* Recall

$$S_z = S_n^{(-)} + R_n R_n^\top.$$

Therefore

$$H_n(\rho) = (S_z + \rho \mathbb{I})^{-1} = (S_n^{(-)} + \rho \mathbb{I} + R_n R_n^\top)^{-1}.$$

Therefore, Woodbury's identity gives

$$H_n = G_n^{(-)} - G_n^{(-)} R_n \left( \mathbb{I}_K + R_n^\top G_n^{(-)} R_n \right)^{-1} R_n^\top G_n^{(-)}.$$

Let

$$T_n = R_n^\top G_n^{(-)} R_n, \qquad L_n = R_n^\top H_n R_n.$$

Multiplying the Woodbury identity by $R_n^\top$ and $R_n$ gives

$$L_n = T_n - T_n (I_K + T_n)^{-1} T_n = T_n (I_K + T_n)^{-1}.$$

By Lemma A.2,

$$L_n \xrightarrow{\mathbb{P}} \frac{d}{1+d} \mathbb{I}_K, \qquad d = \gamma_0 s_{\gamma_0}(\rho).$$

Since

$$T_n = L_n (\mathbb{I}_K - L_n)^{-1},$$

we get

$$T_n \xrightarrow{\mathbb{P}} \frac{d}{1+d} \left( 1 - \frac{d}{1+d} \right)^{-1} \mathbb{I}_K = d \mathbb{I}_K.$$

Hence

$$R_n^\top G_n^{(-)} R_n \xrightarrow{\mathbb{P}} \gamma_0 s_{\gamma_0}(\rho) \mathbb{I}_K.$$

For the mixed bilinear form, multiply the same Woodbury identity by $v^\top$ and $R_n$:

$$v^\top H_n R_n = v^\top G_n^{(-)} R_n - v^\top G_n^{(-)} R_n (\mathbb{I}_K + T_n)^{-1} T_n.$$

Thus

$$v^\top H_n R_n = v^\top G_n^{(-)} R_n \left[ \mathbb{I}_K - (\mathbb{I}_K + T_n)^{-1} T_n \right] = v^\top G_n^{(-)} R_n (\mathbb{I}_K + T_n)^{-1}.$$

Therefore

$$v^\top G_n^{(-)} R_n = v^\top H_n R_n (\mathbb{I}_K + T_n).$$

Lemma A.2 gives $v^\top H_n R_n \xrightarrow{\mathbb{P}} 0$, and $T_n = O_{\mathbb{P}}(1)$. Hence

$$v^\top G_n^{(-)} R_n \xrightarrow{\mathbb{P}} 0.$$

The other statement follows identically. This proves the lemma. □ □



Confidential reviewer copy. This manuscript is submitted to the 40th Conference on Neural Information Processing Systems (NeurIPS 2026). Unauthorized sharing, redistribution, or disclosure is strictly prohibited.

We now complete the proof of the pooled ridge learner using the lemmas. By the anisotropic local law [Knowles and Yin, 2017] and the fact that removing $K$ empirical mean directions is finite rank, for deterministic bounded $v_1, v_2$,

$$v_1^\top G_n^{(-)}(\rho) v_2 - v_1^\top Q_{\gamma_0}(\rho) v_2 \xrightarrow{\mathbb{P}} 0.$$

Consequently,

$$U_n^\top G_n^{(-)}(\rho) U_n \xrightarrow{\mathbb{P}} U_0^\top Q_{\gamma_0}(\rho) U_0,$$

and by Lemma A.3,

$$U_n^\top G_n^{(-)}(\rho) R_n \xrightarrow{\mathbb{P}} 0, \qquad R_n^\top G_n^{(-)}(\rho) U_n \xrightarrow{\mathbb{P}} 0.$$

Next note that, since

$$\widehat{\Sigma}_{\text{pool}} = S_n^{(-)} + (U_n + R_n)(U_n + R_n)^\top,$$

Woodbury's identity gives

$$G_{\text{pool}}(-\rho) = G_n^{(-)}(\rho) - G_n^{(-)}(\rho)(U_n + R_n) D_n(\rho)^{-1} (U_n + R_n)^\top G_n^{(-)}(\rho),$$

where

$$D_n(\rho) = \mathbb{I}_K + (U_n + R_n)^\top G_n^{(-)}(\rho)(U_n + R_n).$$

Therefore,

$$(U_n + R_n)^\top G_n^{(-)}(\rho)(U_n + R_n) \xrightarrow{\mathbb{P}} U_0^\top Q_{\gamma_0}(\rho) U_0 + \gamma_0 s_{\gamma_0}(\rho) \mathbb{I}_K.$$

Thus

$$D_n(\rho) \xrightarrow{\mathbb{P}} \{1 + \gamma_0 s_{\gamma_0}(\rho)\} \mathbb{I}_K + U_0^\top Q_{\gamma_0}(\rho) U_0.$$

Also, for deterministic norm-bounded $v$,

$$v^\top G_n^{(-)}(\rho)(U_n + R_n) \xrightarrow{\mathbb{P}} v^\top Q_{\gamma_0}(\rho) U_0,$$

since

$$v^\top G_n^{(-)}(\rho) R_n \xrightarrow{\mathbb{P}} 0.$$

Therefore

$$v^\top G_{\text{pool}}(-\rho) v \xrightarrow{\mathbb{P}} v^\top Q_{\gamma_0}(\rho) v - v^\top Q_{\gamma_0}(\rho) U_0 \left[\{1 + \gamma_0 s_{\gamma_0}(\rho)\} \mathbb{I}_K + U_0^\top Q_{\gamma_0}(\rho) U_0\right]^{-1} U_0^\top Q_{\gamma_0}(\rho) v.$$

However, by definition,

$$K_{\gamma_0, U_0}(\rho) = Q_{\gamma_0}(\rho) - Q_{\gamma_0}(\rho) U_0 \left[\{1 + \gamma_0 s_{\gamma_0}(\rho)\} \mathbb{I}_K + U_0^\top Q_{\gamma_0}(\rho) U_0\right]^{-1} U_0^\top Q_{\gamma_0}(\rho).$$

Hence

$$v^\top G_{\text{pool}}(\rho) v \xrightarrow{\mathbb{P}} v^\top K_{\gamma_0, U_0}(\rho) v.$$

We now analyze the bias term. We start by recalling

$$\psi_0(\rho) = \beta^\top K_{\gamma_0, U_0}(\rho) \beta.$$

Then as before

$$\boldsymbol{\beta}^\top G_{\text{pool}}(-\rho) \boldsymbol{\beta} \xrightarrow{\mathbb{P}} \psi_0(\rho)$$

locally uniformly for $\rho$ in compact subsets of $(0, \infty)$ since $\boldsymbol{\beta}$ is norm bounded. Next, since

$$\frac{d}{d\rho} G_{\text{pool}}(-\rho) = -G_{\text{pool}}(\rho)^2,$$

we have

$$\boldsymbol{\beta}^\top G_{\text{pool}}(-\rho)^2 \boldsymbol{\beta} = -\frac{d}{d\rho} \left[\boldsymbol{\beta}^\top G_{\text{pool}}(-\rho) \boldsymbol{\beta}\right].$$

Since all the convergences are locally uniform in $\rho$ on compact sets, and all resolvents are uniformly bounded by $\rho^{-1}$ on compact subsets of $(0, \infty)$, differentiating the deterministic equivalent gives

$$\boldsymbol{\beta}^\top G_{\text{pool}}(-\rho_0)^2 \boldsymbol{\beta} \xrightarrow{\mathbb{P}} -\psi_0'(\rho_0).$$

Therefore

$$B_n \xrightarrow{\mathbb{P}} \rho_0^2 [-\psi_0'(\rho_0)].$$

This completes the proof for the fixed-effects pooled ridge estimator.



Confidential reviewer copy. This manuscript is submitted to the 40th Conference on Neural Information Processing Systems (NeurIPS 2026). Unauthorized sharing, redistribution, or disclosure is strictly prohibited.

**Analysis of Pooled Ridgeless Estimator – Fixed Effects:** Recall the notation

$$\widehat{\Sigma}_{\text{pool}} = \frac{1}{n}\mathbb{X}_{\text{pool}}{}^{\top}\mathbb{X}_{\text{pool}}, \qquad G_{\text{pool}}(-\rho) = (\widehat{\Sigma}_{\text{pool}} + \rho I)^{-1}, \qquad \rho > 0.$$

Let the spectral decomposition of $\widehat{\Sigma}_{\text{pool}}$ be

$$\widehat{\Sigma}_{\text{pool}} = \sum_{j=1}^{p} s_j u_j u_j^{\top}, \qquad s_j \geq 0.$$

Also let

$$P_{\mathcal{C}(\mathbb{X}_{\text{pool}}^{\top})} = \widehat{\Sigma}_{\text{pool}}\widehat{\Sigma}_{\text{pool}}^{+}$$

denote the orthogonal projection onto the row space of $\mathbb{X}_{\text{pool}}$. Then

$$I - P_{\mathcal{C}(\mathbb{X}_{\text{pool}}^{\top})} = \sum_{j:s_j=0} u_j u_j^{\top}.$$

By the anisotropic local law [Knowles and Yin, 2017] and smallest singular value bounds for sample covariance matrices with bounded finite-rank deterministic perturbations [Bai et al., 2010], there exists $c_{\gamma_0} > 0$ such that, with probability tending to one, every nonzero eigenvalue of $\widehat{\Sigma}_{\text{pool}}$ lies in

$$[c_{\gamma_0}, 1/c_{\gamma_0}].$$

On this high-probability event, for any deterministic $b$ of bounded norm,

$$\left|\rho^2 b^{\top} G_{\text{pool}}(-\rho)^2 b - b^{\top}(I - P_{\mathcal{C}(\mathbb{X}_{\text{pool}}^{\top})})b\right| \leq \frac{\rho^2}{c_{\gamma_0}^2}\|b\|^2.$$

Since for $s_j \geq c_{\gamma_0}$,

$$0 \leq \frac{\rho^2}{(s_j+\rho)^2} \leq \frac{\rho^2}{c_{\gamma_0}^2},$$

we have that

$$\lim_{n\to\infty} b^{\top}(I - P_{\mathcal{C}(\mathbb{X}_{\text{pool}}^{\top})})b = \lim_{\rho\downarrow 0}\lim_{n\to\infty} \rho^2 b^{\top} G_{\text{pool}}(-\rho)^2 b,$$

since the deterministic equivalent is locally uniform on compact subsets of $(0,\infty)$. Similarly, for the variance term,

$$\frac{1}{n}\text{Tr}(\widehat{\Sigma}^{+}) - \frac{1}{n}\text{Tr}\{\widehat{\Sigma}G_{\text{pool}}(-\rho)^2\} = \frac{1}{n}\sum_{j:s_j>0}\left[\frac{1}{s_j} - \frac{s_j}{(s_j+\rho)^2}\right].$$

For $s_j \geq c_{\gamma_0}$,

$$0 \leq \frac{1}{s_j} - \frac{s_j}{(s_j+\rho)^2} = \frac{2\rho s_j + \rho^2}{s_j(s_j+\rho)^2} \leq C_{\gamma_0}\,\rho,$$

for some constant $C_{\gamma_0} < \infty$ depending only on $c_{\gamma_0}$ (valid for $\rho$ in any compact subset of $(0,\infty)$). Since the number of nonzero eigenvalues is at most $n \wedge p$,

$$\left|\frac{1}{n}\text{Tr}(\widehat{\Sigma}_{\text{pool}}^{+}) - \frac{1}{n}\text{Tr}\{\widehat{\Sigma}_{\text{pool}}G_{\text{pool}}(-\rho)^2\}\right| \leq C_{\gamma_0}\,\rho.$$

Hence

$$\frac{1}{n}\text{Tr}(\widehat{\Sigma}^{+}) = \lim_{\rho\downarrow 0}\lim_{n\to\infty}\frac{1}{n}\text{Tr}\{\widehat{\Sigma}G_{\text{pool}}(-\rho)^2\}.$$

Thus, the ridgeless bias and variance are obtained as the $\rho \downarrow 0$ limits of the corresponding ridge bias and variance deterministic equivalents, i.e.,

$$B_{+} = \lim_{\rho_0\downarrow 0}\rho_0^2[-\psi_0'(\rho_0)]$$

and

$$V_{+} = \lim_{\rho_0\downarrow 0} V_{\gamma}(\rho_0).$$



Confidential reviewer copy. This manuscript is submitted to the 40th Conference on Neural Information Processing Systems (NeurIPS 2026). Unauthorized sharing, redistribution, or disclosure is strictly prohibited.

### A.4 Proof of Theorem 2.3

Here, we only demonstrate the analysis for the random-effects ridge case. All the remaining proofs (random-effects ridgeless and fixed-effects analyses) follow by obvious modifications of the argument below combined with the proof of Theorem 2.5 taken with $K = 1$. For the random-effects ridge calculation, by direct calculations

$$R_{\mathrm{ens},\mathbf{w}} = \sigma_\star^2 + \sum_{\ell=1}^{K} w_\ell^2 \, V_{\gamma_\ell}(\rho_\ell) + o_{\mathbb{P}}(1) \\ + \frac{1}{p}\mathrm{Tr}\left(\sum_{\ell=1}^{K} w_\ell^2 \rho_\ell^2 G_\ell^2(-\rho_\ell)\right) + \frac{1}{p}\mathrm{Tr}\left(\sum_{\ell\neq\ell'} w_\ell w_{\ell'} \rho_\ell \rho_{\ell'} G_\ell(-\rho_\ell) G_{\ell'}(-\rho_{\ell'})\right),$$

where $G_\ell(-\rho_\ell) = (\hat{\Sigma}_\ell + \rho_\ell \mathbb{I})^{-1}$. The analyses of $V_{\gamma_\ell}(\rho_\ell)$ and of $\frac{1}{p}\mathrm{Tr}(\sum w_\ell^2 \rho_\ell^2 G_\ell^2(-\rho_\ell))$ are verbatim the same as in the proof of Theorem 2.5 taken with $K = 1$, so we focus only on the cross-cluster term $\frac{1}{p}\mathrm{Tr}(\sum_{\ell\neq\ell'} w_\ell w_{\ell'} \rho_\ell \rho_{\ell'} G_\ell(-\rho_\ell) G_{\ell'}(-\rho_{\ell'}))$. In the analysis, we drop the argument $(-\rho_\ell)$ from $G_\ell(-\rho_\ell)$ whenever clear from context.

Fix $\ell \neq j$. We claim that

$$\frac{1}{p}\mathrm{Tr}(G_\ell G_j) = \frac{1}{p}\mathrm{Tr}(Q_\ell Q_j) + o_{\mathbb{P}}(1).$$

To that end we write

$$\mathbb{X}_\ell = X_\ell^0 + \mathbf{1}_{n_\ell} \boldsymbol{\mu}_\ell^\top,$$

where $1_{n_\ell}$ is a $n_\ell \times 1$ vector of 1's, and the $i^{\mathrm{th}}$ row of $X_\ell^0$ is $\Sigma^{1/2}\boldsymbol{Z}_{i,\ell}$. Let

$$\widehat{\Sigma}_\ell^0 = \frac{1}{n_\ell}(X_\ell^0)^\top X_\ell^0, \qquad G_\ell^0 = (\widehat{\Sigma}_\ell^0 + \rho_\ell \mathbb{I})^{-1}.$$

Then

$$\widehat{\Sigma}_\ell - \widehat{\Sigma}_\ell^0 = \frac{1}{n_\ell}(X_\ell^0)^\top \mathbf{1}_{n_\ell} \boldsymbol{\mu}_\ell^\top + \frac{1}{n_\ell}\boldsymbol{\mu}_\ell \mathbf{1}_{n_\ell}^\top X_\ell^0 + \boldsymbol{\mu}_\ell \boldsymbol{\mu}_\ell^\top.$$

Each term on the right-hand side has rank at most one, and hence

$$\mathrm{rank}(\widehat{\Sigma}_\ell - \widehat{\Sigma}_\ell^0) \leq 3.$$

By matrix inverse identity,

$$G_\ell - G_\ell^0 = -G_\ell(\widehat{\Sigma}_\ell - \widehat{\Sigma}_\ell^0)G_\ell^0.$$

Therefore

$$\mathrm{rank}(G_\ell - G_\ell^0) \leq 3.$$

Moreover, since $\rho_\ell > 0$,

$$\|G_\ell\|_{\mathrm{op}} \leq \rho_\ell^{-1}, \qquad \|G_\ell^0\|_{\mathrm{op}} \leq \rho_\ell^{-1}.$$

Similarly,

$$\mathrm{rank}(G_j - G_j^0) \leq 3, \qquad \|G_j\|_{\mathrm{op}} \leq \rho_j^{-1}, \qquad \|G_j^0\|_{\mathrm{op}} \leq \rho_j^{-1}.$$

Now

$$G_\ell G_j - G_\ell^0 G_j^0 = (G_\ell - G_\ell^0)G_j + G_\ell^0(G_j - G_j^0).$$

Thus

$$\left|\frac{1}{p}\mathrm{Tr}(G_\ell G_j) - \frac{1}{p}\mathrm{Tr}(G_\ell^0 G_j^0)\right| \leq \frac{1}{p}\left|\mathrm{Tr}\{(G_\ell - G_\ell^0)G_j\}\right| \\ + \frac{1}{p}\left|\mathrm{Tr}\{G_\ell^0(G_j - G_j^0)\}\right|.$$

Using

$$|\mathrm{Tr}(AB)| \leq \mathrm{rank}(A)\|A\|_{\mathrm{op}}\|B\|_{\mathrm{op}},$$

whenever $A$ is finite-rank, together with the uniform boundedness of the resolvents, we get

$$\left|\frac{1}{p}\mathrm{Tr}(G_\ell G_j) - \frac{1}{p}\mathrm{Tr}(G_\ell^0 G_j^0)\right| \leq C'(\rho_\ell^{-1}, \rho_j^{-1})/p,$$



Confidential reviewer copy. This manuscript is submitted to the 40th Conference on Neural Information Processing Systems (NeurIPS 2026). Unauthorized sharing, redistribution, or disclosure is strictly prohibited.

for a finite constant $C'(\rho_\ell^{-1}, \rho_j^{-1})$ depending on $\rho_\ell^{-1}, \rho_j^{-1}$. Therefore

$$\frac{1}{p}\mathrm{Tr}(G_\ell G_j) = \frac{1}{p}\mathrm{Tr}(G_\ell^0 G_j^0) + o_{\mathbb{P}}(1),$$

uniformly in $\ell, j$ (since $K$ is finite). It remains to replace the zero-mean resolvents by their deterministic equivalents. To that end we write

$$\frac{1}{p}\mathrm{Tr}(G_\ell^0 G_j^0) - \frac{1}{p}\mathrm{Tr}(Q_\ell Q_j) = \frac{1}{p}\mathrm{Tr}\{(G_\ell^0 - Q_\ell)G_j^0\} + \frac{1}{p}\mathrm{Tr}\{Q_\ell(G_j^0 - Q_j)\}.$$

Because the clusters are independent, conditional on $G_j^0$, the matrix $G_j^0$ is deterministic with respect to the randomness in $G_\ell^0$. Moreover,

$$\|G_j^0\|_{\mathrm{op}} \le \rho_j^{-1}.$$

Therefore, the anisotropic deterministic-equivalence [Knowles and Yin, 2017] implies that for any deterministic matrix $A_p$ with uniformly bounded operator norm,

$$\frac{1}{p}\mathrm{Tr}\{A_p(G_\ell^0 - Q_\ell)\} \xrightarrow{\mathbb{P}} 0.$$

Applying this conditionally with $A_p = G_j^0$ gives

$$\frac{1}{p}\mathrm{Tr}\{(G_\ell^0 - Q_\ell)G_j^0\} = o_{\mathbb{P}}(1).$$

Similarly, $Q_\ell$ is deterministic and uniformly bounded in operator norm, and hence

$$\frac{1}{p}\mathrm{Tr}\{Q_\ell(G_j^0 - Q_j)\} = o_{\mathbb{P}}(1).$$

Consequently,

$$\frac{1}{p}\mathrm{Tr}(G_\ell^0 G_j^0) = \frac{1}{p}\mathrm{Tr}(Q_\ell Q_j) + o_{\mathbb{P}}(1).$$

Combining this with the finite-rank perturbation bound yields

$$\frac{1}{p}\mathrm{Tr}(G_\ell G_j) = \frac{1}{p}\mathrm{Tr}(Q_\ell Q_j) + o_{\mathbb{P}}(1).$$

Therefore,

$$\frac{1}{p}\mathrm{Tr}\Big(\sum_{\ell \ne \ell'} w_\ell w_{\ell'} \rho_\ell \rho_{\ell'} G_\ell(-\rho_\ell) G_{\ell'}(-\rho_{\ell'})\Big) = \sum_{\ell \ne j} w_\ell w_j \rho_\ell \rho_j \frac{1}{p}\mathrm{Tr}(Q_\ell Q_j) + o_{\mathbb{P}}(1),$$

as promised.

### A.5 Proof of Theorem 2.4

The proof of part (i) follows from Theorem 2.3 together with the optimal tuning result of Dobriban and Wager [2018], since for the random-effects model the cluster mean shifts do not affect the limiting empirical spectral distributions of $\hat{\Sigma}_\ell$ asymptotically. For part (ii), apply the Lagrange-multiplier method to minimize $\mathbf{w}^\top A_{\mathrm{RE}}^\star \mathbf{w}$ over $\{\mathbf{w} \in \mathbb{R}^K : \mathbf{1}^\top \mathbf{w} = 1\}$. The first-order conditions yield $\widetilde{\mathbf{w}} \propto (A_{\mathrm{RE}}^\star)^{-1}\mathbf{1}$ with optimal value $(\mathbf{1}^\top (A_{\mathrm{RE}}^\star)^{-1}\mathbf{1})^{-1}$. When this $\widetilde{\mathbf{w}} \in \mathbb{R}_+^K$, it is also the constrained-positive optimum, and the conclusion follows.

### A.6 Proof of Corollary 2.6

We only verify the second claim and that $\mathcal{R}_{\mathrm{excess}} := \frac{\kappa_1\kappa_2(1-\kappa_1\kappa_2)}{\kappa_0(\kappa_1+\kappa_2-2\kappa_1\kappa_2)} > 1$. The rest of the proof follows from standard algebra by evaluating the expressions in Theorem 2.4 and Theorem 1 through integral w.r.t. classical Marchenko-Pastur distribution [Bai et al., 2010, Section 3.1.1]. We omit the algebraic details.



Confidential reviewer copy. This manuscript is submitted to the 40th Conference on Neural Information Processing Systems (NeurIPS 2026). Unauthorized sharing, redistribution, or disclosure is strictly prohibited.

We now prove the claim that $\mathcal{R}_{\text{excess}} > 1$. First note that $\kappa(\gamma)$ is the positive root of

$$\kappa^2 + \left(\alpha - 1 + \frac{1}{\gamma}\right)\kappa - \alpha = 0.$$

Since the polynomial is negative at 0 and positive at 1,

$$0 < \kappa(\gamma) < 1$$

for every $\gamma > 0$. Hence

$$0 < \kappa_1, \kappa_2, \kappa_0 < 1.$$

Define

$$h(u) = \frac{\alpha}{u} - u - \alpha + 1, \qquad 0 < u < 1.$$

From the defining equation for $\kappa(\gamma)$, dividing by $\kappa(\gamma)$ gives

$$h(\kappa(\gamma)) = \frac{1}{\gamma}.$$

Therefore,

$$h(\kappa_1) = \frac{1}{\gamma_1}, \qquad h(\kappa_2) = \frac{1}{\gamma_2}, \qquad h(\kappa_0) = \frac{1}{\gamma_0}.$$

Since

$$\frac{1}{\gamma_0} = \frac{1}{\gamma_1} + \frac{1}{\gamma_2},$$

we have

$$h(\kappa_0) = h(\kappa_1) + h(\kappa_2).$$

Now write

$$x = \kappa_1, \qquad y = \kappa_2.$$

The optimally weighted ensemble excess-risk contribution, divided by $\tau^2$, is

$$q(x, y) = \frac{xy(1 - xy)}{x + y - 2xy}.$$

Thus

$$\mathcal{R}_{\text{excess}} = \frac{q(x, y)}{\kappa_0}.$$

It is therefore enough to prove

$$q(x, y) > \kappa_0.$$

Because

$$h'(u) = -\frac{\alpha}{u^2} - 1 < 0,$$

the function $h$ is strictly decreasing on $(0, 1)$. Hence

$$q(x, y) > \kappa_0$$

is equivalent to

$$h(q(x, y)) < h(\kappa_0).$$

Since

$$h(\kappa_0) = h(x) + h(y),$$

it is enough to show

$$h(q(x, y)) < h(x) + h(y).$$

A direct algebraic calculation gives

$$h(q(x, y)) - h(x) - h(y) = -\frac{(1 - x)(1 - y)\left[\alpha(x + y - 2xy) + (1 - xy)(x + y - xy)\right]}{(1 - xy)(x + y - 2xy)}.$$

Now $0 < x, y < 1$, so

$$1 - x > 0, \qquad 1 - y > 0, \qquad 1 - xy > 0,$$



Confidential reviewer copy. This manuscript is submitted to the 40th Conference on Neural Information Processing Systems (NeurIPS 2026). Unauthorized sharing, redistribution, or disclosure is strictly prohibited.

and

$$x + y - 2xy = x(1 - y) + y(1 - x) > 0.$$

Also,

$$\alpha(x + y - 2xy) > 0$$

because $\alpha > 0$, and

$$(1 - xy)(x + y - xy) > 0.$$

Therefore the entire fraction is strictly positive, and the leading minus sign implies

$$h(q(x, y)) - h(x) - h(y) < 0.$$

Thus

$$h(q(x, y)) < h(x) + h(y) = h(\kappa_0).$$

Since $h$ is strictly decreasing, this implies

$$q(x, y) > \kappa_0.$$

Therefore

$$\frac{q(x, y)}{\kappa_0} > 1,$$

which is exactly

$$\mathcal{R}_{\text{excess}} > 1.$$



Confidential reviewer copy. This manuscript is submitted to the 40th Conference on Neural Information Processing Systems (NeurIPS 2026). Unauthorized sharing, redistribution, or disclosure is strictly prohibited.

# NeurIPS Paper Checklist

1. **Claims**

   Question: Do the main claims made in the abstract and introduction accurately reflect the paper's contributions and scope?

   Answer: [Yes]

   Justification: The abstract and Section 1 accurately describe the paper's two contributions: (i) the asymptotic comparison of pooled and ensemble ridge regression under covariate shift, formalized in Theorems 2.3, 2.4, 2.5, and Corollary 2.6, and (ii) the simulation study spanning underparameterized, interpolation, and overparameterized regimes presented in Section 3.1 and Figure 2.

2. **Limitations**

   Question: Does the paper discuss the limitations of the work performed by the authors?

   Answer: [Yes]

   Justification: A dedicated "Limitations and future work" paragraph at the end of Section 4 explicitly discusses the well-specified linear model assumption, the shared coefficients and noise variance across clusters, the restriction to mean-shift covariate heterogeneity, the assumption of known cluster memberships, and the random-effects setting under which the optimally weighted optimally tuned comparison is stated. The remark following Assumption 2.1 additionally clarifies the scope of the modeling choices.

3. **Theory assumptions and proofs**

   Question: For each theoretical result, does the paper provide the full set of assumptions and a complete (and correct) proof?

   Answer: [Yes]

   Justification: All assumptions are collected in Assumption 2.1 (Section 2.1), and each theorem statement explicitly references it. Full proofs of Theorems 2.3, 2.4, 2.5, and Corollary 2.6 are provided in Appendix A.3 through A.6, supported by Lemmas A.2 and A.3 and the background result Proposition A.1 from Knowles and Yin [2017].

4. **Experimental result reproducibility**

   Question: Does the paper fully disclose all the information needed to reproduce the main experimental results of the paper to the extent that it affects the main claims and/or conclusions of the paper (regardless of whether the code and data are provided or not)?

   Answer: [Yes]

   Justification: Section 3.1 specifies the full simulation protocol (cluster sizes, covariate distributions, coefficient generation, signal-to-noise ratio, test set construction, range of $p$, and number of repetitions), and Section 3.2 specifies the parameter configurations. The "Base Learners" paragraph in Section 3.1 specifies the numerical routines used for both ridge and ridgeless learners, including the cross-validation grid for the ridge tuning parameter.

5. **Open access to data and code**

   Question: Does the paper provide open access to the data and code, with sufficient instructions to faithfully reproduce the main experimental results, as described in supplemental material?

   Answer: [Yes]

   Justification: We have included all code for replicating analyses and figures as supplementary material.

6. **Experimental setting/details**

   Question: Does the paper specify all the training and test details (e.g., data splits, hyperparameters, how they were chosen, type of optimizer) necessary to understand the results?

   Answer: [Yes]



Confidential reviewer copy. This manuscript is submitted to the 40th Conference on Neural Information Processing Systems (NeurIPS 2026). Unauthorized sharing, redistribution, or disclosure is strictly prohibited.

Justification: Section 3.1 specifies the data-generating process, sample sizes, covariate dimension grid, signal-to-noise ratio, and number of repetitions. The "Base Learners" paragraph in Section 3.1 specifies the use of `numpy.linalg.lstsq` for the ridgeless solution and `RidgeCV` from `scikit-learn` [Pedregosa et al., 2011] with a logarithmic grid of 20 values over $[10^{-6}, 10^{6}]$ tuned via leave-one-out cross-validation independently for every learner.

7. **Experiment statistical significance**

   Question: Does the paper report error bars suitably and correctly defined or other appropriate information about the statistical significance of the experiments?

   Answer: [Yes]

   Justification: As described in Section 3.1, all reported quantities are averaged over 100 independent repetitions, and 95% confidence intervals computed from the $t$-distribution are plotted in Figure 2. Section 4 explicitly references these confidence intervals when discussing the lack of effect of cluster geometry.

8. **Experiments compute resources**

   Question: For each experiment, does the paper provide sufficient information on the computer resources (type of compute workers, memory, time of execution) needed to reproduce the experiments?

   Answer: [Yes]

   Justification: As stated at the beginning of Section 3, all experiments were run using 4 CPU cores on a local machine. The simulations involve ridge and ridgeless least-squares fits on synthetic data with $n = 2000$ and $p$ up to 6000, and complete within standard local-machine time and memory.

9. **Code of ethics**

   Question: Does the research conducted in the paper conform, in every respect, with the NeurIPS Code of Ethics `https://neurips.cc/public/EthicsGuidelines`?

   Answer: [Yes]

   Justification: The research conforms with the NeurIPS Code of Ethics. The work is purely theoretical and methodological, uses only synthetic data generated as described in Section 3.1, involves no human subjects or personally identifiable information, and poses no foreseeable ethical concerns.

10. **Broader impacts**

    Question: Does the paper discuss both potential positive societal impacts and negative societal impacts of the work performed?

    Answer: [N/A]

    Justification: This is foundational theoretical work characterizing the asymptotic risk of two standard estimation strategies for linear regression under covariate heterogeneity. It is not tied to any particular application or deployment, and we do not identify a direct path from the results to specific positive or negative societal impacts.

11. **Safeguards**

    Question: Does the paper describe safeguards that have been put in place for responsible release of data or models that have a high risk for misuse (e.g., pre-trained language models, image generators, or scraped datasets)?

    Answer: [N/A]

    Justification: The paper releases no models, datasets, or other assets that pose a risk for misuse. All data are synthetic and generated by the procedure in Section 3.1.

12. **Licenses for existing assets**

    Question: Are the creators or original owners of assets (e.g., code, data, models), used in the paper, properly credited and are the license and terms of use explicitly mentioned and properly respected?

    Answer: [N/A]



Confidential reviewer copy. This manuscript is submitted to the 40th Conference on Neural Information Processing Systems (NeurIPS 2026). Unauthorized sharing, redistribution, or disclosure is strictly prohibited.

Justification: The paper uses only standard open-source scientific computing libraries (`NumPy` and `scikit-learn` [Pedregosa et al., 2011], cited in the bibliography). No third-party datasets or pretrained models are used.

13. **New assets**

    Question: Are new assets introduced in the paper well documented and is the documentation provided alongside the assets?

    Answer: [N/A]

    Justification: The paper does not introduce new datasets, models, or other released assets. The accompanying simulation code reproduces results from synthetic data only.

14. **Crowdsourcing and research with human subjects**

    Question: For crowdsourcing experiments and research with human subjects, does the paper include the full text of instructions given to participants and screenshots, if applicable, as well as details about compensation (if any)?

    Answer: [N/A]

    Justification: The paper does not involve crowdsourcing or research with human subjects. All data are synthetic, as described in Section 3.1.

15. **Institutional review board (IRB) approvals or equivalent for research with human subjects**

    Question: Does the paper describe potential risks incurred by study participants, whether such risks were disclosed to the subjects, and whether Institutional Review Board (IRB) approvals (or an equivalent approval/review based on the requirements of your country or institution) were obtained?

    Answer: [N/A]

    Justification: The paper does not involve research with human subjects, so IRB review does not apply.

16. **Declaration of LLM usage**

    Question: Does the paper describe the usage of LLMs if it is an important, original, or non-standard component of the core methods in this research? Note that if the LLM is used only for writing, editing, or formatting purposes and does *not* impact the core methodology, scientific rigor, or originality of the research, declaration is not required.

    Answer: [N/A]

    Justification: LLMs were not used at all in the methodology, theoretical analysis, or experimental design for this work.



Confidential reviewer copy. This manuscript is submitted to the 40th Conference on Neural Information Processing Systems (NeurIPS 2026). Unauthorized sharing, redistribution, or disclosure is strictly prohibited.