


Study of the PDO index predictability for 1 to 5 years with INMCM5

M. S. Alexandrov¹, E. M. Volodin²

¹Moscow State University, Moscow, Russia


²Marchuk Institute of Numerical Mathematics of the Russian Academy of Sciences, Moscow, Russia

Introduction




PDO is recurring change in ocean and atmospheric climate in the mid-latitude Pacific Ocean. During positive phase, the western Pacific becomes colder and part of the eastern Pacific becomes warmer; during negative phase, the opposite pattern occurs. The PDO index is defined as the projection of monthly mean sea surface temperature anomalies on their first EOF in the North Pacific.

Task description




First year forecast




We observe correlation decrease from nearly 1.0 in November to 0.355 in October. This indicates a moderate decline in consistency between forecasted and actual PDO index values over the annual forecast period of the model.

1-5 years forecast




Correlation coefficients show that for the first two years of forecast, the statistical relationship between predicted and actual PDO-index is fairly well. This indicates a good predictability of PDO-index for this period.

Inertial forecast



Potential predictability




Conclusion

1. The INMCM5 climate model can predict the PDO index reasonably well for a period of up to two years.
2. The inertial forecast gives a better correlation with the actual PDO index values. However, a significant decrease in correlation is also observed for a period of more than two years.
3. There is potential for improving the INMCM5 model for predicting the PDO index.

References

[1] Volodin E. M., Mortikov E. V., Kostykin S. V. Simulation of the present-day climate with the climate model INMCM5

Questions? aleksandrovm@my.msu.ru



Link to this poster