

08 - “And” Hypothesis

Lean: First Steps

Tariq Rashid

October 27, 2024

“And” Hypotheses

- Previously looked at a “**logical or**” hypothesis.
- Here we’ll look at a “**logical and**” hypothesis.

Task

- Given

$$(x = 5) \wedge (y = x + 3)$$

- where $x, y \in \mathbb{Z}$, show

$$y = 8$$

Conjunction

- The symbol \wedge means “logical and”.
- The statement $P \wedge Q$ means both P and Q are true.
- Statements of the form $P \wedge Q$ are called **conjunctions**.

- A hypothesis $P \wedge Q$ is the same as hypotheses P and Q both being true.
- .. proof proceeds just like earlier examples with two hypotheses.
- Conjunctive hypotheses are **not particularly interesting**,
 - ... but we do need to know how to handle them.

- Structured proof

$(x = 5) \wedge (y = x + 3)$	given fact	(1)
------------------------------	------------	-----

$x = 5$	derived fact (1)	(2)
---------	------------------	-----

$y = x + 3$	derived fact (1)	(3)
-------------	------------------	-----

$y = x + 3$	using fact (3)	
-------------	----------------	--

$= (5) + 3$	using fact (2)	
-------------	----------------	--

$= 8$	using arithmetic	□
-------	------------------	---

- From the single conjunctive hypothesis $(x = 5) \wedge (y = x + 3)$ we derive two smaller hypotheses, both of which are true:
 - $x = 5$
 - $y = x + 3$
- We start with $y = x + 3$ from derived fact (3), then use $x = 5$ from derived fact (2) to finally conclude $y = 8$.

```
-- 08 - Conjunctive "and" Hypothesis
```

```
import Mathlib.Tactic
```

```
example {x y :  $\mathbb{Z}$ } (h : x = 5  $\wedge$  y = x + 3) : y = 8 := by
  obtain ⟨ ha , hb ⟩ := h
  calc
    y = x + 3 := by rw [hb]
    _ = (5) + 3 := by rw [ha]
    _ = 8 := by norm_num
```

- The hypothesis is a conjunction, uses the symbol \wedge for “logical and”.
- The instruction `obtain < ha, hb > := h` splits the conjunctive hypothesis `h` into `ha` and `hb`, and then removes `h`.
- Notice the comma and angle brackets `<>` used to split a conjunction, unlike the vertical bar `|` and no brackets used to split a disjunction.
- The rest of the proof uses the familiar `calc` structure to show `y = 8`.

- Placing the cursor before `obtain` shows only one hypothesis `h`.

$$\begin{array}{l} x \ y : \mathbb{Z} \\ h : x = 5 \wedge y = x + 3 \\ \vdash y = 8 \end{array}$$

- Placing the cursor on the next line after `obtain` shows `h` has been replaced by `ha` and `hb`.

```
x y : ℤ
ha : x = 5
hb : y = x + 3
⊢ y = 8
```

- Notice the hypothesis `h` has been removed.

- Write a Lean program to show $y \geq 8$, where $x, y \in \mathbb{Z}$, given the conjunctive hypothesis

$$(x \geq 5) \wedge (y = x + 3)$$