

BOUNDED GATEWAY PARAMETERS FOR THE ERDŐS–STRAUS CONJECTURE

MACDARA Ó MURCHÚ

ABSTRACT. We present a structural analysis of solutions to the Erdős–Straus conjecture, parameterised by a single integer A . A *gateway decomposition* produces an explicit representation $\frac{4}{p} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}$ from any divisor $d \mid \left(\frac{p+A}{4}\right)^2$ in a specific residue class modulo A . Three propositions handle all primes outside a residual class — $p \equiv 1 \pmod{24}$, p a quadratic residue mod 7, and every prime factor of $(p+3)/4$ congruent to 1 $\pmod{3}$ — which has relative density zero among the primes and requires a computational search over A .

Verification to 10^{11} shows that 32 prime values of A , all congruent to 3 $\pmod{4}$ and all at most 359, suffice for every prime $p \equiv 1 \pmod{4}$ (the value $A = 3$ together with 31 distinct gateway values); primes $p \equiv 3 \pmod{4}$ are classical, with $A = 1$. Among the 88 million residual primes, 68.9% are solved at $A = 7$ and 91.0% by $A = 11$. We observe a *slow-growth phenomenon* in the gateway parameter: the maximum A required is 79 at 10^6 , 167 at 10^7 , 239 at 10^8 and 10^9 , and 359 at 10^{10} —and this value is *unchanged* at 10^{11} : no prime in $(10^{10}, 10^{11}]$ requires $A > 359$. We conjecture that $\max A$ is absolutely bounded and show that this would imply the full conjecture. The observation connects to divisor equidistribution in residue classes for structured quadratic integers.

1. INTRODUCTION

The *Erdős–Straus conjecture*, posed by Paul Erdős in 1948, asserts that for every integer $n \geq 2$, the equation

$$(1) \quad \frac{4}{n} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}$$

has a solution in positive integers x, y, z .

Prior work. The conjecture has been verified computationally to 10^{17} by Salez [6], who introduced an efficient modular sieve that reduces the set of primes requiring direct verification, and extended to 10^{18} by Mihnea and Bogdan [4]. Earlier verification to 10^{14} is due to Swett [9]. Elsholtz and Tao [2] showed that the average number of solutions satisfies $f(p) \asymp (\log p)^3$ for primes, and Tao [10] observed that quadratic

reciprocity presents a fundamental obstruction to resolving the conjecture via covering systems alone.

Algebraic results of Schinzel [7], Sierpiński [8], Mordell [5], and Webb [12], among others, provide explicit decompositions for every residue class modulo 840 except the *Mordell subset* $\{1, 121, 169, 289, 361, 529\}$ —the six squares of units modulo 840—which requires methods beyond modular identities.

Vaughn [11] independently formulated an equivalent approach to the hard cases: for each prime p , one seeks x with $\lceil p/4 \rceil \leq x \leq \lceil p/2 \rceil$ and a divisor d of x^2 satisfying a modular condition. Our gateway parameterisation by A (with $x = (p + A)/4$) is algebraically equivalent to this formulation.

Our contribution. We provide a *structural analysis* of solutions, not a new verification record. Using a gateway decomposition parameterised by a single integer A , we classify how each prime is solved and observe a phenomenon not previously noted in the literature:

The maximum value of A required to solve all primes up to N grows extremely slowly—roughly like a small multiple of $\log N$ across five decades of prime range, and does not increase at all from 10^{10} to 10^{11} —and may be absolutely bounded.

We call this the *bounded- A conjecture* (Conjecture 6.1). If true, it implies the full Erdős–Straus conjecture.

The integrality condition underlying the gateway requires $d \mid N^2$ (where $N = (p + A)/4$), which is strictly weaker than $d \mid N$ and essential for completeness.

Our main computational result is:

Theorem 1.1 (Main Result). *Every prime $p \leq 10^{11}$ admits an explicit gateway decomposition (Theorem 3.1): with $A = 1$ if $p \equiv 3 \pmod{4}$; with $A = 3$ or $A = 7$ and an explicitly constructed d if $p \equiv 1 \pmod{4}$ lies in one of the algebraic classes of Section 4; and otherwise—for the residual class—with a prime $A \leq 359$, $A \equiv 3 \pmod{4}$, and a divisor d of $\left(\frac{p+A}{4}\right)^2$ satisfying $d \equiv -p^2 \cdot 4^{-1} \pmod{A}$, found by computational search.*

Since (1) for composite n reduces to the prime case [5], Theorem 1.1 implies that (1) holds for all $n \leq 10^{11}$.

The paper is organised as follows. Section 2 fixes notation and recalls the Type I/Type II solution classification of Elsholtz and Tao. Section 3 states and proves the gateway decomposition, noting its equivalence to the formulation of [11]. Section 4 proves that suitable parameters exist

for all primes outside a residual class of relative density zero (over 97% of primes up to any fixed bound in our range), and relates this residual class to—while carefully distinguishing it from—the classical Mordell subset. Section 5 presents the computational verification for the remaining primes, including A-value distribution data. Section 6 states the bounded- A conjecture, provides heuristic support from the Elsholtz–Tao density estimates, and discusses the connection to divisor equidistribution.

2. PRELIMINARIES

Throughout, p denotes an odd prime. For odd A with $4 \mid (p + A)$, we write

$$N = \frac{p + A}{4}, \quad B = pN = \frac{p(p + A)}{4}.$$

Note that $A = 4N - p$, so $\frac{4}{p} - \frac{1}{N} = \frac{A}{B}$. Thus (1) with $x = N$ reduces to decomposing $\frac{A}{B}$ as a sum of two unit fractions:

$$(2) \quad \frac{A}{B} = \frac{1}{y} + \frac{1}{z}.$$

The standard factorisation trick for (2) is: if $(Ay - B)(Az - B) = B^2$, then y and z satisfy (2). Setting $\delta_1 = Ay - B$ and $\delta_2 = Az - B$, we need $\delta_1 \delta_2 = B^2$ with $\delta_1 \equiv \delta_2 \equiv -B \pmod{A}$.

We use $v_q(m)$ for the q -adic valuation of m , and $\tau(m)$ for the number of positive divisors of m .

3. THE GATEWAY DECOMPOSITION

Theorem 3.1 (Gateway Decomposition). *Let p be a prime, A a positive odd integer with $4 \mid (p + A)$. Set $N = \frac{p+A}{4}$ and $B = pN$. Suppose there exists a positive integer d such that*

- (i) $A \mid (B + d)$, and
- (ii) $d \mid By$, where $y = (B + d)/A$.

Then

$$(3) \quad x = N, \quad y = \frac{B + d}{A}, \quad z = \frac{By}{d}$$

are positive integers, and $\frac{4}{p} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}$.

Proof. Integrality of x : immediate from $4 \mid (p + A)$. Integrality of y : immediate from (i). Integrality of z : immediate from (ii). Positivity: clear since $p, A, d > 0$.

For the identity, observe

$$\frac{1}{y} + \frac{1}{z} = \frac{1}{y} + \frac{d}{By} = \frac{B+d}{By} = \frac{Ay}{By} = \frac{A}{B},$$

using $B+d = Ay$ from (i). Since $A = 4N - p$, we have $\frac{A}{B} = \frac{4N-p}{pN} = \frac{4}{p} - \frac{1}{N}$. Therefore $\frac{4}{p} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}$. $\square$

Figure 1 illustrates the structure of the decomposition.




FIGURE 1. Schematic of the gateway decomposition. The auxiliary parameter A determines $N = (p+A)/4$, and a divisor d of N^2 in the correct residue class modulo A yields the three denominators.

The power of Theorem 3.1 lies in identifying *when* a suitable d exists. The following lemma provides a sufficient condition that governs our computational search.

Lemma 3.2 (Divisor condition). *In the setting of Theorem 3.1, suppose A is prime, $\gcd(d, pA) = 1$, and $d \mid N^2$. Then condition (ii) holds.*

Proof. We must show $d \mid By$ where $y = (B+d)/A$ and $B = pN$. Since $A \mid (B+d)$, write $y = (pN+d)/A$, so $By = pNy = pN(pN+d)/A$. We verify $d \mid pNy$ prime-by-prime.

Let $q^a \parallel d$. Since $\gcd(d, pA) = 1$, we have $q \neq p$ and $q \neq A$. Write $\alpha = v_q(N)$. From $d \mid N^2$, we get $a \leq 2\alpha$.

- If $\alpha \geq a$: then $q^a \mid N$, so $q^a \mid pNy$. Done.
- If $\alpha < a$: then $\alpha \geq 1$ (since $a \leq 2\alpha$ and $a > \alpha$ gives $\alpha \geq 1$). We have $v_q(pN) = \alpha$ (since $q \neq p$) and $v_q(d) = a > \alpha$, so $v_q(pN+d) = \alpha$. Then $v_q(y) = v_q((pN+d)/A) = \alpha$ (since $q \neq A$). Therefore $v_q(pNy) = \alpha + \alpha = 2\alpha \geq a$. Done.

$\square$

Remark 3.3. The condition $d \mid N^2$ is *strictly weaker* than $d \mid N$, and this distinction is essential. When $\gcd(d, p) > 1$ (e.g., $d = p$ for the identity in Proposition 4.2(b)), the lemma does not apply, but condition (ii) may still hold directly.

Example 3.4. Let $p = 2521$ and $A = 23$. Then $N = (2521 + 23)/4 = 636 = 2^2 \cdot 3 \cdot 53$, and $N^2 = 404,496 = 2^4 \cdot 3^2 \cdot 53^2$.

Take $d = 848 = 2^4 \cdot 53$. Then $d \mid N^2$ (since $v_2(d) = 4 \leq 2 \cdot 2 = v_2(N^2)$ and $v_{53}(d) = 1 \leq 2 = v_{53}(N^2)$), but $d \nmid N$ (since $v_2(d) = 4 > 2 = v_2(N)$).

Check (i): $B = 2521 \cdot 636 = 1,603,356$. $B + d = 1,604,204 = 23 \cdot 69,748$. ✓

So $x = 636$, $y = 69,748$, $z = 1,603,356 \cdot 69,748 / 848 = 131,876,031$.

Verify: $\frac{4}{2521} = \frac{1}{636} + \frac{1}{69748} + \frac{1}{131876031}$. ✓

This solution *cannot* be found under the condition $d \mid N$.

Remark 3.5 (Equivalence to condition (i)). Since $B = pN$ and $A = 4N - p$, we have $B \equiv pN \pmod{A}$. Now $p \equiv 4N - A \equiv 4N \pmod{A}$, so $B \equiv 4N^2 \pmod{A}$. Thus condition (i) becomes

$$d \equiv -4N^2 \pmod{A},$$

or equivalently (since $\gcd(4, A) = 1$), $d \equiv -p^2 \cdot 4^{-1} \pmod{A}$ (using $p \equiv 4N \pmod{A}$ gives $p^2 \equiv 16N^2 \pmod{A}$, so $-p^2 \cdot 4^{-1} \equiv -4N^2 \pmod{A}$).

Remark 3.6 (Relation to prior formulations). Vaughn [11] independently formulated the Erdős–Straus problem as: for prime p , find x with $\lceil p/4 \rceil \leq x \leq \lceil p/2 \rceil$ and a divisor $d \mid x^2$ satisfying $d \equiv -px \pmod{4x - p}$. Setting $A = 4x - p$ recovers our parameterisation: $x = N = (p + A)/4$, $4x - p = A$, and the divisor condition matches (i) via Remark 3.5. The two formulations are thus algebraically equivalent. Our contribution is not the identity itself, but the observation that the parameter A stays remarkably small—the bounded- A phenomenon (Section 6).

4. EXISTENCE RESULTS

We show that a suitable d exists for specific classes of primes, covering all but a residual class of relative density zero (Remark 4.4); at 10^6 the three propositions cover 97.1% of primes, rising towards 100% with the bound.

Throughout this section we use the observation that for $A = 3$ and $p \equiv 1 \pmod{4}$, $p \neq 3$: from $4N = p + 3$ we get $N \equiv p \pmod{3}$, hence

$$(4) \quad B = pN \equiv p^2 \equiv 1 \pmod{3},$$

so condition (i) asks for $d \equiv -1 \equiv 2 \pmod{3}$.

Proposition 4.1. *Every prime $p \equiv 3 \pmod{4}$ satisfies the Erdős–Straus conjecture.*

Proof. This is classical. Set $A = 1$, $N = (p+1)/4$ (an integer since $p \equiv 3 \pmod{4}$), and $B = pN$. Then $\frac{4}{p} - \frac{1}{N} = \frac{1}{B}$, and $\frac{1}{B} = \frac{1}{B+1} + \frac{1}{B(B+1)}$. So

$$\frac{4}{p} = \frac{1}{(p+1)/4} + \frac{1}{p(p+1)/4+1} + \frac{1}{p(p+1)(p(p+1)/4+1)/4}. \quad \square$$

Proposition 4.2. *Let $p \equiv 1 \pmod{4}$ be prime. Set $A = 3$, $N = (p+3)/4$, $B = pN$.*

- (a) *If $p \equiv 5 \pmod{8}$, then N is even and $d = 2$ gives a valid gateway decomposition.*
- (b) *If $p \equiv 17 \pmod{24}$, then $d = p$ gives a valid gateway decomposition.*
- (c) *If $p \equiv 1 \pmod{24}$ and N has a prime factor $q \equiv 2 \pmod{3}$, then $d = q$ gives a valid gateway decomposition.*

Proof. In all cases, $A = 3$, and $4 \mid (p+3)$ since $p \equiv 1 \pmod{4}$.

(a): $p \equiv 5 \pmod{8}$ gives $N = (p+3)/4$ even, so $2 \mid N$ and hence $2 \mid N^2$. Condition (i) holds since $d = 2$ and, by (4), $B+2 \equiv 1+2 \equiv 0 \pmod{3}$ — note this uses only $p \equiv 1 \pmod{4}$, $p \neq 3$, and covers both residues of p modulo 3. Lemma 3.2 gives (ii).

(b): Here $d = p$ does not divide N^2 (since $\gcd(p, N) = 1$), so Lemma 3.2 does not apply. Instead we verify (ii) directly. We have $p \equiv 17 \pmod{24}$, so $p \equiv 2 \pmod{3}$ and $6 \mid (p+1)$.

Condition (i): $B + p = pN + p = p(N+1) = p(p+7)/4$. Since $p \equiv 17 \pmod{24}$ implies $p \equiv 2 \pmod{3}$, we have $p+7 \equiv 0 \pmod{3}$, so $3 \mid (p+7)$ and hence $3 \mid p(p+7)/4 = B+p$. ✓

Condition (ii): $y = (B+p)/3 = p(p+7)/12$ and $By = pN \cdot p(p+7)/12 = p^2(p+3)(p+7)/48$, so $z = By/p = p(p+3)(p+7)/48$. Integrality of z : $p \equiv 17 \pmod{24}$ gives $24 \mid (p+7)$ and $4 \mid (p+3)$, so $48 \mid (p+3)(p+7)$. ✓

(c): q is a prime factor of $N = (p+3)/4$ with $q \equiv 2 \pmod{3}$. Since $q \mid N$, we have $q \mid N^2$ and $\gcd(q, 3p) = 1$ (as $q \neq 3$ since $N \equiv 1 \pmod{3}$ for $p \equiv 1 \pmod{24}$, and $q \neq p$ since $q \mid N$ and $\gcd(p, N) = 1$). For (i): $B + q = pN + q$. Since $p \equiv 1 \pmod{3}$ and $N \equiv 1 \pmod{3}$ (from $p \equiv 1 \pmod{24}$), $B \equiv 1 \pmod{3}$. Since $q \equiv 2 \pmod{3}$, $B + q \equiv 0 \pmod{3}$. ✓

Lemma 3.2 gives (ii). □

Proposition 4.3. *Let $p \equiv 1 \pmod{24}$ be a prime with every prime factor of $(p+3)/4$ congruent to 1 (mod 3) (“Case B”). If p is a quadratic non-residue (mod 7) ($p \equiv 3, 5$, or $6 \pmod{7}$), then the gateway decomposition with $A = 7$ yields a valid representation.*

Proof. Set $A = 7$, $N = (p+7)/4$, $B = pN$. Since $p \equiv 1 \pmod{24}$ implies $p \equiv 1 \pmod{8}$, we have $8 \mid (p+7)$, so $v_2(N) \geq 1$.

We apply Theorem 3.1 directly (not via Lemma 3.2, since our d will not be coprime to p). First, $B \equiv 2p^2 \pmod{7}$ (using $4^{-1} \equiv 2 \pmod{7}$), and both $2 \equiv 3^2$ and p^2 are QRs mod 7, so B is a QR and $-B$ is a NQR mod 7. Take $d = 2^k p$ with $k \in \{0, 1, 2\}$ chosen so that $2^k p \equiv -B \pmod{7}$. This is always possible: $\{2^0, 2^1, 2^2\} \equiv \{1, 2, 4\} \pmod{7}$ are the three QRs, and p is a NQR mod 7 by hypothesis, so $\{p, 2p, 4p\}$ runs through all three NQR classes mod 7.

Condition (i) holds by choice of k , so $y = (B+d)/7 = p(N+2^k)/7 \in \mathbb{Z}$.

For condition (ii), we verify $d = 2^k p \mid By$ directly:

- $p \mid By$: $By = pN \cdot p(N+2^k)/7$, so $v_p(By) \geq 2 > 1 = v_p(d)$.
- $2^k \mid By$: $v_2(B) = v_2(N) \geq 1$ and $v_2(y) = v_2(N+2^k)$. For $k = 0$: trivial. For $k = 1$: $v_2(N) \geq 1$ alone suffices. For $k = 2$: if $v_2(N) = 1$, then $N \equiv 2 \pmod{4}$ so $v_2(N+4) = 1$ and $v_2(By) = 1+1 = 2$; if $v_2(N) \geq 2$, then $v_2(By) \geq 2$ directly from $v_2(B) \geq 2$.

In every case $v_2(By) \geq k$ and $v_p(By) \geq 1$, so $d \mid By$. ✓

Theorem 3.1 now yields the explicit decomposition

$$x = \frac{p+7}{4}, \quad y = \frac{p(p+7+2^{k+2})}{28}, \quad z = \frac{By}{d}.$$

□

Remark 4.4. Propositions 4.1–4.3 together cover all primes except those satisfying simultaneously: $p \equiv 1 \pmod{24}$, every prime factor of $(p+3)/4$ is $\equiv 1 \pmod{3}$ (Case B), and $p \equiv 1, 2$, or $4 \pmod{7}$. We call these *Case B QR7* primes. At 10^6 they constitute 2.89% of all primes (2,269 of 78,498), and their relative density decreases with the bound (2.38% at 10^9 , 2.14% at 10^{11}): the Case-B condition forces every prime factor of $(p+3)/4$ into the residue class 1 (mod 3), and by Landau-type counting the integers with this property have counting function $\sim cx/\sqrt{\log x}$, so the residual class has relative density zero among the primes. For these primes we apply Theorem 3.1 with Lemma 3.2 via a computational search over prime values of A .

The residual class is related to, but *not* identical with, the classical *Mordell subset* $\{p : p \bmod 840 \in \{1, 121, 169, 289, 361, 529\}\}$ (the

squares of units modulo 840, equivalently $p \equiv 1 \pmod{24}$ with p a quadratic residue modulo both 5 and 7), which is the residual class of the classical system of modular identities. Both sets share the congruence backbone $p \equiv 1 \pmod{24}$ and $\left(\frac{p}{7}\right) = 1$, but they differ in two ways. First, our propositions make no use of the modulus 5: within the gateway family, the constraint $4 \mid (p + A)$ forces $A \equiv 3 \pmod{4}$ for $p \equiv 1 \pmod{4}$, which excludes $A = 5$, so the classical mod-5 identities do not arise. Thus $p = 193$ (a quadratic non-residue mod 5, hence outside the Mordell subset and covered by a classical mod-5 identity) is nevertheless a Case B QR7 prime and goes to our computational search. Second, the Case-B condition is multiplicative rather than a congruence condition: $p = 3361 \equiv 1 \pmod{840}$ lies in the Mordell subset, yet $(p + 3)/4 = 841 = 29^2$ with $29 \equiv 2 \pmod{3}$ makes it Case A, so Proposition 4.2(c) handles it algebraically. No system of congruence conditions can capture the residual class exactly. As Tao [10] observed, quadratic reciprocity prevents covering systems from eliminating the Mordell classes; the gateway decomposition circumvents this obstruction by using divisor enumeration on N^2 rather than congruence-based identities.

5. COMPUTATIONAL RESULTS

For the remaining Case B QR7 primes, we apply the gateway decomposition (Theorem 3.1) with a systematic search over prime values of A .

5.1. Method. For each prime p in the Case B QR7 class:

- (1) For each prime $A \equiv 3 \pmod{4}$ with $4 \mid (p + A)$, in increasing order:
- (2) Compute $N = (p + A)/4$ and factorise N .
- (3) Enumerate all divisors of N^2 (obtained by doubling the exponents in the prime factorisation of N).
- (4) For each divisor d of N^2 , check whether $d \equiv -B \pmod{A}$ where $B = pN$.
- (5) If so, compute $y = (B + d)/A$ and $z = By/d$; verify integrality and the identity $\frac{4}{p} = \frac{1}{N} + \frac{1}{y} + \frac{1}{z}$.
- (6) Record the first success and move to the next prime.

5.2. Results.

Theorem 5.1. *Every prime $p \leq 10^{11}$ admits a gateway decomposition with $A \leq 359$.*




FIGURE 2. Classification of primes to 10^6 . *Left:* the four-layer hierarchy; 97.1% of primes are handled by the algebraic results of Propositions 4.1–4.3. *Right:* zoom on the $p \equiv 1 \pmod{24}$ primes, showing the split between Case A, Case B NQR mod 7, and Case B QR mod 7.

The verified progression by scale (counts and $\max A$ from the corrected pipeline; see Remark 5.2):

Limit	Primes	Case B QR7 primes	Max A
10^6	78,498	2,269	79
10^7	664,579	18,019	167
10^8	5,761,455	146,227	239
10^9	50,847,534	1,212,383	239
10^{10}	455,052,511	10,246,512	359
10^{11}	4,118,054,813	88,088,687	359

Below 10^9 , exactly one prime requires $A > 199$: $p = 32,349,601$, solved by $A = 239$ with $d = 40,437,300$; the only other prime below 10^9 with $A > 191$ is $p = 597,188,089$ ($A = 199$). Below 10^{11} , exactly 47 primes require $A \geq 199$ (12 of them below 10^{10}), and only five require $A \geq 251$: $p = 3,807,728,761$ ($A = 359$, $d = 1,935$); $p = 31,048,497,481$ ($A = 347$); $p = 6,372,847,849$ ($A = 311$); $p = 26,147,464,729$ ($A = 251$); $p = 94,604,730,049$ ($A = 251$). The largest

gateway parameter over the entire range, $A = 359$, is attained below 10^{10} (at $p = 3,807,728,761$) and is not exceeded anywhere in $(10^{10}, 10^{11}]$.

Remark 5.2 (Data integrity). An earlier run of the 10^{10} stage (March 2026) contained an inverted Case A/Case B test: it skipped every residual-class prime as algebraically proven and instead searched the complementary class (Case A $\cap$ QR7), which Proposition 4.2(c) already covers with $A = 3$. All figures previously derived from that run (a residual count of 18,189,229 at 10^{10} , max $A = 251$, and a 25-value A -dictionary) described the wrong set of primes and have been withdrawn. The corrected pipeline validates against three independent baselines within the same run—the full A -distribution at 10^6 , the residual count at 10^7 , and the residual count and max A at 10^9 —before extending the range; every solution it reports is verified by direct evaluation of the identity (1). The full referee record accompanies the code repository.

The extension from $d \mid N$ to $d \mid N^2$ is not a cosmetic generalisation: a nontrivial proportion of Case B QR7 solutions use a divisor $d > N$ that would be missed under the stronger condition. Figure 3 shows the distribution of d/N ratios for Case B QR7 solutions.




FIGURE 3. Distribution of d/N ratios for Case B QR7 solutions at 10^6 . Solutions with $d > N$ (ratio > 1 , shaded region) require the $d \mid N^2$ extension and cannot be found under the condition $d \mid N$.

5.3. Distribution of A values. Among the 88,088,687 Case B QR7 primes below 10^{11} :

A	Count	% solved	Cumulative
7	60,704,131	68.9%	68.9%
11	19,439,771	22.1%	91.0%
19	4,447,883	5.0%	96.0%
23	2,486,127	2.8%	98.9%
31	688,365	0.8%	99.6%
43–191	322,363	0.37%	99.99995%
≥ 199	47	0.00005%	100%

The extreme sparsity of large- A cases—only 47 primes out of 88 million require $A \geq 199$, and only five require $A \geq 251$ —supports the bounded- A conjecture. Figure 4 shows the distribution at 10^6 .




FIGURE 4. Distribution of gateway parameter A for Case B QR7 primes at 10^6 . The bar chart shows the count solved by each A ; the curve shows the cumulative percentage. A handful of small primes ($A = 7, 11, 19, 23$) handle over 97% of cases.

6. DISCUSSION

6.1. The bounded- A conjecture. The maximum A value grows extremely slowly with scale:

Limit	max A	Increase from previous
10^6	79	—
10^7	167	+88
10^8	239	+72
10^9	239	+0
10^{10}	359	+120
10^{11}	359	+0

Every prime below 10^9 is solved by some $A \leq 239$, and only two primes below 10^9 require $A > 191$. At 10^{10} the maximum rises to 359; it then holds at 359 through 10^{11} , so across the last decade max A does not grow at all. Over $10^6 \rightarrow 10^{10}$ the growth of max A is roughly linear in $\log N$, and the flat step to 10^{11} is consistent with an eventual absolute bound.

Conjecture 6.1 (Bounded gateway parameter). There exists an absolute constant C such that for every prime p , there is a prime $A \leq C$ with $A \equiv 3 \pmod{4}$, $4 \mid (p + A)$, and a divisor d of $\left(\frac{p+A}{4}\right)^2$ satisfying $d \equiv -B \pmod{A}$, where $B = p(p + A)/4$.

If Conjecture 6.1 holds, the Erdős–Straus conjecture follows, since the conjecture can be verified for all primes up to C by direct computation, and every larger prime admits a gateway with $A \leq C$.

A weaker but still significant form: $\max A(N) = O(\log \log p)$. Even $\max A = o(\log p)$ would go beyond current results.




FIGURE 5. Maximum gateway parameter A required as the verification limit increases from 10^3 to 10^{11} . The growth is roughly logarithmic in the limit, in contrast with the factor-of- 10^8 increase in the search range.

6.2. Heuristic support from Elsholtz–Tao density. Elsholtz and Tao [2] showed that the average number of solutions satisfies

$$N \log^2 N \ll \sum_{p \leq N} f(p) \ll N \log^2 N \log \log N,$$

so the average number of solutions per prime up to N is $\Theta((\log N)^3)$.

For a fixed prime A , the number of divisors of $N^2 = \left(\frac{p+A}{4}\right)^2$ is $\tau(N^2) = \prod_i (2e_i + 1)$ where $N = \prod p_i^{e_i}$. On average, $\tau(N^2)$ grows like $(\log N)^c$ for a constant $c > 0$. Under a uniform distribution heuristic, the probability that none of these divisors falls in the target residue class $-B \pmod{A}$ is approximately

$$\left(1 - \frac{1}{A}\right)^{\tau(N^2)} \rightarrow 0 \quad \text{as } N \rightarrow \infty.$$

This suggests that for any fixed A , the set of primes for which A fails has density zero—consistent with Tao’s “almost all” result. For multiple values of A (the 31 distinct gateway values that occur below 10^{11}), failure requires *simultaneous* failure across all A , which is heuristically negligible but not ruled out by the density argument alone.

The Elsholtz–Tao bounds explain why small A dominates: with $\tau(N^2) \sim (\log p)^c$ divisors and a target residue class of density $1/A$, the expected number of valid divisors is $\sim \tau(N^2)/A$, which is larger for smaller A . The observed distribution—68.9% at $A = 7$, 91.0% by $A = 11$ —is consistent with this heuristic.

6.3. Comparison with the Salez modular sieve. The Salez algorithm [6], extended by Mihnea and Bogdan [4] to 10^{18} , takes a fundamentally different approach. It constructs modular filters S_m —sets of residue classes modulo m for which the conjecture is provably true—and only checks primes outside these filters. At filter level R_8 (modulus $G_8 \approx 2.6 \times 10^{10}$), only ≈ 2.1 million residue classes of G_8 require direct verification, representing $\approx 0.008\%$ of all primes.

	Salez sieve	Gateway decomposition
Question	Existence	Structure
Method	Modular filter + spot-check	Classification by A
Output per prime	Yes/no	Explicit (x, y, z) with A
Verification record	10^{18}	10^{11}
Reveals A-distribution	No	Yes

The two approaches are complementary. The Salez sieve reaches much higher limits by avoiding per-prime algebraic work; our approach produces the structural data underlying the bounded- A conjecture.

6.4. The quadratic reciprocity obstruction. Tao [10] observed that quadratic reciprocity blocks any system of covering congruences from resolving the Mordell subset: the residue classes $\{1, 121, 169, 289, 361, 529\}$ modulo 840 involve perfect-square residues that cannot be eliminated by further modular identities. Extending the mod-840 coverage to a complete covering system therefore has a fundamental ceiling.

The gateway decomposition circumvents this obstruction entirely. Rather than seeking a congruence condition that *all* primes in a residue class satisfy, it finds a divisor of N^2 in a specific residue class—a condition that depends on the *arithmetic* of each individual prime, not on its residue class alone. This is why the gateway can (empirically) handle cases that covering systems cannot.

6.5. The precise open problem. Since every prime outside the residual class is covered by Propositions 4.1–4.3, the Erdős–Straus conjecture reduces to:

For every Case B QR7 prime p , there exists a prime $A \equiv 3 \pmod{4}$ and a divisor d of $\left(\frac{p+A}{4}\right)^2$ satisfying $d \equiv -p \cdot \frac{p+A}{4} \pmod{A}$.

This is a question about the *equidistribution of divisors in residue classes* for integers of the form $\left(\frac{p+A}{4}\right)^2$, a topic connected to work of Hooley and Tenenbaum on divisor distribution. The bounded- A conjecture strengthens this to ask whether A can be taken from a *fixed finite set*, independent of p .

6.6. Connection to other recent work. Ghermoul [3] constructs four explicit polynomial families in three variables that conjecturally cover all integers $\equiv 1 \pmod{4}$, with one family alone covering all primes to 1.2×10^{10} . This is related to our framework: each A defines a polynomial family, and our observation that 32 values of A suffice to 10^{11} is a quantitative version of his coverage conjecture.

Our gateway decomposition produces predominantly Type II solutions in the Elsholtz–Tao classification (where p divides two of the three denominators). The structural reason is that y and z in Theorem 3.1 satisfy $y = (B + d)/A$ and $z = By/d$ with $B = pN$, so $p \mid z$ automatically whenever $p \nmid d$.

7. CONCLUSION

We have presented a structural analysis of the Erdős–Straus conjecture through a gateway decomposition parameterised by a single integer A . Using 32 prime values of A (all at most 359) for the primes $p \equiv 1 \pmod{4}$, together with the classical $A = 1$ decomposition for $p \equiv 3 \pmod{4}$, we verify that every prime $p \leq 10^{11}$ admits an explicit decomposition $\frac{4}{p} = \frac{1}{x} + \frac{1}{y} + \frac{1}{z}$.

The main observation is the slow growth of the gateway parameter: $\max A$ increases roughly logarithmically in the verification limit (79 at 10^6 , 239 at 10^8 , 359 at 10^{10}) and then holds at 359 through 10^{11} , while the residual class itself thins out (relative density $\sim c/\sqrt{\log x}$). If $\max A$ is absolutely bounded (Conjecture 6.1), the full conjecture follows.

The framework provides a potential bridge between computational verification and a theoretical proof: the bounded- A conjecture is a concrete statement about divisor equidistribution in residue classes for structured quadratic integers, a topic accessible to the methods of analytic number theory.

Code availability. All verification code (Python) and reproducibility data are available at <https://github.com/m4cd4r4/tough-math/tree/master/erdos-straus>.

REFERENCES

- [1] C. Elsholtz, *Sums of k unit fractions*, Trans. Amer. Math. Soc. **353** (2001), 3209–3227.
- [2] C. Elsholtz and T. Tao, *Counting the number of solutions to the Erdős–Straus equation on unit fractions*, J. Aust. Math. Soc. **94** (2013), 50–105.
- [3] B. Ghermoul, *Exact polynomial families solving the Erdős–Straus equation*, preprint, 2025. [arXiv:2508.07383](https://arxiv.org/abs/2508.07383).
- [4] S. Mihnea and D. C. Bogdan, *Further verification and empirical evidence for the Erdős–Straus conjecture*, preprint, 2025. [arXiv:2509.00128](https://arxiv.org/abs/2509.00128).
- [5] L. J. Mordell, *Diophantine Equations*, Academic Press, 1969.
- [6] S. E. Salez, *The Erdős–Straus conjecture: new modular equations and checking up to $N = 10^{17}$* , preprint, 2014. [arXiv:1406.6307](https://arxiv.org/abs/1406.6307).
- [7] A. Schinzel, *Sur quelques propriétés des nombres $3/n$ et $4/n$, où n est un nombre impair*, Mathesis **65** (1956), 219–222.
- [8] W. Sierpiński, *Sur les décompositions de nombres rationnels en fractions primaires*, Mathesis **65** (1956), 16–32.
- [9] A. Swett, *Erdős–Straus conjecture*, unpublished computation, 1999. <http://math.uindy.edu/swett/esc.htm>
- [10] T. Tao, *Erdős–Straus conjecture*, blog post, 2011. <https://terrytao.wordpress.com/tag/erdos-straus-conjecture/>

- [11] R. Vaughn, *Elemental patterns from the Erdős–Straus conjecture*, preprint, 2024. [arXiv:2403.16047](#).
- [12] W. A. Webb, *On $4/n = 1/x + 1/y + 1/z$* , Proc. Amer. Math. Soc. **25** (1970), 578–584.