

Lattice

枫聆

2021 年 3 月 11 日

目录

1	The Elements of Universal Algebra	2
1.1	Definition and Examples of Algebras	2
1.2	Isomorphic Algebras and Subalgebras	4
1.3	Algebraic Lattices and Subuniverses	5

The Elements of Universal Algebra

Definition and Examples of Algebras

One of the aims of universal algebra is to extract, whenever possible, the common elements of several seemingly different types of algebraic structures.

Definition 1.1. For A a nonempty set and n a nonnegative integer we define $A^0 = \{\emptyset\}$, and, for $n > 0$, A^n is the set of n -tuples of elements from A . An **n-ary operation** (or function) on A is any function f from A^n to A ; n is the **arity** (or rank) of f . A **finitary operation** is an n -ary operation, for some n . The image of $\langle a_1, \dots, a_n \rangle$ under an n -ary operation f is denoted by $f(a_1, \dots, a_n)$. An operation f on A is called a **nullary operation** (or constant) if its arity is zero; it is completely determined by the image $f(\emptyset)$ in A of the only element \emptyset in A^0 , and as such it is convenient to identify it with the element $f(\emptyset)$. Thus a nullary operation is thought of as an element of A . An operation f on A is **unary, binary, or ternary** if its arity is 1, 2, or 3, respectively.

Definition 1.2. A **language** (or type) of algebras is a set \mathcal{F} of function symbols such a nonnegative integer n is assigned to each member f of \mathcal{F} . This integer is called the arity (or rank) of f , and f is said to be an n -ary function symbol. The subset of n -ary function symbols in \mathcal{F} is denoted by \mathcal{F}_n .

Definition 1.3. If \mathcal{F} is a language of algebras then an **algebra** \mathbf{A} of type \mathcal{F} is an ordered pair $\langle A, F \rangle$ where A is a nonempty set and F is a family of finitary operations on A indexed by the language \mathcal{F} such that corresponding to each n -ary function symbol f in \mathcal{F} there is an n -ary operation $f^{\mathbf{A}}$ on A . The set A is called the **universe** (全域) (or underlying set) of $\mathbf{A} = \langle A, F \rangle$, and the $f^{\mathbf{A}}$'s are called the **fundamental operations** (基本运算) of \mathbf{A} . (In practice we prefer to write just f for $f^{\mathbf{A}}$ —this convention creates an ambiguity which seldom causes a problem. However, in this chapter we will be unusually careful.) If \mathcal{F} is finite, say $\mathcal{F} = \{f_1, \dots, f_k\}$, we often write $\langle A, f_1, \dots, f_k \rangle$ for $\langle A, F \rangle$, usually adopting the convention:

$$\text{arity} f_1 \geq \dots \geq \text{arity} f_k.$$

An algebra \mathbf{A} is **unary** if all of its operations are unary, and it is **mono-unary** if it has just one unary operation.

抽象来说一个 algebra 就是一个集合和一堆 operations 构成的, 在目前已经学到的代数中 operation 的 arity 大多数不会超过 2(够 modern).

Example 1.4. \mathbf{A} is a **groupoid** if it has just one binary operation; this operation is usually denoted by $+$ or \cdot , and we write $a + b$ or $a \cdot b$ for the image of $\langle a, b \rangle$ under this operation, and call it the sum or product of a and b , respectively.

Definition 1.5. An algebra \mathbf{A} is **finite** if $|\mathbf{A}|$ is finite, and **trivial** if $|\mathbf{A}| = 1$.

Example 1.6. Some well-known algebras

- Group
- Semigroup (半群) and Monoid (么半群)
- ...

Isomorphic Algebras and Subalgebras

Definition 1.7. Let \mathbf{A} and \mathbf{B} be two algebras of the same type \mathcal{F} . Then a function $\alpha: A \rightarrow B$ is an **isomorphism** from \mathbf{A} to \mathbf{B} if α is one-to-one and onto, and for every n -ary $f \in \mathcal{F}$, and for $a_1, \dots, a_n \in A$, we have

$$\alpha f^{\mathbf{A}}(a_1, \dots, a_n) = f^{\mathbf{B}}(\alpha a_1, \dots, \alpha a_n).$$

We say \mathbf{A} is isomorphic to B , if there is a isomorphism from \mathbf{A} to \mathbf{B} .

老样子元素上保持 bijective, 对应的运算结果也保持一致.

Definition 1.8. Let \mathbf{A} and \mathbf{B} be two algebras of the same type. Then \mathbf{B} is a **subalgebra** of \mathbf{A} if $B \subseteq A$ and every fundamental operation of \mathbf{B} is the restriction of the corresponding operation of \mathbf{A} , i.e., for each function symbol f , $f^{\mathbf{B}}$ is $f^{\mathbf{A}}$ restricted to B ; we write simply $\mathbf{B} \leq \mathbf{A}$.

A **subuniverse** of \mathbf{A} is a subset B of A which is closed under the fundamental operations of \mathbf{A} , i.e., if f is a fundamental n -ary operation of \mathbf{A} and $a_1, \dots, a_n \in B$ we would required $f(a_1, \dots, a_n) \in B$.

这个 restriction 就是只对定义域做限制的意思, subalgebra 是一种构造新代数的方法, 后面会学到另外几种.

Definition 1.9. Let \mathbf{A} and \mathbf{B} be of the same type. A function $\alpha: A \rightarrow B$ is an embedding of \mathbf{A} into \mathbf{B} if α is one-to-one and satisfies

$$\alpha f^{\mathbf{A}}(a_1, \dots, a_n) = f^{\mathbf{B}}(\alpha a_1, \dots, \alpha a_n).$$

Such an α is also called a monomorphism. We say \mathbf{A} can be embedded in \mathbf{B} if there is an embedding of \mathbf{A} into \mathbf{B} .

单纯地去掉 surjective.

Theorem 1.10. If $\alpha: A \rightarrow B$ is an embedding, then $\alpha(A)$ is a subuniverse of \mathbf{B} .

证明. 我们需要说 $\alpha(A)$ 在 n -ary operation 下保持封闭. 因为 α 是 embedding, 所以对应一个 n -ary operation f 和 $a_1, \dots, a_n \in A$ 有

$$f^{\mathbf{B}}(\alpha a_1, \dots, \alpha a_n) = \alpha f^{\mathbf{A}}(a_1, \dots, a_n) \in \alpha(A).$$

已经证闭. □

Algebraic Lattices and Subuniverses

这一章阐述 algebraic lattice 出现在 universe algebra 的原因.

Definition 1.11. Given an algebra \mathbf{A} define, for every $X \subseteq A$,

$$\text{Sg}(X) = \bigcap \{ B \subseteq A \mid X \subseteq B \text{ and } B \text{ is a subuniverse of } \mathbf{A} \}.$$

We read $\text{Sg}(X)$ as "the subuniverse generated by X ".

Definition 1.12. A closure operator C on the set A is an algebraic closure operator if for $X \subseteq A$

$$C(X) = \bigcup \{ C(Y) \mid Y \subseteq X \text{ and } Y \text{ is finite} \}.$$

Theorem 1.13. If we are given an algebra \mathbf{A} , then Sg is an algebraic closure operator on A .

证明. 很明显任意地 subuniverses 交还是一个 subuniverse, 所有的 subuniverses 构成了一个 closure system, 所以 Sg 是一个 closure operator. 对于任意的 $X \subseteq A$ 我们定义

$$E(X) = X \cup \{ f(a_1, \dots, a_n) \mid f \text{ is a fundamental } n\text{-ary operation on } A \text{ and } a_1, \dots, a_n \in X \}.$$

然后定义它的 n 次复合 $E^n(X)$ 为

$$E^0(X) = X$$

$$E^{n+1}(X) = E(E^n(X)).$$

由于 A 上所有 fundamental operation 都是 finitary, 且有

$$X \subseteq E(X) \subseteq \dots \subseteq E^n(X).$$

接下来我们来证明下面的式子

$$\text{Sg}(X) = X \cup E(X) \cup E^2(X) \cup \dots.$$

思路是 $\text{Sg}(X) \subseteq X \cup E(X) \cup E^2(X) \cup \dots$ 和 $X \cup E(X) \cup E^2(X) \cup \dots \subseteq \text{Sg}(X)$. □