

Genuine Cohomology Theories

Marin Janssen (she/they)

University of Münster

May 6, 2024

Let X be a topological space, CW-complex, simplicial set, anima, G -anima, manifold, ...

Let X be a topological space, CW-complex, simplicial set, anima, G -anima, manifold, ...

Invariants $F(X)$ of X , e.g.: $\chi(X)$, $\beta_i(X)$, $\pi_i(X)$, $H_i(X)$, $H^i(X)$,
..., $i \in \mathbb{Z}$

Let X be a topological space, CW-complex, simplicial set, anima, G -anima, manifold, ...

Invariants $F(X)$ of X , e.g.: $\chi(X)$, $\beta_i(X)$, $\pi_i(X)$, $H_i(X)$, $H^i(X)$,
..., $i \in \mathbb{Z}$

Functoriality: $f : X \rightarrow Y$ induces map $f^* : H^*(Y) \rightarrow H^*(X)$

Let X be a topological space, CW-complex, simplicial set, anima, G -anima, manifold, ...

Invariants $F(X)$ of X , e.g.: $\chi(X)$, $\beta_i(X)$, $\pi_i(X)$, $H_i(X)$, $H^i(X)$,
..., $i \in \mathbb{Z}$

Functoriality: $f : X \rightarrow Y$ induces map $f^* : H^*(Y) \rightarrow H^*(X)$

Homotopy Invariance: homotopic maps induce the same map in cohomology

Let X be a topological space, CW-complex, simplicial set, anima, G -anima, manifold, ...

Invariants $F(X)$ of X , e.g.: $\chi(X)$, $\beta_i(X)$, $\pi_i(X)$, $H_i(X)$, $H^i(X)$,
..., $i \in \mathbb{Z}$

Functoriality: $f : X \rightarrow Y$ induces map $f^* : H^*(Y) \rightarrow H^*(X)$

Homotopy Invariance: homotopic maps induce the same map in cohomology

Suspension: $\tilde{H}^{*+1}(\Sigma X) \simeq \tilde{H}^*(X)$

Let X be a topological space, CW-complex, simplicial set, anima, G -anima, manifold, ...

Invariants $F(X)$ of X , e.g.: $\chi(X)$, $\beta_i(X)$, $\pi_i(X)$, $H_i(X)$, $H^i(X)$,
..., $i \in \mathbb{Z}$

Functoriality: $f : X \rightarrow Y$ induces map $f^* : H^*(Y) \rightarrow H^*(X)$

Homotopy Invariance: homotopic maps induce the same map in cohomology

Suspension: $\tilde{H}^{*+1}(\Sigma X) \simeq \tilde{H}^*(X)$

Additivity: $H^*(\coprod X_\alpha) = \prod H^*(X_\alpha)$

Let X be a topological space, CW-complex, simplicial set, anima, G -anima, manifold, ...

Invariants $F(X)$ of X , e.g.: $\chi(X)$, $\beta_i(X)$, $\pi_i(X)$, $H_i(X)$, $H^i(X)$,
..., $i \in \mathbb{Z}$

Functoriality: $f : X \rightarrow Y$ induces map $f^* : H^*(Y) \rightarrow H^*(X)$

Homotopy Invariance: homotopic maps induce the same map in cohomology

Suspension: $\tilde{H}^{*+1}(\Sigma X) \simeq \tilde{H}^*(X)$

Additivity: $H^*(\coprod X_\alpha) = \prod H^*(X_\alpha)$

Mayer-Vietoris: $H^*(A \amalg_C B) \rightarrow H^*(A) \oplus H^*(B) \rightarrow H^*(C)$

Let X be a topological space, CW-complex, simplicial set, anima, G -anima, manifold, ...

Invariants $F(X)$ of X , e.g.: $\chi(X)$, $\beta_i(X)$, $\pi_i(X)$, $H_i(X)$, $H^i(X)$,
..., $i \in \mathbb{Z}$

Functoriality: $f : X \rightarrow Y$ induces map $f^* : H^*(Y) \rightarrow H^*(X)$

Homotopy Invariance: homotopic maps induce the same map in cohomology

Suspension: $\tilde{H}^{*+1}(\Sigma X) \simeq \tilde{H}^*(X)$

Additivity: $H^*(\coprod X_\alpha) = \prod H^*(X_\alpha)$

Mayer-Vietoris: $H^*(A \coprod_C B) \rightarrow H^*(A) \oplus H^*(B) \rightarrow H^*(C)$

Dimension: $H^i(\text{pt}) = \mathbb{Z}^{\delta_{i,0}}$

Let X be a topological space, CW-complex, simplicial set, anima, G -anima, manifold, ...

Invariants $F(X)$ of X , e.g.: $\chi(X)$, $\beta_i(X)$, $\pi_i(X)$, $H_i(X)$, $H^i(X)$,
..., $i \in \mathbb{Z}$

Eilenberg-Steenrod Axioms for cohomology theories:

Functoriality: $f : X \rightarrow Y$ induces map $f^* : H^*(Y) \rightarrow H^*(X)$

Homotopy Invariance: homotopic maps induce the same map in cohomology

Suspension: $\tilde{H}^{*+1}(\Sigma X) \simeq \tilde{H}^*(X)$

Additivity: $H^*(\coprod X_\alpha) = \prod H^*(X_\alpha)$

Mayer-Vietoris: $H^*(A \amalg_C B) \rightarrow H^*(A) \oplus H^*(B) \rightarrow H^*(C)$

Dimension: $H^i(\text{pt}) = \mathbb{Z}^{\delta_{i,0}}$

Let X be a topological space, CW-complex, simplicial set, anima, G -anima, manifold, ...

Invariants $F(X)$ of X , e.g.: $\chi(X)$, $\beta_i(X)$, $\pi_i(X)$, $H_i(X)$, $H^i(X)$,
..., $i \in \mathbb{Z}$

Eilenberg-Steenrod Axioms for cohomology theories:

Functoriality: $f : X \rightarrow Y$ induces map $f^* : H^*(Y) \rightarrow H^*(X)$

Homotopy Invariance: homotopic maps induce the same map in cohomology

Suspension: $\tilde{H}^{*+1}(\Sigma X) \simeq \tilde{H}^*(X)$

Additivity: $H^*(\coprod X_\alpha) = \prod H^*(X_\alpha)$

Mayer-Vietoris: $H^*(A \coprod_C B) \rightarrow H^*(A) \oplus H^*(B) \rightarrow H^*(C)$

Dimension: $H^i(\text{pt}) = \mathbb{Z}^{\delta_{i,0}}$

Let X be a topological space, CW-complex, simplicial set, anima, G -anima, manifold, ...

Invariants $F(X)$ of X , e.g.: $\chi(X)$, $\beta_i(X)$, $\pi_i(X)$, $H_i(X)$, $H^i(X)$, $K^i(X)$, $\pi_s^i(X)$, $\Omega_G^i(X)$, $K(n, p)^i(X)$, ..., $i \in \mathbb{Z}$

Eilenberg-Steenrod Axioms for cohomology theories:

Functoriality: $f : X \rightarrow Y$ induces map $f^* : H^*(Y) \rightarrow H^*(X)$

Homotopy Invariance: homotopic maps induce the same map in cohomology

Suspension: $\tilde{H}^{*+1}(\Sigma X) \simeq \tilde{H}^*(X)$

Additivity: $H^*(\coprod X_\alpha) = \prod H^*(X_\alpha)$

Mayer-Vietoris: $H^*(A \amalg_C B) \rightarrow H^*(A) \oplus H^*(B) \rightarrow H^*(C)$

Let X be a topological space, CW-complex, simplicial set, anima, G -anima, manifold, ...

Invariants $F(X)$ of X , e.g.: $\chi(X)$, $\beta_i(X)$, $\pi_i(X)$, $H_i(X)$, $H^i(X)$, $K^i(X)$, $\pi_s^i(X)$, $\Omega_G^i(X)$, $K(n, p)^i(X)$, ..., $i \in \mathbb{Z}$

Eilenberg-Steenrod Axioms for cohomology theories:

Functoriality: $f : X \rightarrow Y$ induces map $f^* : H^*(Y) \rightarrow H^*(X)$

Homotopy Invariance: homotopic maps induce the same map in cohomology

Suspension: $\tilde{H}^{*+1}(\Sigma X) \simeq \tilde{H}^*(X)$

Additivity: $H^*(\coprod X_\alpha) = \prod H^*(X_\alpha)$

Mayer-Vietoris: $H^*(A \amalg_C B) \rightarrow H^*(A) \oplus H^*(B) \rightarrow H^*(C)$

Let X be a topological space, CW-complex, simplicial set, anima, G -anima, manifold, ...

Invariants $F(X)$ of X , e.g.: $\chi(X)$, $\beta_i(X)$, $\pi_i(X)$, $H_i(X)$, $H^i(X)$, $K^i(X)$, $\pi_s^i(X)$, $\Omega_G^i(X)$, $K(n, p)^i(X)$, ..., $i \in \mathbb{Z}$

Eilenberg-Steenrod Axioms for (generalized) cohomology theories on compact anima:

Let X be a topological space, CW-complex, simplicial set, anima, G -anima, manifold, ...

Invariants $F(X)$ of X , e.g.: $\chi(X)$, $\beta_i(X)$, $\pi_i(X)$, $H_i(X)$, $H^i(X)$, $K^i(X)$, $\pi_s^i(X)$, $\Omega_G^i(X)$, $K(n, p)^i(X)$, ..., $i \in \mathbb{Z}$

Eilenberg-Steenrod Axioms for (generalized) cohomology theories on compact anima:

Functoriality: $f : X \rightarrow Y$ induces map $f^* : E^*(Y) \rightarrow E^*(X)$

Homotopy Invariance: homotopic maps induce the same map in cohomology

Suspension: $\tilde{E}^{*+1}(\Sigma X) \simeq \tilde{E}^*(X)$

Mayer-Vietoris: $E^*(A \coprod_C B) \rightarrow E^*(A) \oplus E^*(B) \rightarrow E^*(C)$

Let X be a topological space, CW-complex, simplicial set, anima, G -anima, manifold, ...

Invariants $F(X)$ of X , e.g.: $\chi(X)$, $\beta_i(X)$, $\pi_i(X)$, $H_i(X)$, $H^i(X)$, $K^i(X)$, $\pi_s^i(X)$, $\Omega_G^i(X)$, $K(n, p)^i(X)$, ..., $i \in \mathbb{Z}$

Eilenberg-Steenrod Axioms for (generalized) cohomology theories on compact anima:

Functor $E: \mathbf{Sp}^\omega \rightarrow \mathbf{Ab}^{op}$

Mayer-Vietoris: $E(A \coprod_C B) \rightarrow E(A) \oplus E(B) \rightarrow E(C)$ is exact

Motivation

Let M^n closed orientable manifold,

Motivation

Let M^n closed orientable manifold,
 H^* be singular cohomology,

Motivation

Let M^n closed orientable manifold,
 H^* be singular cohomology,
 H_* be singular homology, then

Motivation

Let M^n closed orientable manifold,

H^* be singular cohomology,

H_* be singular homology, then

Poincaré-Duality:

$$H_*(M) \simeq H^{n-*}(M)$$

Motivation

Let M^n closed **not necessarily** orientable manifold,

H^* be singular cohomology,

H_* be singular homology, then

Poincaré-Duality:

$$H_*(M) \simeq H^{n-*}(M)$$

Motivation

Let M^n closed **not necessarily** orientable manifold,

E^* be a generalized cohomology theory,

H_* be singular homology, then

Poincaré-Duality:

$$H_*(M) \simeq H^{n-*}(M)$$

Motivation

Let M^n closed **not necessarily** orientable manifold,

E^* be a generalized cohomology theory,

E_* associated homology theory, then

Poincaré-Duality:

$$H_*(M) \simeq H^{n-*}(M)$$

Motivation

Let M^n closed **not necessarily** orientable manifold,

E^* be a generalized cohomology theory,

E_* associated homology theory, then

Poincaré-Duality:

$$H_*(M) \simeq H^{n-*}(M)$$

Motivation

Let M^n closed **not necessarily** orientable manifold,
 $E^* \simeq [\Sigma_+^\infty(-), \mathcal{E}]_{-*}$ be a generalized cohomology theory,
 E_* associated homology theory, then

Poincaré-Duality:

$$H_*(M) \simeq H^{n-*}(M)$$

Motivation

Let M^n closed **not necessarily** orientable manifold,
 $E^* \simeq [\Sigma_+^\infty(-), \mathcal{E}]_{-*}$ be a generalized cohomology theory,
 $E_* := [\mathbb{S}, \Sigma_+^\infty(-) \otimes \mathcal{E}]_*$ associated homology theory, then
Poincaré-Duality:

$$H_*(M) \simeq H^{n-*}(M)$$

Motivation

Let M^n closed **not necessarily** orientable manifold,
 $E^* \simeq [\Sigma_+^\infty(-), \mathcal{E}]_{-*}$ be a generalized cohomology theory,
 $E_* := [\mathbb{S}, \Sigma_+^\infty(-) \otimes \mathcal{E}]_*$ associated homology theory, then
Poincaré-Duality:

$$E_*(M) \simeq E^{TM-*}(M)$$

Motivation

Let M^n closed **not necessarily** orientable manifold,
 $E^* \simeq [\Sigma_+^\infty(-), \mathcal{E}]_{-*}$ be a generalized cohomology theory,
 $E_* := [\mathbb{S}, \Sigma_+^\infty(-) \otimes \mathcal{E}]_*$ associated homology theory, then
Poincaré-Duality:

$$E_*(M) \simeq E^{TM-*}(M) := [M^{-TM}, \mathcal{E}]_*$$

Motivation

Let M^n closed **not necessarily** orientable manifold,
 $E^* \simeq [\Sigma_+^\infty(-), \mathcal{E}]_{-*}$ be a generalized cohomology theory,
 $E_* := [\mathbb{S}, \Sigma_+^\infty(-) \otimes \mathcal{E}]_*$ associated homology theory, then
Poincaré-Duality:

$$E_*(M) \simeq E^{TM-*}(M) := [M^{-TM}, \mathcal{E}]_*$$

Problem

Extend the ES-axioms to include the groups $E^{V-*}(X) = [X^{-V}, \mathcal{E}]_*$ where (X, V) run through all pairs of $X \in \mathbf{An}^\omega$ and V a vector bundle over X .

Motivation

Let M^n closed **not necessarily** orientable manifold,

$E^* \simeq [\Sigma_+^\infty(-), \mathcal{E}]_{-*}$ be a generalized cohomology theory,

$E_* := [\mathbb{S}, \Sigma_+^\infty(-) \otimes \mathcal{E}]_*$ associated homology theory, then

Poincaré-Duality:

$$E_*(M) \simeq E^{TM-*}(M) := [M^{-TM}, \mathcal{E}]_*$$

Problem

Extend the ES-axioms to include the groups $E^{V-*}(X) = [X^{-V}, \mathcal{E}]_*$ where (X, V) run through all pairs of $X \in \mathbf{An}^\omega$ and V a vector bundle over X .

Inspiration

$RO(G)$ -graded cohomology theories. Graded over G -representation which can be viewed as equivariant vector bundles over G/G .

Genuine Cohomology Theories

Definition

A **genuine Cohomology Theory** E consists of the following data

Genuine Cohomology Theories

Definition

A **genuine Cohomology Theory** E consists of the following data
for each pair (X, V) an abelian group $E^V(X)$,

Genuine Cohomology Theories

Definition

A **genuine Cohomology Theory** E consists of the following data

for each pair (X, V) an abelian group $E^V(X)$,

for each map $f: X \rightarrow Y$ of anima an induced map

$$E^W Y \xrightarrow{E(f)} E^{f^* W} X$$

Genuine Cohomology Theories

Definition

A **genuine Cohomology Theory** E consists of the following data

for each pair (X, V) an abelian group $E^V(X)$,

for each map $f: X \rightarrow Y$ of anima an induced map

$$E^W Y \xrightarrow{E(f)} E^{f^* W} X$$

for each affine linear map $\alpha: U \rightarrow V$ of VB over X an induced map

$$E^U X \xrightarrow{E(\alpha)} E^V X \text{ s.t.}$$

Genuine Cohomology Theories

Definition

A **genuine Cohomology Theory** E consists of the following data

for each pair (X, V) an abelian group $E^V(X)$,

for each map $f: X \rightarrow Y$ of anima an induced map

$$E^W Y \xrightarrow{E(f)} E^{f^* W} X$$

for each affine linear map $\alpha: U \rightarrow V$ of VB over X an induced map

$$E^U X \xrightarrow{E(\alpha)} E^V X \text{ s.t.}$$

Functoriality and Homotopy Invariance

Genuine Cohomology Theories

Definition

A **genuine Cohomology Theory** E consists of the following data

for each pair (X, V) an abelian group $E^V(X)$,

for each map $f: X \rightarrow Y$ of anima an induced map

$$E^W Y \xrightarrow{E(f)} E^{f^* W} X$$

for each affine linear map $\alpha: U \rightarrow V$ of VB over X an induced map

$$E^U X \xrightarrow{E(\alpha)} E^V X \text{ s.t.}$$

Functoriality and Homotopy Invariance

Beck-Chevalley $E(f^* \beta) \circ E(f) = E(f) \circ E(\beta)$

Genuine Cohomology Theories

Definition

A **genuine Cohomology Theory** E consists of the following data

for each pair (X, V) an abelian group $E^V(X)$,

for each map $f: X \rightarrow Y$ of anima an induced map

$$E^W Y \xrightarrow{E(f)} E^{f^* W} X$$

for each affine linear map $\alpha: U \rightarrow V$ of VB over X an induced map

$$E^U X \xrightarrow{E(\alpha)} E^V X \text{ s.t.}$$

Functoriality and Homotopy Invariance

Beck-Chevalley $E(f^* \beta) \circ E(f) = E(f) \circ E(\beta)$

Thom-Isomorphism The map

$E^V(X) \rightarrow E^V(S^W) \rightarrow E^{V \oplus W}(S^W) \rightarrow E^{V \oplus W}(S^W, X)$ is an isomorphism

Genuine Cohomology Theories

Definition

A **genuine Cohomology Theory** E consists of the following data

for each pair (X, V) an abelian group $E^V(X)$,

for each map $f: X \rightarrow Y$ of anima an induced map

$$E^W Y \xrightarrow{E(f)} E^{f^* W} X$$

for each affine linear map $\alpha: U \rightarrow V$ of VB over X an induced map

$$E^U X \xrightarrow{E(\alpha)} E^V X \text{ s.t.}$$

Functoriality and Homotopy Invariance

Beck-Chevalley $E(f^* \beta) \circ E(f) = E(f) \circ E(\beta)$

Thom-Isomorphism The map

$E^V(X) \rightarrow E^V(S^W) \rightarrow E^{V \oplus W}(S^W) \rightarrow E^{V \oplus W}(S^W, X)$ is an isomorphism

Mayer-Vietoris $E^V(B \amalg_A C) \rightarrow E^V(B) \oplus E^V(C) \rightarrow E^V(A)$ exact.

Genuine Cohomology Theories

Definition

A **genuine Cohomology Theory** E consists of the following data

for each pair (X, V) an abelian group $E^V(X)$,

for each map $f: X \rightarrow Y$ of anima an induced map

$$E^W Y \xrightarrow{E(f)} E^{f^* W} X$$

for each affine linear map $\alpha: U \rightarrow V$ of VB over X an induced map

$$E^U X \xrightarrow{E(\alpha)} E^V X \text{ s.t.}$$

Functoriality and Homotopy Invariance

Beck-Chevalley $E(f^* \beta) \circ E(f) = E(f) \circ E(\beta)$

Thom-Isomorphism The map

$E^V(X) \rightarrow E^V(S^W) \rightarrow E^{V \oplus W}(S^W) \rightarrow E^{V \oplus W}(S^W, X)$ is an isomorphism

Mayer-Vietoris $E^V(B \amalg_A C) \rightarrow E^V(B) \oplus E^V(C) \rightarrow E^V(A)$ exact.

Genuine Cohomology Theories

Definition

A **genuine Cohomology Theory** E consists of the following data

Functor $E: \mathcal{C}_{\text{An}, \text{AffLin}} \rightarrow \text{Ab}^{op}$

Thom-Isomorphism The map $E^V(X) \rightarrow E^V(S^W) \rightarrow E^{V \oplus W}(S^W) \rightarrow E^{V \oplus W}(S^W, X)$ is an isomorphism

Mayer-Vietoris $E^V(B \amalg_A C) \rightarrow E^V(B) \oplus E^V(C) \rightarrow E^V(A)$ exact.

Genuine Cohomology Theories

Definition

A **genuine Cohomology Theory** E consists of the following data

Functor $E: \mathcal{C}_{\text{An}, \text{AffLin}} \rightarrow \text{Ab}^{\text{op}}$

Thom-Isomorphism The map $E^V(X) \rightarrow E^V(S^W) \rightarrow E^{V \oplus W}(S^W) \rightarrow E^{V \oplus W}(S^W, X)$ is an isomorphism

Mayer-Vietoris $E^V(B \amalg_A C) \rightarrow E^V(B) \oplus E^V(C) \rightarrow E^V(A)$ exact.

$\text{Gen}(\mathcal{C}_{\text{An}, \text{AffLin}}, \text{Ab}^{\text{op}})$ is the category of genuine cohomology theories.

Genuine Cohomology Theories

Definition

A **genuine Cohomology Theory** E consists of the following data

Functor $E: \mathcal{C}_{\text{An}, \text{AffLin}} \rightarrow \text{Ab}^{\text{op}}$

Thom-Isomorphism The map $E^V(X) \rightarrow E^V(S^W) \rightarrow E^{V \oplus W}(S^W) \rightarrow E^{V \oplus W}(S^W, X)$ is an isomorphism

Mayer-Vietoris $E^V(B \amalg_A C) \rightarrow E^V(B) \oplus E^V(C) \rightarrow E^V(A)$ exact.

$\text{Gen}(\mathcal{C}_{\text{An}, \text{AffLin}}, \text{Ab}^{\text{op}})$ is the category of genuine cohomology theories.

$\text{Gen}^{\text{split}}(\mathcal{C}_{\text{An}, \text{AffLin}}, \mathcal{D})$ is the ∞ -category of split genuine homology theories.

Genuine Cohomology Theories

Definition

Definition: A **split genuine Homology Theory** E with coefficients in a pointed ∞ -category \mathcal{D} consists of the following data

$\mathrm{Gen}(\mathcal{C}_{\mathrm{An}, \mathrm{AffLin}}, \mathrm{Ab}^{\mathrm{op}})$ is the category of genuine cohomology theories.
 $\mathrm{Gen}^{\mathrm{split}}(\mathcal{C}_{\mathrm{An}, \mathrm{AffLin}}, \mathcal{D})$ is the ∞ -category of split genuine homology theories.

Genuine Cohomology Theories

Definition

Definition: A **split genuine Homology Theory** E with coefficients in a pointed ∞ -category \mathcal{D} consists of the following data

Functor $E: \mathcal{C}_{\text{An}, \text{AffLin}} \rightarrow \mathcal{D}$

Thom-Isomorphism The map

$E^V(X) \rightarrow E^V(S^W) \rightarrow E^{V \oplus W}(S^W) \rightarrow E^{V \oplus W}(S^W, X)$ is an isomorphism

$\text{Gen}(\mathcal{C}_{\text{An}, \text{AffLin}}, \text{Ab}^{\text{op}})$ is the category of genuine cohomology theories.

$\text{Gen}^{\text{split}}(\mathcal{C}_{\text{An}, \text{AffLin}}, \mathcal{D})$ is the ∞ -category of split genuine homology theories.

Genuine Cohomology Theories

Definition

Definition: A **split genuine Homology Theory** E with coefficients in a pointed ∞ -category \mathcal{D} consists of the following data

Functor $E: \mathcal{C}_{\text{An}, \text{AffLin}} \rightarrow \mathcal{D}$

Thom-Isomorphism The map $E^V(X) \rightarrow E^V(S^W) \rightarrow E^{V \oplus W}(S^W) \rightarrow E^{V \oplus W}(S^W, X)$ is an isomorphism

Mayer-Vietoris $E^V(B \amalg_A C) \rightarrow E^V(B) \oplus E^V(C) \rightarrow E^V(A)$ exact whenever $A \rightarrow C$ or $A \rightarrow B$ admits a retraction.

$\text{Gen}(\mathcal{C}_{\text{An}, \text{AffLin}}, \text{Ab}^{\text{op}})$ is the category of genuine cohomology theories.

$\text{Gen}^{\text{split}}(\mathcal{C}_{\text{An}, \text{AffLin}}, \mathcal{D})$ is the ∞ -category of split genuine homology theories.

Genuine Cohomology Theories

Definition

Definition: A **split genuine Homology Theory** E with coefficients in a pointed ∞ -category \mathcal{D} consists of the following data

Functor $E: \mathcal{C}_{\text{An}, \text{AffLin}} \rightarrow \mathcal{D}$

Thom-Isomorphism The map $E^V(X) \rightarrow E^V(S^W) \rightarrow E^{V \oplus W}(S^W) \rightarrow E^{V \oplus W}(S^W, X)$ is an isomorphism

Mayer-Vietoris $E^V(B \amalg_A C) \rightarrow E^V(B) \oplus E^V(C) \rightarrow E^V(A)$ exact whenever $A \rightarrow C$ or $A \rightarrow B$ admits a retraction.

$\text{Gen}(\mathcal{C}_{\text{An}, \text{AffLin}}, \text{Ab}^{\text{op}})$ is the category of genuine cohomology theories.

$\text{Gen}^{\text{split}}(\mathcal{C}_{\text{An}, \text{AffLin}}, \mathcal{D})$ is the ∞ -category of split genuine homology theories.

$\text{Gen}(\mathcal{C}_{\text{An}_G, \text{AffLin}}, \text{Ab}^{\text{op}})$ is the category of G -equivariant genuine cohomology theories.

Genuine Cohomology Theories

Definition

Definition: A **split genuine Homology Theory** E with coefficients in a pointed ∞ -category \mathcal{D} consists of the following data

Functor $E: \mathcal{C}_{\text{An}, \text{AffLin}} \rightarrow \mathcal{D}$

Thom-Isomorphism The map $E^V(X) \rightarrow E^V(S^W) \rightarrow E^{V \oplus W}(S^W) \rightarrow E^{V \oplus W}(S^W, X)$ is an isomorphism

Mayer-Vietoris $E^V(B \amalg_A C) \rightarrow E^V(B) \oplus E^V(C) \rightarrow E^V(A)$ exact whenever $A \rightarrow C$ or $A \rightarrow B$ admits a retraction.

$\text{Gen}(\mathcal{C}_{\text{An}, \text{AffLin}}, \text{Ab}^{\text{op}})$ is the category of genuine cohomology theories.

$\text{Gen}^{\text{split}}(\mathcal{C}_{\text{An}, \text{AffLin}}, \mathcal{D})$ is the ∞ -category of split genuine homology theories.

$\text{Gen}(\mathcal{C}_{\text{An}_G, \text{AffLin}}, \text{Ab}^{\text{op}})$ is the category of G -equivariant genuine cohomology theories.

$\text{Gen}^{\text{split}}(\mathcal{C}_{\text{An}_G, \text{AffLin}}, \mathcal{D})$ is the ∞ -category of G -equivariant split genuine homology theories.

Examples and Main Results

$(X, V) \mapsto X^{-V}$ defines a functor $\mathrm{Th}^- : \mathcal{C}_{\mathrm{An}, \mathrm{AffLin}} \rightarrow \mathrm{Sp}^\omega$

Examples and Main Results

$(X, V) \mapsto X^{-V}$ defines a functor $\mathrm{Th}^- : \mathcal{C}_{\mathrm{An}, \mathrm{AffLin}} \rightarrow \mathrm{Sp}^\omega$

Example

Let $E : \mathrm{Sp}^\omega \rightarrow \mathrm{Ab}^{op}$ be a cohomology theory, then

$$E \circ \mathrm{Th}^- \in \mathrm{Gen}(\mathcal{C}_{\mathrm{An}, \mathrm{AffLin}}, \mathrm{Ab}^{op})$$

Examples and Main Results

$(X, V) \mapsto X^{-V}$ defines a functor $\mathrm{Th}^- : \mathcal{C}_{\mathrm{An}, \mathrm{AffLin}} \rightarrow \mathrm{Sp}^\omega$

Example

Let $E : \mathrm{Sp}^\omega \rightarrow \mathrm{Ab}^{\mathrm{op}}$ be a cohomology theory, then
 $E \circ \mathrm{Th}^- \in \mathrm{Gen}(\mathcal{C}_{\mathrm{An}, \mathrm{AffLin}}, \mathrm{Ab}^{\mathrm{op}})$

Theorem

The functor $E \mapsto E \circ \mathrm{Th}^-$ defines an equivalence of categories

$$\{\text{cohomology theories } E : \mathrm{Sp}^\omega \rightarrow \mathrm{Ab}^{\mathrm{op}}\} \simeq \mathrm{Gen}(\mathcal{C}_{\mathrm{An}, \mathrm{AffLin}}, \mathrm{Ab}^{\mathrm{op}})$$

Examples and Main Results

$(X, V) \mapsto X^{-V}$ defines a functor $\mathrm{Th}^- : \mathcal{C}_{\mathrm{An}, \mathrm{AffLin}} \rightarrow \mathrm{Sp}^\omega$

Example

Let $E: \mathrm{Sp}^\omega \rightarrow \mathrm{Ab}^{\mathrm{op}}$ be a cohomology theory, then
 $E \circ \mathrm{Th}^- \in \mathrm{Gen}(\mathcal{C}_{\mathrm{An}, \mathrm{AffLin}}, \mathrm{Ab}^{\mathrm{op}})$

Theorem

The functor $E \mapsto E \circ \mathrm{Th}^-$ defines an equivalence of categories

$$\{\text{cohomology theories } E: \mathrm{Sp}^\omega \rightarrow \mathrm{Ab}^{\mathrm{op}}\} \simeq \mathrm{Gen}(\mathcal{C}_{\mathrm{An}, \mathrm{AffLin}}, \mathrm{Ab}^{\mathrm{op}})$$

Theorem

For a fixed compact Lie group G we have an equivalence of ∞ -categories

$$\{RO(G)\text{-graded cohomology theories}\} \simeq \mathrm{Gen}(\mathcal{C}_{\mathrm{An}_G, \mathrm{AffLin}}, \mathrm{Ab}^{\mathrm{op}})$$

Main Theorem

Main Theorem

Main Theorem

Let \mathcal{D} be a pointed ∞ -category with all split pushouts.

Main Theorem

Main Theorem

Let \mathcal{D} be a pointed ∞ -category with all split pushouts.

The pullback along $\mathrm{Th}^- : \mathcal{C}_{\mathrm{An}, \mathrm{AffLin}} \rightarrow \mathrm{Sp}^\omega$ induces an equivalence of ∞ -categories

$$\mathrm{Exc}^{\mathrm{split}}(\mathrm{Sp}^\omega, \mathcal{D}) \simeq \mathrm{Gen}^{\mathrm{split}}(\mathcal{C}_{\mathrm{An}, \mathrm{AffLin}}, \mathcal{D})$$

Main Theorem

Main Theorem

Let \mathcal{D} be a pointed ∞ -category with all split pushouts.

The pullback along $\mathrm{Th}^- : \mathcal{C}_{\mathrm{An}, \mathrm{AffLin}} \rightarrow \mathrm{Sp}^\omega$ induces an equivalence of ∞ -categories

$$\mathrm{Exc}^{\mathrm{split}}(\mathrm{Sp}^\omega, \mathcal{D}) \simeq \mathrm{Gen}^{\mathrm{split}}(\mathcal{C}_{\mathrm{An}, \mathrm{AffLin}}, \mathcal{D})$$

Slogan

The functor Th^- is the **universal** (split) genuine homology theory.

Main Theorem

Main Theorem

Let \mathcal{D} be a pointed ∞ -category with all split pushouts.

The pullback along $\mathrm{Th}^- : \mathcal{C}_{\mathrm{An}_G, \underline{\mathrm{AffLin}}} \rightarrow \mathrm{Sp}_G^\omega$ induces an equivalence of ∞ -categories

$$\mathrm{Exc}^{\mathrm{split}}(\mathrm{Sp}_G^\omega, \mathcal{D}) \simeq \mathrm{Gen}^{\mathrm{split}}(\mathcal{C}_{\mathrm{An}_G, \underline{\mathrm{AffLin}}}, \mathcal{D})$$

Slogan

The functor Th^- is the **universal** (split) genuine homology theory.

Proof Sketch of the Main Theorem

Prove an unstable version of the theorem.

Proposition

The pullback along $\mathrm{Th}^- : \mathcal{C}_{\mathrm{An}, \mathrm{AffLin}}(V) \rightarrow \mathrm{An}_*^\omega$ induces an equivalence of ∞ -categories

$$\mathrm{Exc}^{\mathrm{split}}(\mathrm{An}_*, \mathcal{D}) \simeq \mathrm{Gen}^{\mathrm{split}}(\mathcal{C}_{\mathrm{An}, \mathrm{AffLin}}(V), \mathcal{D})$$

Proof Sketch of the Main Theorem

Prove an unstable version of the theorem.

Proposition

The pullback along $\mathrm{Th}^- : \mathcal{C}_{\mathrm{An}, \mathrm{AffLin}}(V) \rightarrow \mathrm{An}_*^\omega$ induces an equivalence of ∞ -categories

$$\mathrm{Exc}^{\mathrm{split}}(\mathrm{An}_*, \mathcal{D}) \simeq \mathrm{Gen}^{\mathrm{split}}(\mathcal{C}_{\mathrm{An}, \mathrm{AffLin}}(V), \mathcal{D})$$

Reduce the stable version to the unstable version.

Lemma

We have equivalences of ∞ -categories

$$\mathrm{colim}_V \mathcal{C}_{\mathrm{An}, \mathrm{AffLin}}(V) \simeq \mathcal{C}_{\mathrm{An}, \mathrm{AffLin}}$$

$$\mathrm{colim}_V \mathrm{An}_*^\omega \simeq \mathrm{Sp}^\omega$$

Proof Sketch of the Main Unstable Theorem

We can define the inverse functor of $(\mathrm{Th}^-)^*$ as follows:

$$\begin{aligned} \mathrm{res}: \mathrm{Gen}^{\mathrm{split}}(\mathcal{C}_{\mathrm{An}, \mathrm{AffLin}}(V), \mathcal{D}) &\rightarrow \mathrm{Exc}^{\mathrm{split}}(\mathrm{An}_*^\omega, \mathcal{D}) \\ E &\mapsto \mathrm{res} E: X \mapsto \mathrm{cof}[E^V(\mathrm{pt}) \rightarrow E^{r^*V}(X)] \end{aligned}$$

Proof Sketch of the Main Unstable Theorem

We can define the inverse functor of $(\mathrm{Th}^-)^*$ as follows:

$$\begin{aligned} \mathrm{res}: \mathrm{Gen}^{\mathrm{split}}(\mathcal{C}_{\mathrm{An}, \mathrm{AffLin}}(V), \mathcal{D}) &\rightarrow \mathrm{Exc}^{\mathrm{split}}(\mathrm{An}_*^\omega, \mathcal{D}) \\ E &\mapsto \mathrm{res} E: X \mapsto \mathrm{cof}[E^V(\mathrm{pt}) \rightarrow E^{r^*V}(X)] \end{aligned}$$

We compute:

$$\mathrm{res}(E \circ \mathrm{Th}^-)(X) \simeq \mathrm{cof}[E(\mathrm{pt}_+) \rightarrow E(X_+)]$$

Proof Sketch of the Main Unstable Theorem

We can define the inverse functor of $(\mathrm{Th}^-)^*$ as follows:

$$\begin{aligned} \mathrm{res}: \mathrm{Gen}^{\mathrm{split}}(\mathcal{C}_{\mathrm{An}, \mathrm{AffLin}}(V), \mathcal{D}) &\rightarrow \mathrm{Exc}^{\mathrm{split}}(\mathrm{An}_*^\omega, \mathcal{D}) \\ E &\mapsto \mathrm{res} E: X \mapsto \mathrm{cof}[E^V(\mathrm{pt}) \rightarrow E^{r^*V}(X)] \end{aligned}$$

We compute:

$$\begin{aligned} \mathrm{res}(E \circ \mathrm{Th}^-)(X) &\simeq \mathrm{cof}[E(\mathrm{pt}_+) \rightarrow E(X_+)] \\ &\simeq E(X) \quad (\text{since } E \text{ is split excisive}) \end{aligned}$$

Proof Sketch of the Main Unstable Theorem

We can define the inverse functor of $(\mathrm{Th}^-)^*$ as follows:

$$\begin{aligned} \mathrm{res}: \mathrm{Gen}^{\mathrm{split}}(\mathcal{C}_{\mathrm{An}, \mathrm{AffLin}}(V), \mathcal{D}) &\rightarrow \mathrm{Exc}^{\mathrm{split}}(\mathrm{An}_*^\omega, \mathcal{D}) \\ E &\mapsto \mathrm{res} E: X \mapsto \mathrm{cof}[E^V(\mathrm{pt}) \rightarrow E^{r^*V}(X)] \end{aligned}$$

We compute:

$$\begin{aligned} \mathrm{res}(E \circ \mathrm{Th}^-)(X) &\simeq \mathrm{cof}[E(\mathrm{pt}_+) \rightarrow E(X_+)] \\ &\simeq E(X) \quad (\text{since } E \text{ is split excisive}) \end{aligned}$$

and

$$((\mathrm{res} E) \circ \mathrm{Th}^-)^U(X) \simeq \tilde{E}^V(\mathrm{Th}_X(U^{\perp V}))$$

Proof Sketch of the Main Unstable Theorem

We can define the inverse functor of $(\mathrm{Th}^-)^*$ as follows:

$$\begin{aligned} \mathrm{res}: \mathrm{Gen}^{\mathrm{split}}(\mathcal{C}_{\mathrm{An}, \mathrm{AffLin}}(V), \mathcal{D}) &\rightarrow \mathrm{Exc}^{\mathrm{split}}(\mathrm{An}_*^\omega, \mathcal{D}) \\ E &\mapsto \mathrm{res} E: X \mapsto \mathrm{cof}[E^V(\mathrm{pt}) \rightarrow E^{r^*V}(X)] \end{aligned}$$

We compute:

$$\begin{aligned} \mathrm{res}(E \circ \mathrm{Th}^-)(X) &\simeq \mathrm{cof}[E(\mathrm{pt}_+) \rightarrow E(X_+)] \\ &\simeq E(X) \quad (\text{since } E \text{ is split excisive}) \end{aligned}$$

and

$$\begin{aligned} ((\mathrm{res} E) \circ \mathrm{Th}^-)^U(X) &\simeq \tilde{E}^V(\mathrm{Th}_X(U^{\perp_V})) \\ &\simeq E^V(S_X^{U^{\perp_V}}, X) \quad (\text{Mayer-Vietoris Axiom}) \end{aligned}$$

Proof Sketch of the Main Unstable Theorem

We can define the inverse functor of $(\mathrm{Th}^-)^*$ as follows:

$$\begin{aligned} \mathrm{res}: \mathrm{Gen}^{\mathrm{split}}(\mathcal{C}_{\mathrm{An}, \mathrm{AffLin}}(V), \mathcal{D}) &\rightarrow \mathrm{Exc}^{\mathrm{split}}(\mathrm{An}_*^\omega, \mathcal{D}) \\ E &\mapsto \mathrm{res} E: X \mapsto \mathrm{cof}[E^V(\mathrm{pt}) \rightarrow E^{r^*V}(X)] \end{aligned}$$

We compute:

$$\begin{aligned} \mathrm{res}(E \circ \mathrm{Th}^-)(X) &\simeq \mathrm{cof}[E(\mathrm{pt}_+) \rightarrow E(X_+)] \\ &\simeq E(X) \quad (\text{since } E \text{ is split excisive}) \end{aligned}$$

and

$$\begin{aligned} ((\mathrm{res} E) \circ \mathrm{Th}^-)^U(X) &\simeq \tilde{E}^V(\mathrm{Th}_X(U^{\perp V})) \\ &\simeq E^V(S_X^{U^{\perp V}}, X) \quad (\text{Mayer-Vietoris Axiom}) \\ &\simeq E^{U \oplus U^{\perp V}}(S_X^{U^{\perp V}}, X) \end{aligned}$$

Proof Sketch of the Main Unstable Theorem

We can define the inverse functor of $(\mathrm{Th}^-)^*$ as follows:

$$\begin{aligned} \mathrm{res}: \mathrm{Gen}^{\mathrm{split}}(\mathcal{C}_{\mathrm{An}, \mathrm{AffLin}}(V), \mathcal{D}) &\rightarrow \mathrm{Exc}^{\mathrm{split}}(\mathrm{An}_*^\omega, \mathcal{D}) \\ E &\mapsto \mathrm{res} E: X \mapsto \mathrm{cof}[E^V(\mathrm{pt}) \rightarrow E^{r^*V}(X)] \end{aligned}$$

We compute:

$$\begin{aligned} \mathrm{res}(E \circ \mathrm{Th}^-)(X) &\simeq \mathrm{cof}[E(\mathrm{pt}_+) \rightarrow E(X_+)] \\ &\simeq E(X) \quad (\text{since } E \text{ is split excisive}) \end{aligned}$$

and

$$\begin{aligned} ((\mathrm{res} E) \circ \mathrm{Th}^-)^U(X) &\simeq \tilde{E}^V(\mathrm{Th}_X(U^{\perp V})) \\ &\simeq E^V(S_X^{U^{\perp V}}, X) \quad (\text{Mayer-Vietoris Axiom}) \\ &\simeq E^{U \oplus U^{\perp V}}(S_X^{U^{\perp V}}, X) \\ &\simeq E^U(X) \quad (\text{Thom-Isomorphism Axiom}) \quad \square \end{aligned}$$

Application

Let $\mathcal{D} = \mathbf{An}_*$, then

$$\mathrm{Gen}(\mathcal{C}_{\mathbf{An}, \mathbf{AffLin}}, \mathbf{An}_*) \simeq \mathrm{Exc}(\mathbf{Sp}^\omega, \mathbf{An}_*) \simeq \mathbf{Sp}$$

Application

Let $\mathcal{D} = \mathbf{An}_*$, then

$$\mathrm{Gen}(\mathcal{C}_{\mathbf{An}, \mathbf{AffLin}}, \mathbf{An}_*) \simeq \mathrm{Exc}(\mathbf{Sp}^\omega, \mathbf{An}_*) \simeq \mathbf{Sp}$$

Let $E \in \mathrm{Gen}(\mathcal{C}_{\mathbf{An}, \mathbf{AffLin}}, \mathbf{An}_*)$, then we can compute

Application

Let $\mathcal{D} = \mathbf{An}_*$, then

$$\mathrm{Gen}(\mathcal{C}_{\mathbf{An}, \mathbf{AffLin}}, \mathbf{An}_*) \simeq \mathrm{Exc}(\mathrm{Sp}^\omega, \mathbf{An}_*) \simeq \mathrm{Sp}$$

Let $E \in \mathrm{Gen}(\mathcal{C}_{\mathbf{An}, \mathbf{AffLin}}, \mathbf{An}_*)$, then we can compute

$$E^V(\mathrm{pt}) \simeq E^{V \oplus W}(S^W, \mathrm{pt}) \quad \text{Thom Isomorphism}$$

Application

Let $\mathcal{D} = \mathbf{An}_*$, then

$$\mathrm{Gen}(\mathcal{C}_{\mathbf{An}, \mathbf{AffLin}}, \mathbf{An}_*) \simeq \mathrm{Exc}(\mathbf{Sp}^\omega, \mathbf{An}_*) \simeq \mathbf{Sp}$$

Let $E \in \mathrm{Gen}(\mathcal{C}_{\mathbf{An}, \mathbf{AffLin}}, \mathbf{An}_*)$, then we can compute

$$\begin{aligned} E^V(\mathrm{pt}) &\simeq E^{V \oplus W}(S^W, \mathrm{pt}) && \text{Thom Isomorphism} \\ &\simeq \Omega^W E^{V \oplus W}(\mathrm{pt}) && \text{Mayer-Vietoris} \end{aligned}$$

Application

Let $\mathcal{D} = \mathbf{An}_*$, then

$$\mathrm{Gen}(\mathcal{C}_{\mathbf{An}, \mathbf{AffLin}}, \mathbf{An}_*) \simeq \mathrm{Exc}(\mathbf{Sp}^\omega, \mathbf{An}_*) \simeq \mathbf{Sp}$$

Let $E \in \mathrm{Gen}(\mathcal{C}_{\mathbf{An}, \mathbf{AffLin}}, \mathbf{An}_*)$, then we can compute

$$\begin{aligned} E^V(\mathrm{pt}) &\simeq E^{V \oplus W}(S^W, \mathrm{pt}) && \text{Thom Isomorphism} \\ &\simeq \Omega^W E^{V \oplus W}(\mathrm{pt}) && \text{Mayer-Vietoris} \end{aligned}$$

$X(V) := E^V(\mathrm{pt})$ defines an **orthogonal spectrum**.