Matematika pro fyziky 1: The X-Files ••



Michal Grňo

4. ledna 2020

1 ???

$$oldsymbol{p},oldsymbol{q}\in\mathbb{R}^{2} \hspace{0.5cm}H=\left\|oldsymbol{p}
ight\|^{2}+\left\|oldsymbol{q}
ight\|^{2} \ \dot{q}_{j}=rac{\partial H}{\partial p_{j}}\hspace{0.5cm}\dot{p}_{j}=-rac{\partial H}{\partial q_{j}}\hspace{0.5cm}-\zeta\left(rac{\partial H}{\partial p_{1}}-rac{\partial H}{\partial p_{2}}
ight)$$

2

2.1 p, q = ?

$$\boldsymbol{v} = \begin{pmatrix} \boldsymbol{q} \\ \boldsymbol{p} \end{pmatrix} = \begin{pmatrix} q_1 \\ q_2 \\ p_1 \\ p_2 \end{pmatrix} \qquad \qquad \dot{\boldsymbol{v}} = \begin{pmatrix} -\frac{\partial H}{\partial q_1} - \zeta \left(\frac{\partial H}{\partial p_1} - \frac{\partial H}{\partial p_2} \right) \\ -\frac{\partial H}{\partial q_2} - \zeta \left(\frac{\partial H}{\partial p_1} - \frac{\partial H}{\partial p_2} \right) \\ \frac{\partial H}{\partial p_1} \\ \frac{\partial H}{\partial p_2} \end{pmatrix} = 2 \underbrace{\begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & -\zeta & \zeta \\ 0 & 1 & -\zeta & \zeta \end{pmatrix}}_{M} \boldsymbol{v}$$

$$\dot{\boldsymbol{v}} = M \boldsymbol{v} \implies \boldsymbol{v} = \exp(tM) \boldsymbol{v_0}, \quad \boldsymbol{v_0} \in \mathbb{R}^2$$

$\exp(tM) = ?$ 2.2

$$2t \begin{pmatrix} 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ 1 & 0 & -\zeta & \zeta \\ 0 & 1 & -\zeta & \zeta \end{pmatrix} = \begin{pmatrix} t\zeta & \frac{\zeta}{2} - 1 & -t\zeta & \frac{\zeta}{2} + 1 \\ t\zeta & \frac{\zeta}{2} & -t\zeta & \frac{\zeta}{2} \\ -t\zeta & 1 & -t\zeta & 1 \\ -t\zeta & 0 & -t\zeta & 0 \end{pmatrix} \begin{pmatrix} -2t & 1 & 0 & 0 \\ 0 & -2t & 0 & 0 \\ 0 & 0 & 2t & 1 \\ 0 & 0 & 0 & 2t \end{pmatrix} \begin{pmatrix} 0 & \frac{1}{2t\zeta} & -\frac{1}{4t} & \frac{\zeta-2}{4t\zeta} \\ -\frac{1}{2} & \frac{1}{2} & -\frac{1}{2} \\ 0 & -\frac{1}{2t\zeta} & \frac{1}{4t} & -\frac{\zeta+2}{4t\zeta} \\ \frac{1}{2} & -\frac{1}{2} & \frac{1}{2} & -\frac{1}{2} \end{pmatrix}$$

$$\exp(tM) = \exp(P^{-1}JP) = P^{-1}\exp(J)P$$

$$\exp(tM) = P^{-1} \begin{pmatrix} e^{-2t} & e^{-2t} & 0 & 0\\ 0 & e^{-2t} & 0 & 0\\ 0 & 0 & e^{2t} & e^{2t}\\ 0 & 0 & 0 & e^{2t} \end{pmatrix} P$$

$$\exp(tM) = \begin{pmatrix} \frac{\left(-2t\zeta - \zeta + \left(-2t\zeta + \zeta + 2\right)e^{4t} + 2\right)e^{-2t}}{4} & \frac{\zeta(2t + (2t - 1)e^{4t} + 1)e^{-2t}}{4} & \frac{(t\zeta + (-t\zeta + 1)e^{4t} - 1)e^{-2t}}{2} & t\zeta \sinh(2t) \\ \frac{\zeta(-2t + (1 - 2t)e^{4t} - 1)e^{-2t}}{4} & \frac{(2t\zeta + \zeta + (2t\zeta - \zeta + 2)e^{4t} + 2)e^{-2t}}{4} & -t\zeta \sinh(2t) \\ \frac{\zeta(t\zeta + (-t\zeta + 1)e^{4t} - 1)e^{-2t}}{2} & t\zeta \sinh(2t) & \frac{(-t\zeta + (t\zeta + 1)e^{4t} - 1)e^{-2t}}{4} \\ -t\zeta \sinh(2t) & \frac{(-t\zeta + (t\zeta + 1)e^{4t} - 1)e^{-2t}}{2} & \frac{\zeta(2t + (2t\zeta - \zeta + 2)e^{4t} + 2)e^{-2t}}{4} & \frac{(-2t\zeta + \zeta + (-2t\zeta - \zeta + 2)e^{4t} + 2)e^{-2t}}{4} & \frac{\zeta(2t + (2t\tau + 1)e^{4t} - 1)e^{-2t}}{4} \\ -t\zeta \sinh(2t) & \frac{(-t\zeta + (t\zeta + 1)e^{4t} - 1)e^{-2t}}{2} & \frac{\zeta(2t + (2t\zeta + 1)e^{4t} - 1)e^{-2t}}{4} & \frac{\zeta(2t + (2t\zeta + 1)e^{4t} - 1)e^{-2t}}{4} & \frac{\zeta(2t + (2t\zeta + 1)e^{4t} - 1)e^{-2t}}{4} \end{pmatrix}$$

$$\exp(tM)|_{\zeta=0} = \begin{pmatrix} \cosh(2t) & 0 & \sinh(2t) & 0\\ 0 & \cosh(2t) & 0 & \sinh(2t)\\ \sinh(2t) & 0 & \cosh(2t) & 0\\ 0 & \sinh(2t) & 0 & \cosh(2t) \end{pmatrix}$$

2.3 $\det \exp(tM) = ?$

$$\det \exp(tM) = \exp \operatorname{Tr}(tM) = \exp \left(t\operatorname{Tr}(M) \right) = \exp \left(t\left(z-z \right) \right) = \exp (0) = 1$$