

3.5

$$f(x) = \arctan\left(\frac{2+x}{2-x}\right) - \arctan\left(\frac{x}{2}\right)$$

$$D_f = \mathbb{R} \setminus \{2\}$$

$$\begin{aligned} \frac{df}{dx} &= \frac{1}{1 + \left(\frac{2+x}{2-x}\right)^2} \cdot \frac{(1)(2-x) - (-1)(2+x)}{(2-x)^2} - \frac{1}{1 + \left(\frac{x}{2}\right)^2} \cdot \frac{1}{2} \\ &= \frac{1}{\frac{4 - 4x + x^2 + 4 + 4x + x^2}{(2-x)^2}} \cdot \frac{2-x+2+x}{(2-x)^2} - \frac{1}{\frac{4+x^2}{4}} \cdot \frac{1}{2} \\ &= \frac{(2-x)^2}{2x^2+8} \cdot \frac{4}{(2-x)^2} - \frac{2}{x^2+4} = \frac{2}{x^2+4} - \frac{2}{x^2+4} = 0 \end{aligned}$$


$$\lim_{x \rightarrow 2^+} f(x) = \lim_{x \rightarrow 2^+} \arctan\left(\frac{2+x}{2-x}\right) - \arctan\left(\frac{x}{2}\right) = -\frac{\pi}{2} - \frac{\pi}{4} = -\frac{3\pi}{4}$$

$$\lim_{x \rightarrow 2^-} f(x) = \lim_{x \rightarrow 2^-} \arctan\left(\frac{2+x}{2-x}\right) - \arctan\left(\frac{x}{2}\right) = \frac{\pi}{2} - \frac{\pi}{4} = \frac{\pi}{4}$$

$$\lim_{x \rightarrow +\infty} f(x) = \lim_{x \rightarrow +\infty} \arctan\left(\frac{x+2}{-x+2}\right) - \arctan\left(\frac{x}{2}\right) = -\frac{\pi}{4} - \frac{\pi}{2} = -\frac{3\pi}{4}$$

$$\lim_{x \rightarrow -\infty} f(x) = \lim_{x \rightarrow -\infty} \arctan\left(\frac{x+2}{-x+2}\right) - \arctan\left(\frac{x}{2}\right) = -\frac{\pi}{4} - \left(-\frac{\pi}{2}\right) = \frac{\pi}{4}$$

brak asymptot pionowych

asymptoty poziome $y = -\frac{3\pi}{4}$ $y = \frac{\pi}{4}$ 

3.4a

$$f(x) = \sqrt{1+x^2} \quad f(x) \approx f(0) + \frac{df(0)}{1!}x + \frac{d^2f(0)}{2!}x^2 + \frac{d^3f(0)}{3!}x^3 + \frac{d^4f(0)}{4!}x^4$$

$$\frac{df}{dx} = \frac{1}{2\sqrt{1+x^2}} \cdot 2x = \frac{x}{\sqrt{1+x^2}}$$

$$\frac{d^2f}{dx^2} = \frac{(1)\sqrt{1+x^2} - x \cdot \frac{x}{\sqrt{1+x^2}}}{1+x^2} = \frac{1+x^2 - x^2}{(1+x^2)\sqrt{1+x^2}} = \frac{1}{(1+x^2)^{3/2}} = (1+x^2)^{-3/2}$$

$$\frac{d^3f}{dx^3} = -\frac{3}{2}(1+x^2)^{-5/2} \cdot 2x = -3x(1+x^2)^{-5/2}$$

$$\frac{d^4f}{dx^4} = -3(1+x^2)^{-5/2} + (-3x) \cdot \left(-\frac{5}{2}\right)(1+x^2)^{-7/2} \cdot 2x = -3(1+x^2)^{-5/2} + 15x^2(1+x^2)^{-7/2}$$

$$f(0) = \sqrt{1} = 1$$

$$\frac{df}{dx}(0) = \frac{0}{1} = 0$$

$$\frac{d^2f}{dx^2}(0) = 1^{-3/2} = 1$$

$$\frac{d^3f}{dx^3}(0) = -3 \cdot 0 \cdot 1^{-5/2} = 0$$

$$\frac{d^4f}{dx^4}(0) = -3(1)^{-5/2} + 0 = -3$$

$$f(x) \approx 1 + \frac{1}{2}x^2 - \frac{1}{8}x^4$$

6-7.1c

$$\int \sqrt{x} \arctan(\sqrt{x}) dx = \left| \begin{array}{l} f = \arctan(\sqrt{x}) \quad g' = \sqrt{x} \\ f' = \frac{1}{1+x} \cdot \frac{1}{2\sqrt{x}} \quad g = \frac{2}{3}x^{3/2} \end{array} \right|$$

$$= \frac{2}{3}x\sqrt{x} \arctan(\sqrt{x}) - \int \frac{2}{3} \cdot \frac{x\sqrt{x}}{2\sqrt{x}(x+1)} dx = \frac{2}{3}x\sqrt{x} \arctan(\sqrt{x}) - \frac{1}{3} \int \frac{x}{x+1} dx$$

$$\int \frac{x}{x+1} dx = \int \frac{x+1-1}{x+1} dx = \int 1 dx - \int \frac{1}{x+1} dx = x - \ln|x+1| + C$$

$$= \frac{2}{3}x\sqrt{x} \arctan(\sqrt{x}) - \frac{1}{3}x + \frac{1}{3}\ln|x+1| + C$$

6-7.2

$$b) \int e^{\sqrt{x}} dx = \left| \begin{array}{l} t = \sqrt{x} \\ \frac{dt}{dx} = \frac{1}{2\sqrt{x}} \\ dx = 2t dt \end{array} \right| = 2 \int t e^t dt = \left| \begin{array}{l} f = t \quad g' = e^t \\ f' = 1 \quad g = e^t \end{array} \right| = 2te^t - 2 \int e^t dt = 2te^t - 2e^t + C = 2e^{\sqrt{x}}(\sqrt{x} - 1) + C$$

$$c) \int \arcsin(x) dx = \left| \begin{array}{l} x = \sin(t) \\ dx = \cos(t) dt \\ t = \arcsin(x) \end{array} \right| = \int \arcsin(\sin(t)) \cos(t) dt = \int t \cos(t) dt = \left| \begin{array}{l} f = t \quad g' = \cos(t) \\ f' = 1 \quad g = \sin(t) \end{array} \right| = t \sin(t) - \int \sin(t) dt = t \sin(t) + \cos(t) + C$$

$$= x \arcsin(x) + \cos(\arcsin(x)) + C = x \arcsin(x) + \sqrt{1-x^2} + C$$

$$\cos(\arcsin(x)) = \sqrt{1 - \sin^2(\arcsin(x))} = \sqrt{1-x^2}$$

$$f) \int x^3 e^{x^2} dx = \left| \begin{array}{l} t = x^2 \\ dt = 2x dx \end{array} \right| = \frac{1}{2} \int t e^t dt = \left| \begin{array}{l} f = t \quad g' = e^t \\ f' = 1 \quad g = e^t \end{array} \right| = \frac{1}{2} t e^t - \frac{1}{2} e^t + C = \frac{1}{2} x^2 e^{x^2} - \frac{1}{2} e^{x^2} + C = \frac{1}{2} e^{x^2}(x^2 - 1) + C$$

6-7.3

$$b) \int \sin(4x) \cos(6x) dx = \int \frac{1}{2} [\sin((4-6)x) + \sin((4+6)x)] dx$$

$$= \frac{1}{2} \int \sin(-2x) dx + \frac{1}{2} \int \sin(10x) dx$$

$$= -\frac{1}{2} \int \sin(2x) dx + \frac{1}{2} \int \sin(10x) dx$$

$$= -\frac{1}{2} \left[-\frac{1}{2} \cos(2x) - \left(-\frac{1}{10} \cos(10x) \right) \right] + C$$

$$= \frac{1}{4} \cos(2x) - \frac{1}{20} \cos(10x) + C$$

$$c) \int \sin^4(x) \cos^3(x) dx = \int \sin^4(x) \cdot (1 - \sin^2(x)) \cdot \cos(x) dx = \left| \begin{array}{l} t = \sin(x) \\ dt = \cos(x) dx \end{array} \right| = \int t^4 (1 - t^2) dt = \int [-t^6 + t^4] dt = -\frac{1}{7} \sin^7(x) + \frac{1}{5} \sin^5(x) + C$$

6-7.4

$$b) \int \frac{x+1}{x^2+8x+25} dx = \frac{1}{2} \int \frac{2x+8-6}{x^2+8x+25} dx = \frac{1}{2} \int \frac{2x+8}{x^2+8x+25} dx - 3 \int \frac{dx}{x^2+8x+25}$$

$$\int \frac{2x+8}{x^2+8x+25} dx = \left| \begin{array}{l} t = x^2+8x+25 \\ dt = (2x+8) dx \end{array} \right| = \int \frac{dt}{t} = \ln|x^2+8x+25| + C$$

$$\int \frac{dx}{x^2+8x+25} = \int \frac{dx}{(x+4)^2+9} = \left| \begin{array}{l} t = \frac{1}{3}(x+4) \\ dt = \frac{1}{3} dx \end{array} \right| = 3 \int \frac{dt}{9t^2+9} = \frac{1}{3} \int \frac{dt}{t^2+1} = \frac{1}{3} \arctan\left(\frac{x+4}{3}\right) + C$$

$$\int \frac{x+1}{x^2+8x+25} dx = \frac{1}{2} \ln|x^2+8x+25| - \arctan\left(\frac{x+4}{3}\right) + C$$

6-7.5 e

$$\int x \sqrt{6x-x^2} dx = \int \frac{x(6x-x^2)}{\sqrt{-x^2+6x}} dx = \int \frac{-x^3+6x^2}{\sqrt{-x^2+6x}} dx = [Ax^2+Bx+C] \sqrt{-x^2+6x} + \lambda \int \frac{dx}{\sqrt{-x^2+6x}}$$

$$\frac{-x^3+6x^2}{\sqrt{-x^2+6x}} = \frac{[2Ax+B]\sqrt{-x^2+6x}}{2\sqrt{-x^2+6x}} + \frac{[Ax^2+Bx+C]}{2\sqrt{-x^2+6x}} \cdot (-2x+6) + \frac{\lambda}{\sqrt{-x^2+6x}}$$

$$-x^3+6x^2 = (2Ax+B)(-x^2+6x) + (Ax^2+Bx+C)(-x+3) + \lambda$$

$$-x^3+6x^2 = -2Ax^3+12Ax^2-Bx^2+6Bx-Ax^3+3Ax^2-Bx^2+3Bx-Cx+3C+\lambda$$

$$-x^3+6x^2 = (-3A)x^3 + (15A-2B)x^2 + (3B-C)x + (3C+\lambda)$$

$$\begin{cases} -1 = -3A & A = \frac{1}{3} \\ 6 = 15A - 2B & B = -\frac{1}{2} \\ 0 = 3B - C & C = -\frac{3}{2} \\ 0 = 3C + \lambda & \lambda = \frac{27}{2} \end{cases} \quad \int \frac{dx}{\sqrt{-x^2+6x}} = \int \frac{dx}{\sqrt{-x^2+6x-9+9}} = \int \frac{dx}{\sqrt{9-(x-3)^2}} = \left| \begin{array}{l} t = \frac{1}{3}(x-3) \\ dt = \frac{1}{3} dx \\ dx = 3dt \end{array} \right| = \int \frac{3dt}{\sqrt{9-9t^2}} = \int \frac{dt}{\sqrt{1-t^2}} = \arcsin\left(\frac{x-3}{3}\right) + C$$

$$\int x \sqrt{6x-x^2} dx = \left(\frac{1}{3}x^2 - \frac{1}{2}x - \frac{3}{2}\right)\sqrt{6x-x^2} + \frac{27}{2} \arcsin\left(\frac{x-3}{3}\right) + C$$

6-7.6

$$d) \int \frac{1}{\sin(x) + 2\cos(x) + 3} dx = \int \frac{\frac{2}{1+t^2}}{\frac{2t}{1+t^2} + 2 \frac{1-t^2}{1+t^2} + 3 \frac{1+t^2}{1+t^2}} dt = \int \frac{\frac{2}{1+t^2}}{\frac{2t + 2(1-t^2) + 3(1+t^2)}{1+t^2}} dt = 2 \int \frac{dt}{2t + 2 - 2t^2 + 3 + 3t^2} = 2 \int \frac{dt}{t^2 + 2t + 5}$$

$$\left| \begin{array}{l} t = \tan\left(\frac{x}{2}\right) \\ dx = \frac{2}{1+t^2} dt \\ \sin(x) = \frac{2t}{1+t^2} \\ \cos(x) = \frac{1-t^2}{1+t^2} \end{array} \right|$$

$$= 2 \int \frac{dt}{(t+1)^2 + 4} = \left| u = \frac{1}{2}(t+1) \right| = 4 \int \frac{du}{4u^2 + 4} = \int \frac{du}{u^2 + 1} = \arctan(u) + C = \arctan\left(\frac{t+1}{2}\right) + C = \arctan\left(\frac{\tan\left(\frac{x}{2}\right) + 1}{2}\right) + C$$

$$e) \int \frac{e^{2x} + e^x}{\sqrt{4 - e^{2x}}} dx = \left| \begin{array}{l} t = e^x \\ dt = e^x dx \end{array} \right| = \int \frac{t+1}{\sqrt{4-t^2}} dt = \int \frac{t}{\sqrt{4-t^2}} dt + \int \frac{dt}{\sqrt{4-t^2}}$$

$$\int \frac{t}{\sqrt{4-t^2}} dt = \left| u = 4-t^2 \right| = -\frac{1}{2} \int \frac{du}{\sqrt{u}} = -\frac{1}{2} \cdot \frac{1}{\frac{1}{2}} \cdot u^{\frac{1}{2}} + C = -\sqrt{u} + C = -\sqrt{4-t^2} + C$$

$$\int \frac{dt}{\sqrt{4-t^2}} = \left| u = \frac{1}{2}t \right| = 2 \int \frac{du}{\sqrt{4-4u^2}} = \int \frac{du}{\sqrt{1-u^2}} = \arcsin(u) + C = \arcsin\left(\frac{t}{2}\right) + C$$

$$\int \frac{e^{2x} + e^x}{\sqrt{4 - e^{2x}}} dx = \arcsin\left(\frac{1}{2}e^x\right) - \sqrt{4 - e^{2x}} + C$$

8-9.1c

$$-f(x) = f(-x)$$

$$\int_{-a}^a f(x) dx = \int_{-a}^0 f(x) dx + \int_0^a f(x) dx = -\int_0^a f(x) dx + \int_0^a f(x) dx = 0$$

$$\int_{-a}^0 f(x) dx = \left| \begin{array}{l} t = -x \\ dt = -dx \end{array} \right| = \int_a^0 -f(-t) dt = \int_0^a f(t) dt = -\int_0^a f(t) dt$$

8-9.2b


$$F(x) = \int_x^{2x} \frac{t}{t} dt = G(2x) - G(x)$$

$$G(t) = \int \frac{e^t}{t} dt$$

$$\frac{dF}{dx} = G'(2x) \cdot 2 - G'(x) = \frac{e^{2x}}{2x} \cdot 2 - \frac{e^x}{x} = \frac{e^{2x} - e^x}{x} = e^x \frac{e^x - 1}{x}$$

$$\frac{dF}{dx} > 0 \Leftrightarrow x e^x (e^x - 1) > 0$$

$\downarrow \begin{matrix} x > 0 \\ x < 0 \end{matrix}$
 $\downarrow \begin{matrix} x > 0 \\ x < 0 \end{matrix}$



$$F \nearrow \cup (-\infty, 0), (0, \infty)$$

8-9.3c

$$\lim_{x \rightarrow \infty} \frac{\int_x^{3x} \frac{dt}{\ln^3(t)}}{e^{3x}} \stackrel{H}{=} \lim_{x \rightarrow \infty} \frac{\frac{e^{3x} - e^x}{3x^3}}{e^{3x}} = \lim_{x \rightarrow \infty} \frac{1 - e^{-2x}}{27x^3} = 0$$

$$G(t) = \int \frac{dt}{\ln^3(t)}$$

$$\frac{d}{dx} \int_x^{3x} \frac{dt}{\ln^3(t)} = \frac{d}{dx} G(e^{3x}) - \frac{d}{dx} G(e^x) = G'(e^{3x}) \cdot e^{3x} \cdot 3 - G'(e^x) e^x$$

$$= \frac{3e^{3x}}{\ln^3(e^{3x})} - \frac{e^x}{\ln^3(e^x)} = \frac{3e^{3x}}{(3x)^3} - \frac{e^x}{x^3} = \frac{e^{3x}}{3x^3} - \frac{e^x}{x^3} = \frac{e^{3x} - e^x}{3x^3}$$

8-9.4 a

$$2y = -x^2 \rightarrow y = -\frac{1}{2}x^2$$

$$2x = -y^2 \rightarrow y = -\sqrt{-2x}$$

$$-\sqrt{-2x} = -\frac{1}{2}x^2$$


$$-2x = \frac{1}{4}x^4$$

$$\frac{1}{4}x^4 + 2x = 0$$

$$\frac{1}{4}x(x^3 + 8) = 0 \quad \frac{1}{4}x(x+2)(x^2 - 2x + 4) = 0$$

$$x = 0 \quad x = -2$$

$$y = 0 \quad y = -2$$



$$|S| = \int_{-2}^0 \left[-\frac{1}{2}x^2 - (-\sqrt{-2x}) \right] dx$$

$$= \int_{-2}^0 \left[-\frac{1}{2}x^2 + \sqrt{-2x} \right] dx = -\frac{1}{2} \int_{-2}^0 x^2 dx + \int_{-2}^0 \sqrt{-2x} dx$$

$$\int x^2 dx = \frac{1}{3}x^3 + C \quad \int \sqrt{-2x} dx = \left| \frac{t^{-2+1}}{-2+1} \right| = -\frac{1}{2} \int \sqrt{t} dt = -\frac{1}{2} \cdot \frac{2}{3} t^{\frac{3}{2}} = -\frac{1}{3}(-2x)\sqrt{-2x} = \frac{2}{3}x\sqrt{-2x}$$

$$= -\frac{1}{2} \cdot \frac{1}{3} x^3 \Big|_{-2}^0 + \frac{2}{3} x\sqrt{-2x} \Big|_{-2}^0$$

$$= -\frac{1}{6} \cdot (0 - (-8)) + \frac{2}{3} (0 - (-2\sqrt{4})) = -\frac{8}{6} + \frac{2}{3} \cdot 4 = \frac{16}{6} - \frac{8}{6} = \frac{8}{6} = \frac{4}{3}$$

$$\int \frac{dx}{\sqrt{x^2+k}} = \ln|x + \sqrt{x^2+k}| + C$$

$$\int \frac{u(x)}{\sqrt{ax^2+bx+c}} dx = u|_{x=-1}(x) \sqrt{ax^2+bx+c} + \lambda \int \frac{dx}{\sqrt{ax^2+bx+c}}$$

$$\int R(\sin(x), \cos(x)) dx = \left| \begin{array}{l} t = \tan(\frac{x}{2}) \\ dx = \frac{2}{1+t^2} dt \\ \sin(x) = \frac{2t}{1+t^2} \\ \cos(x) = \frac{1-t^2}{1+t^2} \end{array} \right|$$

$$\int f(x)g(x) dx = f(x)g(x) - \int f'(x)g(x) dx$$

$$\sin(ax)\cos(bx) = \frac{1}{2} [\sin((a-b)x) + \sin((a+b)x)]$$

$$\sin(ax)\sin(bx) = \frac{1}{2} [\cos((a-b)x) - \cos((a+b)x)]$$

$$\cos(ax)\cos(bx) = \frac{1}{2} [\cos((a-b)x) + \cos((a+b)x)]$$

$$\cos(2x) = 2\cos^2(x) - 1 = 1 - 2\sin^2(x)$$

$$\sin^2(x) = \frac{1 - \cos(2x)}{2}$$

$$\cos^2(x) = \frac{1 + \cos(2x)}{2}$$

$$\int \ln(x) dx = x\ln(x) - x$$

$$\int \frac{f'(x)}{f(x)} dx = \ln|f(x)|$$

$$\int_{h(x)}^{g(x)} f(t) dt = F(g(x)) - F(h(x))$$

$$F(t) = \int f(t) dt$$

$$\frac{d}{dx} \int_{h(x)}^{g(x)} f(t) dt = F'(g(x))g'(x) - F'(h(x))h'(x) = f(g(x))g'(x) - f(h(x))h'(x)$$

$$|V| = \pi \int_a^b f(x) dx$$

$$|L| = \int_a^b \sqrt{1 + [f'(x)]^2} dx$$

$$|S| = 2\pi \int_a^b f(x) \sqrt{1 + [f'(x)]^2} dx$$

$$1. \quad \int \frac{3x^2 + 3x + 1}{\sqrt{x^2 + 4x + 8}} dx = (Ax + B)\sqrt{x^2 + 4x + 8} + \lambda \int \frac{dx}{\sqrt{x^2 + 4x + 8}}$$

$$\frac{3x^2 + 3x + 1}{\sqrt{x^2 + 4x + 8}} = \frac{(Ax + B)(2x + 4)}{2\sqrt{x^2 + 4x + 8}} + A\sqrt{x^2 + 4x + 8} + \frac{\lambda}{\sqrt{x^2 + 4x + 8}}$$

$$3x^2 + 3x + 1 = (Ax + B)(x + 2) + A(x^2 + 4x + 8) + \lambda$$

$$3x^2 + 3x + 1 = Ax^2 + 2Ax + Bx + 2B + Ax^2 + 4Ax + 8A + \lambda$$

$$3x^2 + 3x + 1 = 2Ax^2 + (6A + B)x + (8A + 2B + \lambda)$$

$$\begin{cases} 3 = 2A \\ 3 = 6A + B \\ 1 = 8A + 2B + \lambda \end{cases} \quad \begin{cases} A = \frac{3}{2} \\ B = -6 \\ \lambda = 1 - 12 + 12 = 1 \end{cases}$$

$$\int \frac{dx}{\sqrt{x^2 + 4x + 8}} = \int \frac{dx}{\sqrt{x^2 + 4x + 4 + 4}} = \int \frac{dx}{\sqrt{(x+2)^2 + 4}} = \left| t = \frac{1}{2}(x+2) \right| = \int \frac{2dt}{\sqrt{4t^2 + 4}} = \int \frac{dt}{\sqrt{t^2 + 1}} =$$

$$\ln|t + \sqrt{t^2 + 1}| + C = \ln\left|\frac{x+2}{2} + \sqrt{\left(\frac{x+2}{2}\right)^2 + 1}\right| + C = \ln|x+2 + \sqrt{x^2 + 4x + 8}| + C$$

$$\int \frac{3x^2 + 3x + 1}{\sqrt{x^2 + 4x + 8}} dx = \left(\frac{3}{2}x - 6\right)\sqrt{x^2 + 4x + 8} + \ln|x+2 + \sqrt{x^2 + 4x + 8}| + C$$

$$2? \quad \int_0^{+\infty} \frac{e^x}{1 + e^{2x}} dx = \lim_{T \rightarrow +\infty} \int_0^T \frac{e^x}{1 + e^{2x}} dx \quad \left| \begin{array}{l} t = e^x \\ dt = e^x dx \end{array} \right| = \lim_{T \rightarrow +\infty} \int_1^{e^T} \frac{dt}{t^2 + 1} = \lim_{T \rightarrow +\infty} \arctan(t) \Big|_1^{e^T} = \lim_{T \rightarrow +\infty} \arctan(e^T) - \arctan(1) = \frac{\pi}{2} - \frac{\pi}{4} = \frac{\pi}{4}$$