MIPT & YSDA · AUTUMN 2026

Deep Generative Models

Lecture 7

Roman Isachenko

Moscow Institute of Physics and Technology
Yandex School of Data Analysis

Recap of Previous Lecture


Define **score function** $\mathbf{s}_{\boldsymbol{\theta}}(\mathbf{x}) = \nabla_{\mathbf{x}} \log p_{\boldsymbol{\theta}}(\mathbf{x})$.

Fisher Divergence

$$D_F(p_{\text{data}}, p_{\boldsymbol{\theta}}) = \frac{1}{2}\mathbb{E}_{p_{\text{data}}} \left\| \mathbf{s}_{\boldsymbol{\theta}}(\mathbf{x}) - \nabla_{\mathbf{x}} \log p_{\text{data}}(\mathbf{x}) \right\|_2^2 \to \min_{\boldsymbol{\theta}}$$

Langevin Dynamics

$$\mathbf{x}_{l+1} = \mathbf{x}_l + \frac{\eta}{2} \cdot \mathbf{s}_{\boldsymbol{\theta}}(\mathbf{x}_l) + \sqrt{\eta} \cdot \boldsymbol{\epsilon}_l, \quad \boldsymbol{\epsilon}_l \sim \mathcal{N}(0, \mathbf{I})$$



Recap of Previous Lecture

Let us perturb the original data with Gaussian noise $q(\mathbf{x}_\sigma|\mathbf{x}) = \mathcal{N}(\mathbf{x}, \sigma^2 \cdot \mathbf{I})$.


$$q(\mathbf{x}_\sigma) = \int q(\mathbf{x}_\sigma|\mathbf{x})p_{\text{data}}(\mathbf{x})d\mathbf{x}.$$

Assumption

The solution to

$$\frac{1}{2}\mathbb{E}_{q(\mathbf{x}_\sigma)}\left\|\mathbf{s}_{\boldsymbol{\theta},\sigma}(\mathbf{x}_\sigma) - \nabla_{\mathbf{x}_\sigma}\log q(\mathbf{x}_\sigma)\right\|_2^2 \to \min_{\boldsymbol{\theta}}$$

satisfies $\mathbf{s}_{\boldsymbol{\theta},\sigma}(\mathbf{x}_\sigma) \approx \mathbf{s}_{\boldsymbol{\theta},0}(\mathbf{x}_0) = \mathbf{s}_{\boldsymbol{\theta}}(\mathbf{x})$ if σ is sufficiently small.



Song Y. Generative Modeling by Estimating Gradients of the Data Distribution, blog post, 2021
Vincent P. A Connection Between Score Matching and Denoising Autoencoders, 2010

Recap of Previous Lecture

Theorem

Under mild regularity conditions, the last term is independent of $\boldsymbol{\theta}$:

$$\begin{aligned}&\mathbb{E}_{q(\mathbf{x}_\sigma)}\left\|\mathbf{s}_{\boldsymbol{\theta},\sigma}(\mathbf{x}_\sigma) - \nabla_{\mathbf{x}_\sigma}\log q(\mathbf{x}_\sigma)\right\|_2^2 \\ &\quad= \mathbb{E}_{p_{\text{data}}(\mathbf{x})}\mathbb{E}_{q(\mathbf{x}_\sigma|\mathbf{x})}\left\|\mathbf{s}_{\boldsymbol{\theta},\sigma}(\mathbf{x}_\sigma) - \nabla_{\mathbf{x}_\sigma}\log q(\mathbf{x}_\sigma|\mathbf{x})\right\|_2^2 + \text{const}\end{aligned}$$

Here,

$$\nabla_{\mathbf{x}_\sigma}\log q(\mathbf{x}_\sigma|\mathbf{x}) = -\frac{\mathbf{x}_\sigma - \mathbf{x}}{\sigma^2} = -\frac{\boldsymbol{\epsilon}}{\sigma}.$$

Tweedie's Formula

From Lecture 6, the MSE-optimal clean-data prediction is

$$\mathbb{E}_{q(\mathbf{x}|\mathbf{x}_\sigma)}[\mathbf{x}] = \mathbf{x}_\sigma + \sigma^2\nabla_{\mathbf{x}_\sigma}\log q(\mathbf{x}_\sigma).$$


Vincent P. A Connection Between Score Matching and Denoising Autoencoders, 2010
Efron B. Tweedie's Formula and Selection Bias, 2011

Recap of Previous Lecture

$$\mathbb{E}_{p_{\text{data}}(\mathbf{x})}\mathbb{E}_{\mathcal{N}(0,\mathbf{I})}\left\|\mathbf{s}_{\boldsymbol{\theta},\sigma}(\mathbf{x}+\sigma\boldsymbol{\epsilon})+\frac{\boldsymbol{\epsilon}}{\sigma}\right\|_2^2 \to \min_{\boldsymbol{\theta}}$$

$$\mathbf{x}_{l+1} = \mathbf{x}_l + \frac{\eta}{2}\cdot\mathbf{s}_{\boldsymbol{\theta},\sigma}(\mathbf{x}_l) + \sqrt{\eta}\cdot\boldsymbol{\epsilon}_l$$

- For **small** σ, $\mathbf{s}_{\boldsymbol{\theta},\sigma}(\mathbf{x})$ becomes inaccurate and Langevin dynamics fails to traverse modes
- For **large** σ, robustness in low-density regions is achieved, but the model learns a distribution that is overly corrupted




Recap of Previous Lecture

Noise-Conditioned Score Network (NCSN)

- Define a sequence of noise levels: $\sigma_1 < \sigma_2 < \cdots < \sigma_T$.
- Train the denoising score function $\mathbf{s}_{\boldsymbol{\theta},\sigma_t}(\mathbf{x}_t)$ for each noise level:

$$\sum_{t=1}^{T} \sigma_t^2 \cdot \mathbb{E}_{p_{\text{data}}(\mathbf{x})} \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x})} \left\| \mathbf{s}_{\boldsymbol{\theta},\sigma_t}(\mathbf{x}_t) - \nabla_{\mathbf{x}_t} \log q(\mathbf{x}_t|\mathbf{x}) \right\|_2^2 \to \min_{\boldsymbol{\theta}}$$

- Sample using **annealed** Langevin dynamics (for $t = T, \ldots, 1$).



Song Y. et al. Generative Modeling by Estimating Gradients of the Data Distribution, 2019

Recap of Previous Lecture

Training

1. Sample $\mathbf{x}_0 \sim p_{\text{data}}(\mathbf{x})$, $t \sim U\{1, T\}$, $\boldsymbol{\epsilon} \sim \mathcal{N}(0, \mathbf{I})$.
2. Compute noisy image $\mathbf{x}_t = \mathbf{x}_0 + \sigma_t \boldsymbol{\epsilon}$.
3. Compute loss $\mathcal{L} = \sigma_t^2 \left\| \mathbf{s}_{\boldsymbol{\theta}, \sigma_t}(\mathbf{x}_t) + \frac{\boldsymbol{\epsilon}}{\sigma_t} \right\|^2$.
4. Update $\boldsymbol{\theta}$ via stochastic gradient descent.

Sampling (Annealed Langevin Dynamics)

1. Sample $\mathbf{x}_T^0 \sim \mathcal{N}(0, \sigma_T^2 \mathbf{I}) \approx q(\mathbf{x}_T)$.
2. Update $\mathbf{x}_t^l$ via L steps of Langevin dynamics at each noise level σ_t:

$$\mathbf{x}_t^l = \mathbf{x}_t^{l-1} + \frac{\eta_t}{2} \mathbf{s}_{\boldsymbol{\theta}, \sigma_t}(\mathbf{x}_t^{l-1}) + \sqrt{\eta_t} \boldsymbol{\epsilon}_t^l.$$

3. Update $\mathbf{x}_{t-1}^0 := \mathbf{x}_t^L$ and proceed to the next noise level σ_{t-1}.

Song Y. et al. Generative Modeling by Estimating Gradients of the Data Distribution, 2019
Song Y. et al. Improved Techniques for Training Score-Based Generative Models, 2020

Recap of Previous Lecture

$$\mathbf{x}_t = \sqrt{1-\beta_t}\mathbf{x}_{t-1} + \sqrt{\beta_t}\boldsymbol{\epsilon}_t, \quad \boldsymbol{\epsilon}_t \sim \mathcal{N}(0, \mathbf{I})$$

$$q(\mathbf{x}_t|\mathbf{x}_{t-1}) = \mathcal{N}(\sqrt{1-\beta_t}\mathbf{x}_{t-1}, \beta_t\mathbf{I})$$


Statement 1

Let $\alpha_t = 1 - \beta_t$ and $\bar{\alpha}_t = \prod_{s=1}^{t} \alpha_s = \prod_{s=1}^{t}(1-\beta_s)$. Then

$$q(\mathbf{x}_t|\mathbf{x}_0) = \mathcal{N}(\sqrt{\bar{\alpha}_t}\,\mathbf{x}_0, (1-\bar{\alpha}_t)\mathbf{I})$$

Recap of Previous Lecture

Diffusion describes the migration of particles from regions of high density to those of low density.



1. $\mathbf{x}_0 = \mathbf{x} \sim p_{\text{data}}(\mathbf{x})$
2. $\mathbf{x}_t = \sqrt{1-\beta_t}\mathbf{x}_{t-1} + \sqrt{\beta_t}\boldsymbol{\epsilon}_t$, $\boldsymbol{\epsilon}_t \sim \mathcal{N}(0, \mathbf{I})$, $t \geq 1$
3. After $T \gg 1$ steps: $\mathbf{x}_T \sim p_\infty(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$

If this process can be reversed, we can sample from $p_{\text{data}}(\mathbf{x})$ by starting from noise $p_\infty(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$.
Our goal now becomes inverting this diffusion.

Outline

01 Reverse Gaussian Diffusion Process

02 Gaussian Diffusion Model as VAE

03 Diffusion ELBO Derivation

04 Gaussian Diffusion Reparametrization

05 Denoising Diffusion Probabilistic Model (DDPM)

Outline

01 **Reverse Gaussian Diffusion Process**


02 Gaussian Diffusion Model as VAE

03 Diffusion ELBO Derivation


04 Gaussian Diffusion Reparametrization

05 Denoising Diffusion Probabilistic Model (DDPM)

Reverse Gaussian Diffusion Process




Reverse Gaussian Diffusion Process



Forward Process

$$q(\mathbf{x}_t|\mathbf{x}_{t-1}) = \mathcal{N}\left(\sqrt{1-\beta_t}\mathbf{x}_{t-1}, \beta_t\mathbf{I}\right)$$

Reverse Gaussian Diffusion Process




Forward Process

$$q(\mathbf{x}_t|\mathbf{x}_{t-1}) = \mathcal{N}\left(\sqrt{1-\beta_t}\mathbf{x}_{t-1}, \beta_t\mathbf{I}\right)$$

Reverse Process

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) = \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1})q(\mathbf{x}_{t-1})}{q(\mathbf{x}_t)} \approx p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)$$

Reverse Gaussian Diffusion Process



Forward Process

$$q(\mathbf{x}_t|\mathbf{x}_{t-1}) = \mathcal{N}\left(\sqrt{1-\beta_t}\mathbf{x}_{t-1}, \beta_t\mathbf{I}\right)$$

Reverse Process

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) = \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1})q(\mathbf{x}_{t-1})}{q(\mathbf{x}_t)} \approx p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)$$

$q(\mathbf{x}_{t-1})$ and $q(\mathbf{x}_t)$ are intractable:

$$q(\mathbf{x}_t) = \int q(\mathbf{x}_t|\mathbf{x}_0)p_{\text{data}}(\mathbf{x}_0)d\mathbf{x}_0$$

Reverse Gaussian Diffusion Process

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) = \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1})q(\mathbf{x}_{t-1})}{q(\mathbf{x}_t)}$$

Theorem (Feller, 1949)

If β_t is sufficiently small, $q(\mathbf{x}_{t-1}|\mathbf{x}_t)$ is approximately Gaussian.

The original DDPM used $T = 1000$ steps. Faster sampling methods will be discussed later.

Feller W. On the theory of stochastic processes, with particular reference to applications, 1949
Ho J. et al. Denoising Diffusion Probabilistic Models, 2020
Xiao Z., Kreis K., Vahdat A. Tackling the generative learning trilemma with denoising diffusion GANs, 2021


Reverse Gaussian Diffusion Process

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) = \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1})q(\mathbf{x}_{t-1})}{q(\mathbf{x}_t)}$$

Theorem (Feller, 1949)


If β_t is sufficiently small, $q(\mathbf{x}_{t-1}|\mathbf{x}_t)$ is approximately Gaussian.

The original DDPM used $T = 1000$ steps. Faster sampling methods will be discussed later.



Feller W. On the theory of stochastic processes, with particular reference to applications, 1949
Ho J. et al. Denoising Diffusion Probabilistic Models, 2020
Xiao Z., Kreis K., Vahdat A. Tackling the generative learning trilemma with denoising diffusion GANs, 2021

Reverse Gaussian Diffusion Process (Ancestral Sampling)




Define the reverse process as:

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) \approx p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t) = \mathcal{N}\left(\boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t), \boldsymbol{\sigma}^2_{\boldsymbol{\theta},t}(\mathbf{x}_t)\right)$$

If β_t is sufficiently small, $q(\mathbf{x}_{t-1}|\mathbf{x}_t)$ is approximately Gaussian.


Reverse Gaussian Diffusion Process (Ancestral Sampling)



Forward Process

1. $\mathbf{x}_0 = \mathbf{x} \sim p_{\text{data}}(\mathbf{x})$
2. $\mathbf{x}_t = \sqrt{1-\beta_t}\mathbf{x}_{t-1} + \sqrt{\beta_t}\boldsymbol{\epsilon}_t$
3. $\mathbf{x}_T \sim p_\infty(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$

Reverse Gaussian Diffusion Process (Ancestral Sampling)




Forward Process

1. $\mathbf{x}_0 = \mathbf{x} \sim p_{\text{data}}(\mathbf{x})$
2. $\mathbf{x}_t = \sqrt{1-\beta_t}\mathbf{x}_{t-1} + \sqrt{\beta_t}\boldsymbol{\epsilon}_t$
3. $\mathbf{x}_T \sim p_\infty(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$

Reverse Process

1. $\mathbf{x}_T \sim p_\infty(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$
2. $\mathbf{x}_{t-1} = \boldsymbol{\sigma}_{\boldsymbol{\theta},t}(\mathbf{x}_t)\boldsymbol{\epsilon} + \boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t)$
3. $\mathbf{x}_0 = \mathbf{x} \sim p_{\text{data}}(\mathbf{x})$

Reverse Gaussian Diffusion Process (Ancestral Sampling)



Forward Process

1. $\mathbf{x}_0 = \mathbf{x} \sim p_{\text{data}}(\mathbf{x})$
2. $\mathbf{x}_t = \sqrt{1-\beta_t}\mathbf{x}_{t-1} + \sqrt{\beta_t}\boldsymbol{\epsilon}_t$
3. $\mathbf{x}_T \sim p_\infty(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$

Reverse Process

1. $\mathbf{x}_T \sim p_\infty(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$
2. $\mathbf{x}_{t-1} = \boldsymbol{\sigma}_{\boldsymbol{\theta},t}(\mathbf{x}_t)\boldsymbol{\epsilon} + \boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t)$
3. $\mathbf{x}_0 = \mathbf{x} \sim p_{\text{data}}(\mathbf{x})$

Note: The forward process is non-learnable, i.e., it does not involve trainable parameters

Weng L. What are Diffusion Models?, blog post, 2021

Conditioned Reverse Distribution

Reverse Kernel (Intractable)

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) = \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1})q(\mathbf{x}_{t-1})}{q(\mathbf{x}_t)}$$

Conditioned Reverse Distribution

Reverse Kernel (Intractable)

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) = \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1})q(\mathbf{x}_{t-1})}{q(\mathbf{x}_t)}$$

Conditioned Reverse Kernel (Tractable)

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) = \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1}, \mathbf{x}_0)q(\mathbf{x}_{t-1}|\mathbf{x}_0)}{q(\mathbf{x}_t|\mathbf{x}_0)}$$

Conditioned Reverse Distribution

Reverse Kernel (Intractable)

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) = \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1})q(\mathbf{x}_{t-1})}{q(\mathbf{x}_t)}$$

Conditioned Reverse Kernel (Tractable)

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) = \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1}, \mathbf{x}_0)q(\mathbf{x}_{t-1}|\mathbf{x}_0)}{q(\mathbf{x}_t|\mathbf{x}_0)}$$

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) = \frac{\mathcal{N}(\sqrt{1-\beta_t}\cdot\mathbf{x}_{t-1}, \beta_t\mathbf{I})\cdot\mathcal{N}(\sqrt{\bar{\alpha}_{t-1}}\cdot\mathbf{x}_0, (1-\bar{\alpha}_{t-1})\cdot\mathbf{I})}{\mathcal{N}(\sqrt{\bar{\alpha}_t}\cdot\mathbf{x}_0, (1-\bar{\alpha}_t)\cdot\mathbf{I})}$$

Conditioned Reverse Distribution

Reverse Kernel (Intractable)

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) = \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1})q(\mathbf{x}_{t-1})}{q(\mathbf{x}_t)}$$

Conditioned Reverse Kernel (Tractable)

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) = \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1}, \mathbf{x}_0)q(\mathbf{x}_{t-1}|\mathbf{x}_0)}{q(\mathbf{x}_t|\mathbf{x}_0)}$$

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) = \frac{\mathcal{N}(\sqrt{1-\beta_t}\cdot\mathbf{x}_{t-1}, \beta_t\mathbf{I})\cdot\mathcal{N}(\sqrt{\bar{\alpha}_{t-1}}\cdot\mathbf{x}_0, (1-\bar{\alpha}_{t-1})\cdot\mathbf{I})}{\mathcal{N}(\sqrt{\bar{\alpha}_t}\cdot\mathbf{x}_0, (1-\bar{\alpha}_t)\cdot\mathbf{I})}$$

$$= \mathcal{N}(\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0), \tilde{\beta}_t\cdot\mathbf{I})$$

$$\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) = \frac{\sqrt{\alpha_t}(1-\bar{\alpha}_{t-1})}{1-\bar{\alpha}_t}\cdot\mathbf{x}_t + \frac{\sqrt{\bar{\alpha}_{t-1}}(1-\alpha_t)}{1-\bar{\alpha}_t}\cdot\mathbf{x}_0$$

$$\tilde{\beta}_t = \frac{(1-\alpha_t)(1-\bar{\alpha}_{t-1})}{1-\bar{\alpha}_t} = \text{const}$$

What Does Knowing the Clean Data Change?

Same observation $x_t = 0$: unknown clean data versus each known clean value.

Two possible clean values

$\mathbb{P}(x_0 = -2) = \mathbb{P}(x_0 = 2) = \frac{1}{2}$

$x_0 = -2$


Posterior weight 50.0%

$x_0 = 2$

Posterior weight 50.0%

Unknown x_0: posterior-weighted mixture.
Known x_0: the corresponding Gaussian.

$q(x_{t-1}|x_t)$ and $q(x_{t-1}|x_t, x_0)$



$$q(x_{t-1}|x_t) = \sum_{x_0 \in \{-2,2\}} q(x_{t-1}|x_t, x_0)\, q(x_0|x_t)$$

Exact toy example. During training x_0 is available; during generation it is unknown.

Distribution Summary

Forward process maps any distribution $p_{\text{data}}(\mathbf{x})$ to $\mathcal{N}(0, \mathbf{I})$ by injection of noise:

$$q(\mathbf{x}_t|\mathbf{x}_{t-1}) = \mathcal{N}(\sqrt{1-\beta_t} \cdot \mathbf{x}_{t-1}, \beta_t \cdot \mathbf{I});$$
$$q(\mathbf{x}_t|\mathbf{x}_0) = \mathcal{N}(\sqrt{\bar{\alpha}_t} \cdot \mathbf{x}_0, (1-\bar{\alpha}_t) \cdot \mathbf{I}).$$

Distribution Summary

Forward process maps any distribution $p_{\text{data}}(\mathbf{x})$ to $\mathcal{N}(0, \mathbf{I})$ by injection of noise:

$$q(\mathbf{x}_t|\mathbf{x}_{t-1}) = \mathcal{N}(\sqrt{1-\beta_t} \cdot \mathbf{x}_{t-1}, \beta_t \cdot \mathbf{I});$$
$$q(\mathbf{x}_t|\mathbf{x}_0) = \mathcal{N}(\sqrt{\bar{\alpha}_t} \cdot \mathbf{x}_0, (1-\bar{\alpha}_t) \cdot \mathbf{I}).$$

Reverse process refers to an intractable distribution that can be approximated by a normal distribution (with unknown parameters) for small β_t:

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) = \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1})q(\mathbf{x}_{t-1})}{q(\mathbf{x}_t)} \approx \mathcal{N}\left(\boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t), \boldsymbol{\sigma}^2_{\boldsymbol{\theta},t}(\mathbf{x}_t)\right)$$

Distribution Summary

Forward process maps any distribution $p_{\text{data}}(\mathbf{x})$ to $\mathcal{N}(0, \mathbf{I})$ by injection of noise:

$$q(\mathbf{x}_t|\mathbf{x}_{t-1}) = \mathcal{N}(\sqrt{1-\beta_t} \cdot \mathbf{x}_{t-1}, \beta_t \cdot \mathbf{I});$$
$$q(\mathbf{x}_t|\mathbf{x}_0) = \mathcal{N}(\sqrt{\bar{\alpha}_t} \cdot \mathbf{x}_0, (1-\bar{\alpha}_t) \cdot \mathbf{I}).$$

Reverse process refers to an intractable distribution that can be approximated by a normal distribution (with unknown parameters) for small β_t:

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) = \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1})q(\mathbf{x}_{t-1})}{q(\mathbf{x}_t)} \approx \mathcal{N}\left(\boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t), \boldsymbol{\sigma}^2_{\boldsymbol{\theta},t}(\mathbf{x}_t)\right)$$

Conditioned reverse process is a normal distribution with known parameters, describing how to denoise a noisy image $\mathbf{x}_t$ when we know the clean image $\mathbf{x}_0$.

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) = \mathcal{N}(\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0), \tilde{\beta}_t \cdot \mathbf{I})$$

Noise Schedule through Signal-to-Noise Ratio

Signal and Noise

$$\mathbf{x}_t = \sqrt{\bar{\alpha}_t}\mathbf{x}_0 + \sqrt{1 - \bar{\alpha}_t}\boldsymbol{\epsilon}.$$

For $\boldsymbol{\epsilon} \sim \mathcal{N}(0, \mathbf{I})$, compare squared scales:

$$\mathrm{SNR}(t) = \frac{\bar{\alpha}_t}{1 - \bar{\alpha}_t}.$$

Kingma D. et al. Variational Diffusion Models, 2021
Nichol A., Dhariwal P. Improved Denoising Diffusion Probabilistic Models, 2021
Hoogeboom E. et al. Simple diffusion: End-to-end diffusion for high resolution images, 2023

Noise Schedule through Signal-to-Noise Ratio

Signal and Noise

$$\mathbf{x}_t = \sqrt{\bar{\alpha}_t}\mathbf{x}_0 + \sqrt{1-\bar{\alpha}_t}\boldsymbol{\epsilon}.$$


For $\boldsymbol{\epsilon} \sim \mathcal{N}(0, \mathbf{I})$, compare squared scales:

$$\mathrm{SNR}(t) = \frac{\bar{\alpha}_t}{1-\bar{\alpha}_t}.$$

A Single Noise-Level Coordinate

$$\lambda_t = \log \mathrm{SNR}(t), \qquad \bar{\alpha}_t = \frac{1}{1+e^{-\lambda_t}}.$$

$\lambda_t > 0$: signal dominates.
$\lambda_t = 0$: equal signal and noise scales.
$\lambda_t < 0$: noise dominates.



Same clean image and Gaussian noise.
Fixed display scale; log-SNR axis clipped to [−12, 10].

Same timestep, different noise level.

Kingma D. et al. Variational Diffusion Models, 2021
Nichol A., Dhariwal P. Improved Denoising Diffusion Probabilistic Models, 2021
Hoogeboom E. et al. Simple diffusion: End-to-end diffusion for high resolution images, 2023

Noise Schedule through Signal-to-Noise Ratio

Signal and Noise

$$\mathbf{x}_t = \sqrt{\bar{\alpha}_t}\mathbf{x}_0 + \sqrt{1-\bar{\alpha}_t}\boldsymbol{\epsilon}.$$


For $\boldsymbol{\epsilon} \sim \mathcal{N}(0, \mathbf{I})$, compare squared scales:

$$\mathrm{SNR}(t) = \frac{\bar{\alpha}_t}{1-\bar{\alpha}_t}.$$

A Single Noise-Level Coordinate

$$\lambda_t = \log \mathrm{SNR}(t), \qquad \bar{\alpha}_t = \frac{1}{1+e^{-\lambda_t}}.$$

$\lambda_t > 0$: signal dominates.
$\lambda_t = 0$: equal signal and noise scales.
$\lambda_t < 0$: noise dominates.



Same clean image and Gaussian noise.
Fixed display scale; log-SNR axis clipped to [−12, 10].

Same timestep, different noise level.

> **Higher pixel-space resolution:** more noise can help; shifting log-SNR down increases corruption.

Kingma D. et al. Variational Diffusion Models, 2021
Nichol A., Dhariwal P. Improved Denoising Diffusion Probabilistic Models, 2021
Hoogeboom E. et al. Simple diffusion: End-to-end diffusion for high resolution images, 2023

Outline

01 Reverse Gaussian Diffusion Process

02 Gaussian Diffusion Model as VAE

03 Diffusion ELBO Derivation

04 Gaussian Diffusion Reparametrization

05 Denoising Diffusion Probabilistic Model (DDPM)

Gaussian Diffusion Model as VAE

Let's treat $\mathbf{z} = (\mathbf{x}_1, \dots, \mathbf{x}_T)$ as a latent variable (**note:** each $\mathbf{x}_t$ has the same dimension), and $\mathbf{x} = \mathbf{x}_0$ as the observed variable.

Latent Variable Model


$$p_{\boldsymbol{\theta}}(\mathbf{x}, \mathbf{z}) = p_{\boldsymbol{\theta}}(\mathbf{x}|\mathbf{z})p_{\boldsymbol{\theta}}(\mathbf{z})$$

Gaussian Diffusion Model as VAE

Let's treat $\mathbf{z} = (\mathbf{x}_1, \dots, \mathbf{x}_T)$ as a latent variable (**note:** each $\mathbf{x}_t$ has the same dimension), and $\mathbf{x} = \mathbf{x}_0$ as the observed variable.

Latent Variable Model

$$p_{\boldsymbol{\theta}}(\mathbf{x}, \mathbf{z}) = p_{\boldsymbol{\theta}}(\mathbf{x}|\mathbf{z})p_{\boldsymbol{\theta}}(\mathbf{z})$$




Gaussian Diffusion Model as VAE

Let's treat $\mathbf{z} = (\mathbf{x}_1, \dots, \mathbf{x}_T)$ as a latent variable (**note:** each $\mathbf{x}_t$ has the same dimension), and $\mathbf{x} = \mathbf{x}_0$ as the observed variable.

Latent Variable Model

$$p_{\boldsymbol{\theta}}(\mathbf{x}, \mathbf{z}) = p_{\boldsymbol{\theta}}(\mathbf{x}|\mathbf{z})p_{\boldsymbol{\theta}}(\mathbf{z})$$



Forward Diffusion

Variational posterior (encoder), with **no learnable parameters**:

$$q(\mathbf{z}|\mathbf{x}) = q(\mathbf{x}_1, \dots, \mathbf{x}_T|\mathbf{x}_0) = \prod_{t=1}^{T} q(\mathbf{x}_t|\mathbf{x}_{t-1}).$$


Gaussian Diffusion Model as VAE

$$p_{\boldsymbol{\theta}}(\mathbf{x}, \mathbf{z}) = p_{\boldsymbol{\theta}}(\mathbf{x}|\mathbf{z})p_{\boldsymbol{\theta}}(\mathbf{z})$$

Gaussian Diffusion Model as VAE

$$p_{\theta}(\mathbf{x}, \mathbf{z}) = p_{\theta}(\mathbf{x}|\mathbf{z})p_{\theta}(\mathbf{z})$$

$\mathbf{z} = (\mathbf{x}_1, \ldots, \mathbf{x}_T)$ latent trajectory




observed data

Gaussian Diffusion Model as VAE

$$p_{\theta}(\mathbf{x}, \mathbf{z}) = p_{\theta}(\mathbf{x}|\mathbf{z})p_{\theta}(\mathbf{z})$$

$\mathbf{z} = (\mathbf{x}_1, \ldots, \mathbf{x}_T)$ latent trajectory



Reverse Diffusion


Generative distribution (decoder)

$$p_{\theta}(\mathbf{x}|\mathbf{z}) = p_{\theta}(\mathbf{x}_0|\mathbf{x}_1).$$

Gaussian Diffusion Model as VAE

$$p_{\theta}(\mathbf{x}, \mathbf{z}) = p_{\theta}(\mathbf{x}|\mathbf{z})p_{\theta}(\mathbf{z})$$

$\mathbf{z} = (\mathbf{x}_1, \ldots, \mathbf{x}_T)$ latent trajectory



observed data

$p_{\theta}(\mathbf{z})$ learned reverse transitions + terminal $p(\mathbf{x}_T)$

Reverse Diffusion

Generative distribution (decoder)


$$p_{\theta}(\mathbf{x}|\mathbf{z}) = p_{\theta}(\mathbf{x}_0|\mathbf{x}_1).$$

Prior distribution

$$p_{\theta}(\mathbf{z}) = p_{\theta}(\mathbf{x}_1, \ldots, \mathbf{x}_T) = \prod_{t=2}^{T} p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t) \cdot p(\mathbf{x}_T).$$

Gaussian Diffusion Model as VAE

$$p_{\theta}(\mathbf{x}, \mathbf{z}) = p_{\theta}(\mathbf{x}|\mathbf{z})p_{\theta}(\mathbf{z})$$



Reverse Diffusion

Generative distribution (decoder)

$$p_{\theta}(\mathbf{x}|\mathbf{z}) = p_{\theta}(\mathbf{x}_0|\mathbf{x}_1).$$

Prior distribution

$$p_{\theta}(\mathbf{z}) = p_{\theta}(\mathbf{x}_1, \ldots, \mathbf{x}_T) = \prod_{t=2}^{T} p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t) \cdot p(\mathbf{x}_T).$$

Unlike the vanilla VAE: a **simple decoder** $p_{\theta}(\mathbf{x}|\mathbf{z})$ and a **Markov-chain prior** $p_{\theta}(\mathbf{z})$.

Outline

01 Reverse Gaussian Diffusion Process

02 Gaussian Diffusion Model as VAE

03 Diffusion ELBO Derivation

04 Gaussian Diffusion Reparametrization

05 Denoising Diffusion Probabilistic Model (DDPM)

ELBO for Gaussian Diffusion Model

Standard ELBO

$$\log p_{\boldsymbol{\theta}}(\mathbf{x}) \geq \mathbb{E}_{q(\mathbf{z}|\mathbf{x})} \log \frac{p_{\boldsymbol{\theta}}(\mathbf{x}, \mathbf{z})}{q(\mathbf{z}|\mathbf{x})} = \mathcal{L}_{\boldsymbol{\phi},\boldsymbol{\theta}}(\mathbf{x}) \to \max_{\boldsymbol{\phi},\boldsymbol{\theta}}$$

ELBO for Gaussian Diffusion Model

Standard ELBO

$$\log p_{\boldsymbol{\theta}}(\mathbf{x}) \geq \mathbb{E}_{q(\mathbf{z}|\mathbf{x})} \log \frac{p_{\boldsymbol{\theta}}(\mathbf{x}, \mathbf{z})}{q(\mathbf{z}|\mathbf{x})} = \mathcal{L}_{\boldsymbol{\phi},\boldsymbol{\theta}}(\mathbf{x}) \to \max_{\boldsymbol{\phi},\boldsymbol{\theta}}$$

Derivation

$$\mathcal{L}_{\boldsymbol{\phi},\boldsymbol{\theta}}(\mathbf{x}) = \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \log \frac{p_{\boldsymbol{\theta}}(\mathbf{x}_0, \mathbf{x}_{1:T})}{q(\mathbf{x}_{1:T}|\mathbf{x}_0)}$$

ELBO for Gaussian Diffusion Model

Standard ELBO

$$\log p_{\boldsymbol{\theta}}(\mathbf{x}) \geq \mathbb{E}_{q(\mathbf{z}|\mathbf{x})} \log \frac{p_{\boldsymbol{\theta}}(\mathbf{x}, \mathbf{z})}{q(\mathbf{z}|\mathbf{x})} = \mathcal{L}_{\boldsymbol{\phi},\boldsymbol{\theta}}(\mathbf{x}) \to \max_{\boldsymbol{\phi},\boldsymbol{\theta}}$$

Derivation

$$\begin{aligned}\mathcal{L}_{\boldsymbol{\phi},\boldsymbol{\theta}}(\mathbf{x}) &= \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \log \frac{p_{\boldsymbol{\theta}}(\mathbf{x}_0, \mathbf{x}_{1:T})}{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \\ &= \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \log \frac{p(\mathbf{x}_T) \prod_{t=1}^{T} p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{\prod_{t=1}^{T} q(\mathbf{x}_t|\mathbf{x}_{t-1})}\end{aligned}$$

ELBO for Gaussian Diffusion Model

Standard ELBO

$$\log p_{\boldsymbol{\theta}}(\mathbf{x}) \geq \mathbb{E}_{q(\mathbf{z}|\mathbf{x})} \log \frac{p_{\boldsymbol{\theta}}(\mathbf{x}, \mathbf{z})}{q(\mathbf{z}|\mathbf{x})} = \mathcal{L}_{\boldsymbol{\phi},\boldsymbol{\theta}}(\mathbf{x}) \to \max_{\boldsymbol{\phi},\boldsymbol{\theta}}$$

Derivation

$$\begin{aligned}\mathcal{L}_{\boldsymbol{\phi},\boldsymbol{\theta}}(\mathbf{x}) &= \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \log \frac{p_{\boldsymbol{\theta}}(\mathbf{x}_0, \mathbf{x}_{1:T})}{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \\ &= \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \log \frac{p(\mathbf{x}_T) \prod_{t=1}^{T} p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{\prod_{t=1}^{T} q(\mathbf{x}_t|\mathbf{x}_{t-1})}\end{aligned}$$

- Let's try to decompose the ELBO into individual KL divergence terms.
- We need to replace $q(\mathbf{x}_t|\mathbf{x}_{t-1})$ with $q(\mathbf{x}_{t-1}|\mathbf{x}_t)$ in the denominator.
- Let's condition on $\mathbf{x}_0$ to make the reverse $q(\mathbf{x}_{t-1}|\mathbf{x}_t)$ tractable.

ELBO for Gaussian Diffusion Model

Bayes' Rule ($t \geq 2$)

$$q(\mathbf{x}_t|\mathbf{x}_{t-1}, \mathbf{x}_0) = \frac{q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0)q(\mathbf{x}_t|\mathbf{x}_0)}{q(\mathbf{x}_{t-1}|\mathbf{x}_0)}$$

Markov Property

$$q(\mathbf{x}_t|\mathbf{x}_{t-1}, \mathbf{x}_0) = q(\mathbf{x}_t|\mathbf{x}_{t-1})$$

ELBO for Gaussian Diffusion Model

Bayes' Rule ($t \geq 2$)

$$q(\mathbf{x}_t|\mathbf{x}_{t-1}, \mathbf{x}_0) = \frac{q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0)q(\mathbf{x}_t|\mathbf{x}_0)}{q(\mathbf{x}_{t-1}|\mathbf{x}_0)}$$

$$\mathcal{L}_{\phi,\theta}(\mathbf{x}) = \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \log \frac{p(\mathbf{x}_T)\prod_{t=1}^{T} p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{\prod_{t=1}^{T} q(\mathbf{x}_t|\mathbf{x}_{t-1}, \mathbf{x}_0)}$$

Markov Property

$$q(\mathbf{x}_t|\mathbf{x}_{t-1}, \mathbf{x}_0) = q(\mathbf{x}_t|\mathbf{x}_{t-1})$$

Conditioning on $\mathbf{x}_0$ leaves the forward transitions unchanged.

ELBO for Gaussian Diffusion Model

Bayes' Rule ($t \geq 2$)

$$q(\mathbf{x}_t|\mathbf{x}_{t-1}, \mathbf{x}_0) = \frac{q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0)q(\mathbf{x}_t|\mathbf{x}_0)}{q(\mathbf{x}_{t-1}|\mathbf{x}_0)}$$

$$\mathcal{L}_{\phi,\theta}(\mathbf{x}) = \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \log \frac{p(\mathbf{x}_T)\prod_{t=1}^{T} p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{\prod_{t=1}^{T} q(\mathbf{x}_t|\mathbf{x}_{t-1}, \mathbf{x}_0)}$$

$$= \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \log \frac{p(\mathbf{x}_T)p_{\theta}(\mathbf{x}_0|\mathbf{x}_1)\prod_{t=2}^{T} p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{q(\mathbf{x}_1|\mathbf{x}_0)\prod_{t=2}^{T} q(\mathbf{x}_t|\mathbf{x}_{t-1}, \mathbf{x}_0)}$$

Markov Property

$$q(\mathbf{x}_t|\mathbf{x}_{t-1}, \mathbf{x}_0) = q(\mathbf{x}_t|\mathbf{x}_{t-1})$$

Conditioning on $\mathbf{x}_0$ leaves the forward transitions unchanged.

Separate the decoder step $t = 1$.

ELBO for Gaussian Diffusion Model

Bayes' Rule ($t \geq 2$)

$$q(\mathbf{x}_t|\mathbf{x}_{t-1}, \mathbf{x}_0) = \frac{q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0)q(\mathbf{x}_t|\mathbf{x}_0)}{q(\mathbf{x}_{t-1}|\mathbf{x}_0)}$$

$$\mathcal{L}_{\phi,\theta}(\mathbf{x}) = \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \log \frac{p(\mathbf{x}_T)\prod_{t=1}^{T} p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{\prod_{t=1}^{T} q(\mathbf{x}_t|\mathbf{x}_{t-1}, \mathbf{x}_0)}$$

$$= \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \log \frac{p(\mathbf{x}_T)p_{\boldsymbol{\theta}}(\mathbf{x}_0|\mathbf{x}_1)\prod_{t=2}^{T} p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{q(\mathbf{x}_1|\mathbf{x}_0)\prod_{t=2}^{T} q(\mathbf{x}_t|\mathbf{x}_{t-1}, \mathbf{x}_0)}$$

$$= \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \log \frac{p(\mathbf{x}_T)p_{\boldsymbol{\theta}}(\mathbf{x}_0|\mathbf{x}_1)\prod_{t=2}^{T} p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{q(\mathbf{x}_1|\mathbf{x}_0)\prod_{t=2}^{T} \frac{q(\mathbf{x}_{t-1}|\mathbf{x}_t,\mathbf{x}_0)q(\mathbf{x}_t|\mathbf{x}_0)}{q(\mathbf{x}_{t-1}|\mathbf{x}_0)}}$$

Markov Property

$$q(\mathbf{x}_t|\mathbf{x}_{t-1}, \mathbf{x}_0) = q(\mathbf{x}_t|\mathbf{x}_{t-1})$$

Conditioning on $\mathbf{x}_0$ leaves the forward transitions unchanged.

Separate the decoder step $t = 1$.

Apply Bayes' rule for each $t \geq 2$.

ELBO for Gaussian Diffusion Model

Bayes' Rule ($t \geq 2$)

$$q(\mathbf{x}_t|\mathbf{x}_{t-1}, \mathbf{x}_0) = \frac{q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0)q(\mathbf{x}_t|\mathbf{x}_0)}{q(\mathbf{x}_{t-1}|\mathbf{x}_0)}$$

$$\mathcal{L}_{\phi,\theta}(\mathbf{x}) = \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \log \frac{p(\mathbf{x}_T)\prod_{t=1}^{T} p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{\prod_{t=1}^{T} q(\mathbf{x}_t|\mathbf{x}_{t-1}, \mathbf{x}_0)}$$

$$= \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \log \frac{p(\mathbf{x}_T)p_{\theta}(\mathbf{x}_0|\mathbf{x}_1)\prod_{t=2}^{T} p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{q(\mathbf{x}_1|\mathbf{x}_0)\prod_{t=2}^{T} q(\mathbf{x}_t|\mathbf{x}_{t-1}, \mathbf{x}_0)}$$

$$= \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \log \frac{p(\mathbf{x}_T)p_{\theta}(\mathbf{x}_0|\mathbf{x}_1)\prod_{t=2}^{T} p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{q(\mathbf{x}_1|\mathbf{x}_0)\prod_{t=2}^{T} \frac{q(\mathbf{x}_{t-1}|\mathbf{x}_t,\mathbf{x}_0)q(\mathbf{x}_t|\mathbf{x}_0)}{q(\mathbf{x}_{t-1}|\mathbf{x}_0)}}$$

$$= \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \log \frac{p(\mathbf{x}_T)p_{\theta}(\mathbf{x}_0|\mathbf{x}_1)\prod_{t=2}^{T} p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{q(\mathbf{x}_T|\mathbf{x}_0)\prod_{t=2}^{T} q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0)}$$

Markov Property

$$q(\mathbf{x}_t|\mathbf{x}_{t-1}, \mathbf{x}_0) = q(\mathbf{x}_t|\mathbf{x}_{t-1})$$

Conditioning on $\mathbf{x}_0$ leaves the forward transitions unchanged.

Separate the decoder step $t = 1$.

Apply Bayes' rule for each $t \geq 2$.

Intermediate marginals telescope; the $q(\mathbf{x}_1|\mathbf{x}_0)$ factor cancels too.

ELBO for Gaussian Diffusion Model

Derivation (continued)

$$\mathcal{L}_{\phi,\theta}(\mathbf{x}) = \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)}\Bigg[\log p_{\theta}(\mathbf{x}_0|\mathbf{x}_1) + \log \frac{p(\mathbf{x}_T)}{q(\mathbf{x}_T|\mathbf{x}_0)} + \sum_{t=2}^{T} \log\left(\frac{p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0)}\right)\Bigg]$$

Expand the logarithm: products become sums.

ELBO for Gaussian Diffusion Model

Derivation (continued)

$$\mathcal{L}_{\phi,\theta}(\mathbf{x}) = \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)}\Bigg[\log p_{\boldsymbol{\theta}}(\mathbf{x}_0|\mathbf{x}_1) + \log\frac{p(\mathbf{x}_T)}{q(\mathbf{x}_T|\mathbf{x}_0)} + \sum_{t=2}^{T}\log\left(\frac{p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{q(\mathbf{x}_{t-1}|\mathbf{x}_t,\mathbf{x}_0)}\right)\Bigg]$$

Expand the logarithm: products become sums.

$$= \mathbb{E}_{q(\mathbf{x}_1|\mathbf{x}_0)}\log p_{\boldsymbol{\theta}}(\mathbf{x}_0|\mathbf{x}_1) + \mathbb{E}_{q(\mathbf{x}_T|\mathbf{x}_0)}\log\frac{p(\mathbf{x}_T)}{q(\mathbf{x}_T|\mathbf{x}_0)} + \sum_{t=2}^{T}\mathbb{E}_{q(\mathbf{x}_{t-1},\mathbf{x}_t|\mathbf{x}_0)}\log\left(\frac{p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{q(\mathbf{x}_{t-1}|\mathbf{x}_t,\mathbf{x}_0)}\right)$$

Integrate out unused variables. Each term depends on one state or one adjacent pair.

ELBO for Gaussian Diffusion Model

Derivation (continued)

$$\mathcal{L}_{\phi,\theta}(\mathbf{x}) = \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)}\Bigg[\log p_{\boldsymbol{\theta}}(\mathbf{x}_0|\mathbf{x}_1) + \log\frac{p(\mathbf{x}_T)}{q(\mathbf{x}_T|\mathbf{x}_0)} + \sum_{t=2}^{T}\log\left(\frac{p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{q(\mathbf{x}_{t-1}|\mathbf{x}_t,\mathbf{x}_0)}\right)\Bigg]$$

Expand the logarithm: products become sums.

$$= \mathbb{E}_{q(\mathbf{x}_1|\mathbf{x}_0)}\log p_{\boldsymbol{\theta}}(\mathbf{x}_0|\mathbf{x}_1) + \mathbb{E}_{q(\mathbf{x}_T|\mathbf{x}_0)}\log\frac{p(\mathbf{x}_T)}{q(\mathbf{x}_T|\mathbf{x}_0)} + \sum_{t=2}^{T}\mathbb{E}_{q(\mathbf{x}_{t-1},\mathbf{x}_t|\mathbf{x}_0)}\log\left(\frac{p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{q(\mathbf{x}_{t-1}|\mathbf{x}_t,\mathbf{x}_0)}\right)$$

Integrate out unused variables. Each term depends on one state or one adjacent pair.


$$= \mathbb{E}_{q(\mathbf{x}_1|\mathbf{x}_0)}\log p_{\boldsymbol{\theta}}(\mathbf{x}_0|\mathbf{x}_1) - D_{\mathrm{KL}}\big(q(\mathbf{x}_T|\mathbf{x}_0)\|p(\mathbf{x}_T)\big) - \sum_{t=2}^{T}\underbrace{\mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)}D_{\mathrm{KL}}\big(q(\mathbf{x}_{t-1}|\mathbf{x}_t,\mathbf{x}_0)\|p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)\big)}_{\mathcal{L}_t}$$

Use the definition of KL. For each $\mathbf{x}_t$, average over $q(\mathbf{x}_{t-1}|\mathbf{x}_t,\mathbf{x}_0)$; then average over $q(\mathbf{x}_t|\mathbf{x}_0)$.

Ho J. Denoising Diffusion Probabilistic Models, 2020


ELBO for Gaussian Diffusion Model

$$\mathcal{L}_{\phi,\theta}(\mathbf{x}) = \mathbb{E}_{q(\mathbf{x}_1|\mathbf{x}_0)} \log p_{\boldsymbol{\theta}}(\mathbf{x}_0|\mathbf{x}_1) - D_{\mathrm{KL}}\big(q(\mathbf{x}_T|\mathbf{x}_0)\|p(\mathbf{x}_T)\big) - \sum_{t=2}^{T} \underbrace{\mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} D_{\mathrm{KL}}\big(q(\mathbf{x}_{t-1}|\mathbf{x}_t,\mathbf{x}_0)\|p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)\big)}_{\mathcal{L}_t}$$



ELBO for Gaussian Diffusion Model

$$\mathcal{L}_{\phi,\theta}(\mathbf{x}) = \mathbb{E}_{q(\mathbf{x}_1|\mathbf{x}_0)} \log p_{\boldsymbol{\theta}}(\mathbf{x}_0|\mathbf{x}_1) - D_{\mathrm{KL}}\big(q(\mathbf{x}_T|\mathbf{x}_0)\|p(\mathbf{x}_T)\big) - \sum_{t=2}^{T} \underbrace{\mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} D_{\mathrm{KL}}\big(q(\mathbf{x}_{t-1}|\mathbf{x}_t,\mathbf{x}_0)\|p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)\big)}_{\mathcal{L}_t}$$




- First term is the decoder distribution, with $\mathbf{x}_1 \sim q(\mathbf{x}_1|\mathbf{x}_0)$:

$$\log p_{\boldsymbol{\theta}}(\mathbf{x}_0|\mathbf{x}_1) = \log \mathcal{N}\big(\mathbf{x}_0|\boldsymbol{\mu}_{\boldsymbol{\theta},1}(\mathbf{x}_1), \boldsymbol{\sigma}^2_{\boldsymbol{\theta},1}(\mathbf{x}_1)\big).$$

ELBO for Gaussian Diffusion Model

$$\mathcal{L}_{\phi,\theta}(\mathbf{x}) = \mathbb{E}_{q(\mathbf{x}_1|\mathbf{x}_0)} \log p_{\boldsymbol{\theta}}(\mathbf{x}_0|\mathbf{x}_1) - D_{\mathrm{KL}}\big(q(\mathbf{x}_T|\mathbf{x}_0)\|p(\mathbf{x}_T)\big) - \sum_{t=2}^{T} \underbrace{\mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} D_{\mathrm{KL}}\big(q(\mathbf{x}_{t-1}|\mathbf{x}_t,\mathbf{x}_0)\|p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)\big)}_{\mathcal{L}_t}$$



- First term is the decoder distribution, with $\mathbf{x}_1 \sim q(\mathbf{x}_1|\mathbf{x}_0)$:


$$\log p_{\boldsymbol{\theta}}(\mathbf{x}_0|\mathbf{x}_1) = \log \mathcal{N}\big(\mathbf{x}_0|\boldsymbol{\mu}_{\boldsymbol{\theta},1}(\mathbf{x}_1), \boldsymbol{\sigma}^2_{\boldsymbol{\theta},1}(\mathbf{x}_1)\big).$$

- Second term is constant:

$$p(\mathbf{x}_T) = \mathcal{N}(0, \mathbf{I}), \qquad q(\mathbf{x}_T|\mathbf{x}_0) = \mathcal{N}(\sqrt{\bar{\alpha}_T} \cdot \mathbf{x}_0, (1 - \bar{\alpha}_T) \cdot \mathbf{I}).$$

ELBO for Gaussian Diffusion Model

$$\mathcal{L}_{\phi,\theta}(\mathbf{x}) = \mathbb{E}_{q(\mathbf{x}_1|\mathbf{x}_0)} \log p_{\boldsymbol{\theta}}(\mathbf{x}_0|\mathbf{x}_1) - D_{\mathrm{KL}}\big(q(\mathbf{x}_T|\mathbf{x}_0)\|p(\mathbf{x}_T)\big) - \sum_{t=2}^{T} \underbrace{\mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} D_{\mathrm{KL}}\big(q(\mathbf{x}_{t-1}|\mathbf{x}_t,\mathbf{x}_0)\|p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)\big)}_{\mathcal{L}_t}$$



- First term is the decoder distribution, with $\mathbf{x}_1 \sim q(\mathbf{x}_1|\mathbf{x}_0)$:


$$\log p_{\boldsymbol{\theta}}(\mathbf{x}_0|\mathbf{x}_1) = \log \mathcal{N}\big(\mathbf{x}_0|\boldsymbol{\mu}_{\boldsymbol{\theta},1}(\mathbf{x}_1), \boldsymbol{\sigma}^2_{\boldsymbol{\theta},1}(\mathbf{x}_1)\big).$$

- Second term is constant:

$$p(\mathbf{x}_T) = \mathcal{N}(0, \mathbf{I}), \qquad q(\mathbf{x}_T|\mathbf{x}_0) = \mathcal{N}(\sqrt{\bar{\alpha}_T} \cdot \mathbf{x}_0, (1 - \bar{\alpha}_T) \cdot \mathbf{I}).$$

- Third term is the main contributor to the ELBO.

ELBO for Gaussian Diffusion Model



$$\mathcal{L}_t = \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} D_{\mathrm{KL}}\left(q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) \| p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)\right)$$

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) = \mathcal{N}(\mathbf{x}_{t-1}|\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0), \tilde{\beta}_t \mathbf{I}),$$

$$p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t) = \mathcal{N}\left(\mathbf{x}_{t-1}|\boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t), \boldsymbol{\sigma}^2_{\boldsymbol{\theta},t}(\mathbf{x}_t)\right)$$

ELBO for Gaussian Diffusion Model

$$\mathcal{L}_t = \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} D_{\mathrm{KL}}\left(q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) \| p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)\right)$$

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) = \mathcal{N}(\mathbf{x}_{t-1}|\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0), \tilde{\beta}_t \mathbf{I}),$$

$$p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t) = \mathcal{N}\left(\mathbf{x}_{t-1}|\boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t), \boldsymbol{\sigma}^2_{\boldsymbol{\theta},t}(\mathbf{x}_t)\right)$$

ELBO for Gaussian Diffusion Model

$$\mathcal{L}_t = \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} D_{\mathrm{KL}}\big(q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) \| p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)\big)$$

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) = \mathcal{N}(\mathbf{x}_{t-1}|\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0), \tilde{\beta}_t \mathbf{I}),$$

$$p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t) = \mathcal{N}\big(\mathbf{x}_{t-1}|\boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t), \boldsymbol{\sigma}^2_{\boldsymbol{\theta},t}(\mathbf{x}_t)\big)$$

Let's assume that

$$\boldsymbol{\sigma}^2_{\boldsymbol{\theta},t}(\mathbf{x}_t) = \tilde{\beta}_t \mathbf{I}$$

$$\Rightarrow \quad p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t) = \mathcal{N}\big(\mathbf{x}_{t-1}|\boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t), \tilde{\beta}_t \mathbf{I}\big).$$

Ho J. Denoising Diffusion Probabilistic Models, 2020

ELBO for Gaussian Diffusion Model

$$\mathcal{L}_t = \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} D_{\mathrm{KL}}\big(q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) \| p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)\big)$$

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) = \mathcal{N}(\mathbf{x}_{t-1}|\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0), \tilde{\beta}_t \mathbf{I}),$$

$$p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t) = \mathcal{N}\big(\mathbf{x}_{t-1}|\boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t), \boldsymbol{\sigma}^2_{\boldsymbol{\theta},t}(\mathbf{x}_t)\big)$$

Let's assume that

$$\boldsymbol{\sigma}^2_{\boldsymbol{\theta},t}(\mathbf{x}_t) = \tilde{\beta}_t \mathbf{I}$$

$$\Rightarrow \quad p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t) = \mathcal{N}\big(\mathbf{x}_{t-1}|\boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t), \tilde{\beta}_t \mathbf{I}\big).$$

Theoretically, the optimal $\boldsymbol{\sigma}^2_{\boldsymbol{\theta},t}(\mathbf{x}_t)$ lies in $[\tilde{\beta}_t, \beta_t]$:

- β_t is optimal for $\mathbf{x}_0 \sim \mathcal{N}(0, \mathbf{I})$;
- $\tilde{\beta}_t$ is optimal for $\mathbf{x}_0 \sim \delta(\mathbf{x}_0 - \mathbf{x}^*)$.

ELBO for Gaussian Diffusion Model

$$\mathcal{L}_t = \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} D_{\mathrm{KL}}\big(q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) \| p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)\big)$$

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) = \mathcal{N}(\mathbf{x}_{t-1}|\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0), \tilde{\beta}_t \mathbf{I}),$$

$$p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t) = \mathcal{N}\big(\mathbf{x}_{t-1}|\boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t), \boldsymbol{\sigma}^2_{\boldsymbol{\theta},t}(\mathbf{x}_t)\big)$$

Let's assume that

$$\boldsymbol{\sigma}^2_{\boldsymbol{\theta},t}(\mathbf{x}_t) = \tilde{\beta}_t \mathbf{I}$$

$$\Rightarrow \quad p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t) = \mathcal{N}\big(\mathbf{x}_{t-1}|\boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t), \tilde{\beta}_t \mathbf{I}\big).$$

Theoretically, the optimal $\boldsymbol{\sigma}^2_{\boldsymbol{\theta},t}(\mathbf{x}_t)$ lies in $[\tilde{\beta}_t, \beta_t]$:

- β_t is optimal for $\mathbf{x}_0 \sim \mathcal{N}(0, \mathbf{I})$;
- $\tilde{\beta}_t$ is optimal for $\mathbf{x}_0 \sim \delta(\mathbf{x}_0 - \mathbf{x}^*)$.

$$\mathcal{L}_t = \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} D_{\mathrm{KL}}\big(\mathcal{N}\big(\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0), \tilde{\beta}_t \mathbf{I}\big) \| \mathcal{N}\big(\boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t), \tilde{\beta}_t \mathbf{I}\big)\big)$$

Ho J. Denoising Diffusion Probabilistic Models, 2020

ELBO for Gaussian Diffusion Model

$$\mathcal{L}_t = \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} D_{\mathrm{KL}}\big(q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) \| p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)\big)$$

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) = \mathcal{N}(\mathbf{x}_{t-1}|\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0), \tilde{\beta}_t \mathbf{I}),$$

$$p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t) = \mathcal{N}\big(\mathbf{x}_{t-1}|\boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t), \boldsymbol{\sigma}^2_{\boldsymbol{\theta},t}(\mathbf{x}_t)\big)$$

Let's assume that

$$\boldsymbol{\sigma}^2_{\boldsymbol{\theta},t}(\mathbf{x}_t) = \tilde{\beta}_t \mathbf{I}$$

$$\Rightarrow \quad p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t) = \mathcal{N}\big(\mathbf{x}_{t-1}|\boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t), \tilde{\beta}_t \mathbf{I}\big).$$

Theoretically, the optimal $\boldsymbol{\sigma}^2_{\boldsymbol{\theta},t}(\mathbf{x}_t)$ lies in $[\tilde{\beta}_t, \beta_t]$:

- β_t is optimal for $\mathbf{x}_0 \sim \mathcal{N}(0, \mathbf{I})$;
- $\tilde{\beta}_t$ is optimal for $\mathbf{x}_0 \sim \delta(\mathbf{x}_0 - \mathbf{x}^*)$.

$$\mathcal{L}_t = \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} D_{\mathrm{KL}}\big(\mathcal{N}\big(\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0), \tilde{\beta}_t \mathbf{I}\big) \| \mathcal{N}\big(\boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t), \tilde{\beta}_t \mathbf{I}\big)\big)$$

$$= \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} \left[\frac{1}{2\tilde{\beta}_t} \left\| \tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) - \boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t) \right\|^2\right]$$

ELBO for Gaussian Diffusion Model

Training

1. Sample $\mathbf{x}_0 \sim p_{\text{data}}(\mathbf{x})$, $\boldsymbol{\epsilon} \sim \mathcal{N}(0, \mathbf{I})$.
2. Compute noisy image $\mathbf{x}_t = \sqrt{\bar{\alpha}_t} \cdot \mathbf{x}_0 + \sqrt{1 - \bar{\alpha}_t} \cdot \boldsymbol{\epsilon}$.
3. Compute the ELBO:

$$\mathcal{L}_{\boldsymbol{\phi},\boldsymbol{\theta}}(\mathbf{x}) = \mathbb{E}_{q(\mathbf{x}_1|\mathbf{x}_0)} \log p_{\boldsymbol{\theta}}(\mathbf{x}_0|\mathbf{x}_1) - D_{\text{KL}}\left(q(\mathbf{x}_T|\mathbf{x}_0) \| p(\mathbf{x}_T)\right) - \sum_{t=2}^{T} \underbrace{\mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} \left[\frac{1}{2\tilde{\beta}_t} \left\| \tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) - \boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t) \right\|^2 \right]}_{\mathcal{L}_t}.$$

ELBO for Gaussian Diffusion Model

Training

1. Sample $\mathbf{x}_0 \sim p_{\text{data}}(\mathbf{x})$, $\boldsymbol{\epsilon} \sim \mathcal{N}(0, \mathbf{I})$.
2. Compute noisy image $\mathbf{x}_t = \sqrt{\bar{\alpha}_t} \cdot \mathbf{x}_0 + \sqrt{1 - \bar{\alpha}_t} \cdot \boldsymbol{\epsilon}$.
3. Compute the ELBO:

$$\mathcal{L}_{\boldsymbol{\phi},\boldsymbol{\theta}}(\mathbf{x}) = \mathbb{E}_{q(\mathbf{x}_1|\mathbf{x}_0)} \log p_{\boldsymbol{\theta}}(\mathbf{x}_0|\mathbf{x}_1) - D_{\text{KL}}\left(q(\mathbf{x}_T|\mathbf{x}_0) \| p(\mathbf{x}_T)\right) - \sum_{t=2}^{T} \underbrace{\mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} \left[\frac{1}{2\tilde{\beta}_t} \left\| \tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) - \boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t) \right\|^2 \right]}_{\mathcal{L}_t}.$$

Sampling

1. Sample $\mathbf{x}_T \sim \mathcal{N}(0, \mathbf{I})$.
2. Denoise $\mathbf{x}_{t-1} = \boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t) + \sqrt{\tilde{\beta}_t} \cdot \boldsymbol{\epsilon}$, $\boldsymbol{\epsilon} \sim \mathcal{N}(0, \mathbf{I})$.

Outline

01 Reverse Gaussian Diffusion Process

02 Gaussian Diffusion Model as VAE

03 Diffusion ELBO Derivation

04 Gaussian Diffusion Reparametrization

05 Denoising Diffusion Probabilistic Model (DDPM)

Reparametrization of DDPM

$$\mathcal{L}_t = \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} \left[\frac{1}{2\tilde{\beta}_t} \left\| \tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) - \boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t) \right\|^2 \right]$$

$$\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) = \frac{\sqrt{\alpha_t}(1 - \bar{\alpha}_{t-1})}{1 - \bar{\alpha}_t} \cdot \mathbf{x}_t + \frac{\sqrt{\bar{\alpha}_{t-1}}(1 - \alpha_t)}{1 - \bar{\alpha}_t} \cdot \mathbf{x}_0$$

Reparametrization of DDPM

$$\mathcal{L}_t = \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} \left[\frac{1}{2\tilde{\beta}_t} \left\| \tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) - \boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t) \right\|^2 \right]$$

$$\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) = \frac{\sqrt{\alpha_t}(1 - \bar{\alpha}_{t-1})}{1 - \bar{\alpha}_t} \cdot \mathbf{x}_t + \frac{\sqrt{\bar{\alpha}_{t-1}}(1 - \alpha_t)}{1 - \bar{\alpha}_t} \cdot \mathbf{x}_0$$

$$\mathbf{x}_t = \sqrt{\bar{\alpha}_t} \cdot \mathbf{x}_0 + \sqrt{1 - \bar{\alpha}_t} \cdot \boldsymbol{\epsilon}$$

Reparametrization of DDPM

$$\mathcal{L}_t = \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)}\left[\frac{1}{2\tilde{\beta}_t}\left\|\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) - \boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t)\right\|^2\right]$$

$$\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) = \frac{\sqrt{\alpha_t}(1-\bar{\alpha}_{t-1})}{1-\bar{\alpha}_t}\cdot\mathbf{x}_t + \frac{\sqrt{\bar{\alpha}_{t-1}}(1-\alpha_t)}{1-\bar{\alpha}_t}\cdot\mathbf{x}_0$$

$$\mathbf{x}_t = \sqrt{\bar{\alpha}_t}\cdot\mathbf{x}_0 + \sqrt{1-\bar{\alpha}_t}\cdot\boldsymbol{\epsilon} \quad\Rightarrow\quad \mathbf{x}_0 = \frac{\mathbf{x}_t - \sqrt{1-\bar{\alpha}_t}\cdot\boldsymbol{\epsilon}}{\sqrt{\bar{\alpha}_t}}$$

Reparametrization of DDPM

$$\mathcal{L}_t = \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)}\left[\frac{1}{2\tilde{\beta}_t}\left\|\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) - \boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t)\right\|^2\right]$$

$$\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) = \frac{\sqrt{\alpha_t}(1-\bar{\alpha}_{t-1})}{1-\bar{\alpha}_t}\cdot\mathbf{x}_t + \frac{\sqrt{\bar{\alpha}_{t-1}}(1-\alpha_t)}{1-\bar{\alpha}_t}\cdot\mathbf{x}_0$$

$$\mathbf{x}_t = \sqrt{\bar{\alpha}_t}\cdot\mathbf{x}_0 + \sqrt{1-\bar{\alpha}_t}\cdot\boldsymbol{\epsilon} \quad\Rightarrow\quad \mathbf{x}_0 = \frac{\mathbf{x}_t - \sqrt{1-\bar{\alpha}_t}\cdot\boldsymbol{\epsilon}}{\sqrt{\bar{\alpha}_t}}$$

- There is a linear relationship between $\boldsymbol{\epsilon}$, $\mathbf{x}_t$, and $\mathbf{x}_0$.
- Let's try to rewrite this mean using only $\mathbf{x}_t$ and $\boldsymbol{\epsilon}$.

Reparametrization of DDPM

$$\mathcal{L}_t = \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)}\left[\frac{1}{2\tilde{\beta}_t}\left\|\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) - \boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t)\right\|^2\right]$$

$$\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) = \frac{\sqrt{\alpha_t}(1-\bar{\alpha}_{t-1})}{1-\bar{\alpha}_t}\cdot\mathbf{x}_t + \frac{\sqrt{\bar{\alpha}_{t-1}}(1-\alpha_t)}{1-\bar{\alpha}_t}\cdot\mathbf{x}_0$$

$$\mathbf{x}_t = \sqrt{\bar{\alpha}_t}\cdot\mathbf{x}_0 + \sqrt{1-\bar{\alpha}_t}\cdot\boldsymbol{\epsilon} \quad\Rightarrow\quad \mathbf{x}_0 = \frac{\mathbf{x}_t - \sqrt{1-\bar{\alpha}_t}\cdot\boldsymbol{\epsilon}}{\sqrt{\bar{\alpha}_t}}$$

- There is a linear relationship between $\boldsymbol{\epsilon}$, $\mathbf{x}_t$, and $\mathbf{x}_0$.
- Let's try to rewrite this mean using only $\mathbf{x}_t$ and $\boldsymbol{\epsilon}$.

$$\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \boldsymbol{\epsilon}) = \frac{\sqrt{\alpha_t}(1-\bar{\alpha}_{t-1})}{1-\bar{\alpha}_t}\cdot\mathbf{x}_t + \frac{\sqrt{\bar{\alpha}_{t-1}}(1-\alpha_t)}{1-\bar{\alpha}_t}\cdot\left(\frac{\mathbf{x}_t - \sqrt{1-\bar{\alpha}_t}\cdot\boldsymbol{\epsilon}}{\sqrt{\bar{\alpha}_t}}\right)$$

Reparametrization of DDPM

$$\mathcal{L}_t = \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)}\left[\frac{1}{2\tilde{\beta}_t}\left\|\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) - \boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t)\right\|^2\right]$$

$$\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) = \frac{\sqrt{\alpha_t}(1-\bar{\alpha}_{t-1})}{1-\bar{\alpha}_t}\cdot\mathbf{x}_t + \frac{\sqrt{\bar{\alpha}_{t-1}}(1-\alpha_t)}{1-\bar{\alpha}_t}\cdot\mathbf{x}_0$$

$$\mathbf{x}_t = \sqrt{\bar{\alpha}_t}\cdot\mathbf{x}_0 + \sqrt{1-\bar{\alpha}_t}\cdot\boldsymbol{\epsilon} \quad\Rightarrow\quad \mathbf{x}_0 = \frac{\mathbf{x}_t - \sqrt{1-\bar{\alpha}_t}\cdot\boldsymbol{\epsilon}}{\sqrt{\bar{\alpha}_t}}$$

- There is a linear relationship between $\boldsymbol{\epsilon}$, $\mathbf{x}_t$, and $\mathbf{x}_0$.
- Let's try to rewrite this mean using only $\mathbf{x}_t$ and $\boldsymbol{\epsilon}$.

$$\begin{aligned}\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \boldsymbol{\epsilon}) &= \frac{\sqrt{\alpha_t}(1-\bar{\alpha}_{t-1})}{1-\bar{\alpha}_t}\cdot\mathbf{x}_t + \frac{\sqrt{\bar{\alpha}_{t-1}}(1-\alpha_t)}{1-\bar{\alpha}_t}\cdot\left(\frac{\mathbf{x}_t - \sqrt{1-\bar{\alpha}_t}\cdot\boldsymbol{\epsilon}}{\sqrt{\bar{\alpha}_t}}\right) \\ &= \frac{1}{\sqrt{\alpha_t}}\cdot\mathbf{x}_t - \frac{1-\alpha_t}{\sqrt{\alpha_t(1-\bar{\alpha}_t)}}\cdot\boldsymbol{\epsilon}\end{aligned}$$

Ho J. Denoising Diffusion Probabilistic Models, 2020

Reparametrization of DDPM

$$\mathcal{L}_t = \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} \left[\frac{1}{2\tilde{\beta}_t} \left\| \tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) - \boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t) \right\|^2 \right]$$

Reparametrization

$$\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) = \frac{1}{\sqrt{\alpha_t}} \cdot \mathbf{x}_t - \frac{1 - \alpha_t}{\sqrt{\alpha_t(1 - \bar{\alpha}_t)}} \cdot \boldsymbol{\epsilon}$$

Reparametrization of DDPM

$$\mathcal{L}_t = \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} \left[\frac{1}{2\tilde{\beta}_t} \left\| \tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) - \boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t) \right\|^2 \right]$$

Reparametrization

$$\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) = \frac{1}{\sqrt{\alpha_t}} \cdot \mathbf{x}_t - \frac{1 - \alpha_t}{\sqrt{\alpha_t(1 - \bar{\alpha}_t)}} \cdot \boldsymbol{\epsilon}$$

$$\boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t) = \frac{1}{\sqrt{\alpha_t}} \cdot \mathbf{x}_t - \frac{1 - \alpha_t}{\sqrt{\alpha_t(1 - \bar{\alpha}_t)}} \cdot \boldsymbol{\epsilon}_{\boldsymbol{\theta},t}(\mathbf{x}_t)$$

Reparametrization of DDPM

$$\mathcal{L}_t = \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} \left[\frac{1}{2\tilde{\beta}_t} \left\| \tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) - \boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t) \right\|^2 \right]$$

Reparametrization

$$\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) = \frac{1}{\sqrt{\alpha_t}} \cdot \mathbf{x}_t - \frac{1 - \alpha_t}{\sqrt{\alpha_t(1 - \bar{\alpha}_t)}} \cdot \boldsymbol{\epsilon}$$

$$\boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t) = \frac{1}{\sqrt{\alpha_t}} \cdot \mathbf{x}_t - \frac{1 - \alpha_t}{\sqrt{\alpha_t(1 - \bar{\alpha}_t)}} \cdot \boldsymbol{\epsilon}_{\boldsymbol{\theta},t}(\mathbf{x}_t)$$

$$\mathcal{L}_t = \mathbb{E}_{\boldsymbol{\epsilon} \sim \mathcal{N}(0,\mathbf{I})} \left[\frac{(1 - \alpha_t)^2}{2\tilde{\beta}_t \alpha_t (1 - \bar{\alpha}_t)} \left\| \boldsymbol{\epsilon} - \boldsymbol{\epsilon}_{\boldsymbol{\theta},t}(\mathbf{x}_t) \right\|^2 \right]$$

Reparametrization of DDPM

$$\mathcal{L}_t = \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} \left[\frac{1}{2\tilde{\beta}_t} \left\| \tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) - \boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t) \right\|^2 \right]$$

Reparametrization

$$\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) = \frac{1}{\sqrt{\alpha_t}} \cdot \mathbf{x}_t - \frac{1 - \alpha_t}{\sqrt{\alpha_t(1 - \bar{\alpha}_t)}} \cdot \boldsymbol{\epsilon}$$

$$\boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t) = \frac{1}{\sqrt{\alpha_t}} \cdot \mathbf{x}_t - \frac{1 - \alpha_t}{\sqrt{\alpha_t(1 - \bar{\alpha}_t)}} \cdot \boldsymbol{\epsilon}_{\boldsymbol{\theta},t}(\mathbf{x}_t)$$

$$\begin{aligned}
\mathcal{L}_t &= \mathbb{E}_{\boldsymbol{\epsilon} \sim \mathcal{N}(0,\mathbf{I})} \left[\frac{(1 - \alpha_t)^2}{2\tilde{\beta}_t \alpha_t (1 - \bar{\alpha}_t)} \left\| \boldsymbol{\epsilon} - \boldsymbol{\epsilon}_{\boldsymbol{\theta},t}(\mathbf{x}_t) \right\|^2 \right] \\
&= \mathbb{E}_{\boldsymbol{\epsilon} \sim \mathcal{N}(0,\mathbf{I})} \left[\frac{(1 - \alpha_t)^2}{2\tilde{\beta}_t \alpha_t (1 - \bar{\alpha}_t)} \left\| \boldsymbol{\epsilon} - \boldsymbol{\epsilon}_{\boldsymbol{\theta},t}\left(\sqrt{\bar{\alpha}_t}\mathbf{x}_0 + \sqrt{1 - \bar{\alpha}_t}\boldsymbol{\epsilon} \right) \right\|^2 \right]
\end{aligned}$$

Reparametrization of DDPM

$$\mathcal{L}_t = \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)}\left[\frac{1}{2\tilde{\beta}_t}\left\|\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) - \boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t)\right\|^2\right]$$

Reparametrization

$$\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) = \frac{1}{\sqrt{\alpha_t}} \cdot \mathbf{x}_t - \frac{1-\alpha_t}{\sqrt{\alpha_t(1-\bar{\alpha}_t)}} \cdot \boldsymbol{\epsilon}$$

$$\boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t) = \frac{1}{\sqrt{\alpha_t}} \cdot \mathbf{x}_t - \frac{1-\alpha_t}{\sqrt{\alpha_t(1-\bar{\alpha}_t)}} \cdot \boldsymbol{\epsilon}_{\boldsymbol{\theta},t}(\mathbf{x}_t)$$


$$\begin{aligned}\mathcal{L}_t &= \mathbb{E}_{\boldsymbol{\epsilon}\sim\mathcal{N}(0,\mathbf{I})}\left[\frac{(1-\alpha_t)^2}{2\tilde{\beta}_t\alpha_t(1-\bar{\alpha}_t)}\left\|\boldsymbol{\epsilon} - \boldsymbol{\epsilon}_{\boldsymbol{\theta},t}(\mathbf{x}_t)\right\|^2\right] \\ &= \mathbb{E}_{\boldsymbol{\epsilon}\sim\mathcal{N}(0,\mathbf{I})}\left[\frac{(1-\alpha_t)^2}{2\tilde{\beta}_t\alpha_t(1-\bar{\alpha}_t)}\left\|\boldsymbol{\epsilon} - \boldsymbol{\epsilon}_{\boldsymbol{\theta},t}\left(\sqrt{\bar{\alpha}_t}\mathbf{x}_0 + \sqrt{1-\bar{\alpha}_t}\boldsymbol{\epsilon}\right)\right\|^2\right]\end{aligned}$$

At every step of the reverse process, we attempt to predict the noise $\boldsymbol{\epsilon}$ that was used in the forward diffusion process!

One Prediction, Three Parameterizations

One coordinate. Keep the input and timestep fixed: $x_t = 1, \quad \alpha_t = 0.6, \quad \bar{\alpha}_t = 0.39$

Choose one network output; compute the other two. Example: predict noise.



$$x_t = \sqrt{\bar{\alpha}_t}\hat{x}_0 + \sqrt{1-\bar{\alpha}_t}\hat{\epsilon}$$

$$\mu_{\theta,t} = \frac{x_t}{\sqrt{\alpha_t}} - \frac{1-\alpha_t}{\sqrt{\alpha_t(1-\bar{\alpha}_t)}}\hat{\epsilon}$$

Switch the network output: the values stay the same; only the parameterization changes.

All three are predicted quantities, linked by known conversions.

Diffusion Parameterizations

$$\mathbf{x}_t = \sqrt{\bar{\alpha}_t}\,\mathbf{x}_0 + \sqrt{1-\bar{\alpha}_t}\,\boldsymbol{\epsilon}.$$

Diffusion Parameterizations

$$\mathbf{x}_t = \sqrt{\bar{\alpha}_t}\,\mathbf{x}_0 + \sqrt{1-\bar{\alpha}_t}\,\boldsymbol{\epsilon}.$$

Network output at $\mathbf{x}_t$	Regression target	Clean-data estimate
Noise $\boldsymbol{\epsilon}_{\boldsymbol{\theta},t}$	$\boldsymbol{\epsilon}$	$\dfrac{\mathbf{x}_t - \sqrt{1-\bar{\alpha}_t}\,\boldsymbol{\epsilon}_{\boldsymbol{\theta},t}}{\sqrt{\bar{\alpha}_t}}$
Clean data $\hat{\mathbf{x}}_{\boldsymbol{\theta},t}$	$\mathbf{x}_0$	$\hat{\mathbf{x}}_{\boldsymbol{\theta},t}$
Score $\mathbf{s}_{\boldsymbol{\theta},t}$	$-\dfrac{\boldsymbol{\epsilon}}{\sqrt{1-\bar{\alpha}_t}}$	$\dfrac{\mathbf{x}_t + (1-\bar{\alpha}_t)\,\mathbf{s}_{\boldsymbol{\theta},t}}{\sqrt{\bar{\alpha}_t}}$
v-prediction $\mathbf{v}_{\boldsymbol{\theta},t}$	$\sqrt{\bar{\alpha}_t}\,\boldsymbol{\epsilon} - \sqrt{1-\bar{\alpha}_t}\,\mathbf{x}_0$	$\sqrt{\bar{\alpha}_t}\,\mathbf{x}_t - \sqrt{1-\bar{\alpha}_t}\,\mathbf{v}_{\boldsymbol{\theta},t}$

Salimans T., Ho J. Progressive Distillation for Fast Sampling of Diffusion Models, 2022

Diffusion Parameterizations

$$\mathbf{x}_t = \sqrt{\bar{\alpha}_t}\,\mathbf{x}_0 + \sqrt{1-\bar{\alpha}_t}\,\boldsymbol{\epsilon}.$$

Network output at $\mathbf{x}_t$	Regression target	Clean-data estimate
Noise $\boldsymbol{\epsilon}_{\boldsymbol{\theta},t}$	$\boldsymbol{\epsilon}$	$\dfrac{\mathbf{x}_t - \sqrt{1-\bar{\alpha}_t}\,\boldsymbol{\epsilon}_{\boldsymbol{\theta},t}}{\sqrt{\bar{\alpha}_t}}$
Clean data $\hat{\mathbf{x}}_{\boldsymbol{\theta},t}$	$\mathbf{x}_0$	$\hat{\mathbf{x}}_{\boldsymbol{\theta},t}$
Score $\mathbf{s}_{\boldsymbol{\theta},t}$	$-\dfrac{\boldsymbol{\epsilon}}{\sqrt{1-\bar{\alpha}_t}}$	$\dfrac{\mathbf{x}_t + (1-\bar{\alpha}_t)\,\mathbf{s}_{\boldsymbol{\theta},t}}{\sqrt{\bar{\alpha}_t}}$
v-prediction $\mathbf{v}_{\boldsymbol{\theta},t}$	$\sqrt{\bar{\alpha}_t}\,\boldsymbol{\epsilon} - \sqrt{1-\bar{\alpha}_t}\,\mathbf{x}_0$	$\sqrt{\bar{\alpha}_t}\,\mathbf{x}_t - \sqrt{1-\bar{\alpha}_t}\,\mathbf{v}_{\boldsymbol{\theta},t}$

Convertible predictions; different losses. Plain MSE assigns different weights to different noise levels.

Salimans T., Ho J. Progressive Distillation for Fast Sampling of Diffusion Models, 2022

Reparametrization of DDPM

$$\mathcal{L}_{\phi,\theta}(\mathbf{x}) = \mathbb{E}_{q(\mathbf{x}_1|\mathbf{x}_0)} \log p_{\theta}(\mathbf{x}_0|\mathbf{x}_1) - D_{\mathrm{KL}}\left(q(\mathbf{x}_T|\mathbf{x}_0)||p(\mathbf{x}_T)\right) - \sum_{t=2}^{T} \underbrace{\mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} D_{\mathrm{KL}}\left(q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0)||p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t)\right)}_{\mathcal{L}_t}$$

Reparametrization of DDPM

$$\mathcal{L}_{\phi,\theta}(\mathbf{x}) = \mathbb{E}_{q(\mathbf{x}_1|\mathbf{x}_0)} \log p_{\boldsymbol{\theta}}(\mathbf{x}_0|\mathbf{x}_1) - D_{\mathrm{KL}}\big(q(\mathbf{x}_T|\mathbf{x}_0)\|p(\mathbf{x}_T)\big) - \sum_{t=2}^{T} \underbrace{\mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} D_{\mathrm{KL}}\big(q(\mathbf{x}_{t-1}|\mathbf{x}_t,\mathbf{x}_0)\|p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)\big)}_{\mathcal{L}_t}$$

$$\mathcal{L}_t = \mathbb{E}_{\boldsymbol{\epsilon}\sim\mathcal{N}(0,\mathbf{I})}\left[\frac{(1-\alpha_t)^2}{2\tilde{\beta}_t\alpha_t(1-\bar{\alpha}_t)}\left\|\boldsymbol{\epsilon} - \boldsymbol{\epsilon}_{\boldsymbol{\theta},t}\big(\sqrt{\bar{\alpha}_t}\mathbf{x}_0 + \sqrt{1-\bar{\alpha}_t}\boldsymbol{\epsilon}\big)\right\|^2\right]$$

Reparametrization of DDPM

$$\mathcal{L}_{\phi,\boldsymbol{\theta}}(\mathbf{x}) = \mathbb{E}_{q(\mathbf{x}_1|\mathbf{x}_0)} \log p_{\boldsymbol{\theta}}(\mathbf{x}_0|\mathbf{x}_1) - D_{\mathrm{KL}}\big(q(\mathbf{x}_T|\mathbf{x}_0)\|p(\mathbf{x}_T)\big) - \sum_{t=2}^{T} \underbrace{\mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} D_{\mathrm{KL}}\big(q(\mathbf{x}_{t-1}|\mathbf{x}_t,\mathbf{x}_0)\|p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)\big)}_{\mathcal{L}_t}$$

$$\mathcal{L}_t = \mathbb{E}_{\boldsymbol{\epsilon}\sim\mathcal{N}(0,\mathbf{I})}\left[\frac{(1-\alpha_t)^2}{2\tilde{\beta}_t\alpha_t(1-\bar{\alpha}_t)}\left\|\boldsymbol{\epsilon} - \boldsymbol{\epsilon}_{\boldsymbol{\theta},t}\big(\sqrt{\bar{\alpha}_t}\mathbf{x}_0 + \sqrt{1-\bar{\alpha}_t}\boldsymbol{\epsilon}\big)\right\|^2\right]$$

Let's drop the scaling coefficient; the reconstruction term ($t=1$) has the same form.

Simplified Objective

$$\mathcal{L}_{\text{simple}} = \mathbb{E}_{t\sim U\{1,T\}}\mathbb{E}_{\boldsymbol{\epsilon}\sim\mathcal{N}(0,\mathbf{I})}\left\|\boldsymbol{\epsilon} - \boldsymbol{\epsilon}_{\boldsymbol{\theta},t}\big(\sqrt{\bar{\alpha}_t}\cdot\mathbf{x}_0 + \sqrt{1-\bar{\alpha}_t}\cdot\boldsymbol{\epsilon}\big)\right\|^2$$

Outline

01 Reverse Gaussian Diffusion Process


02 Gaussian Diffusion Model as VAE

03 Diffusion ELBO Derivation

04 Gaussian Diffusion Reparametrization

05 Denoising Diffusion Probabilistic Model (DDPM)

Generative Models Taxonomy



Denoising Diffusion Probabilistic Model (DDPM)

DDPM is a VAE Model

- The encoder is a fixed Gaussian Markov chain $q(\mathbf{x}_1, \dots, \mathbf{x}_T | \mathbf{x}_0)$.
- The latent variable is hierarchical (at each step, its dimension equals the input's).
- The decoder is a simple Gaussian model $p_{\boldsymbol{\theta}}(\mathbf{x}_0 | \mathbf{x}_1)$.
- The prior distribution is given by a parametric Gaussian Markov chain $p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1} | \mathbf{x}_t)$.

Denoising Diffusion Probabilistic Model (DDPM)

DDPM is a VAE Model

- The encoder is a fixed Gaussian Markov chain $q(\mathbf{x}_1, \ldots, \mathbf{x}_T | \mathbf{x}_0)$.
- The latent variable is hierarchical (at each step, its dimension equals the input's).
- The decoder is a simple Gaussian model $p_{\boldsymbol{\theta}}(\mathbf{x}_0 | \mathbf{x}_1)$.
- The prior distribution is given by a parametric Gaussian Markov chain $p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1} | \mathbf{x}_t)$.

Forward Process

1. $\mathbf{x}_0 = \mathbf{x} \sim p_{\text{data}}(\mathbf{x})$;
2. $\mathbf{x}_t = \sqrt{1 - \beta_t} \cdot \mathbf{x}_{t-1} + \sqrt{\beta_t} \cdot \boldsymbol{\epsilon}_t$;
3. $\mathbf{x}_T \sim p_\infty(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$.

Denoising Diffusion Probabilistic Model (DDPM)

DDPM is a VAE Model

- The encoder is a fixed Gaussian Markov chain $q(\mathbf{x}_1, \dots, \mathbf{x}_T | \mathbf{x}_0)$.
- The latent variable is hierarchical (at each step, its dimension equals the input's).
- The decoder is a simple Gaussian model $p_{\boldsymbol{\theta}}(\mathbf{x}_0 | \mathbf{x}_1)$.
- The prior distribution is given by a parametric Gaussian Markov chain $p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1} | \mathbf{x}_t)$.

Forward Process

1. $\mathbf{x}_0 = \mathbf{x} \sim p_{\text{data}}(\mathbf{x})$;
2. $\mathbf{x}_t = \sqrt{1 - \beta_t} \cdot \mathbf{x}_{t-1} + \sqrt{\beta_t} \cdot \boldsymbol{\epsilon}_t$;
3. $\mathbf{x}_T \sim p_\infty(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$.

Reverse Process

1. $\mathbf{x}_T \sim p_\infty(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$;
2. $\mathbf{x}_{t-1} = \boldsymbol{\sigma}_{\boldsymbol{\theta}, t}(\mathbf{x}_t) \cdot \boldsymbol{\epsilon} + \boldsymbol{\mu}_{\boldsymbol{\theta}, t}(\mathbf{x}_t)$;
3. $\mathbf{x}_0 = \mathbf{x} \sim p_{\text{data}}(\mathbf{x})$.

Denoising Diffusion Probabilistic Model (DDPM)

Training

1. Sample $\mathbf{x}_0 \sim p_{\text{data}}(\mathbf{x})$, $t \sim U\{1, T\}$, $\boldsymbol{\epsilon} \sim \mathcal{N}(0, \mathbf{I})$.
2. Compute noisy image $\mathbf{x}_t = \sqrt{\bar{\alpha}_t} \cdot \mathbf{x}_0 + \sqrt{1 - \bar{\alpha}_t} \cdot \boldsymbol{\epsilon}$.
3. Compute loss $\mathcal{L}_{\text{simple}} = \|\boldsymbol{\epsilon} - \boldsymbol{\epsilon}_{\boldsymbol{\theta}, t}(\mathbf{x}_t)\|^2$.

Denoising Diffusion Probabilistic Model (DDPM)

Training

1. Sample $\mathbf{x}_0 \sim p_{\text{data}}(\mathbf{x})$, $t \sim U\{1, T\}$, $\boldsymbol{\epsilon} \sim \mathcal{N}(0, \mathbf{I})$.
2. Compute noisy image $\mathbf{x}_t = \sqrt{\bar{\alpha}_t} \cdot \mathbf{x}_0 + \sqrt{1 - \bar{\alpha}_t} \cdot \boldsymbol{\epsilon}$.
3. Compute loss $\mathcal{L}_{\text{simple}} = \|\boldsymbol{\epsilon} - \boldsymbol{\epsilon}_{\boldsymbol{\theta},t}(\mathbf{x}_t)\|^2$.

Sampling (Ancestral)

1. Sample $\mathbf{x}_T \sim \mathcal{N}(0, \mathbf{I})$.
2. Compute the mean of $p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t) = \mathcal{N}(\boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t), \tilde{\beta}_t \cdot \mathbf{I})$:

$$\boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t) = \frac{1}{\sqrt{\alpha_t}} \cdot \mathbf{x}_t - \frac{1 - \alpha_t}{\sqrt{\alpha_t(1 - \bar{\alpha}_t)}} \cdot \boldsymbol{\epsilon}_{\boldsymbol{\theta},t}(\mathbf{x}_t).$$

3. Denoise $\mathbf{x}_{t-1} = \boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t) + \sqrt{\tilde{\beta}_t} \cdot \boldsymbol{\epsilon}$, $\boldsymbol{\epsilon} \sim \mathcal{N}(0, \mathbf{I})$.

Summary

- The noise schedule sets the signal-to-noise ratio at each diffusion step.
- The reverse diffusion process reconstructs data from noise; we approximate its intractable transitions with Gaussian distributions.
- Conditioning on the clean image gives a tractable Gaussian reverse distribution.
- Gaussian diffusion is a VAE with a hierarchy of latent variables and a fixed encoder.
- Its ELBO decomposes into reconstruction, prior matching, and denoising terms; Gaussian denoising terms reduce to squared error.
- The reverse mean can be parameterized through clean data, noise, score or v; plain MSE induces different noise-level weights.
- DDPM combines a simplified noise-prediction objective with ancestral sampling.