MIPT & YSDA · AUTUMN 2026

# Deep Generative Models

## Lecture 7

Roman Isachenko

Moscow Institute of Physics and Technology
Yandex School of Data Analysis

# Recap of Previous Lecture


Define **score function** $\mathbf{s}_{\boldsymbol{\theta}}(\mathbf{x}) = \nabla_{\mathbf{x}} \log p_{\boldsymbol{\theta}}(\mathbf{x})$.

## Fisher Divergence

$$D_F(p_{\text{data}}, p_{\boldsymbol{\theta}}) = \frac{1}{2}\mathbb{E}_{p_{\text{data}}} \left\| \mathbf{s}_{\boldsymbol{\theta}}(\mathbf{x}) - \nabla_{\mathbf{x}} \log p_{\text{data}}(\mathbf{x}) \right\|_2^2 \to \min_{\boldsymbol{\theta}}$$

## Langevin Dynamics

$$\mathbf{x}_{l+1} = \mathbf{x}_l + \frac{\eta}{2} \cdot \mathbf{s}_{\boldsymbol{\theta}}(\mathbf{x}_l) + \sqrt{\eta} \cdot \boldsymbol{\epsilon}_l, \quad \boldsymbol{\epsilon}_l \sim \mathcal{N}(0, \mathbf{I})$$



# Recap of Previous Lecture

Let us perturb the original data with Gaussian noise $q(\mathbf{x}_\sigma|\mathbf{x}) = \mathcal{N}(\mathbf{x}, \sigma^2 \cdot \mathbf{I})$.


$$q(\mathbf{x}_\sigma) = \int q(\mathbf{x}_\sigma|\mathbf{x})p_{\text{data}}(\mathbf{x})d\mathbf{x}.$$

## Assumption

The solution to

$$\frac{1}{2}\mathbb{E}_{q(\mathbf{x}_\sigma)}\left\|\mathbf{s}_{\boldsymbol{\theta},\sigma}(\mathbf{x}_\sigma) - \nabla_{\mathbf{x}_\sigma}\log q(\mathbf{x}_\sigma)\right\|_2^2 \to \min_{\boldsymbol{\theta}}$$

satisfies $\mathbf{s}_{\boldsymbol{\theta},\sigma}(\mathbf{x}_\sigma) \approx \mathbf{s}_{\boldsymbol{\theta},0}(\mathbf{x}_0) = \mathbf{s}_{\boldsymbol{\theta}}(\mathbf{x})$ if $\sigma$ is sufficiently small.




Song Y. Generative Modeling by Estimating Gradients of the Data Distribution, blog post, 2021
Vincent P. A Connection Between Score Matching and Denoising Autoencoders, 2010

# Recap of Previous Lecture

## Theorem

Under mild regularity conditions, the last term is independent of $\boldsymbol{\theta}$:

$$\begin{aligned}&\mathbb{E}_{q(\mathbf{x}_\sigma)}\left\|\mathbf{s}_{\boldsymbol{\theta},\sigma}(\mathbf{x}_\sigma) - \nabla_{\mathbf{x}_\sigma}\log q(\mathbf{x}_\sigma)\right\|_2^2 \\ &\quad= \mathbb{E}_{p_{\text{data}}(\mathbf{x})}\mathbb{E}_{q(\mathbf{x}_\sigma|\mathbf{x})}\left\|\mathbf{s}_{\boldsymbol{\theta},\sigma}(\mathbf{x}_\sigma) - \nabla_{\mathbf{x}_\sigma}\log q(\mathbf{x}_\sigma|\mathbf{x})\right\|_2^2 + \text{const}\end{aligned}$$

Here,

$$\nabla_{\mathbf{x}_\sigma}\log q(\mathbf{x}_\sigma|\mathbf{x}) = -\frac{\mathbf{x}_\sigma - \mathbf{x}}{\sigma^2} = -\frac{\boldsymbol{\epsilon}}{\sigma}.$$

## Tweedie's Formula

From Lecture 6, the MSE-optimal clean-data prediction is

$$\mathbb{E}_{q(\mathbf{x}|\mathbf{x}_\sigma)}[\mathbf{x}] = \mathbf{x}_\sigma + \sigma^2\nabla_{\mathbf{x}_\sigma}\log q(\mathbf{x}_\sigma).$$


Vincent P. A Connection Between Score Matching and Denoising Autoencoders, 2010
Efron B. Tweedie's Formula and Selection Bias, 2011

# Recap of Previous Lecture

$$\mathbb{E}_{p_{\text{data}}(\mathbf{x})}\mathbb{E}_{\mathcal{N}(0,\mathbf{I})}\left\|\mathbf{s}_{\boldsymbol{\theta},\sigma}(\mathbf{x}+\sigma\boldsymbol{\epsilon})+\frac{\boldsymbol{\epsilon}}{\sigma}\right\|_2^2 \to \min_{\boldsymbol{\theta}}$$

$$\mathbf{x}_{l+1} = \mathbf{x}_l + \frac{\eta}{2}\cdot\mathbf{s}_{\boldsymbol{\theta},\sigma}(\mathbf{x}_l) + \sqrt{\eta}\cdot\boldsymbol{\epsilon}_l$$

- For **small** $\sigma$, $\mathbf{s}_{\boldsymbol{\theta},\sigma}(\mathbf{x})$ becomes inaccurate and Langevin dynamics fails to traverse modes
- For **large** $\sigma$, robustness in low-density regions is achieved, but the model learns a distribution that is overly corrupted




# Recap of Previous Lecture

## Noise-Conditioned Score Network (NCSN)

- Define a sequence of noise levels: $\sigma_1 < \sigma_2 < \cdots < \sigma_T$.
- Balance the score targets $-\boldsymbol{\epsilon}/\sigma_t$ using weights $\sigma_t^2$:

$$\sum_{t=1}^{T} \sigma_t^2 \cdot \mathbb{E}_{p_{\text{data}}(\mathbf{x})} \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x})} \left\| \mathbf{s}_{\boldsymbol{\theta},\sigma_t}(\mathbf{x}_t) - \nabla_{\mathbf{x}_t} \log q(\mathbf{x}_t|\mathbf{x}) \right\|_2^2 \to \min_{\boldsymbol{\theta}}$$

- Sample using **annealed** Langevin dynamics (for $t = T, \ldots, 1$).




Song Y. et al. Generative Modeling by Estimating Gradients of the Data Distribution, 2019

# Recap of Previous Lecture

## Training

1. Sample $\mathbf{x}_0 \sim p_{\text{data}}(\mathbf{x})$, $t \sim U\{1, T\}$, $\boldsymbol{\epsilon} \sim \mathcal{N}(0, \mathbf{I})$.
2. Compute noisy image $\mathbf{x}_t = \mathbf{x}_0 + \sigma_t \boldsymbol{\epsilon}$.
3. Compute loss $\mathcal{L} = \sigma_t^2 \left\| \mathbf{s}_{\boldsymbol{\theta}, \sigma_t}(\mathbf{x}_t) + \frac{\boldsymbol{\epsilon}}{\sigma_t} \right\|^2$.
4. Update $\boldsymbol{\theta}$ via stochastic gradient descent.

## Sampling (Annealed Langevin Dynamics)

1. Sample $\mathbf{x}_T^0 \sim \mathcal{N}(0, \sigma_T^2 \mathbf{I}) \approx q(\mathbf{x}_T)$.
2. Update $\mathbf{x}_t^l$ via $L$ steps of Langevin dynamics at each noise level $\sigma_t$:

$$\mathbf{x}_t^l = \mathbf{x}_t^{l-1} + \frac{\eta_t}{2} \mathbf{s}_{\boldsymbol{\theta}, \sigma_t}(\mathbf{x}_t^{l-1}) + \sqrt{\eta_t} \boldsymbol{\epsilon}_t^l.$$

3. Update $\mathbf{x}_{t-1}^0 := \mathbf{x}_t^L$ and proceed to the next noise level $\sigma_{t-1}$.

Song Y. et al. Generative Modeling by Estimating Gradients of the Data Distribution, 2019
Song Y. et al. Improved Techniques for Training Score-Based Generative Models, 2020

# Recap of Previous Lecture

$$\mathbf{x}_t = \sqrt{1-\beta_t}\mathbf{x}_{t-1} + \sqrt{\beta_t}\boldsymbol{\epsilon}_t, \quad \boldsymbol{\epsilon}_t \sim \mathcal{N}(0, \mathbf{I})$$

$$q(\mathbf{x}_t|\mathbf{x}_{t-1}) = \mathcal{N}(\sqrt{1-\beta_t}\mathbf{x}_{t-1}, \beta_t\mathbf{I})$$


## Statement 1

Let $\alpha_t = 1 - \beta_t$ and $\bar{\alpha}_t = \prod_{s=1}^{t} \alpha_s = \prod_{s=1}^{t}(1-\beta_s)$. Then

$$q(\mathbf{x}_t|\mathbf{x}_0) = \mathcal{N}(\sqrt{\bar{\alpha}_t}\,\mathbf{x}_0, (1-\bar{\alpha}_t)\mathbf{I})$$

# Recap of Previous Lecture

**Diffusion** describes the migration of particles from regions of high density to those of low density.




1. $\mathbf{x}_0 = \mathbf{x} \sim p_{\text{data}}(\mathbf{x})$
2. $\mathbf{x}_t = \sqrt{1-\beta_t}\mathbf{x}_{t-1} + \sqrt{\beta_t}\boldsymbol{\epsilon}_t$, $\boldsymbol{\epsilon}_t \sim \mathcal{N}(0, \mathbf{I})$, $t \geq 1$
3. After $T \gg 1$ steps: $\mathbf{x}_T \sim p_\infty(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$

If this process can be reversed, we can sample from $p_{\text{data}}(\mathbf{x})$ by starting from noise $p_\infty(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$.
Our goal now becomes inverting this diffusion.

# Outline

01 Reverse Gaussian Diffusion Process

02 Gaussian Diffusion Model as VAE

03 Diffusion ELBO Derivation

04 Gaussian Diffusion Reparametrization

05 Denoising Diffusion Probabilistic Model (DDPM)

# Outline

01 **Reverse Gaussian Diffusion Process**


02 Gaussian Diffusion Model as VAE

03 Diffusion ELBO Derivation

04 Gaussian Diffusion Reparametrization

05 Denoising Diffusion Probabilistic Model (DDPM)

# Reverse Gaussian Diffusion Process




## Forward Process

$$q(\mathbf{x}_t|\mathbf{x}_{t-1}) = \mathcal{N}\left(\sqrt{1-\beta_t}\mathbf{x}_{t-1}, \beta_t\mathbf{I}\right)$$

## Reverse Process

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) = \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1})q(\mathbf{x}_{t-1})}{q(\mathbf{x}_t)} \approx p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)$$

$q(\mathbf{x}_{t-1})$ and $q(\mathbf{x}_t)$ are intractable:

$$q(\mathbf{x}_t) = \int q(\mathbf{x}_t|\mathbf{x}_0)p_{\text{data}}(\mathbf{x}_0)d\mathbf{x}_0$$


# Reverse Gaussian Diffusion Process

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) = \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1})q(\mathbf{x}_{t-1})}{q(\mathbf{x}_t)}$$

## Theorem (Feller, 1949)


If $\beta_t$ is sufficiently small, $q(\mathbf{x}_{t-1}|\mathbf{x}_t)$ is approximately Gaussian.

The original DDPM used $T = 1000$ steps. Faster sampling methods will be discussed later.




Feller W. On the theory of stochastic processes, with particular reference to applications, 1949
Ho J. et al. Denoising Diffusion Probabilistic Models, 2020
Xiao Z., Kreis K., Vahdat A. Tackling the generative learning trilemma with denoising diffusion GANs, 2021

# Reverse Gaussian Diffusion Process (Ancestral Sampling)





Define the reverse process as:

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) \approx p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t) = \mathcal{N}\left(\boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t), \boldsymbol{\sigma}^2_{\boldsymbol{\theta},t}(\mathbf{x}_t)\right)$$

If $\beta_t$ is sufficiently small, $q(\mathbf{x}_{t-1}|\mathbf{x}_t)$ is approximately Gaussian.

# Reverse Gaussian Diffusion Process (Ancestral Sampling)




## Forward Process

1. $\mathbf{x}_0 = \mathbf{x} \sim p_{\text{data}}(\mathbf{x})$
2. $\mathbf{x}_t = \sqrt{1-\beta_t}\mathbf{x}_{t-1} + \sqrt{\beta_t}\boldsymbol{\epsilon}_t$
3. $\mathbf{x}_T \sim p_\infty(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$

## Reverse Process

1. $\mathbf{x}_T \sim p_\infty(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$
2. $\mathbf{x}_{t-1} = \boldsymbol{\sigma}_{\boldsymbol{\theta},t}(\mathbf{x}_t)\boldsymbol{\epsilon} + \boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t)$
3. $\mathbf{x}_0 = \mathbf{x} \sim p_{\text{data}}(\mathbf{x})$

**Note:** The forward process is non-learnable, i.e., it does not involve trainable parameters

Weng L. What are Diffusion Models?, blog post, 2021

# Conditioned Reverse Distribution

## Reverse Kernel (Intractable)

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) = \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1})q(\mathbf{x}_{t-1})}{q(\mathbf{x}_t)}$$

## Conditioned Reverse Kernel (Tractable)

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) = \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1}, \mathbf{x}_0)q(\mathbf{x}_{t-1}|\mathbf{x}_0)}{q(\mathbf{x}_t|\mathbf{x}_0)}$$

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) = \frac{\mathcal{N}(\sqrt{1-\beta_t}\cdot\mathbf{x}_{t-1}, \beta_t\mathbf{I})\cdot\mathcal{N}(\sqrt{\bar{\alpha}_{t-1}}\cdot\mathbf{x}_0, (1-\bar{\alpha}_{t-1})\cdot\mathbf{I})}{\mathcal{N}(\sqrt{\bar{\alpha}_t}\cdot\mathbf{x}_0, (1-\bar{\alpha}_t)\cdot\mathbf{I})}$$

$$= \mathcal{N}(\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0), \tilde{\beta}_t\cdot\mathbf{I})$$

$$\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) = \frac{\sqrt{\alpha_t}(1-\bar{\alpha}_{t-1})}{1-\bar{\alpha}_t}\cdot\mathbf{x}_t + \frac{\sqrt{\bar{\alpha}_{t-1}}(1-\alpha_t)}{1-\bar{\alpha}_t}\cdot\mathbf{x}_0$$

$$\tilde{\beta}_t = \frac{(1-\alpha_t)(1-\bar{\alpha}_{t-1})}{1-\bar{\alpha}_t} = \text{const}$$

# What Does Knowing the Clean Data Change?

Same observation $x_t = 0$: unknown clean data versus each known clean value.

### Two possible clean values

$\mathbb{P}(x_0 = -2) = \mathbb{P}(x_0 = 2) = \frac{1}{2}$

$x_0 = -2$


Posterior weight 50.0%

$x_0 = 2$

Posterior weight 50.0%

Unknown $x_0$: posterior-weighted mixture.
Known $x_0$: the corresponding Gaussian.

$q(x_{t-1}|x_t)$ and $q(x_{t-1}|x_t, x_0)$




$$q(x_{t-1}|x_t) = \sum_{x_0 \in \{-2,2\}} q(x_{t-1}|x_t, x_0)\, q(x_0|x_t)$$

Exact toy example. During training $x_0$ is available; during generation it is unknown.

# Distribution Summary

**Forward process** maps any distribution $p_{\text{data}}(\mathbf{x})$ to $\mathcal{N}(0, \mathbf{I})$ by injection of noise:

$$q(\mathbf{x}_t|\mathbf{x}_{t-1}) = \mathcal{N}(\sqrt{1-\beta_t} \cdot \mathbf{x}_{t-1}, \beta_t \cdot \mathbf{I});$$
$$q(\mathbf{x}_t|\mathbf{x}_0) = \mathcal{N}(\sqrt{\bar{\alpha}_t} \cdot \mathbf{x}_0, (1-\bar{\alpha}_t) \cdot \mathbf{I}).$$

**Reverse process** refers to an intractable distribution that can be approximated by a normal distribution (with unknown parameters) for small $\beta_t$:

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t) = \frac{q(\mathbf{x}_t|\mathbf{x}_{t-1})q(\mathbf{x}_{t-1})}{q(\mathbf{x}_t)} \approx \mathcal{N}\left(\boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t), \boldsymbol{\sigma}^2_{\boldsymbol{\theta},t}(\mathbf{x}_t)\right)$$

**Conditioned reverse process** is a normal distribution with known parameters, describing how to denoise a noisy image $\mathbf{x}_t$ when we know the clean image $\mathbf{x}_0$.

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) = \mathcal{N}(\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0), \tilde{\beta}_t \cdot \mathbf{I})$$

# Noise Schedule through Signal-to-Noise Ratio

## Signal and Noise

$$\mathbf{x}_t = \sqrt{\bar{\alpha}_t}\mathbf{x}_0 + \sqrt{1-\bar{\alpha}_t}\boldsymbol{\epsilon}.$$


For $\boldsymbol{\epsilon} \sim \mathcal{N}(0, \mathbf{I})$, compare squared scales:

$$\mathrm{SNR}(t) = \frac{\bar{\alpha}_t}{1-\bar{\alpha}_t}.$$

## A Single Noise-Level Coordinate

$$\lambda_t = \log \mathrm{SNR}(t), \qquad \bar{\alpha}_t = \frac{1}{1+e^{-\lambda_t}}.$$

$\lambda_t > 0$: signal dominates.
$\lambda_t = 0$: equal signal and noise scales.
$\lambda_t < 0$: noise dominates.




Same clean image and Gaussian noise.
Fixed display scale; log-SNR axis clipped to [−12, 10].

**Same timestep, different noise level.**

> **Higher pixel-space resolution:** more noise can help; shifting log-SNR down increases corruption.

Kingma D. et al. Variational Diffusion Models, 2021
Nichol A., Dhariwal P. Improved Denoising Diffusion Probabilistic Models, 2021
Hoogeboom E. et al. Simple diffusion: End-to-end diffusion for high resolution images, 2023

# Outline

01 Reverse Gaussian Diffusion Process

**02 Gaussian Diffusion Model as VAE**

03 Diffusion ELBO Derivation

04 Gaussian Diffusion Reparametrization


05 Denoising Diffusion Probabilistic Model (DDPM)

# Gaussian Diffusion Model as VAE

Let's treat $\mathbf{z} = (\mathbf{x}_1, \dots, \mathbf{x}_T)$ as a latent variable (**note:** each $\mathbf{x}_t$ has the same dimension), and $\mathbf{x} = \mathbf{x}_0$ as the observed variable.

## Latent Variable Model

$$p_{\boldsymbol{\theta}}(\mathbf{x}, \mathbf{z}) = p_{\boldsymbol{\theta}}(\mathbf{x}|\mathbf{z})p_{\boldsymbol{\theta}}(\mathbf{z})$$





## Forward Diffusion

Variational posterior (encoder), with **no learnable parameters**:

$$q(\mathbf{z}|\mathbf{x}) = q(\mathbf{x}_1, \dots, \mathbf{x}_T|\mathbf{x}_0) = \prod_{t=1}^{T} q(\mathbf{x}_t|\mathbf{x}_{t-1}).$$

# Gaussian Diffusion Model as VAE

$$p_{\theta}(\mathbf{x}, \mathbf{z}) = p_{\theta}(\mathbf{x}|\mathbf{z})p_{\theta}(\mathbf{z})$$




## Reverse Diffusion

Generative distribution (decoder)

$$p_{\theta}(\mathbf{x}|\mathbf{z}) = p_{\theta}(\mathbf{x}_0|\mathbf{x}_1).$$

Prior distribution

$$p_{\theta}(\mathbf{z}) = p_{\theta}(\mathbf{x}_1, \ldots, \mathbf{x}_T) = \prod_{t=2}^{T} p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t) \cdot p(\mathbf{x}_T).$$

Unlike the vanilla VAE: a **simple decoder** $p_{\theta}(\mathbf{x}|\mathbf{z})$ and a **Markov-chain prior** $p_{\theta}(\mathbf{z})$.

# Outline

01 Reverse Gaussian Diffusion Process

02 Gaussian Diffusion Model as VAE

**03 Diffusion ELBO Derivation**

04 Gaussian Diffusion Reparametrization

05 Denoising Diffusion Probabilistic Model (DDPM)

# ELBO for Gaussian Diffusion Model

## Standard ELBO

$$\log p_{\boldsymbol{\theta}}(\mathbf{x}) \geq \mathbb{E}_{q(\mathbf{z}|\mathbf{x})} \log \frac{p_{\boldsymbol{\theta}}(\mathbf{x}, \mathbf{z})}{q(\mathbf{z}|\mathbf{x})} = \mathcal{L}_{\boldsymbol{\phi},\boldsymbol{\theta}}(\mathbf{x}) \to \max_{\boldsymbol{\phi},\boldsymbol{\theta}}$$

## Derivation

$$\begin{aligned}\mathcal{L}_{\boldsymbol{\phi},\boldsymbol{\theta}}(\mathbf{x}) &= \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \log \frac{p_{\boldsymbol{\theta}}(\mathbf{x}_0, \mathbf{x}_{1:T})}{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \\ &= \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \log \frac{p(\mathbf{x}_T) \prod_{t=1}^{T} p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{\prod_{t=1}^{T} q(\mathbf{x}_t|\mathbf{x}_{t-1})}\end{aligned}$$

- Let's try to decompose the ELBO into individual KL divergence terms.
- We need to replace $q(\mathbf{x}_t|\mathbf{x}_{t-1})$ with $q(\mathbf{x}_{t-1}|\mathbf{x}_t)$ in the denominator.
- Let's condition on $\mathbf{x}_0$ to make the reverse $q(\mathbf{x}_{t-1}|\mathbf{x}_t)$ tractable.

# ELBO for Gaussian Diffusion Model

## Bayes' Rule ($t \geq 2$)

$$q(\mathbf{x}_t|\mathbf{x}_{t-1}, \mathbf{x}_0) = \frac{q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0)q(\mathbf{x}_t|\mathbf{x}_0)}{q(\mathbf{x}_{t-1}|\mathbf{x}_0)}$$

$$\mathcal{L}_{\phi,\theta}(\mathbf{x}) = \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \log \frac{p(\mathbf{x}_T)\prod_{t=1}^{T} p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{\prod_{t=1}^{T} q(\mathbf{x}_t|\mathbf{x}_{t-1}, \mathbf{x}_0)}$$

$$= \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \log \frac{p(\mathbf{x}_T)p_{\theta}(\mathbf{x}_0|\mathbf{x}_1)\prod_{t=2}^{T} p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{q(\mathbf{x}_1|\mathbf{x}_0)\prod_{t=2}^{T} q(\mathbf{x}_t|\mathbf{x}_{t-1}, \mathbf{x}_0)}$$

$$= \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \log \frac{p(\mathbf{x}_T)p_{\theta}(\mathbf{x}_0|\mathbf{x}_1)\prod_{t=2}^{T} p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{q(\mathbf{x}_1|\mathbf{x}_0)\prod_{t=2}^{T} \frac{q(\mathbf{x}_{t-1}|\mathbf{x}_t,\mathbf{x}_0)q(\mathbf{x}_t|\mathbf{x}_0)}{q(\mathbf{x}_{t-1}|\mathbf{x}_0)}}$$

$$= \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)} \log \frac{p(\mathbf{x}_T)p_{\theta}(\mathbf{x}_0|\mathbf{x}_1)\prod_{t=2}^{T} p_{\theta}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{q(\mathbf{x}_T|\mathbf{x}_0)\prod_{t=2}^{T} q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0)}$$

## Markov Property

$$q(\mathbf{x}_t|\mathbf{x}_{t-1}, \mathbf{x}_0) = q(\mathbf{x}_t|\mathbf{x}_{t-1})$$

Conditioning on $\mathbf{x}_0$ leaves the forward transitions unchanged.

Separate the decoder step $t = 1$.

Apply Bayes' rule for each $t \geq 2$.

Intermediate marginals telescope; the $q(\mathbf{x}_1|\mathbf{x}_0)$ factor cancels too.

# ELBO for Gaussian Diffusion Model

## Derivation (continued)

$$\mathcal{L}_{\phi,\theta}(\mathbf{x}) = \mathbb{E}_{q(\mathbf{x}_{1:T}|\mathbf{x}_0)}\Bigg[\log p_{\boldsymbol{\theta}}(\mathbf{x}_0|\mathbf{x}_1) + \log\frac{p(\mathbf{x}_T)}{q(\mathbf{x}_T|\mathbf{x}_0)} + \sum_{t=2}^{T}\log\left(\frac{p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{q(\mathbf{x}_{t-1}|\mathbf{x}_t,\mathbf{x}_0)}\right)\Bigg]$$

Expand the logarithm: products become sums.

$$= \mathbb{E}_{q(\mathbf{x}_1|\mathbf{x}_0)}\log p_{\boldsymbol{\theta}}(\mathbf{x}_0|\mathbf{x}_1) + \mathbb{E}_{q(\mathbf{x}_T|\mathbf{x}_0)}\log\frac{p(\mathbf{x}_T)}{q(\mathbf{x}_T|\mathbf{x}_0)} + \sum_{t=2}^{T}\mathbb{E}_{q(\mathbf{x}_{t-1},\mathbf{x}_t|\mathbf{x}_0)}\log\left(\frac{p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)}{q(\mathbf{x}_{t-1}|\mathbf{x}_t,\mathbf{x}_0)}\right)$$

Integrate out unused variables. Each term depends on one state or one adjacent pair.


$$= \mathbb{E}_{q(\mathbf{x}_1|\mathbf{x}_0)}\log p_{\boldsymbol{\theta}}(\mathbf{x}_0|\mathbf{x}_1) - D_{\mathrm{KL}}\big(q(\mathbf{x}_T|\mathbf{x}_0)\|p(\mathbf{x}_T)\big) - \sum_{t=2}^{T}\underbrace{\mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)}D_{\mathrm{KL}}\big(q(\mathbf{x}_{t-1}|\mathbf{x}_t,\mathbf{x}_0)\|p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)\big)}_{\mathcal{L}_t}$$

Use the definition of KL. For each $\mathbf{x}_t$, average over $q(\mathbf{x}_{t-1}|\mathbf{x}_t,\mathbf{x}_0)$; then average over $q(\mathbf{x}_t|\mathbf{x}_0)$.

Ho J. Denoising Diffusion Probabilistic Models, 2020

# ELBO for Gaussian Diffusion Model

$$\mathcal{L}_{\phi,\theta}(\mathbf{x}) = \mathbb{E}_{q(\mathbf{x}_1|\mathbf{x}_0)} \log p_{\boldsymbol{\theta}}(\mathbf{x}_0|\mathbf{x}_1) - D_{\mathrm{KL}}\big(q(\mathbf{x}_T|\mathbf{x}_0)\|p(\mathbf{x}_T)\big) - \sum_{t=2}^{T} \underbrace{\mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} D_{\mathrm{KL}}\big(q(\mathbf{x}_{t-1}|\mathbf{x}_t,\mathbf{x}_0)\|p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)\big)}_{\mathcal{L}_t}$$




- First term is the decoder distribution, with $\mathbf{x}_1 \sim q(\mathbf{x}_1|\mathbf{x}_0)$:


$$\log p_{\boldsymbol{\theta}}(\mathbf{x}_0|\mathbf{x}_1) = \log \mathcal{N}\big(\mathbf{x}_0|\boldsymbol{\mu}_{\boldsymbol{\theta},1}(\mathbf{x}_1), \boldsymbol{\sigma}^2_{\boldsymbol{\theta},1}(\mathbf{x}_1)\big).$$

- Second term is constant:

$$p(\mathbf{x}_T) = \mathcal{N}(0, \mathbf{I}), \qquad q(\mathbf{x}_T|\mathbf{x}_0) = \mathcal{N}(\sqrt{\bar{\alpha}_T} \cdot \mathbf{x}_0, (1 - \bar{\alpha}_T) \cdot \mathbf{I}).$$

- Third term is the main contributor to the ELBO.

# ELBO for Gaussian Diffusion Model




$$\mathcal{L}_t = \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} D_{\mathrm{KL}}\left(q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) \| p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)\right)$$

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) = \mathcal{N}(\mathbf{x}_{t-1}|\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0), \tilde{\beta}_t \mathbf{I}),$$

$$p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t) = \mathcal{N}\left(\mathbf{x}_{t-1}|\boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t), \boldsymbol{\sigma}^2_{\boldsymbol{\theta},t}(\mathbf{x}_t)\right)$$

# ELBO for Gaussian Diffusion Model

$$\mathcal{L}_t = \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} D_{\mathrm{KL}}\big(q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) \| p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)\big)$$

$$q(\mathbf{x}_{t-1}|\mathbf{x}_t, \mathbf{x}_0) = \mathcal{N}(\mathbf{x}_{t-1}|\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0), \tilde{\beta}_t \mathbf{I}),$$

$$p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t) = \mathcal{N}\big(\mathbf{x}_{t-1}|\boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t), \boldsymbol{\sigma}^2_{\boldsymbol{\theta},t}(\mathbf{x}_t)\big)$$

Let's assume that

$$\boldsymbol{\sigma}^2_{\boldsymbol{\theta},t}(\mathbf{x}_t) = \tilde{\beta}_t \mathbf{I}$$

$$\Rightarrow \quad p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t) = \mathcal{N}\big(\mathbf{x}_{t-1}|\boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t), \tilde{\beta}_t \mathbf{I}\big).$$

Theoretically, the optimal $\boldsymbol{\sigma}^2_{\boldsymbol{\theta},t}(\mathbf{x}_t)$ lies in $[\tilde{\beta}_t, \beta_t]$:

- $\beta_t$ is optimal for $\mathbf{x}_0 \sim \mathcal{N}(0, \mathbf{I})$;
- $\tilde{\beta}_t$ is optimal for $\mathbf{x}_0 \sim \delta(\mathbf{x}_0 - \mathbf{x}^*)$.

$$\mathcal{L}_t = \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} D_{\mathrm{KL}}\big(\mathcal{N}\big(\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0), \tilde{\beta}_t \mathbf{I}\big) \| \mathcal{N}\big(\boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t), \tilde{\beta}_t \mathbf{I}\big)\big)$$

$$= \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} \left[\frac{1}{2\tilde{\beta}_t} \left\| \tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) - \boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t) \right\|^2\right]$$

# ELBO for Gaussian Diffusion Model

## Training

1. Sample $\mathbf{x}_0 \sim p_{\text{data}}(\mathbf{x})$, $\boldsymbol{\epsilon} \sim \mathcal{N}(0, \mathbf{I})$.
2. Compute noisy image $\mathbf{x}_t = \sqrt{\bar{\alpha}_t} \cdot \mathbf{x}_0 + \sqrt{1 - \bar{\alpha}_t} \cdot \boldsymbol{\epsilon}$.
3. Compute the ELBO:

$$\mathcal{L}_{\boldsymbol{\phi},\boldsymbol{\theta}}(\mathbf{x}) = \mathbb{E}_{q(\mathbf{x}_1|\mathbf{x}_0)} \log p_{\boldsymbol{\theta}}(\mathbf{x}_0|\mathbf{x}_1) - D_{\text{KL}}\left(q(\mathbf{x}_T|\mathbf{x}_0) \| p(\mathbf{x}_T)\right) - \sum_{t=2}^{T} \underbrace{\mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} \left[ \frac{1}{2\tilde{\beta}_t} \left\| \tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) - \boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t) \right\|^2 \right]}_{\mathcal{L}_t}.$$

## Sampling

1. Sample $\mathbf{x}_T \sim \mathcal{N}(0, \mathbf{I})$.
2. Denoise $\mathbf{x}_{t-1} = \boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t) + \sqrt{\tilde{\beta}_t} \cdot \boldsymbol{\epsilon}$, $\boldsymbol{\epsilon} \sim \mathcal{N}(0, \mathbf{I})$.

# Outline

01 Reverse Gaussian Diffusion Process

02 Gaussian Diffusion Model as VAE

03 Diffusion ELBO Derivation

**04 Gaussian Diffusion Reparametrization**

05 Denoising Diffusion Probabilistic Model (DDPM)

# Reparametrization of DDPM

$$\mathcal{L}_t = \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)}\left[\frac{1}{2\tilde{\beta}_t}\left\|\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) - \boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t)\right\|^2\right]$$

$$\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) = \frac{\sqrt{\alpha_t}(1-\bar{\alpha}_{t-1})}{1-\bar{\alpha}_t}\cdot\mathbf{x}_t + \frac{\sqrt{\bar{\alpha}_{t-1}}(1-\alpha_t)}{1-\bar{\alpha}_t}\cdot\mathbf{x}_0$$

$$\mathbf{x}_t = \sqrt{\bar{\alpha}_t}\cdot\mathbf{x}_0 + \sqrt{1-\bar{\alpha}_t}\cdot\boldsymbol{\epsilon} \quad\Rightarrow\quad \mathbf{x}_0 = \frac{\mathbf{x}_t - \sqrt{1-\bar{\alpha}_t}\cdot\boldsymbol{\epsilon}}{\sqrt{\bar{\alpha}_t}}$$

- There is a linear relationship between $\boldsymbol{\epsilon}$, $\mathbf{x}_t$, and $\mathbf{x}_0$.
- Let's try to rewrite this mean using only $\mathbf{x}_t$ and $\boldsymbol{\epsilon}$.

$$\begin{aligned}\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \boldsymbol{\epsilon}) &= \frac{\sqrt{\alpha_t}(1-\bar{\alpha}_{t-1})}{1-\bar{\alpha}_t}\cdot\mathbf{x}_t + \frac{\sqrt{\bar{\alpha}_{t-1}}(1-\alpha_t)}{1-\bar{\alpha}_t}\cdot\left(\frac{\mathbf{x}_t - \sqrt{1-\bar{\alpha}_t}\cdot\boldsymbol{\epsilon}}{\sqrt{\bar{\alpha}_t}}\right) \\ &= \frac{1}{\sqrt{\alpha_t}}\cdot\mathbf{x}_t - \frac{1-\alpha_t}{\sqrt{\alpha_t(1-\bar{\alpha}_t)}}\cdot\boldsymbol{\epsilon}\end{aligned}$$

Ho J. Denoising Diffusion Probabilistic Models, 2020

# Reparametrization of DDPM

$$\mathcal{L}_t = \mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)}\left[\frac{1}{2\tilde{\beta}_t}\left\|\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) - \boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t)\right\|^2\right]$$

## Reparametrization

$$\tilde{\boldsymbol{\mu}}_t(\mathbf{x}_t, \mathbf{x}_0) = \frac{1}{\sqrt{\alpha_t}} \cdot \mathbf{x}_t - \frac{1-\alpha_t}{\sqrt{\alpha_t(1-\bar{\alpha}_t)}} \cdot \boldsymbol{\epsilon}$$

$$\boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t) = \frac{1}{\sqrt{\alpha_t}} \cdot \mathbf{x}_t - \frac{1-\alpha_t}{\sqrt{\alpha_t(1-\bar{\alpha}_t)}} \cdot \boldsymbol{\epsilon}_{\boldsymbol{\theta},t}(\mathbf{x}_t)$$


$$\begin{aligned}\mathcal{L}_t &= \mathbb{E}_{\boldsymbol{\epsilon}\sim\mathcal{N}(0,\mathbf{I})}\left[\frac{(1-\alpha_t)^2}{2\tilde{\beta}_t\alpha_t(1-\bar{\alpha}_t)}\left\|\boldsymbol{\epsilon} - \boldsymbol{\epsilon}_{\boldsymbol{\theta},t}(\mathbf{x}_t)\right\|^2\right] \\ &= \mathbb{E}_{\boldsymbol{\epsilon}\sim\mathcal{N}(0,\mathbf{I})}\left[\frac{(1-\alpha_t)^2}{2\tilde{\beta}_t\alpha_t(1-\bar{\alpha}_t)}\left\|\boldsymbol{\epsilon} - \boldsymbol{\epsilon}_{\boldsymbol{\theta},t}\left(\sqrt{\bar{\alpha}_t}\mathbf{x}_0 + \sqrt{1-\bar{\alpha}_t}\boldsymbol{\epsilon}\right)\right\|^2\right]\end{aligned}$$

At every step of the reverse process, we attempt to predict the noise $\boldsymbol{\epsilon}$ that was used in the forward diffusion process!

# One Prediction, Three Parameterizations

One coordinate. Keep the input and timestep fixed: $x_t = 1, \quad \alpha_t = 0.6, \quad \bar{\alpha}_t = 0.39$

Choose one network output; compute the other two. Example: predict noise.




$$x_t = \sqrt{\bar{\alpha}_t}\hat{x}_0 + \sqrt{1-\bar{\alpha}_t}\hat{\epsilon}$$

$$\mu_{\theta,t} = \frac{x_t}{\sqrt{\alpha_t}} - \frac{1-\alpha_t}{\sqrt{\alpha_t(1-\bar{\alpha}_t)}}\hat{\epsilon}$$

Switch the network output: the values stay the same; only the parameterization changes.

All three are predicted quantities, linked by known conversions.

# Diffusion Parameterizations

$$\mathbf{x}_t = \sqrt{\bar{\alpha}_t}\,\mathbf{x}_0 + \sqrt{1-\bar{\alpha}_t}\,\boldsymbol{\epsilon}.$$

| Network output at $\mathbf{x}_t$ | Regression target | Clean-data estimate |
|---|---|---|
| Noise $\boldsymbol{\epsilon}_{\boldsymbol{\theta},t}$ | $\boldsymbol{\epsilon}$ | $\dfrac{\mathbf{x}_t - \sqrt{1-\bar{\alpha}_t}\,\boldsymbol{\epsilon}_{\boldsymbol{\theta},t}}{\sqrt{\bar{\alpha}_t}}$ |
| Clean data $\hat{\mathbf{x}}_{\boldsymbol{\theta},t}$ | $\mathbf{x}_0$ | $\hat{\mathbf{x}}_{\boldsymbol{\theta},t}$ |
| Score $\mathbf{s}_{\boldsymbol{\theta},t}$ | $-\dfrac{\boldsymbol{\epsilon}}{\sqrt{1-\bar{\alpha}_t}}$ | $\dfrac{\mathbf{x}_t + (1-\bar{\alpha}_t)\,\mathbf{s}_{\boldsymbol{\theta},t}}{\sqrt{\bar{\alpha}_t}}$ |
| $v$-prediction $\mathbf{v}_{\boldsymbol{\theta},t}$ | $\sqrt{\bar{\alpha}_t}\,\boldsymbol{\epsilon} - \sqrt{1-\bar{\alpha}_t}\,\mathbf{x}_0$ | $\sqrt{\bar{\alpha}_t}\,\mathbf{x}_t - \sqrt{1-\bar{\alpha}_t}\,\mathbf{v}_{\boldsymbol{\theta},t}$ |

**Convertible predictions; different losses.** Plain MSE assigns different weights to different noise levels.

Salimans T., Ho J. Progressive Distillation for Fast Sampling of Diffusion Models, 2022

# Reparametrization of DDPM

$$\mathcal{L}_{\phi,\theta}(\mathbf{x}) = \mathbb{E}_{q(\mathbf{x}_1|\mathbf{x}_0)} \log p_{\boldsymbol{\theta}}(\mathbf{x}_0|\mathbf{x}_1) - D_{\mathrm{KL}}\big(q(\mathbf{x}_T|\mathbf{x}_0)\|p(\mathbf{x}_T)\big) - \sum_{t=2}^{T} \underbrace{\mathbb{E}_{q(\mathbf{x}_t|\mathbf{x}_0)} D_{\mathrm{KL}}\big(q(\mathbf{x}_{t-1}|\mathbf{x}_t,\mathbf{x}_0)\|p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t)\big)}_{\mathcal{L}_t}$$

$$\mathcal{L}_t = \mathbb{E}_{\boldsymbol{\epsilon}\sim\mathcal{N}(0,\mathbf{I})}\left[\frac{(1-\alpha_t)^2}{2\tilde{\beta}_t\alpha_t(1-\bar{\alpha}_t)}\left\|\boldsymbol{\epsilon} - \boldsymbol{\epsilon}_{\boldsymbol{\theta},t}\big(\sqrt{\bar{\alpha}_t}\mathbf{x}_0 + \sqrt{1-\bar{\alpha}_t}\boldsymbol{\epsilon}\big)\right\|^2\right]$$

Dropping the time-dependent coefficient **changes the relative weights of noise levels**. The reconstruction term ($t = 1$) also uses noise prediction.

## Simplified Objective

$$\mathcal{L}_{\text{simple}} = \mathbb{E}_{t\sim U\{1,T\}}\mathbb{E}_{\boldsymbol{\epsilon}\sim\mathcal{N}(0,\mathbf{I})}\left\|\boldsymbol{\epsilon} - \boldsymbol{\epsilon}_{\boldsymbol{\theta},t}\big(\sqrt{\bar{\alpha}_t}\cdot\mathbf{x}_0 + \sqrt{1-\bar{\alpha}_t}\cdot\boldsymbol{\epsilon}\big)\right\|^2$$

# Outline

01 Reverse Gaussian Diffusion Process


02 Gaussian Diffusion Model as VAE

03 Diffusion ELBO Derivation

04 Gaussian Diffusion Reparametrization

**05 Denoising Diffusion Probabilistic Model (DDPM)**

# Generative Models Taxonomy



# Denoising Diffusion Probabilistic Model (DDPM)

## DDPM is a VAE Model

- The encoder is a fixed Gaussian Markov chain $q(\mathbf{x}_1, \dots, \mathbf{x}_T | \mathbf{x}_0)$.
- The latent variable is hierarchical (at each step, its dimension equals the input's).
- The decoder is a simple Gaussian model $p_{\boldsymbol{\theta}}(\mathbf{x}_0 | \mathbf{x}_1)$.
- The prior distribution is given by a parametric Gaussian Markov chain $p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1} | \mathbf{x}_t)$.

## Forward Process

1. $\mathbf{x}_0 = \mathbf{x} \sim p_{\text{data}}(\mathbf{x})$;
2. $\mathbf{x}_t = \sqrt{1 - \beta_t} \cdot \mathbf{x}_{t-1} + \sqrt{\beta_t} \cdot \boldsymbol{\epsilon}_t$;
3. $\mathbf{x}_T \sim p_\infty(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$.

## Reverse Process

1. $\mathbf{x}_T \sim p_\infty(\mathbf{x}) = \mathcal{N}(0, \mathbf{I})$;
2. $\mathbf{x}_{t-1} = \boldsymbol{\sigma}_{\boldsymbol{\theta}, t}(\mathbf{x}_t) \cdot \boldsymbol{\epsilon} + \boldsymbol{\mu}_{\boldsymbol{\theta}, t}(\mathbf{x}_t)$;
3. $\mathbf{x}_0 = \mathbf{x} \sim p_{\text{data}}(\mathbf{x})$.

# Denoising Diffusion Probabilistic Model (DDPM)

## Training

1. Sample $\mathbf{x}_0 \sim p_{\text{data}}(\mathbf{x})$, $t \sim U\{1, T\}$, $\boldsymbol{\epsilon} \sim \mathcal{N}(0, \mathbf{I})$.
2. Compute noisy image $\mathbf{x}_t = \sqrt{\bar{\alpha}_t} \cdot \mathbf{x}_0 + \sqrt{1 - \bar{\alpha}_t} \cdot \boldsymbol{\epsilon}$.
3. Compute loss $\mathcal{L}_{\text{simple}} = \|\boldsymbol{\epsilon} - \boldsymbol{\epsilon}_{\boldsymbol{\theta},t}(\mathbf{x}_t)\|^2$.

## Sampling (Ancestral)

1. Sample $\mathbf{x}_T \sim \mathcal{N}(0, \mathbf{I})$.
2. Compute the mean of $p_{\boldsymbol{\theta}}(\mathbf{x}_{t-1}|\mathbf{x}_t) = \mathcal{N}(\boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t), \tilde{\beta}_t \cdot \mathbf{I})$:

$$\boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t) = \frac{1}{\sqrt{\alpha_t}} \cdot \mathbf{x}_t - \frac{1 - \alpha_t}{\sqrt{\alpha_t(1 - \bar{\alpha}_t)}} \cdot \boldsymbol{\epsilon}_{\boldsymbol{\theta},t}(\mathbf{x}_t).$$

3. Denoise $\mathbf{x}_{t-1} = \boldsymbol{\mu}_{\boldsymbol{\theta},t}(\mathbf{x}_t) + \sqrt{\tilde{\beta}_t} \cdot \boldsymbol{\epsilon}$, $\boldsymbol{\epsilon} \sim \mathcal{N}(0, \mathbf{I})$.

# Summary

- The noise schedule sets the signal-to-noise ratio at each diffusion step.
- The reverse diffusion process reconstructs data from noise; we approximate its intractable transitions with Gaussian distributions.
- Conditioning on the clean image gives a tractable Gaussian reverse distribution.
- Gaussian diffusion is a VAE with a hierarchy of latent variables and a fixed encoder.
- Its ELBO decomposes into reconstruction, prior matching, and denoising terms; Gaussian denoising terms reduce to squared error.
- The reverse mean can be parameterized through clean data, noise, score or $v$; plain MSE induces different noise-level weights.
- DDPM reweights the ELBO denoising terms into a simple noise-prediction loss and uses ancestral sampling.