

# Extinction, Fixation, and Invasion in an Ecological Niche

MattheW Badali

Final defense  
performed as a requirement for  
the degree of Doctor of Philosophy  
5 September 2019

# Biodiversity: Number And Distribution Of Species

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction


Fixation

Invasion

Discussion

Extra Slides

## Paradox of the Plankton a problem of biodiversity



corp2365, NOAA Corps Collection

# Biodiversity: Number And Distribution Of Species

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction


Fixation

Invasion


Discussion

Extra Slides

Paradox of the Plankton    Gut Microbiome  
a problem of biodiversity important to health



corp2365, NOAA Corps Collection



E. coli, Rocky Mountain  
Laboratories, NIAID, NIH

# Biodiversity: Number And Distribution Of Species

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction


Fixation

Invasion


Discussion

Extra Slides

Paradox of the Plankton      Gut Microbiome  
a problem of biodiversity important to health



corp2365, NOAA Corps Collection



E. coli, Rocky Mountain  
Laboratories, NIAID, NIH

Conservationism



# Biodiversity: Number And Distribution Of Species

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction


Fixation

Invasion

Discussion


Extra Slides

Paradox of the Plankton      Gut Microbiome  
a problem of biodiversity important to health



corp2365, NOAA Corps Collection

Coalescent Trees



E. coli, Rocky Mountain  
Laboratories, NIAID, NIH

Conservationism



# Biodiversity: Number And Distribution Of Species

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction


Fixation

Invasion

Discussion


Extra Slides

Paradox of the Plankton      Gut Microbiome  
a problem of biodiversity important to health



corp2365, NOAA Corps Collection

Coalescent Trees




*E. coli*, Rocky Mountain  
Laboratories, NIAID, NIH

Conservationism



Small Populations  
e.g. microfluidics



Microfluidic Device, Cooksey/NIST

# Niche Theories - Each Species In Its Own Niche

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation

Invasion

Discussion

Extra Slides

- niche/resource apportionment explains abundance curve



# Niche Theories - Each Species In Its Own Niche

- niche/resource apportionment explains abundance curve



- Competitive Exclusion Principle: “two species cannot coexist if they share a single [ecological] niche”
- niche: survivable values of those factors which affect the birth and death rates
- use logistic equation  $\dot{x} = rx(1 - x/K)$

# Niche Theories - Each Species In Its Own Niche

- niche/resource apportionment explains abundance curve



- Competitive Exclusion Principle: “two species cannot coexist if they share a single [ecological] niche”
- niche: survivable values of those factors which affect the birth and death rates
- use logistic equation  $\dot{x} = rx(1 - x/K)$
- with stochasticity, mean time to extinction  $\tau \sim e^K$

# Niche Theories - Each Species In Its Own Niche

- niche/resource apportionment explains abundance curve



- Competitive Exclusion Principle: “two species cannot coexist if they share a single [ecological] niche”
- niche: survivable values of those factors which affect the birth and death rates
- use logistic equation  $\dot{x} = rx(1 - x/K)$
- with stochasticity, mean time to extinction  $\tau \sim e^K$
- problem: too many resources, parameters required

# Neutral Theories - All Species In One Niche

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation

Invasion

Discussion

Extra Slides

- better prediction of abundance curves (Hubbell),  
also allele frequencies (Kimura), fixation (Moran)
- inherently stochastic

# Neutral Theories - All Species In One Niche

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction


Fixation

Invasion

Discussion

Extra Slides

- better prediction of abundance curves (Hubbell), also allele frequencies (Kimura), fixation (Moran)
- inherently stochastic
- Moran model



- mean time to extinction  $\tau \sim K$

# Neutral Theories - All Species In One Niche

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction


Fixation

Invasion

Discussion

Extra Slides

- better prediction of abundance curves (Hubbell), also allele frequencies (Kimura), fixation (Moran)
- inherently stochastic
- Moran model



- mean time to extinction  $\tau \sim K$
- problem: all species in one niche seems unphysical

# Main Questions Motivating My Research

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation

Invasion

Discussion

Extra Slides

# Main Questions Motivating My Research

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation

Invasion

Discussion

Extra Slides

It is known that stochasticity connects niche and neutral theories: how do the qualitative changes occur between these extremes?

# Main Questions Motivating My Research

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation

Invasion

Discussion

Extra Slides

It is known that stochasticity connects niche and neutral theories: how do the qualitative changes occur between these extremes?

Biodiversity comes from a balance of extinction, fixation (species exiting the system) and invasion (species entering the system): what are the timescales of these processes?

# Main Questions Motivating My Research

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation

Invasion

Discussion

Extra Slides

It is known that stochasticity connects niche and neutral theories: how do the qualitative changes occur between these extremes?

Biodiversity comes from a balance of extinction, fixation (species exiting the system) and invasion (species entering the system): what are the timescales of these processes?

→ **The relevant parameter is niche overlap.**

# How To Calculate The Mean Time To Extinction

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background


Extinction

Fixation

Invasion

Discussion

Extra Slides



Master equation:  $\partial_t P_n = b_{n-1}P_{n-1} + d_{n+1}P_{n+1} - (b_n + d_n)P_n$ ,  
in vector form  $\partial_t \vec{P} = \hat{M}\vec{P}$ , is solved by  $\vec{P}(t) = e^{\hat{M}t}\vec{P}(0)$ .

# How To Calculate The Mean Time To Extinction

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background


Extinction

Fixation

Invasion

Discussion

Extra Slides




Master equation:  $\partial_t P_n = b_{n-1}P_{n-1} + d_{n+1}P_{n+1} - (b_n + d_n)P_n$ ,  
in vector form  $\partial_t \vec{P} = \hat{M}\vec{P}$ , is solved by  $\vec{P}(t) = e^{\hat{M}t}\vec{P}(0)$ .

mean time to extinction  $\tau_n = \frac{1}{b_n+d_n} + \frac{b_n}{b_n+d_n}\tau_{n+1} + \frac{d_n}{b_n+d_n}\tau_{n-1}$ ,

in vector form  $\hat{M}^T \vec{T} = -\vec{1}$ , is solved by  $\boxed{\tau_{n^0} = -\sum_n \hat{M}_{n^0,n}^{-1}}$ .

# Extinction within a Niche

deterministic logistic equation  $\dot{x} = r x \left(1 - \frac{x}{K}\right)$



# Rates Of The Stochastic 1D Logistic Model

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation

Invasion

Discussion

Extra Slides

deterministic logistic equation  $\dot{x} = rx \left(1 - \frac{x}{K}\right)$

$$b_n = r(1 + \delta)n - \frac{r q}{K} n^2$$

$$d_n = r\delta n + \frac{r(1 - q)}{K} n^2$$

# Rates Of The Stochastic 1D Logistic Model

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation

Invasion

Discussion

Extra Slides

deterministic logistic equation  $\dot{x} = rx \left(1 - \frac{x}{K}\right)$

$$b_n = r(1 + \delta)n - \frac{r q}{K} n^2$$

$$d_n = r\delta n + \frac{r(1 - q)}{K} n^2$$

- $\delta$  gives magnitude of birth or death (rather than their average difference, the growth rate  $r$ )
- $q$  shifts intraspecies interactions from increasing death rate ( $q \sim 0$ ) to reducing birth rate ( $q \sim 1$ )



# Mean Time To Extinction Depends On Interactions

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background  
Extinction  
Fixation  
Invasion  
Discussion  
Extra Slides

(inset is heat map of mean time to extinction)



Carrying capacity  $K = 100$ , growth rate  $r = 1$ . The mean time to extinction (MTE) decreases with increased  $\delta$  or decreased  $q$ .



# Mean Time To Extinction Depends On Interactions

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background  
Extinction  
Fixation  
Invasion  
Discussion  
Extra Slides


(inset is heat map of mean time to extinction)



Carrying capacity  $K = 100$ , growth rate  $r = 1$ . The mean time to extinction (MTE) decreases with increased  $\delta$  or decreased  $q$ .

→ **increasing birth and death rates (e.g. competition increasing death rate) reduces the mean time to extinction**


# Fixation versus Coexistence of Two Competing Species



$$\dot{x}_1 = r_1 x_1 \left(1 - \frac{x_1 + a_{12}x_2}{K_1}\right)$$

$$\dot{x}_2 = r_2 x_2 \left(1 - \frac{a_{21}x_1 + x_2}{K_2}\right)$$

# Coupled Logistic Has Niche And Neutral Limits



- for niche theory (independent limit):  
 $a = 0$ ,  
 $\tau \sim e^K$
- for neutral theory (Moran limit):  
 $a = 1$ ,  
 $\tau \sim K$

# How The System Transitions To Neutrality

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation

Invasion

Discussion

Extra Slides

niche theory

$$\tau \sim e^K$$


neutral theory

$$\tau \sim K$$

# How The System Transitions To Neutrality

niche theory  
 $\tau \sim e^K$

neutral theory  
 $\tau \sim K$



# How The System Transitions To Neutrality


niche theory

$$\tau \sim e^K$$

ansatz:  $\tau(a, K) = e^{h(a)} K^{g(a)} e^{f(a)K}$

neutral theory

$$\tau \sim K$$



# How The System Transitions To Neutrality



niche theory

$$\tau \sim e^K$$

ansatz:  $\tau(a, K) = e^{h(a)} K^{g(a)} e^{f(a)K}$

neutral theory

$$\tau \sim K$$



# How The System Transitions To Neutrality



niche theory

$$\tau \sim e^K$$

neutral theory


$$\text{ansatz: } \tau(a, K) = e^{h(a)} K^{g(a)} e^{f(a)K}$$

$$\tau \sim K$$




→ Effective coexistence except for complete niche overlap

# The Timescale of Invasion of a Second Species





- invasion balances extinction to maintain biodiversity

# The Timescale of Invasion of a Second Species



- invasion balances extinction to maintain biodiversity
- invasion is going from one organism to half the population
- invasion into a niche is deterministic, fast (logarithmic)
- invasion into Moran is linear


# Invasion Less Probable As Niche Overlap Increases



→ **Invasion probability lessens as niche overlap increases.**  
The trend is more stark for large  $K$ . Dependence on  $K$  is lesser than  $a$  dependence.

# Invasion Times Are Fast


mean time of successful invasion




Scaling varies from linear in neutral limit to logarithmic in niche limit. → **Successful invasion times are as expected.**

# Invasion Times Are Fast

mean time of successful invasion



mean time of failed attempt



Scaling varies from linear in neutral limit to logarithmic in niche limit. → **Successful invasion times are as expected.**

Except for  $a = 1$  the time has an asymptote of a fast time.  
→ **Transients die quickly.**

# Conclusions

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation

Invasion

Discussion

Extra Slides

Particular scientific contributions:

- mechanism of competition matters: faster extinction when competition increases death rate (lower  $q$ );

# Conclusions

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation

Invasion

Discussion

Extra Slides

Particular scientific contributions:

- mechanism of competition matters: faster extinction when competition increases death rate (lower  $q$ );
- two species will effectively coexist unless they have exactly the same niche;

# Conclusions

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation

Invasion

Discussion

Extra Slides

Particular scientific contributions:

- mechanism of competition matters: faster extinction when competition increases death rate (lower  $q$ );
- two species will effectively coexist unless they have exactly the same niche;
- arbitrarily precise technique captures  $e^{fK}$  and  $K^g$ ;

# Conclusions

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation

Invasion

Discussion

Extra Slides

Particular scientific contributions:

- mechanism of competition matters: faster extinction when competition increases death rate (lower  $q$ );
- two species will effectively coexist unless they have exactly the same niche;
- arbitrarily precise technique captures  $e^{fK}$  and  $K^g$ ;
- greater niche overlap leads to longer invasion times, and less likelihood of success of an attempt;

# Conclusions

Particular scientific contributions:

- mechanism of competition matters: faster extinction when competition increases death rate (lower  $q$ );
- two species will effectively coexist unless they have exactly the same niche;
- arbitrarily precise technique captures  $e^{fK}$  and  $K^g$ ;
- greater niche overlap leads to longer invasion times, and less likelihood of success of an attempt;

More generally:

- with partial niche overlap, extinction is slow and invasion is fast so we expect high biodiversity;

# Conclusions

Particular scientific contributions:

- mechanism of competition matters: faster extinction when competition increases death rate (lower  $q$ );
- two species will effectively coexist unless they have exactly the same niche;
- arbitrarily precise technique captures  $e^{fK}$  and  $K^g$ ;
- greater niche overlap leads to longer invasion times, and less likelihood of success of an attempt;

More generally:

- with partial niche overlap, extinction is slow and invasion is fast so we expect high biodiversity;
- partial niche overlap gives a niche theory with carrying capacities modified by niche overlaps ( $K' = f(a)K$ )

# Potential Future Research

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation

Invasion

Discussion

Extra Slides

# Thank You!

- experimental systems (microfluidics, plasmids, mitochondria)
- predator-prey model (centre fixed point)
- rock-paper-scissors model (limit cycle)
- other 3D models (e.g. chaos)
- SIR model (epidemics)
- evolving parameters (ecology and evolutionary biology)

# Failure Of Fokker-Planck

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation

Invasion

Discussion

Extra Slides

- [101] Charles R Doering, Khachik V Sargsyan, and Leonard M Sander. Extinction times for birth-death processes: exact results, continuum asymptotics, and the failure of the Fokker-Planck approximation. *Multiscale Model. Simul.*, 3(2):283–299, 2005.
- [103] Michael Assaf and Baruch Meerson. Wkb theory of large deviations in stochastic populations. *Journal of Physics A: Mathematical and Theoretical*, 50(26):263001, 2017.
- [117] J. Grasman and D. Ludwig. The accuracy of the diffusion approximation to the expected time to extinction for some discrete stochastic processes. *J. Appl. Probab.*, 20(2):305–321, 1983.

# Failure Of Fokker-Planck

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background  
Extinction

Fixation  
Invasion

Discussion

Extra Slides

- [101] Charles R Doering, Khachik V Sargsyan, and Leonard M Sander. Extinction times for birth-death processes: exact results, continuum asymptotics, and the failure of the Fokker-Planck approximation. *Multiscale Model. Simul.*, 3(2):283–299, 2005.
- [103] Michael Assaf and Baruch Meerson. Wkb theory of large deviations in stochastic populations. *Journal of Physics A: Mathematical and Theoretical*, 50(26):263001, 2017.
- [117] J. Grasman and D. Ludwig. The accuracy of the diffusion approximation to the expected time to extinction for some discrete stochastic processes. *J. Appl. Probab.*, 20(2):305–321, 1983.

with  $V(x) = -2 \int dy \frac{b(y)-d(y)}{b(y)+d(y)}$  the Fokker-Planck approach gives

$$\tau = 2K \int ds \int dt \frac{1}{b(s) + d(s)} \exp [2K (V(s) - V(t))]$$

# Failure Of Fokker-Planck

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background  
Extinction

Fixation  
Invasion

Discussion

Extra Slides

- [101] Charles R Doering, Khachik V Sargsyan, and Leonard M Sander. Extinction times for birth-death processes: exact results, continuum asymptotics, and the failure of the Fokker-Planck approximation. *Multiscale Model. Simul.*, 3(2):283–299, 2005.
- [103] Michael Assaf and Baruch Meerson. Wkb theory of large deviations in stochastic populations. *Journal of Physics A: Mathematical and Theoretical*, 50(26):263001, 2017.
- [117] J. Grasman and D. Ludwig. The accuracy of the diffusion approximation to the expected time to extinction for some discrete stochastic processes. *J. Appl. Probab.*, 20(2):305–321, 1983.

with  $V(x) = -2 \int dy \frac{b(y)-d(y)}{b(y)+d(y)}$  the Fokker-Planck approach gives

$$\tau = 2K \int ds \int dt \frac{1}{b(s) + d(s)} \exp [2K(V(s) - V(t))]$$


$$\begin{aligned}\tau &\approx \frac{2}{V'(0)} \sqrt{\frac{2\pi}{K V''(x^*)}} \frac{e^{-K V(x^*)}}{b(x^*) + d(x^*)} \\ &= \sqrt{\frac{4\pi(1+2\delta)^2}{K(1+\delta-q)}} e^{K(4(1+\delta-q)\ln 2(1+\delta-q)-2(1-2q))/(1-2q)^2} \\ &\sim e^{(4\ln 2-2)K}/\sqrt{K} \text{ and not } \sim e^K/K\end{aligned}$$

# Failure Of Fokker-Planck

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali


Background  
Extinction  
Fixation  
Invasion  
Discussion  
Extra Slides




with  $\delta = 0.4$  and  $q = 0.2$

# Approximation Methods

low  $q$  ( $q = 0.2$ )  
competition in death rate



high  $q$  ( $q = 0.7$ )  
competition in birth rate



appears that FP works for high  $\delta$ , and WKB works for low  $\delta$

# Stochastic Analysis

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation

Invasion

Discussion

Extra Slides

## Master equation

$$\frac{dP_n}{dt} = b_{n-1}P_{n-1}(t) + d_{n+1}P_{n+1}(t) - (b_n + d_n)P_n(t).$$

# Stochastic Analysis

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background  
Extinction

Fixation

Invasion

Discussion

Extra Slides

## Master equation

$$\frac{dP_n}{dt} = b_{n-1}P_{n-1}(t) + d_{n+1}P_{n+1}(t) - (b_n + d_n)P_n(t).$$

$$\dot{\vec{P}}(t) = \hat{M}\vec{P}(t) \text{ is solved by } \vec{P}(t) = \exp\left(\hat{M}t\right)\vec{P}(0)$$

# Stochastic Analysis

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation

Invasion

Discussion

Extra Slides

## Master equation

$$\frac{dP_n}{dt} = b_{n-1}P_{n-1}(t) + d_{n+1}P_{n+1}(t) - (b_n + d_n)P_n(t).$$

$\vec{P}(t) = \hat{M}\vec{P}(t)$  is solved by  $\vec{P}(t) = \exp(\hat{M}t)\vec{P}(0)$

Residence time is  $\langle t(s^0) \rangle_s = -\int_0^\infty dt P(s, t | s^0, 0) = -\hat{M}_{s,s^0}^{-1}$   
so MTE given by  $\hat{M}^T \vec{T} = -\vec{1}$

# Stochastic Analysis

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation

Invasion

Discussion

Extra Slides

## Master equation

$$\frac{dP_n}{dt} = b_{n-1}P_{n-1}(t) + d_{n+1}P_{n+1}(t) - (b_n + d_n)P_n(t).$$

$\vec{P}(t) = \hat{M}\vec{P}(t)$  is solved by  $\vec{P}(t) = \exp(\hat{M}t)\vec{P}(0)$

Residence time is  $\langle t(s^0) \rangle_s = -\int_0^\infty dt P(s, t | s^0, 0) = -\hat{M}_{s,s^0}^{-1}$

so MTE given by  $\hat{M}^T \vec{T} = -\vec{1}$

With a stable fixed point  $\tau \sim e^K$  (actually  $e^K/K$ )

# How To Calculate The Mean Time To Extinction

Master equation  $\dot{\vec{P}}(t) = \hat{M}\vec{P}(t)$  is solved by  $\vec{P}(t) = e^{\hat{M}t}\vec{P}(0)$ .

# How To Calculate The Mean Time To Extinction

Master equation  $\dot{\vec{P}}(t) = \hat{M}\vec{P}(t)$  is solved by  $\vec{P}(t) = e^{\hat{M}t}\vec{P}(0)$ .

Residence time  $= \int_0^\infty dt P(s|s^0) = -\hat{M}_{s,s^0}^{-1}$  so  $\boxed{\hat{M}\vec{T} = -\vec{1}}$ .

# How To Calculate The Mean Time To Extinction

Master equation  $\dot{\vec{P}}(t) = \hat{M}\vec{P}(t)$  is solved by  $\vec{P}(t) = e^{\hat{M}t}\vec{P}(0)$ .

Residence time  $= \int_0^\infty dt P(s|s^0) = -\hat{M}_{s,s^0}^{-1}$  so  $\hat{M}\vec{T} = -\vec{1}$ .

---

## Approximations

- $\tau \approx \frac{1}{d_1 P_1^c}$

# How To Calculate The Mean Time To Extinction

Master equation  $\dot{\vec{P}}(t) = \hat{M}\vec{P}(t)$  is solved by  $\vec{P}(t) = e^{\hat{M}t}\vec{P}(0)$ .

Residence time  $= \int_0^\infty dt P(s|s^0) = -\hat{M}_{s,s^0}^{-1}$  so  $\boxed{\hat{M}\vec{T} = -\vec{1}}$ .

## Approximations

- $\tau \approx \frac{1}{d_1 P_1^c}$

### Fokker-Planck:

$$\partial_t P_x = -\partial_x ((b_x - d_x)P_x) + \frac{1}{2K} \partial_x^2 ((b_x + d_x)P_x)$$

- $P_n \approx \frac{1}{\sqrt{2\pi\sigma^2}} \exp \left\{ -\frac{(n-n^*)^2}{2\sigma^2} \right\}$  with  $\sigma^2 = \frac{-(b_n+d_n)|_{n^*}}{2\partial_n(b_n-d_n)|_{n^*}}$

# How To Calculate The Mean Time To Extinction

Master equation  $\dot{\vec{P}}(t) = \hat{M}\vec{P}(t)$  is solved by  $\vec{P}(t) = e^{\hat{M}t}\vec{P}(0)$ .

Residence time  $= \int_0^\infty dt P(s|s^0) = -\hat{M}_{s,s^0}^{-1}$  so  $\boxed{\hat{M}\vec{T} = -\vec{1}}$ .

---

## Approximations

- $\tau \approx \frac{1}{d_1 P_1^c}$

---

- Fokker-Planck:

$$\partial_t P_x = -\partial_x ((b_x - d_x)P_x) + \frac{1}{2K} \partial_x^2 ((b_x + d_x)P_x)$$

- $P_n \approx \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left\{-\frac{(n-n^*)^2}{2\sigma^2}\right\}$  with  $\sigma^2 = \frac{-(b_n+d_n)|_{n^*}}{2\partial_n(b_n-d_n)|_{n^*}}$

---

- WKB ansatz:  $P_n \propto \exp\left\{K \sum_i \frac{1}{K^i} S_i(n)\right\}$  with  $S_0(n) = \frac{1}{K} \int_0^n dx \ln(b_x/d_x)$  along extinction trajectory

# Mean Time to Extinction

## Approximations

- larger fluctuations lead to shorter MTE:  $\tau \approx \frac{1}{d_1 P_1}$

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation

Invasion

Discussion

Extra Slides

# Mean Time to Extinction

## Approximations

- larger fluctuations lead to shorter MTE:  $\tau \approx \frac{1}{d_1 P_1}$
- $\hat{M} \vec{T} = -\vec{1}$  is equivalent to  $\tau(n) = \sum_{i=1}^N \frac{1}{d_i} \prod_{k=1}^{i-1} \frac{b_k}{d_k} + \sum_{j=1}^{n-1} \prod_{l=1}^j \frac{d_l}{b_l} \sum_{i=j+1}^N \frac{1}{d_i} \prod_{k=1}^{i-1} \frac{b_k}{d_k}$

# Mean Time to Extinction

## Approximations

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation

Invasion

Discussion

Extra Slides

- larger fluctuations lead to shorter MTE:  $\tau \approx \frac{1}{d_1 P_1}$
- $\hat{M} \vec{T} = -\vec{1}$  is equivalent to  $\tau(n) = \sum_{i=1}^N \frac{1}{d_i} \prod_{k=1}^{i-1} \frac{b_k}{d_k} + \sum_{j=1}^{n-1} \prod_{l=1}^j \frac{d_l}{b_l} \sum_{i=j+1}^N \frac{1}{d_i} \prod_{k=1}^{i-1} \frac{b_k}{d_k}$
- Fokker-Planck equation  $\partial_t P(x, t) = -\partial_x((b(x) - d(x))P(x, t)) + \frac{1}{2K} \partial_x^2((b(x) + d(x))P(x, t))$
- Gaussian approximation<sup>†</sup>  $p(n) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left\{-\frac{(n-n^*)^2}{2\sigma^2}\right\}$   
with  $\sigma^2 = \frac{-(b_n+d_n)|_{n=n^*}}{2\partial_n(b_n-d_n)|_{n=n^*}}$

# Mean Time to Extinction

## Approximations

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation

Invasion

Discussion

Extra Slides

- larger fluctuations lead to shorter MTE:  $\tau \approx \frac{1}{d_1 P_1}$
- $\hat{M} \vec{T} = -\vec{1}$  is equivalent to  $\tau(n) = \sum_{i=1}^N \frac{1}{d_i} \prod_{k=1}^{i-1} \frac{b_k}{d_k} + \sum_{j=1}^{n-1} \prod_{l=1}^j \frac{d_l}{b_l} \sum_{i=j+1}^N \frac{1}{d_i} \prod_{k=1}^{i-1} \frac{b_k}{d_k}$
- Fokker-Planck equation  $\partial_t P(x, t) = -\partial_x((b(x) - d(x))P(x, t)) + \frac{1}{2K} \partial_x^2((b(x) + d(x))P(x, t))$ 
  - Gaussian approximation<sup>†</sup>  $p(n) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left\{-\frac{(n-n^*)^2}{2\sigma^2}\right\}$   
with  $\sigma^2 = \frac{-(b_n+d_n)|_{n=n^*}}{2\partial_n(b_n-d_n)|_{n=n^*}}$
  - WKB ansatz  $P_n \propto \exp\left\{K \sum_i \frac{1}{K^i} S_i(n)\right\}$   
with  $S_0(n) = \int_{n=0}^K dn \ln\left(\frac{b_n}{d_n}\right)$  along extinction trajectory

<sup>†</sup>Gaussian approximation was written incorrectly in thesis.

# Mean Time to Extinction Approximations

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background



Extinction

Fixation

Invasion

Discussion



Extra Slides



*Approximations of the MTE in various regimes of parameter space. WKB is good for low  $\delta$ , is otherwise poor as FP.*

# Extinction within a Niche

deterministic logistic equation  $\dot{x} = rx(1 - \frac{x}{K})$



# Quasi-Steady State

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background


Extinction

Fixation

Invasion

Discussion

Extra Slides



*Characterizing the quasi-stationary probability distribution function for varying  $\delta$  and  $q$ .* Lightness indicates an increased mean or variance in left and right respectively. Carrying capacity  $K = 100$ . The QSD has decreasing mean and increasing variance with increased  $\delta$  or decreased  $q$ .

**Coexistence  
and  
Extinction of  
Competing  
Species**

**M.A.Badali**

Background

Extinction

Fixation

Invasion

Discussion

**Extra Slides**

# Fixation

# Coupled Logistic Equations

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background


Extinction

Fixation

Invasion

Discussion

Extra Slides



# Coupled Logistic Equations

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background


Extinction

Fixation

Invasion

Discussion

Extra Slides



$$\dot{x}_1 = (\beta_1 - \mu_1 - e_{11}t_1 - e_{12}t_2)x_1$$

$$\dot{x}_2 = (\beta_2 - \mu_2 - e_{21}t_1 - e_{22}t_2)x_2$$

$$\dot{t}_1 = g_{11}x_1 + g_{12}x_2 - \lambda_1 t_1$$

$$\dot{t}_2 = g_{21}x_1 + g_{22}x_2 - \lambda_2 t_2$$

# Coupled Logistic Equations

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background


Extinction

Fixation

Invasion

Discussion

Extra Slides



$$\dot{x}_1 = (\beta_1 - \mu_1 - e_{11}t_1 - e_{12}t_2)x_1$$

$$\dot{x}_2 = (\beta_2 - \mu_2 - e_{21}t_1 - e_{22}t_2)x_2$$

$$\dot{t}_1 = g_{11}x_1 + g_{12}x_2 - \lambda_1 t_1$$

$$\dot{t}_2 = g_{21}x_1 + g_{22}x_2 - \lambda_2 t_2$$

$$\dot{\vec{x}} = \hat{R}\hat{X} \left( \vec{1} - (\hat{E}\hat{G})\vec{x} \right)$$

# Coupled Logistic Equations

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background


Extinction

Fixation

Invasion

Discussion

Extra Slides



$$\dot{x}_1 = (\beta_1 - \mu_1 - e_{11}t_1 - e_{12}t_2)x_1$$

$$\dot{x}_2 = (\beta_2 - \mu_2 - e_{21}t_1 - e_{22}t_2)x_2$$

$$\dot{t}_1 = g_{11}x_1 + g_{12}x_2 - \lambda_1 t_1$$

$$\dot{t}_2 = g_{21}x_1 + g_{22}x_2 - \lambda_2 t_2$$

$$\dot{\vec{x}} = \hat{R}\hat{X} \left( \vec{1} - (\hat{E}\hat{G})\vec{x} \right)$$

When the matrix  $(\hat{E}\hat{G})$  is singular ( $a_{12}a_{21} = 1$ , complete niche overlap), the coexistence fixed point  $\vec{x}^* = (EG)^{-1}\vec{1}$  does not exist. Coexistence is allowed only when neutral:  $K_1/K_2 = a_{12} = 1/a_{21}$ .

# Coupled Logistic Includes Competitive Exclusion

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background


Extinction

Fixation

Invasion

Discussion

Extra Slides



2,6 = parasitism/predation/antagonism, 3,7 = mutualism,  
4,5 = competitive exclusion, 1 = (weak) competition

# Coupled Logistic Includes Competitive Exclusion

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background



Extinction

Fixation

Invasion

Discussion

Extra Slides




$2, 6 =$  parasitism/predation/antagonism,  $3, 7 =$  mutualism,

$4, 5 =$  competitive exclusion,  $1 =$  (weak) competition



# Exponential Scaling With $K$ Is Long

■ ansatz:  $\tau(a, K) = e^{h(a)} K^{g(a)} e^{f(a)K}$



# Route to Fixation

$$\text{Residence time } \langle t(s^0) \rangle_s = \int_0^\infty dt P(s, t | s^0, 0) = \hat{M}_{s, s^0}^{-1}$$



*The system samples multiple trajectories on its way to fixation.*  
Left: Complete niche overlap limit,  $a = 1$ , for  $K = 64$ .  
Right: Independent limit with  $a = 0$  and  $K = 32$ .

# Discussion

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation

Invasion

Discussion

Extra Slides

$f(a)$  (exponential dependence of MTE) approaches zero monotonically as niche overlap reaches Moran limit  $a = 1$

- only for complete niche overlap will there be no exponential dependence: fixation will be rapid

# Discussion

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation

Invasion

Discussion

Extra Slides

$f(a)$  (exponential dependence of MTE) approaches zero monotonically as niche overlap reaches Moran limit  $a = 1$

- only for complete niche overlap will there be no exponential dependence: fixation will be rapid
- any niche mismatch allows for exponential dependence on  $K$ , which is typically large
  - any niche mismatch implies effective coexistence

# Discussion

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation

Invasion

Discussion

Extra Slides

$f(a)$  (exponential dependence of MTE) approaches zero monotonically as niche overlap reaches Moran limit  $a = 1$

- only for complete niche overlap will there be no exponential dependence: fixation will be rapid
- any niche mismatch allows for exponential dependence on  $K$ , which is typically large
  - any niche mismatch implies effective coexistence
- small departure from neutrality gives a niche theory

**Coexistence  
and  
Extinction of  
Competing  
Species**

**M.A.Badali**

Background

Extinction

Fixation

Invasion

Discussion

**Extra Slides**

# Invasion

# Invasion - Definition And Expectations

Invasion is going from one organism to half the population.

- invasion balances extinction to maintain biodiversity

# Invasion - Definition And Expectations

Invasion is going from one organism to half the population.



- invasion balances extinction to maintain biodiversity
- invasion in niche/independent limit should be fast (logarithmic)

# Invasion - Definition And Expectations

Invasion is going from one organism to half the population.

- invasion balances extinction to maintain biodiversity
- invasion in niche/independent limit should be fast (logarithmic)
- invasion in neutral limit should be slower (linear)

# Invasion Probability



*Probability of a successful invasion.* Left: Numerical results, from  $a = 0$  at the top to  $a = 1$  at the bottom. The purple solid line is the expected analytical solution in the independent limit. The green solid line is the prediction of the Moran model in the complete niche overlap case. Right: The red data show the results for carrying capacity  $K = 4$ , and suggest the solid black line  $\frac{b_{mut}}{b_{mut} + d_{mut}}$  is an appropriate small carrying capacity limit. Successive lines are at larger system size, and approach the solid magenta line of  $1 - d_{mut}/b_{mut} \approx 1 - a$ .

# Invasion Probability

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction


Fixation

Invasion

Discussion

Extra Slides

Invasion probability  
approaches  $1 - a$ .



# Invasion Probability

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation


Invasion

Discussion



Extra Slides

Invasion probability  
approaches  $1 - a$ .

Successful invasion  
goes from logarithmic  
to linear in  $K$ .



Failed invasion  
attempts go  
from constant to  
logarithmic in  $K$ .



# Invasion Times

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background



Extinction

Fixation

Invasion

Discussion

Extra Slides



Mean time of a successful or failed invasion attempt. Left: Mean time vs  $K$ . Right: Mean time vs  $a$ . Upper: Mean time conditioned on eventual invasion success. Lower: Mean time conditioned on failed attempt.

# Discussion

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation

Invasion

Discussion

Extra Slides

- we can rationalize most of the behaviour
- some questions remain (why is there a max time for failed attempts, why do probabilities remain intermediate for large  $K$ )
- implication is that any invasion attempt (whether successful or not) is faster than fixation times
- comparison of interest is invasion attempt times with immigration rate

# Maintenance of a Species with Repeated Immigration

# Moran Model With Immigration

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation

Invasion

Discussion

Extra Slides

Immigration comes from a constant reservoir of focal species fraction  $g = n_{\text{reservoir}}/K_{\text{reservoir}}$  at a rate  $\nu$ . Defining  $f = n/K$ , we have the following transition rates.

transition	function	value
$n \rightarrow n + 1$	$b(n)$	$f(1 - f)(1 - \nu) + \nu g(1 - f)$
$n \rightarrow n - 1$	$d(n)$	$f(1 - f)(1 - \nu) + \nu(1 - g)f$
$n \rightarrow n$	$1 - b - d$	$(f^2 + (1 - f)^2)(1 - \nu) + \nu(gf + (1 - g)(1 - f))$

# Moran Model With Immigration

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation

Invasion

Discussion

Extra Slides

Immigration comes from a constant reservoir of focal species fraction  $g = n_{\text{reservoir}}/K_{\text{reservoir}}$  at a rate  $\nu$ . Defining  $f = n/K$ , we have the following transition rates.

transition	function	value
$n \rightarrow n + 1$	$b(n)$	$f(1 - f)(1 - \nu) + \nu g(1 - f)$
$n \rightarrow n - 1$	$d(n)$	$f(1 - f)(1 - \nu) + \nu(1 - g)f$
$n \rightarrow n$	$1 - b - d$	$(f^2 + (1 - f)^2)(1 - \nu) + \nu(gf + (1 - g)(1 - f))$

The crucial comparison for biodiversity is between  $1/\nu$  and the invasion times previously described.

# Steady State Results

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background



Extinction

Fixation

Invasion

Discussion

Extra Slides



Metapopulation focal fraction is  $g = 0.4$ , local system size  $N = 100$ , immigration rate  $\nu$  is given by the colour.

For high immigration rate the distribution should be centered near the metapopulation fraction  $g N$  whereas for low immigration the system spends most of its time fixated. What is “high” and “low” depends on  $g$ .

# Infrequent Immigration

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation


Invasion

Discussion

Extra Slides

The model recovers qualitative experimental results.  
(See Vega and Gore, *PLoS Biology*, 2017.)

C



# First Passage Results

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background



Extinction

Fixation

Invasion

Discussion

Extra Slides



Metapopulation focal fraction is  $g = 0.4$ , local system size  $N = 100$ , immigration rate  $\nu$  is given by the colour. The black line is the regular Moran result without immigration.

When the immigrant is mostly not from the focal species ( $g < 0.5$ ) immigration increases the likelihood of the focal species going extinct before fixating. Conditioned first passage times are longer when immigration is more frequent, rare events take even longer still.

# Discussion

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation

Invasion

Discussion

Extra Slides

- when immigration is uncommon  
 $(N\nu < \min(1/g, 1/(1-g)))$ , focal species either fixated or extinct most of the time
- when immigration is common  
 $(N\nu > \max(1/g, 1/(1-g)))$ , focal species is maintained at moderate abundance in the system, specifically  $gN$
- immigration increases the times to (temporary) fixation or extinction

**Coexistence  
and  
Extinction of  
Competing  
Species**

**M.A.Badali**

Background

Extinction

Fixation

Invasion

Discussion

**Extra Slides**

# Discussion

# Conclusions

- higher commensurate birth and death rates (*i.e.* higher  $\delta$ , lower  $q$ ) leads to faster extinction;

# Conclusions

- higher commensurate birth and death rates (*i.e.* higher  $\delta$ , lower  $q$ ) leads to faster extinction;
- WKB is fine for exponential scaling of the MTE, FP fails;

# Conclusions

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation

Invasion

Discussion

Extra Slides

- higher commensurate birth and death rates (*i.e.* higher  $\delta$ , lower  $q$ ) leads to faster extinction;
- WKB is fine for exponential scaling of the MTE, FP fails;
- two species will effectively coexist unless they have exactly the same niche;
- similarly, greater niche overlap leads to longer invasion times, and less likelihood of success of an attempt;

# Conclusions

- higher commensurate birth and death rates (*i.e.* higher  $\delta$ , lower  $q$ ) leads to faster extinction;
- WKB is fine for exponential scaling of the MTE, FP fails;
- two species will effectively coexist unless they have exactly the same niche;
- similarly, greater niche overlap leads to longer invasion times, and less likelihood of success of an attempt;
- in Moran model with immigration, a focal species at moderate size if  $K\nu > 1/g$ ;

# Conclusions

- higher commensurate birth and death rates (*i.e.* higher  $\delta$ , lower  $q$ ) leads to faster extinction;
- WKB is fine for exponential scaling of the MTE, FP fails;
- two species will effectively coexist unless they have exactly the same niche;
- similarly, greater niche overlap leads to longer invasion times, and less likelihood of success of an attempt;
- in Moran model with immigration, a focal species at moderate size if  $K\nu > 1/g$ ;
- incomplete niche overlap is a niche theory with carrying capacities modified by niche overlaps;
- complete niche overlap (neutralism) on an island with immigration has abundance curve like mainland for species with  $g_i > 1/K\nu$ ; other species are transients.

# Utility Of My Results

Coexistence  
and  
Extinction of  
Competing  
Species

M.A.Badali

Background

Extinction

Fixation

Invasion

Discussion

Extra Slides

To reiterate:

- human health (gut microbiome)
- planet health (conservation)
- minimal working models
- coalescent theory
- small population systems like:
  - microfluidics
  - plasmids
  - mitochondria