

Extinction, Fixation, and Invasion in an Ecological Niche

MattheW Badali

Internal defense
performed as a requirement for
the degree of Doctor of Philosophy
4 July 2019

Motivation and Background

Coexistence
and
Extinction of
Competing
Species

M.A.Badali

Introduction

Extinction


Fixation

Invasion

Maintenance

Discussion

Paradox of the Plankton - a problem of biodiversity



corp2365, NOAA Corps Collection

Motivation and Background

Coexistence
and
Extinction of
Competing
Species

M.A.Badali

Introduction

Extinction


Fixation

Invasion

Maintenance

Discussion

Paradox of the Plankton - a problem of biodiversity



corp2365, NOAA Corps Collection

- human health (gut)
- planet health (conservation)
- minimal working models
- coalescent theory

Niche Theories

Coexistence and Extinction of Competing Species

M.A.Badali

Introduction

Extinction

Fixation

Invasion

Maintenance

Discussion

- Competitive Exclusion: “two species cannot coexist if they share a single [ecological] niche”¹

¹Gause. *Science*, 1934

Niche Theories

- Competitive Exclusion: “two species cannot coexist if they share a single [ecological] niche”¹
- Lotka-Volterra

$$\frac{\dot{x}_1}{r_1 x_1} = 1 - \frac{(x_1 + a_{12}x_2)}{K_1}$$
$$\frac{\dot{x}_2}{r_2 x_2} = 1 - \frac{(a_{21}x_1 + x_2)}{K_2}.$$

¹Gause. *Science*, 1934

Niche Theories

- Competitive Exclusion: “two species cannot coexist if they share a single [ecological] niche”¹
- Lotka-Volterra

$$\frac{\dot{x}_1}{r_1 x_1} = 1 - \frac{(x_1 + a_{12}x_2)}{K_1}$$
$$\frac{\dot{x}_2}{r_2 x_2} = 1 - \frac{(a_{21}x_1 + x_2)}{K_2}.$$

- Niche Apportionment

¹Gause. *Science*, 1934

Stochastic Analysis

Coexistence
and
Extinction of
Competing
Species

M.A.Badali

Introduction


Extinction

Fixation

Invasion

Maintenance

Discussion



- Demographic Stochasticity: fluctuations

Stochastic Analysis

Coexistence
and
Extinction of
Competing
Species

M.A.Badali

Introduction


Extinction

Fixation

Invasion

Maintenance

Discussion



- Demographic Stochasticity: fluctuations
- Probability of being in state, Extinction

Stochastic Analysis

Coexistence
and
Extinction of
Competing
Species

M.A.Badali

Introduction

Extinction

Fixation

Invasion

Maintenance

Discussion

Master equation

$$\frac{dP_n}{dt} = b_{n-1}P_{n-1}(t) + d_{n+1}P_{n+1}(t) - (b_n + d_n)P_n(t).$$

Stochastic Analysis

Coexistence
and
Extinction of
Competing
Species

M.A.Badali

Introduction

Extinction

Fixation

Invasion

Maintenance

Discussion

Master equation

$$\frac{dP_n}{dt} = b_{n-1}P_{n-1}(t) + d_{n+1}P_{n+1}(t) - (b_n + d_n)P_n(t).$$

$$\dot{\vec{P}}(t) = \hat{M}\vec{P}(t) \text{ is solved by } \vec{P}(t) = \exp(\hat{M}t) \vec{P}(0)$$

Stochastic Analysis

Coexistence
and
Extinction of
Competing
Species

M.A.Badali

Introduction

Extinction

Fixation

Invasion

Maintenance

Discussion

Master equation

$$\frac{dP_n}{dt} = b_{n-1}P_{n-1}(t) + d_{n+1}P_{n+1}(t) - (b_n + d_n)P_n(t).$$

$\dot{\vec{P}}(t) = \hat{M}\vec{P}(t)$ is solved by $\vec{P}(t) = \exp(\hat{M}t) \vec{P}(0)$

Residence time is $\langle t(s^0) \rangle_s = \int_0^\infty dt P(s, t | s^0, 0) = \hat{M}_{s, s^0}^{-1}$

so MTE given by $\hat{M}\vec{T} = -\vec{1}$

Stochastic Analysis

Coexistence
and
Extinction of
Competing
Species

M.A.Badali

Introduction

Extinction

Fixation

Invasion

Maintenance

Discussion

Master equation

$$\frac{dP_n}{dt} = b_{n-1}P_{n-1}(t) + d_{n+1}P_{n+1}(t) - (b_n + d_n)P_n(t).$$

$\dot{\vec{P}}(t) = \hat{M}\vec{P}(t)$ is solved by $\vec{P}(t) = \exp(\hat{M}t) \vec{P}(0)$


Residence time is $\langle t(s^0) \rangle_s = \int_0^\infty dt P(s, t | s^0, 0) = \hat{M}_{s, s^0}^{-1}$

so MTE given by $\hat{M}\vec{T} = -\vec{1}$

With a stable fixed point $\tau \sim e^K$ (actually e^K/K)

Neutral Theories

- inherently stochastic
- used for allele frequencies (Kimura), fixation (Moran), abundance curves (Hubbell)
- Moran model



$$\tau(n) = -\Delta t K^2 \left(\frac{n}{K} \ln \left(\frac{n}{K} \right) + \frac{K-n}{K} \ln \left(\frac{K-n}{K} \right) \right) \sim K.$$

Structure of Thesis

Coexistence
and
Extinction of
Competing
Species

M.A.Badali

Introduction

Extinction

Fixation

Invasion

Maintenance

Discussion

- Biodiversity is balance of species out and species in to system

Structure of Thesis

Coexistence
and
Extinction of
Competing
Species

M.A.Badali

Introduction

Extinction

Fixation

Invasion

Maintenance

Discussion

- Biodiversity is balance of species out and species in to system
- Extinction - Single Logistic System

Structure of Thesis

Coexistence
and
Extinction of
Competing
Species

M.A.Badali

Introduction

Extinction

Fixation

Invasion

Maintenance

Discussion

- Biodiversity is balance of species out and species in to system
- Extinction - Single Logistic System
- Fixation - Coupled Logistic System/LV

Structure of Thesis

Coexistence
and
Extinction of
Competing
Species

M.A.Badali

Introduction

Extinction

Fixation

Invasion

Maintenance

Discussion

- Biodiversity is balance of species out and species in to system
- Extinction - Single Logistic System
- Fixation - Coupled Logistic System/LV
- Invasion - Coupled Logistic System/LV

Structure of Thesis

Coexistence
and
Extinction of
Competing
Species

M.A.Badali

Introduction

Extinction

Fixation

Invasion

Maintenance

Discussion

- Biodiversity is balance of species out and species in to system
- Extinction - Single Logistic System
- Fixation - Coupled Logistic System/LV
- Invasion - Coupled Logistic System/LV
- Maintenance - Moran with Immigration

Structure of Thesis

Coexistence
and
Extinction of
Competing
Species

M.A.Badali

Introduction

Extinction

Fixation

Invasion

Maintenance

Discussion

- Biodiversity is balance of species out and species in to system
- Extinction - Single Logistic System
- Fixation - Coupled Logistic System/LV
- Invasion - Coupled Logistic System/LV
- Maintenance - Moran with Immigration
- Discussion

Extinction

Logistic Equation

The deterministic logistic equation $\dot{x} = r x \left(1 - \frac{x}{K}\right)$

Coexistence
and
Extinction of
Competing
Species

M.A.Badali

Introduction

Extinction

Fixation


Invasion

Maintenance

Discussion

Logistic Equation

The deterministic logistic equation $\dot{x} = r x \left(1 - \frac{x}{K}\right)$




can be derived from a stochastic model with birth/death rates

$$b_n = (1 + \delta) r n - \frac{q r}{K} n^2$$

$$d_n = \delta r n + \frac{(1 - q) r}{K} n^2$$

Logistic Equation

The deterministic logistic equation $\dot{x} = r x \left(1 - \frac{x}{K}\right)$



can be derived from a stochastic model with birth/death rates

$$b_n = (1 + \delta) r n - \frac{q r}{K} n^2$$

$$d_n = \delta r n + \frac{(1 - q) r}{K} n^2$$

- 4 terms (2nd order in birth/death) so 4 total parameters
- note that $b_n > 0$ implies a maximum population size, N

Quasi-Steady State

Coexistence
and
Extinction of
Competing
Species

M.A.Badali

Introduction


Extinction

Fixation

Invasion



Maintenance

Discussion



Characterizing the quasi-stationary probability distribution function for varying δ and q . Lightness indicates an increased mean or variance in left and right respectively. Carrying capacity $K = 100$. The QSD has decreasing mean and increasing variance with increased δ or decreased q .

Mean Time to Extinction



Mean time to extinction for varying δ and q . Lightness of the line indicates an increase of q or δ in left and right respectively. Carrying capacity $K = 100$. The MTE decreases with increased δ or decreased q .

Mean Time to Extinction

Approximations

- larger fluctuations lead to shorter MTE: $\tau \approx \frac{1}{d_1 P_1}$

Mean Time to Extinction

Approximations

- larger fluctuations lead to shorter MTE: $\tau \approx \frac{1}{d_1 P_1}$
- $\hat{M} \vec{T} = -\vec{1}$ is equivalent to $\tau(n) = \sum_{i=1}^N \frac{1}{d_i} \prod_{k=1}^{i-1} \frac{b_k}{d_k} + \sum_{j=1}^{n-1} \prod_{l=1}^j \frac{d_l}{b_l} \sum_{i=j+1}^N \frac{1}{d_i} \prod_{k=1}^{i-1} \frac{b_k}{d_k}$

Mean Time to Extinction

Approximations

- larger fluctuations lead to shorter MTE: $\tau \approx \frac{1}{d_1 P_1}$
- $\hat{M} \vec{T} = -\vec{1}$ is equivalent to $\tau(n) = \sum_{i=1}^N \frac{1}{d_i} \prod_{k=1}^{i-1} \frac{b_k}{d_k} + \sum_{j=1}^{n-1} \prod_{l=1}^j \frac{d_l}{b_l} \sum_{i=j+1}^N \frac{1}{d_i} \prod_{k=1}^{i-1} \frac{b_k}{d_k}$
- Fokker-Planck equation $\partial_t P(x, t) = -\partial_x((b(x) - d(x))P(x, t)) + \frac{1}{2K} \partial_x^2((b(x) + d(x))P(x, t))$
 - Gaussian approximation[†] $p(n) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left\{-\frac{(n-n^*)^2}{2\sigma^2}\right\}$
with $\sigma^2 = \frac{-(b_n + d_n)|_{n=n^*}}{2\partial_n(b_n - d_n)|_{n=n^*}}$

Mean Time to Extinction

Approximations

- larger fluctuations lead to shorter MTE: $\tau \approx \frac{1}{d_1 P_1}$
- $\hat{M}\vec{T} = -\vec{1}$ is equivalent to $\tau(n) = \sum_{i=1}^N \frac{1}{d_i} \prod_{k=1}^{i-1} \frac{b_k}{d_k} + \sum_{j=1}^{n-1} \prod_{l=1}^j \frac{d_l}{b_l} \sum_{i=j+1}^N \frac{1}{d_i} \prod_{k=1}^{i-1} \frac{b_k}{d_k}$
- Fokker-Planck equation $\partial_t P(x, t) = -\partial_x((b(x) - d(x))P(x, t)) + \frac{1}{2K} \partial_x^2((b(x) + d(x))P(x, t))$
 - Gaussian approximation[†] $p(n) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left\{-\frac{(n-n^*)^2}{2\sigma^2}\right\}$
with $\sigma^2 = \frac{-(b_n + d_n)|_{n=n^*}}{2\partial_n(b_n - d_n)|_{n=n^*}}$
- WKB ansatz $P_n \propto \exp\left\{K \sum_i \frac{1}{K^i} S_i(n)\right\}$
with $S_0(n) = \int_{n=0}^K dn \ln\left(\frac{b_n}{d_n}\right)$ along extinction trajectory

[†]Gaussian approximation was written incorrectly in thesis.

Mean Time to Extinction

Approximations

Coexistence
and
Extinction of
Competing
Species

M.A.Badali

Introduction





Extinction

Fixation

Invasion

Maintenance

Discussion



Approximations of the MTE in various regimes of parameter space. WKB is good for low δ , is otherwise poor as FP.

Fixation

Coupled Logistic Equations

Coexistence
and
Extinction of
Competing
Species

M.A.Badali

Introduction


Extinction

Fixation

Invasion

Maintenance

Discussion



Each of the two species reproduces (arrows to self) and produces a toxin (arrows to limiting factors) which inhibits its own growth (square-ending lines to self) and the growth of the other (square-ending lines to other colour).

Coupled Logistic Equations

Coexistence
and
Extinction of
Competing
Species

M.A.Badali

Introduction


Extinction

Fixation

Invasion

Maintenance

Discussion



Each of the two species reproduces (arrows to self) and produces a toxin (arrows to limiting factors) which inhibits its own growth (square-ending lines to self) and the growth of the other (square-ending lines to other colour).

The deterministic coupled logistic equations are $\dot{x}_1 = r_1 x_1 \left(1 - \frac{x_1 + a_{12} x_2}{K_1} \right)$ and

$$\dot{x}_2 = r_2 x_2 \left(1 - \frac{a_{21} x_1 + x_2}{K_2} \right)$$

Coupled Logistic Equations

Coexistence
and
Extinction of
Competing
Species

M.A.Badali

Introduction

Extinction



Fixation

Invasion

Maintenance

Discussion

$$\dot{x}_1 = r_1 x_1 \left(1 - \frac{x_1 + a_{12} x_2}{K_1} \right) \text{ and } \dot{x}_2 = r_2 x_2 \left(1 - \frac{a_{21} x_1 + x_2}{K_2} \right)$$



$$O = (0, 0), A = (0, K_2), B = (K_1, 0), C = \left(\frac{K_1 - a_{12} K_2}{1 - a_{12} a_{21}}, \frac{K_2 - a_{21} K_1}{1 - a_{12} a_{21}} \right)$$

2,6 = parasitism/predation/antagonism, 3,7 = mutualism,

4,5 = competitive exclusion, 1 = (weak) competition


Transition to Neutrality

- Recall for niches $\tau \sim e^K$
- $a_{12} = a_{21} = 1$ limit recovers Moran results² $\tau \sim K$

²Lin, Kim, and Doering. *J. Stat. Phys.*, 2012.

Transition to Neutrality



- Recall for niches $\tau \sim e^K$
- $a_{12} = a_{21} = 1$ limit recovers Moran results² $\tau \sim K$



²Lin, Kim, and Doering. *J. Stat. Phys.*, 2012.

Transition to Neutrality



- Recall for niches $\tau \sim e^K$
- $a_{12} = a_{21} = 1$ limit recovers Moran results² $\tau \sim K$



²Lin, Kim, and Doering. *J. Stat. Phys.*, 2012.

Transition to Neutrality

- Recall for niches $\tau \sim e^K$
- $a_{12} = a_{21} = 1$ limit recovers Moran results² $\tau \sim K$




Effective coexistence except with complete niche overlap!

²Lin, Kim, and Doering. *J. Stat. Phys.*, 2012.

Route to Fixation

$$\text{Residence time } \langle t(s^0) \rangle_s = \int_0^\infty dt P(s, t | s^0, 0) = \hat{M}_{s, s^0}^{-1}$$



The system samples multiple trajectories on its way to fixation.

Left: Complete niche overlap limit, $a = 1$, for $K = 64$.

Right: Independent limit with $a = 0$ and $K = 32$.

Discussion

Coexistence and Extinction of Competing Species

M.A.Badali

Introduction

Extinction

Fixation

Invasion

Maintenance

Discussion

$f(a)$ (exponential dependence of MTE) approaches zero monotonically as niche overlap reaches Moran limit $a = 1$

- only for complete niche overlap will there be no exponential dependence: fixation will be rapid

Discussion

$f(a)$ (exponential dependence of MTE) approaches zero monotonically as niche overlap reaches Moran limit $a = 1$

- only for complete niche overlap will there be no exponential dependence: fixation will be rapid
- any niche mismatch allows for exponential dependence on K , which is typically large
 - any niche mismatch implies effective coexistence

Discussion

$f(a)$ (exponential dependence of MTE) approaches zero monotonically as niche overlap reaches Moran limit $a = 1$

- only for complete niche overlap will there be no exponential dependence: fixation will be rapid
- any niche mismatch allows for exponential dependence on K , which is typically large
 - any niche mismatch implies effective coexistence
- small departure from neutrality gives a niche theory

Invasion

Invasion

Coexistence
and
Extinction of
Competing
Species

M.A.Badali

Introduction

Extinction

Fixation

Invasion

Maintenance

Discussion

Invasion is going from one organism to half the population

Invasion

Coexistence
and
Extinction of
Competing
Species

M.A.Badali

Introduction

Extinction

Fixation

Invasion

Maintenance

Discussion

Invasion is going from one organism to half the population

- invasion is the other part of maintenance of biodiversity

Invasion

Coexistence
and
Extinction of
Competing
Species

M.A.Badali

Introduction

Extinction

Fixation

Invasion

Maintenance

Discussion

- Invasion is going from one organism to half the population
- invasion is the other part of maintenance of biodiversity
 - invasion with a fixed point should be fast (logarithmic)

Invasion

Invasion is going from one organism to half the population

- invasion is the other part of maintenance of biodiversity
- invasion with a fixed point should be fast (logarithmic)
- invasion on the line should be slower (linear)

Invasion

Invasion is going from one organism to half the population

- invasion is the other part of maintenance of biodiversity
- invasion with a fixed point should be fast (logarithmic)
- invasion on the line should be slower (linear)
- effects of a and K are not trivial

Invasion

Coexistence
and
Extinction of
Competing
Species

M.A.Badali

Introduction

Extinction

Fixation

Invasion

Maintenance

Discussion

Invasion is going from one organism to half the population

- invasion is the other part of maintenance of biodiversity
- invasion with a fixed point should be fast (logarithmic)
- invasion on the line should be slower (linear)
- effects of a and K are not trivial
- invasion attempts characterized by invasion probability E_s , successful invasion time τ_s , and failed invasion time τ_f

Invasion Probability

Coexistence
and
Extinction of
Competing
Species

M.A.Badali

Introduction



Extinction

Fixation

Invasion

Maintenance

Discussion



Probability of a successful invasion. **Left:** Numerical results, from $a = 0$ at the top to $a = 1$ at the bottom. The purple solid line is the expected analytical solution in the independent limit. The green solid line is the prediction of the Moran model in the complete niche overlap case. **Right:** The red data show the results for carrying capacity $K = 4$, and suggest the solid black line $\frac{b_{mut}}{b_{mut} + d_{mut}}$ is an appropriate small carrying capacity limit. Successive lines are at larger system size, and approach the solid magenta line of $1 - d_{mut}/b_{mut} \approx 1 - a$.

Invasion Times

Coexistence
and
Extinction of
Competing
Species

M.A.Badali

Introduction





Extinction

Fixation

Invasion

Maintenance

Discussion



Mean time of a successful or failed invasion attempt. Left: Mean time vs K . Right: Mean time vs a . Upper: Mean time conditioned on eventual invasion success. Lower: Mean time conditioned on failed attempt.

Maintenance

Moran Model with Immigration

Immigration comes from a constant reservoir of focal species fraction $g = n_{\text{reservoir}}/K_{\text{reservoir}}$ at a rate ν . Defining $f = n/K$, we have the following transition rates.

transition	function	value
$n \rightarrow n + 1$	$b(n)$	$f(1 - f)(1 - \nu) + \nu g(1 - f)$
$n \rightarrow n - 1$	$d(n)$	$f(1 - f)(1 - \nu) + \nu(1 - g)f$
$n \rightarrow n$	$1 - b - d$	$(f^2 + (1 - f)^2)(1 - \nu) + \nu(gf + (1 - g)(1 - f))$

Moran Model with Immigration

Immigration comes from a constant reservoir of focal species fraction $g = n_{\text{reservoir}}/K_{\text{reservoir}}$ at a rate ν . Defining $f = n/K$, we have the following transition rates.

transition	function	value
$n \rightarrow n + 1$	$b(n)$	$f(1 - f)(1 - \nu) + \nu g(1 - f)$
$n \rightarrow n - 1$	$d(n)$	$f(1 - f)(1 - \nu) + \nu(1 - g)f$
$n \rightarrow n$	$1 - b - d$	$(f^2 + (1 - f)^2)(1 - \nu) + \nu(gf + (1 - g)(1 - f))$

The crucial comparison is between $1/\nu$ and the invasion times previously described.

Steady State Results

Coexistence
and
Extinction of
Competing
Species

M.A.Badali

Introduction

Extinction



Fixation

Invasion

Maintenance

Discussion


$P(n)$ at steady state




PDF of stationary Moran process with immigration. Metapopulation focal fraction is $g = 0.4$, local system size $N = 100$, immigration rate ν is given by the colour. For high immigration rate the distribution should be centered near the metapopulation fraction $g N$ whereas for low immigration the system spends most of its time fixated.

First Passage Results

probability of extinction from n



mean conditional first passage time



Probability and conditional times of the focal species reaching temporary extinction before fixation, as a function of initial population. Metapopulation focal fraction is $g = 0.4$, local system size $N = 100$, immigration rate ν is given by the colour. The black line is the regular Moran result without immigration. When the immigrant is mostly not from the focal species ($g < 0.5$) immigration increases the likelihood of the focal species going extinct before fixating. Conditioned first passage times are longer when immigration is more frequent. Rare events take even longer still.

Discussion

- when immigration is uncommon
($N\nu < \min(1/g, 1/(1 - g))$), focal species either fixated or extinct most of the time
- when immigration is common
($N\nu > \max(1/g, 1/(1 - g))$), focal species is maintained at moderate abundance in the system, specifically gN

Discussion

- when immigration is uncommon ($N\nu < \min(1/g, 1/(1 - g))$), focal species either fixated or extinct most of the time
- when immigration is common ($N\nu > \max(1/g, 1/(1 - g))$), focal species is maintained at moderate abundance in the system, specifically gN
- immigration increases the times to (temporary) fixation or extinction

Discussion

Testable Systems

Coexistence
and
Extinction of
Competing
Species

M.A.Badali

Introduction

Extinction

Fixation

Invasion

Maintenance

Discussion

■ microfluidics

Testable Systems

Coexistence and Extinction of Competing Species

M.A.Badali

- microfluidics
- plasmids

Introduction

Extinction

Fixation

Invasion

Maintenance

Discussion

Testable Systems

Coexistence and Extinction of Competing Species

M.A.Badali

Introduction

Extinction

Fixation

Invasion

Maintenance

Discussion

- microfluidics
- plasmids
- mitochondria

Testable Systems

Coexistence
and
Extinction of
Competing
Species

M.A.Badali

Introduction

Extinction

Fixation

Invasion

Maintenance

Discussion

- microfluidics
- plasmids
- mitochondria
- nematode gut

Testable Systems

Coexistence
and
Extinction of
Competing
Species

M.A.Badali

Introduction

Extinction

Fixation


Invasion

Maintenance

Discussion

- microfluidics
- plasmids
- mitochondria
- nematode gut

Vega and Gore, *PLoS Biology*, 2017.



Conclusions

- higher commensurate birth and death rates (*i.e.* higher δ , lower q) leads to faster extinction;

Conclusions

- higher commensurate birth and death rates (*i.e.* higher δ , lower q) leads to faster extinction;
- WKB is fine for exponential scaling of the MTE, FP fails;

Conclusions

- higher commensurate birth and death rates (*i.e.* higher δ , lower q) leads to faster extinction;
- WKB is fine for exponential scaling of the MTE, FP fails;
- two species will effectively coexist unless they have exactly the same niche;
- similarly, greater niche overlap leads to longer invasion times, and less likelihood of success of an attempt;

Conclusions

- higher commensurate birth and death rates (*i.e.* higher δ , lower q) leads to faster extinction;
- WKB is fine for exponential scaling of the MTE, FP fails;
- two species will effectively coexist unless they have exactly the same niche;
- similarly, greater niche overlap leads to longer invasion times, and less likelihood of success of an attempt;
- in Moran model with immigration, a focal species at moderate size if $K\nu > 1/g$;

Conclusions

- higher commensurate birth and death rates (*i.e.* higher δ , lower q) leads to faster extinction;
- WKB is fine for exponential scaling of the MTE, FP fails;
- two species will effectively coexist unless they have exactly the same niche;
- similarly, greater niche overlap leads to longer invasion times, and less likelihood of success of an attempt;
- in Moran model with immigration, a focal species at moderate size if $K\nu > 1/g$;
- incomplete niche overlap is a niche theory with carrying capacities modified by niche overlaps;
- complete niche overlap (neutrality) on an island with immigration has abundance curve like mainland for species with $g_i > 1/K\nu$; other species are transients.

Next Steps

Coexistence and Extinction of Competing Species

M.A.Badali

Introduction

Extinction

Fixation

Invasion

Maintenance

Discussion

- predator-prey model (centre fixed point)

Next Steps

Coexistence and Extinction of Competing Species

M.A.Badali

Introduction

Extinction

Fixation

Invasion

Maintenance

Discussion

- predator-prey model (centre fixed point)
- rock-paper-scissors model (limit cycle)

Next Steps

Coexistence and Extinction of Competing Species

M.A.Badali

Introduction

Extinction

Fixation

Invasion

Maintenance

Discussion

- predator-prey model (centre fixed point)
- rock-paper-scissors model (limit cycle)
- other 3D models (chaos)

Next Steps

Coexistence and Extinction of Competing Species

M.A.Badali

Introduction

Extinction

Fixation

Invasion

Maintenance

Discussion

- predator-prey model (centre fixed point)
- rock-paper-scissors model (limit cycle)
- other 3D models (chaos)
- SIR model (epidemics)

Next Steps

Coexistence and Extinction of Competing Species

M.A.Badali

Introduction

Extinction

Fixation

Invasion

Maintenance

Discussion

- predator-prey model (centre fixed point)
- rock-paper-scissors model (limit cycle)
- other 3D models (chaos)
- SIR model (epidemics)
- evolving parameters (ecology and evolutionary biology)