



Master in Business Analytics & Data Science
Research Capstone in Operations Research

**The Price of Sophistication: A Pre-Registered
Falsification Test for When Spatial and Robust Modelling Pay
in Carbon-Aware Data-Center Scheduling**

A Day-Ahead Migration Scheduler and a Complexity–Value Decision Rule

Marco Ortiz Togashi

Supervisor: Prof. Bissan Ghaddar

June 2026

AI Use Declaration. Generative AI tools were used as a research and engineering assistant under the author’s direction: for code implementation support, experiment orchestration, literature triage, and drafting assistance. All research questions, modelling decisions, experimental designs, and interpretations are the author’s, developed with the supervisor; all results were produced by version-controlled code reviewed by the author (with the Phase 1–2 falsification protocol pre-registered; the Part 3 transfer and emergency-crossover sweeps are exploratory extensions, not part of that pre-registration), and all text was reviewed and edited by the author.

Ethics Statement. This research involves no human subjects, personal data, or animal experimentation, and required no ethics-board approval. It uses only grid-level carbon-intensity estimates (Electricity Maps, under an academic licence that prohibits redistribution of the raw series, respected here) and public reanalysis weather data (Open-Meteo, CC-BY). The work is motivated by an environmental aim, reducing the carbon footprint of flexible compute, and the author reports the negative and adverse results (the dependence-modelling null and the cases where robustness loses) as candidly as the positive ones, to avoid overstating the value of the proposed methods. The author declares no conflicts of interest.

Abstract

Data-center operators can shift compute toward cleaner hours and regions, and recent work applies distributionally robust optimization (DRO) to carbon-aware scheduling. The unexamined question is which modelling layers actually pay for their complexity. This thesis builds a working day-ahead carbon-aware migration scheduler and characterises the value of each layer. Over the honest baseline (carbon-aware scheduling with no inter-region transfer, $\Phi = 0$, on the same feasible set), active migration cuts daily $\text{CVaR}_{0.95}$ emissions by 4.0–9.9% (Western 4.0%, Eastern 9.9%, Diversified 9.0%), and this deterministic transfer channel is the dominant lever; the carbon-computing literature independently finds spatial shifting dominates temporal shifting. This thesis introduces a pre-registered, falsification-based methodology for *pricing* each modelling layer, separating the *validity* of the spatial-dependence assumption (the cross-region carbon correlations are real) from its *value* to the decision (a replicated null). We then isolate the value of *passive* sophistication. Using a Mahalanobis–Wasserstein DRO solved as a second-order cone program, a pre-registered shuffled-marginals falsification test across three real US/Canada grids spanning the dependence spectrum (the US West, an Ontario-anchored Eastern belt, and an engineered solar/wind/hydro portfolio), with an Iberia–France anchor, returns a replicated **null**. The encoded cross-region covariance adds below 0.4% of $\text{CVaR}_{0.95}$, and the null survives Ledoit–Wolf shrinkage, removing each region’s average daily pattern (residualization), Benjamini–Hochberg correction, walk-forward validation, and a per-cell equivalence test. A mean-leveling test shows the covariance signal is genuine but masked by the mean field (a mean-dominance bound), and a tail-dependence analysis shows the residual dependence is non-elliptical and upper-tail independent, hence invisible to covariance-based ambiguity sets; a one-paragraph copula appendix confirms even the maximal coupling recovers nothing. Finally we ask when robustness itself pays. Following the price of robustness of Bertsimas and Sim, a DRO over day-ahead forecast error buys a small worst-day ($\text{CVaR}_{0.95}$) tail reduction at a mean premium, with a value of the stochastic solution near zero, and pays over deterministic transfer only past an emergency-severity crossover $M^* \approx 3$ (with M the multiple by which a region’s carbon intensity spikes in a rare grid emergency, so $M = 3$ is a tripling); tested against real data-grounded worst-tail emergencies across 17 zones, that crossover does not activate. The thesis contributes a working scheduler with honest savings, a screening rule for when passive dependence modelling fails, and a price-of-robustness decision rule with a tested, data-grounded bound, all under 202 unit tests, with the Phase 1–2 falsification protocol pre-registered (the stylised-emergency crossover of RQ3 is a calibrated stress test, not part of the pre-registration).

Contents

Nomenclature	7
1 Introduction	8
1.1 Context	8
1.2 Problem statement and research questions	8
1.3 Contributions	9
1.4 Thesis structure	10
2 Literature Review	10
2.1 Carbon-aware computing	10
2.2 Distributionally robust optimization and the price of robustness	11
2.3 Modelling cross-region dependence	11
2.4 The gap: value has been assumed, not measured	12
3 Methodology	12
3.1 Research design overview	12
3.2 The scheduling problem	12
3.3 Distributionally robust formulation: day-ahead forecast-error hedging	13
3.4 A ladder of modelling layers	13
3.5 Feasible set and constraint regimes	14
3.6 Model validation on a small instance	15
3.7 The falsification test: shuffled marginals	16
3.8 Data	17
3.9 Validation, diagnostics, and the mean-dominance bound	17
3.10 Phase 2: richer dependence models (summary)	18
3.11 Reproducibility	20
4 Results	20
4.1 The spatial assumption is valid (RQ1)	20
4.2 The day-ahead scheduler and its savings (RQ1)	21
4.3 Where the value comes from: the complexity–value frontier (RQ1)	22
4.4 The screening rule: passive covariance adds nothing (RQ2)	22
4.5 The null survives a pre-registered robustness battery (RQ2)	24
4.6 Why the covariance is masked: the mean-leveling test (RQ2)	28
4.7 Tail dependence: the structure is non-elliptical (RQ2)	29
4.8 The null is not an artifact of the coherent objective (RQ2)	29

4.9	Richer dependence models confirm the screening rule (RQ2)	31
4.10	The price of robustness: when does the crossover pay? (RQ3)	31
5	Discussion	34
5.1	What the null means, and what it does not	34
5.2	Why the mean dominates: a mechanism	34
5.3	Relation to prior work	35
5.4	The decision rule: when does each layer pay?	35
5.5	Limitations	36
6	Conclusions	37
6.1	Answers to the research questions	37
6.2	Contributions	38
6.3	Future work	39
A	Reproducibility Details	41
B	Richer dependence models: confirming the screening rule	43
C	Extended outlook: online and multistage robust transfer	44
D	Glossary: plain-language terminology	47
	Annex A: Individual Contribution Statement	48

List of Figures

- 1 Full-model validation on a two-region instance. The complete feasible set (capacity, work conservation, ramp, deferral deadline) plus a shared aggregate hourly cap is solved jointly; stacked bars show how the two regions share each hour, the dashed line is the aggregate cap, the curves are each region’s carbon field, the green band the deadline window. **A:** baseline, the regions split the contested clean hours (4–5) under the cap. **B:** after dirtying region 1’s clean window, it cedes those hours to region 2 and spreads out. **C:** after tightening the shared cap, both regions spread further in time. Each response is the one an analyst anticipates, confirming the model encodes the intended cross-region behaviour. 15
- 2 The pre-registered falsification pipeline. Cross-validation and schedule fitting touch only the training years; the 2025 test set is read once, after the configuration is commit-locked, and the spatial gap is read off a paired day-bootstrap. 16
- 3 Residual (hour-of-day-removed) cross-region correlation of hourly carbon intensity, 2021–2025. Left: US West (CISO, BANC, LDWP, NEVP, AZPS). Right: Ontario–Eastern belt (CA-ON, NYISO, MISO, PJM). Cross-region correlation is strong and genuine in both, the assumption motivating spatial DRO is empirically valid. 21
- 4 The transfer lever. Out-of-sample CVaR_{0.95} saving against the inter-region transfer budget Φ , relative to the carbon-aware no-transfer baseline ($\Phi = 0$). Active migration captures this spatial lever; the curve saturates as the feasible mean-based gain is exhausted. The robust layer is analyzed in Section 4.10. 22
- 5 The complexity–value frontier. Cumulative out-of-sample emissions saved versus a carbon-blind scheduler, as each modelling layer is added. The value lives in deterministic transfer; passive dependence sophistication is flat; robust transfer is conditional on tail severity (Section 4.10). The transfer jump here is the same lever the headline states as a 4.0–9.9% reduction in tail-risk CVaR_{0.95} over the like-for-like $\Phi = 0$ baseline, a stricter metric than the cumulative-versus-blind axis shown. 23
- 6 The DRO genuinely engages. Blocked 5-fold cross-validation CVaR_{0.95} (relative to the nominal $\varepsilon = 0$ problem) against the Wasserstein radius ε , for the joint (solid) and shuffled (dashed) covariances. Cross-validation selects an interior ε^* (circled), not $\varepsilon = 0$, so the robustification is on; the two curves coincide, so the cross-region covariance changes nothing. Large ε is over-conservative. 24

7	The Phase 1 finding. A: spatial gap vs. mean residual cross-region correlation; value does not track correlation anywhere on the spectrum. B: mean-leveling test; in the covariance-only world (constant scheduling mean) the joint covariance pays off by up to +1.46%, proving the spatial signal is real but <i>masked</i> by the mean carbon field, not absent.	26
8	Robustness of the null. For each grid and each robustness arm (sample covariance, Ledoit–Wolf, seasonal and AR(1) residuals, walk-forward to 2024, and a 3× tighter ramp), dots are the spatial gap for the nine (regime, α) cells and the tick marks the median. The shaded band is the $\pm 0.1\%$ no-value threshold. No arm produces a sign-stable, material gap; the few Diversified excursions are negative (the joint covariance mildly hurts).	27
9	Upper-tail dependence $\chi_U(q)$ for the Ontario–Eastern belt region pairs (solid) against the Gaussian-copula benchmark at the matched correlation (dashed). The empirical curves sit at or below the benchmark as $q \rightarrow 1$: in the dirtiest hours the regions decouple, so there is no exploitable upper-tail co-movement for a richer ambiguity set to capture. The complementary radial asymmetry ($\chi_L > \chi_U$) is quantified in Table 6.	30
10	The price-of-robustness crossover. Robust-minus-risk-neutral CVaR _{0.95} advantage against synthetic single-region emergency severity M . The materially relevant crossover is $M^* \approx 3$ (grid-specific); the plotted curves are single representative seeds, and under four independent seeds the crossover is sign-stable only on the US West grid (+4.4% at $M = 4$). Realized joint severity stays at 1.3–1.9× and the robust commitment does not pay on real top-5% emergency days, so robustness does not activate on observed data. Rule: deploy deterministic transfer; robustify only if you expect $M > M^*$	33
11	The comonotone gap is scenario-sampling noise. Mean (line) and range over five seeds (band) of the comonotone-vs-independence CVaR _{0.95} gap on Eastern US–Canada, against the scenario count S . The band straddles zero and narrows with S ; the $\pm 0.1\%$ no-value band is shaded.	41
12	Transfer module validation. A: active transfer reduces out-of-sample CVaR _{0.95} by 4.0–9.9% over the $\Phi = 0$ baseline. B: the severity crossover, the robust (CVaR-hedged) commitment’s advantage over the risk-neutral one as grid-emergency severity M grows; it turns materially positive only past $M^* \approx 3$. The same numbers are reported and interpreted in the body (Section 4.10).	46

List of Tables

1	Spatial gap (% of test $\text{CVaR}_{0.95}$; positive = joint covariance better) for the three US/Canada cases. R3 = reference regime, R1 = +deadline, R2 = +variable carbon-coupled capacity. “det.” = bootstrap 95% CI excludes zero.	25
2	Spatial gap range (min, max over the nine cells, % of test CVaR_β) at three tail levels. The DRO is engaged ($\varepsilon^\star = 1$ almost everywhere) throughout. No level yields a sign-stable positive gap reaching the 0.4% materiality margin.	26
3	Statistical power of the spatial-gap test. SE is the paired-bootstrap standard error of the gap; $\text{MDE}_{80} = 2.80 \text{ SE}$ is the smallest true gap detectable at 80% power ($\alpha = 0.05$, two-sided). All are well below the $\approx 1\%$ materiality threshold.	28
4	Robustness battery: maximum $ \text{gap} $ (%) per covariance estimator, and temporal walk-forward (test year 2024). No cell passes the pre-registered agreement rule (positive and sign-stable under both seasonal and AR(1) residualization).	29
5	Mean-leveling test: spatial gap range (%) across the nine cells per case. LEVEL (equalize regional time-averages) leaves the null untouched; FLAT (constant scheduling mean = covariance-only world) reveals genuine spatial value except where correlation is common-mode (Western US).	29
6	Residual tail dependence at $q = 0.95$ for selected pairs. “Excess” = empirical χ_U minus the Gaussian-copula value at the matched correlation ρ . Throughout, χ_U shows no exploitable upper-tail excess while $\chi_L > \chi_U$ signals non-elliptical (radially asymmetric) dependence.	31
7	Risk-measure swap. Re-running the shuffled-marginals test under a non-coherent mean-variance objective, at the weight where the variance term balances the mean. The schedule moves materially, but the out-of-sample $\text{CVaR}_{0.95}$ gap (modelling joint covariance minus shuffling it; positive means covariance helps) stays immaterial and is adverse on two grids. Paired bootstrap, 1000 resamples.	31
8	Online robust vs. deterministic transfer, rolling 2025 evaluation with no injected emergency. Out-of-sample $\text{CVaR}_{0.95}$ reduction from robustifying (positive means robust is better); only the Diversified grid’s interval excludes zero, and there the robust controller is significantly worse.	32
9	Mapping of reported results to archived summary tables.	42
10	Copula gaps (% of test $\text{CVaR}_{0.95}$): out-of-sample improvement of each dependence model over the independence (shuffled) arm, median over the nine (regime, α) cells. No model reaches the 0.4% materiality margin.	44

Nomenclature

Symbols used throughout; plain-language glosses are in Appendix D.

Symbol	Meaning
R, T	number of regions; hours in the day-ahead horizon ($T = 24$)
$x \in \mathbb{R}^{R \times T}$	schedule (compute allocated per region and hour) [MW]
$y_{r,t}$	executed load after inter-region migration [MW]
W_r	day-ahead workload to be served by region r [MWh]
$\rho \in \mathbb{R}^{R \times T}$	uncertain carbon-intensity field [gCO ₂ eq/kWh]
$\bar{\rho}$	mean (forecast) carbon field; $\rho = \bar{\rho} + \xi$
ξ	zero-mean carbon residual (the dependence-bearing part)
Σ	cross-region space–time covariance of ρ ($RT \times RT$)
L	Cholesky factor, $LL^\top = \Sigma$
ε	Wasserstein ambiguity radius (DRO ball size)
η	scale-adaptive ridge factor for Σ
α_r	inflexible fraction of region r 's workload
Φ	inter-region transfer budget ($\Phi = 0$ is the no-transfer baseline)
$f_{r \rightarrow s,t}$	migrated load from region r to s at hour t [MW]
CVaR_β	Conditional Value-at-Risk at level β ($\beta = 0.95$)
τ	Value-at-Risk threshold in the CVaR program
$M, M^\star$	emergency severity multiple; the crossover severity
χ_L, χ_U	lower / upper tail-dependence coefficients
Λ	dependence leverage (comonotone minus diversified tail spread)
$\text{OPT}(\Sigma)$	optimal robust value of the SOCP under covariance Σ

1 Introduction

1.1 Context

Data centers consumed an estimated 1–2% of global electricity, and demand is growing rapidly with AI workloads [10]. Unlike most industrial loads, compute is unusually *flexible*: many batch and training workloads can be delayed by hours or moved between sites with little loss of utility. This movement is *logical*. Jobs relocate over the network between an operator’s data centers and do not require the regions’ power grids to be physically interconnected, so cross-region correlation of carbon intensity matters as information for the scheduler, not as an electrical constraint.¹ The carbon intensity of grid electricity varies strongly by hour and region with wind, solar, hydro availability, and demand, and this flexibility lets an operator act as a grid-aware participant that defers work from dirty hours to clean ones [20, 23]. Google’s carbon-intelligent computing platform demonstrated such temporal shifting at production scale [20], and recent systems work extends it to variable-capacity operation [24].

Scheduling against *tomorrow’s* carbon intensity, however, means scheduling under uncertainty. Distributionally robust optimization (DRO) addresses this by optimizing against the worst distribution in an ambiguity ball around the empirical distribution [8], and has been applied to carbon-aware scheduling with carbon intensity as the uncertain quantity [9]. Existing formulations treat each region’s carbon intensity essentially in isolation, yet the carbon intensities of electrically interconnected regions co-move: weather fronts traverse neighbouring balancing areas, and shared solar fleets create synchronized clean periods. We therefore model carbon intensity as a stochastic *vector* with a full cross-region covariance.

1.2 Problem statement and research questions

Each modelling layer (cross-region covariance, distributional robustness, richer copulas) adds estimation and computational cost and is worthwhile only if it buys measurably better schedules. The literature has assumed, rather than measured, that this sophistication pays. This thesis builds the scheduler and measures the value of each layer directly.

RQ1. How much can a day-ahead carbon-aware migration scheduler cut daily CVaR_{0.95} emissions relative to carbon-aware scheduling without inter-region transfer, and which lever drives the saving?

¹We use “spatial” throughout in the sense of *across the operator’s geographically separate regions* (sites), as in the standard spatial-versus-temporal load-shifting distinction. The dependence we study is the cross-region statistical correlation of carbon intensity, driven by shared weather and generation; it is not distance-decay spatial autocorrelation (e.g. Moran’s *I* or a variogram) and not an electrical coupling of the grids, which need not be interconnected.

RQ2. When does adding passive modelling complexity (cross-region covariance, or a non-elliptical copula) improve out-of-sample tail emissions, and at what cost?

RQ3. When does robustifying against day-ahead forecast error pay over deterministic transfer, and do real grids ever reach that regime? What decision rule follows?

The value question for RQ2 was committed as a falsifiable, pre-registered hypothesis test before the held-out year was touched (Section 3.7):

Pre-registered hypotheses (Phase 1–2, RQ2).

H_1 (modelling pays): encoding cross-region carbon dependence in the ambiguity set lowers out-of-sample $\text{CVaR}_{0.95}$ by more than a 0.4% materiality margin, relative to the dependence-destroyed (shuffled-marginals) model on the same feasible set.

H_0 (the spatial null): it does not; the out-of-sample spatial gap lies within $\pm 0.4\%$.

Decision rule: a per-cell equivalence test (TOST) that affirmatively rejects any effect beyond the margin, across the pre-registered grids and constraint regimes, supports H_0 ; a mean-leveling positive control must fire (detect value where value is constructed to exist) to license the null as a true negative rather than an underpowered test.

1.3 Contributions

The thesis makes five contributions: two pre-registered and confirmatory (the day-ahead savings, RQ1, and the dependence null, RQ2), one methodological, one exploratory (the robustness crossover, RQ3), and one infrastructural. (i) *A working day-ahead scheduler with honest, baseline-anchored savings:* active inter-region migration cuts out-of-sample $\text{CVaR}_{0.95}$ by 4.0–9.9% over the no-transfer ($\Phi = 0$) baseline on the same feasible set across three real grids (a gross spatial ceiling, reported before migration cost). The transfer mechanism follows established carbon-aware practice; the contribution is the $\Phi = 0$ -anchored measurement of its spatial value. (ii) *A pre-registered methodology for pricing modelling layers:* holding the optimization fixed and manipulating one object at a time, it measures the marginal out-of-sample CVaR value of each layer, and separates the *validity* of the spatial-dependence assumption (correlations are genuine, 0.7–0.8) from its *value* to the decision (null). To our knowledge this is the first treatment to place carbon intensity itself in a Mahalanobis–Wasserstein ball as a correlated cross-region vector and isolate the marginal value of that coupling via a shuffled-marginals falsification test. (iii) *A screening rule* for when passive dependence modelling cannot help, supported from four independent directions: an a-priori SOCP gap-bound (Proposition 1) before any data is touched; a mean-leveling positive control showing the covariance signal is genuine but masked by the mean field; a tail-dependence/comonotone-copula argument showing even the maximal coupling recovers nothing; and a risk-measure swap showing

the null survives replacing the coherent objective with a non-coherent mean-variance one (Section 4.8). (iv) A *price-of-robustness decision rule*, reported as an *exploratory* extension outside the pre-registration, with a data-grounded, falsified crossover: robustification pays only past an emergency-severity crossover $M^* \approx 3$ that realized severity (1.3–1.9 $\times$; Winter Storm Uri 1.3 $\times$) does not reach on observed data, stated as the rule plus its tested bound, not a universal impossibility. (v) A *reproducible, validated pipeline*: pre-registration, 202 unit tests, a small-instance model validation, a robustness battery, Benjamini–Hochberg correction, and per-cell TOST equivalence tests that affirmatively reject any effect above the 0.4% materiality margin.

1.4 Thesis structure

Section 2 reviews the carbon-aware scheduling and DRO literature and locates the gap: the value of modelling sophistication has been assumed, not measured. Section 3 develops the day-ahead migration scheduler, the DRO model, the falsification test, and the evaluation protocol. Section 4 reports the scheduler savings and the value of each modelling layer, the screening rule for passive dependence, and the price-of-robustness decision rule with the severity crossover. Section 5 interprets the findings and states limitations, and Section 6 concludes with the practitioner decision rule and future work. Appendix D gives a plain-language glossary of the main technical terms.

2 Literature Review

2.1 Carbon-aware computing

Carbon-aware scheduling exploits temporal and spatial variation in grid carbon intensity. In production at Google, Radovanović et al. [20] describe a system that computes day-ahead “virtual capacity curves” to throttle flexible compute during high-carbon hours; the measured impact on latency-sensitive workloads is negligible while load shifts measurably toward cleaner hours. Wiesner et al. [23] analyse the potential of temporal shifting across regions and workloads and find the benefit depends strongly on the grid’s marginal mix and the workload’s flexibility. Wijayawardana and Chien [24] study cloud-VM scheduling on datacenters whose capacity varies with grid conditions, including carbon-coupled capacity, where a constant carbon budget forces capacity to fall as intensity rises, and document the goodput cost of capacity variation for long-running jobs. This systems literature establishes operational levers (deferral, throttling, capacity coupling) that we adopt as constraints, but it treats future carbon intensity as a point forecast and handles uncertainty implicitly, by re-planning. That geographic migration itself lowers carbon, moving compute toward whichever region is cleaner, is established by this line of work [20, 23], and we take the

mechanism as given. What remains assumed rather than measured is whether *modelling* the joint spatial structure, or hedging forecast error robustly, adds anything on top.

2.2 Distributionally robust optimization and the price of robustness

DRO optimizes against the worst-case distribution within an ambiguity set, and the Wasserstein ball around the empirical distribution has emerged as the canonical data-driven choice, with tractable finite-dimensional reformulations [8, 18]. For costs linear in the uncertain vector, type-2 Wasserstein DRO with a Mahalanobis ground metric reduces to a mean term plus a covariance-weighted regularizer, an interpretable “mean plus $\varepsilon \sqrt{x^\top \Sigma x}$ ” objective solvable as a second-order cone program (SOCP) [2]. Hall et al. [9] apply Wasserstein DRO to carbon-aware data-center scheduling with virtual capacity curves, treating each region’s carbon intensity as the uncertain quantity and demonstrating robustness gains over sample-average approximation. The risk functional we target, CVaR_β , is standard for tail-focused evaluation [21]. Beyond second moments, recent work robustifies the *dependence structure* itself: Fan et al. [5] place both the marginals and the *copula* within a Wasserstein ball, the natural home for the non-elliptical structure our copula schedulers target.

Robustness is never free. Bertsimas and Sim [3] framed this as the *price of robustness*: protecting against worst-case realizations trades a guaranteed degradation in nominal performance for a bounded improvement in the tail, and the operator chooses where on that frontier to sit. This lens is central to our RQ3: we do not ask whether robustification *can* help (it can, in the tail) but whether the price it charges is paid back under conditions real grids actually reach.

2.3 Modelling cross-region dependence

Cross-region dependence enters a Mahalanobis–Wasserstein model only through the off-diagonal blocks of the covariance matrix. Estimating large covariance matrices from short panels is noisy; shrinkage estimators [19] are the standard remedy and serve here as a robustness arm. Beyond second moments, dependence is fully described by the copula [11, 22]; tail-dependence coefficients quantify joint extremes that correlation cannot see, and the Gaussian copula has zero asymptotic tail dependence for any correlation below one [7], a property we use as a diagnostic benchmark. Vine (pair-copula) constructions make high-dimensional non-elliptical dependence tractable [11, 12, 14], with established sequential structure-selection procedures [13], and copula-based scenario generation for stochastic programs is well developed [15]. Whether richer non-elliptical structure (vine copulas, or copula-ambiguity DRO [5]) improves a *stochastic program* depends on whether it changes the optimal decision out-of-sample; we test this directly in Appendix B.

2.4 The gap: value has been assumed, not measured

The carbon-aware scheduling literature establishes that *temporal* shifting (deferral to clean hours) delivers measurable savings, confirmed operationally at Google and in recent studies [20, 23]. The natural extension is *spatial* shifting (migration toward clean regions), and the distributed-computing literature finds spatial shifting dominates temporal shifting in potential [23], independently motivating the transfer channel studied here. Yet the value of *modelling* that spatial structure (via cross-region covariance, copulas, or distributional robustness) has been *assumed*, not measured. The DRO literature applies sophisticated ambiguity sets to uncertain vectors without isolating the marginal value of the cross-coordinate dependence they encode, and neither literature offers a controlled mechanism for *why* spatial structure should or should not matter. Adjacent work finds simple carbon-aware heuristics capture most attainable savings, with added sophistication offering little incremental benefit [4], and the role of dependence structure in Wasserstein DRO has been studied abstractly [6]; our wedge is the controlled, pre-registered, triangulated *measurement* of the dependence layer’s value. This thesis fills that gap: we build a working day-ahead scheduler that actively migrates compute and measure the savings (RQ1), isolate the value of passive modelling sophistication via a falsification test with a mechanistic explanation (RQ2), and frame robustification as a price-of-robustness trade-off, testing when that price is paid in real data (RQ3).

3 Methodology

3.1 Research design overview

The design tests whether passive (covariance-based) spatial modelling of carbon intensity adds value to robust day-ahead scheduling, and measures the value of active transfer separately. We hold the optimization machinery fixed and manipulate one object, the cross-region blocks of the covariance supplied to the scheduler, then measure the consequence on held-out tail emissions. Three primary region sets spanning the dependence spectrum (two earlier panels corroborating) serve as replications; three nested constraint regimes guard against a result driven by a particular feasible-set geometry; a robustness battery (estimators, multiple-testing correction, walk-forward) guards against statistical artifacts; and two further experiments (mean-leveling test; tail dependence) identify the mechanism. A pre-registration discipline removes analyst degrees of freedom.

3.2 The scheduling problem

Compute is scheduled across R regions over a $T = 24$ -hour day. The decision is $x \in \mathbb{R}_{\geq 0}^{R \times T}$, where $x_{r,t}$ is compute power (MW) in region r at hour t . Carbon intensity $\rho_{r,t}$ (gCO₂eq/kWh) is uncertain;

realized emissions are linear, $\langle \rho, x \rangle = \sum_{r,t} \rho_{r,t} x_{r,t}$.

3.3 Distributionally robust formulation: day-ahead forecast-error hedging

The robust layer hedges *day-ahead forecast error*. Each day the scheduler commits against a day-ahead point forecast $\bar{\rho}$ of the carbon field (the hour-of-day climatological mean on the training window, updated daily); the realized field departs by a residual $\xi = \rho - \bar{\rho}$ whose empirical distribution $\hat{\mathbb{P}}$, pooled over 2021–2024, defines the ambiguity. We minimize worst-case expected emissions over a type-2 Wasserstein ball $\mathcal{B}_\varepsilon(\hat{\mathbb{P}})$ of radius ε centred on that distribution, with Mahalanobis ground metric induced by the empirical residual covariance $\hat{\Sigma} \in \mathbb{R}^{RT \times RT}$:

$$\min_{x \in \mathcal{X}} \sup_{\mathbb{Q} \in \mathcal{B}_\varepsilon(\hat{\mathbb{P}})} \mathbb{E}_{\mathbb{Q}}[\langle \rho, x \rangle] = \min_{x \in \mathcal{X}} \langle \bar{\rho}, x \rangle + \varepsilon \sqrt{x^\top \hat{\Sigma} x}, \quad (1)$$

by Wasserstein duality for linear costs [2, 8]. With the Cholesky factorization $\hat{\Sigma} = LL^\top$ the penalty becomes $\varepsilon \|L^\top \text{vec}(x)\|_2$. Introducing an epigraph variable η (distinct from the hour index t), (1) is the second-order cone program

$$\min_{x \in \mathcal{X}, \eta \geq 0} \langle \bar{\rho}, x \rangle + \varepsilon \eta \quad \text{s.t.} \quad \|L^\top \text{vec}(x)\|_2 \leq \eta, \quad (2)$$

solved to global optimality (CLARABEL). The radius ε interpolates from the deterministic LP ($\varepsilon = 0$) to increasingly conservative schedules.

Where cross-region correlation enters. Partitioning $\hat{\Sigma}$ into $T \times T$ blocks Σ_{rs} by region pair, the quadratic form splits as

$$x^\top \hat{\Sigma} x = \sum_r x_r^\top \Sigma_{rr} x_r + \sum_{r \neq s} x_r^\top \Sigma_{rs} x_s, \quad (3)$$

so the off-diagonal (spatial) blocks are the *only* channel by which cross-region dependence can alter the schedule. The experiment manipulates exactly these blocks.

3.4 A ladder of modelling layers

The thesis studies schedulers arranged by sophistication: **(0)** carbon-blind; **(1)** per-region deterministic (each region by its own mean forecast); **(2)** per-region robust (single-region Wasserstein DRO, no cross-region structure); **(3)** joint covariance (the SOCP (1) with the full spatial Σ); and **(4)** joint copula (non-elliptical dependence; Appendix B). The screening test (Section 3.7) measures the marginal value of (3) over (2). Active transfer ($\Phi > 0$, inter-region flows) is orthogonal to this passive hierarchy and is the dominant lever, studied in Sections 4.2 and 4.10.

3.5 Feasible set and constraint regimes

The feasible set $\mathcal{X}$ collects the linear constraints below; introducing an auxiliary flexible-load variable $x^f \in \mathbb{R}_{\geq 0}^{R \times T}$, the scheduler solves (1) over

$$0 \leq x_{r,t} \leq \bar{x}_{r,t}, \quad \forall r, t \quad (\text{C0: capacity}) \quad (4a)$$

$$x_{r,t} = \alpha_r W_r p_{r,t} + x_{r,t}^f, \quad \forall r, t \quad (\text{C2a: flex/inflex split}) \quad (4b)$$

$$\sum_t x_{r,t}^f = (1 - \alpha_r) W_r, \quad \forall r \quad (\text{C2b: work conservation}) \quad (4c)$$

$$|x_{r,t} - x_{r,t-1}| \leq \Delta_r, \quad \forall r, t \quad (\text{C3: ramp}) \quad (4d)$$

$$\sum_{t \in \mathcal{W}} x_{r,t}^f \geq \gamma (1 - \alpha_r) W_r, \quad \forall r \quad (3a: \text{deferral deadline}) \quad (4e)$$

$$\pi_{r,t} x_{r,t} \leq \bar{P}, \quad \forall r, t \quad (3b: \text{thermal/PUE}) \quad (4f)$$

$$\sum_{r,t} \bar{\rho}_{r,t} x_{r,t} \leq B \quad (3d: \text{carbon budget}) \quad (4g)$$

where W_r is region r 's daily work (utilization fixed at 80%, $W_r = 960$ MWh); $\alpha_r \in \{0.30, 0.50, 0.75\}$ is the inflexible fraction following the fixed intraday shape $p_{r,t}$ ($\sum_t p_{r,t} = 1$); $\Delta_r = 15$ MW/h is the ramp limit; $\mathcal{W} = [0, 7]$ and $\gamma = 0.20$ set the deferral deadline; and $\pi_{r,t} = \text{PUE}_0 + \kappa \max(\mathcal{T}_{r,t} - \mathcal{T}_{\text{set}}, 0)$ is the temperature-coupled power-usage multiplier (data, hence a constant per cell), capped at effective power $\bar{P}$. Every constraint is linear, so $\mathcal{X}$ is polyhedral and the program (1) remains an SOCP. (We follow the source labelling: C0–C3 are the base operational constraints and the **3**· series is adapted from the SoCC'25 formulation [24]; a further aggregate hourly cap $\sum_r x_{r,t} \leq p_{\text{max}}$ is implemented but inactive in the reported regimes.)

The capacity ceiling is constant ($\bar{x}_{r,t} = 50$ MW) except under constraint **3c**, the carbon-coupled variable capacity adapted from Wijayawardana and Chien [24], which replaces it with

$$\bar{x}_{r,t} = x_{\min} + (x_{\max} - x_{\min}) \text{CFE}_{r,t}/100, \quad (5)$$

where $\text{CFE}_{r,t}$ is the training-mean carbon-free-energy share, capacity falls as the grid gets dirtier. The bounds $(x_{\min}, x_{\max}) = (50, 75)$ were calibrated on training data to a loosely binding (“Goldilocks”) regime: the constraint binds on 28–37% of region-hours, leaving slack and capacity margin so it shapes the schedule without freezing it.

Three nested regimes report results as a function of operational tightness: **R3** (reference) imposes C0 + C2 + C3; **R1** (lean) adds the deferral deadline 3a; and **R2** adds the thermal ceiling 3b (climate cases) or the variable-capacity ceiling 3c (interconnection cases). The carbon budget 3d is available but inactive in the reported regimes. Each regime is solved as a *complete* model with all its constraints active jointly; the nesting probes the null's robustness to operational tightness, not the effect of any single constraint in isolation.




Figure 1: Full-model validation on a two-region instance. The complete feasible set (capacity, work conservation, ramp, deferral deadline) plus a shared aggregate hourly cap is solved jointly; stacked bars show how the two regions share each hour, the dashed line is the aggregate cap, the curves are each region’s carbon field, the green band the deadline window. **A:** baseline, the regions split the contested clean hours (4–5) under the cap. **B:** after dirtying region 1’s clean window, it cedes those hours to region 2 and spreads out. **C:** after tightening the shared cap, both regions spread further in time. Each response is the one an analyst anticipates, confirming the model encodes the intended cross-region behaviour.

3.6 Model validation on a small instance

Before running the falsification experiment we check that the optimization model behaves correctly. We validate the full model *as a whole*, with all constraints of (4) active simultaneously, on a small human-checkable *two-region* instance (eight hours, 10 MWh per-cell capacity, $W = 30$ MWh each), rather than each constraint in isolation: correctness is a property of the constraints’ *interaction*, and a set that every constraint satisfies alone can still be jointly binding in unexpected ways. Two regions exercise the *cross-region* interaction, since a shared aggregate hourly cap $\sum_r x_{r,t} \leq P_{\text{agg}}$ couples the regions and forces them to compete for the same clean hours. We fix an analyst-anticipated outcome, perturb one input at a time, and verify the solved schedule moves the anticipated way (Figure 1). At baseline the two regions’ clean windows overlap, and the aggregate cap forces them to split the contested hours (panel A). Making region 1’s clean window dirty drives it to cede those hours to region 2 and spread to its own next-cleanest hours, with each region’s work still conserved (panel B). Tightening the shared cap removes joint headroom, so both regions spread further in time rather than co-locating in the clean window (panel C). All three responses match the hand-anticipated solution and conserve work per region to machine precision. This model-level check is distinct from the software test suite (Section 3.11, which pins component behaviour such as feasible-set equivalence) and from the falsification experiment below, which probes a scientific hypothesis rather than model correctness.




Figure 2: The pre-registered falsification pipeline. Cross-validation and schedule fitting touch only the training years; the 2025 test set is read once, after the configuration is commit-locked, and the spatial gap is read off a paired day-bootstrap.

3.7 The falsification test: shuffled marginals

The scheduler is fitted twice, with covariances differing in exactly one respect:

$$(\hat{\Sigma}^{\text{shuf}})_{rs} = \begin{cases} \Sigma_{rr}, & r = s, \\ \mathbf{0}, & r \neq s, \end{cases} \quad (6)$$

preserving every region’s marginal distribution and within-region autocorrelation while destroying only cross-region dependence. Each fitted schedule x is evaluated by the out-of-sample conditional value-at-risk of daily emissions $e_i = \langle \rho_i, x \rangle$ realized on test day i ; CVaR is a coherent risk measure [16] standard in stochastic programming [17]. We use the Rockafellar–Uryasev form [21]

$$\text{CVaR}_{\beta}(e) = \min_{\tau \in \mathbb{R}} \left\{ \tau + \frac{1}{1-\beta} \mathbb{E}[(e - \tau)_+] \right\}, \quad \beta = 0.95, \quad (7)$$

estimated as the empirical mean of the worst 5% of test days. The *spatial gap* is the difference between the two arms,

$$\text{gap} = \text{CVaR}_{0.95}(e^{\text{shuf}}) - \text{CVaR}_{0.95}(e^{\text{joint}}), \quad (8)$$

positive when the joint covariance helps. The test is a screening tool built to detect a spatial benefit if one exists; it separates the *value* of passive covariance modelling from the *validity* of cross-region correlation in the data, and checks the mean-dominance bound (Proposition 1). Figure 2 sketches the pipeline and Algorithm 1 gives the full protocol. The recipe is generic: hold the optimizer fixed, perturb exactly one modelling object, pre-register the analysis, and read off that object’s marginal out-of-sample tail value; only the perturbed object (here the cross-region covariance) changes when the same protocol is applied to another stochastic program.

Algorithm 1 Shuffled-marginals experiment (one region set)

- 1: **Input:** daily panels $\rho_{1:N}$ (train 2021–2024, test 2025); regimes $\{R3, R1, R2\}$; α -levels $\{0.30, 0.50, 0.75\}$; grid $\mathcal{E} = \{0, 0.1, 1, 10, 100, 1000\}$
 - 2: **for** each regime, each α , each arm $\Sigma \in \{\text{joint}, \text{shuf}\}$ **do**
 - 3: **for** each $\varepsilon \in \mathcal{E}$ **do** ▷ blocked 5-fold time-series CV
 - 4: **for** each contiguous fold k **do**
 - 5: fit $\bar{\rho}, L$ on training folds; solve SOCP (1); record validation $\text{CVaR}_{0.95}$
 - 6: $\varepsilon^* \leftarrow \arg \min_{\varepsilon} \text{mean validation CVaR}$ ▷ per arm, independently
 - 7: refit on full training set at ε^* ; evaluate *once* on 2025 ▷ single test read
 - 8: paired bootstrap (1000 resamples, fixed seed) on test days $\Rightarrow$ 95% CI for the gap (8)
 - 9: **Pre-registration:** configuration commit-locked before any test read; –dry-run gate runs CV only
-

3.8 Data

Carbon intensity is hourly lifecycle intensity from Electricity Maps (academic licence), 2021–2025, 43,824 hourly observations per zone; panels are built on a common local clock per region set, trained on 2021–2024 and tested once on 2025. Three primary US/Canada region sets span the dependence spectrum: **(1)** the *US West*, California, Nevada and Phoenix (CISO, BANC, LDWP, NEVP, AZPS); **(2)** an *Ontario–Eastern belt* (CA-ON, NYISO, MISO, PJM); and **(3)** an *engineered heterogeneous portfolio* (CISO (solar), ERCOT (wind), BPAT (hydro), spanning *different* interconnections) deliberately chosen so clean periods do *not* align: the adversarial best case for diversification. The spread is deliberate on both ends. The two correlated, within-interconnection grids are the *best case for the spatial hypothesis*, the most cross-region correlation a covariance model could exploit, and double as a stress test rather than a realistic deployment (no operator clusters data centers within one interconnection). A realistic, globally dispersed fleet looks instead like the heterogeneous portfolio, with low and mixed correlation, so the practically relevant case is the one where diversification *should* help most. The null holding at both ends is therefore stronger than a null on either alone. Two earlier panels corroborate: an *Iberia–France* European set (ES, PT, FR), where residual correlation collapses to the diurnal cycle, and the California/Nevada subset of the US West before Phoenix was added at the supervisor’s request. Temperature for constraint 3b is hourly ERA5 2 m air temperature (Open-Meteo), one load-weighted point per zone. The licensed raw data is never committed; derived summary tables are archived in the repository for traceability.

3.9 Validation, diagnostics, and the mean-dominance bound

Correlation validation (RQ1). Raw Pearson correlation conflates co-movement with a shared diurnal cycle. We therefore report correlations of *residuals* after removing each zone’s hour-of-day mean (local clock), with overlay plots of standardized intensity for summer and winter weeks.

Robustness battery. Every experiment is re-run with (i) Ledoit–Wolf shrinkage covariance [19]; (ii) covariance estimated from seasonal (monthly-mean) residuals; (iii) covariance from AR(1) residuals. A pre-registered agreement rule counts a spatial effect only if positive and sign-stable across both residual estimators. Bootstrap p -values across all cells receive Benjamini–Hochberg correction at $q = 0.05$ [1]; a walk-forward refit (train 2021–2023, test 2024) checks temporal stability; a log-denser ε grid checks grid sensitivity.

The mean-leveling test (RQ2, mechanism). To test whether the mean field masks the covariance, the mean used for *scheduling* is transformed while evaluation stays on real emissions: `LEVEL` equalizes regional time-averages (keeping hour-of-day shape); `FLAT` sets a constant mean, making the mean term constant under fixed demand, a covariance-only world in which any joint-vs-shuf difference is attributable purely to the spatial blocks.

Tail dependence (RQ2, structural). For each region pair we estimate the upper and lower tail-dependence coefficients $\chi_U(q) = \Pr(U > q, V > q)/(1 - q)$ and $\chi_L(p) = \Pr(U < p, V < p)/p$ on rank pseudo-observations of residual intensity, comparing χ_U at $q = 0.95$ with the Gaussian-copula value at the matched correlation. The Gaussian copula, the dependence family a covariance ball can represent, is asymptotically tail-independent [7], so a positive empirical excess would be exploitable tail structure invisible to the DRO; and since elliptical copulas force $\chi_L = \chi_U$, radial asymmetry diagnoses non-elliptical dependence.

3.10 Phase 2: richer dependence models (summary)

The covariance ball represents only *elliptical* dependence. To foreclose the “wrong object” objection, Appendix B re-runs the falsification with Gaussian, lower-tail Clayton, and the maximal (comonotone) copula, fitted to the empirical rank structure on the same feasible set. The result confirms the screening rule and is summarized in one paragraph there; Proposition 1 below explains why it must hold.

Why the dependence model is second-order. Two facts make the dependence model second-order to the mean field. The first is a decomposition. By the translation invariance of CVaR, writing the carbon field as mean plus zero-mean residual $\rho = \bar{\rho} + \xi$ and noting $\langle \bar{\rho}, x \rangle$ is deterministic,

$$\text{CVaR}_\beta(\langle \rho, x \rangle) = \underbrace{\langle \bar{\rho}, x \rangle}_{\text{dependence-independent}} + \text{CVaR}_\beta(\langle \xi, x \rangle), \quad (9)$$

so for a *fixed* schedule the dependence model moves only the residual term. The second is an a-priori bound on how far it can move the *optimal* value.

Proposition 1 (Spatial gap control, SOCP). *Let $\text{OPT}(\Sigma)$ be the optimal value of the SOCP (2) for a covariance Σ , and let Σ^{shuf} be its block-diagonal (cross-region-zeroed) counterpart with $\Sigma^{\text{off}} = \Sigma - \Sigma^{\text{shuf}}$. Then the spatial gap obeys*

$$|\text{OPT}(\Sigma) - \text{OPT}(\Sigma^{\text{shuf}})| \leq \varepsilon \left(\max_{x \in \mathcal{X}} \|x\|_2 \right) \|\Sigma^{\text{off}}\|_2^{1/2}, \quad (10)$$

which vanishes as $\varepsilon \rightarrow 0$ (no robustification) or $\Sigma^{\text{off}} \rightarrow 0$ (no cross-region structure).

Proof. The two programs share the feasible set $\mathcal{X}$ and differ only in the conic penalty. For fixed x , using $|\sqrt{a} - \sqrt{b}| \leq \sqrt{|a - b|}$ and the Rayleigh bound $|x^\top \Sigma^{\text{off}} x| \leq \|\Sigma^{\text{off}}\|_2 \|x\|_2^2$,

$$|\varepsilon \sqrt{x^\top \Sigma x} - \varepsilon \sqrt{x^\top \Sigma^{\text{shuf}} x}| \leq \varepsilon \sqrt{|x^\top \Sigma^{\text{off}} x|} \leq \varepsilon \|x\|_2 \|\Sigma^{\text{off}}\|_2^{1/2}.$$

Since the infimum is 1-Lipschitz in the sup-norm of the objective, two objectives that differ pointwise by at most δ over a common feasible set have optimal values differing by at most δ . The feasible set $\mathcal{X}$ is a bounded polytope (the capacity bounds C0), hence compact, so the maximum $\max_{x \in \mathcal{X}} \|x\|_2$ is attained. Its value follows from combining the capacity bound $x_{r,t} \leq \bar{x}$ (C0) with work conservation (C2b): $\|x\|_2^2 = \sum_{r,t} x_{r,t}^2 \leq \bar{x} \sum_{r,t} x_{r,t} = \bar{x} \sum_r W_r = \bar{x}^2 R T u$, so $\max_{x \in \mathcal{X}} \|x\|_2 \leq \bar{x} \sqrt{R T u}$; taking that maximum gives (10). $\square$

The bound is a conservative *certificate*, not a tight prediction. Evaluated on each grid’s training covariance (with $\varepsilon^* = 1$, feasible bound $\max_x \|x\|_2 = \bar{x} \sqrt{R T u}$ and $\|\Sigma^{\text{off}}\|_2$ the spectral norm of the destroyed cross-region blocks), its right-hand side is 7.4% (Eastern US–Canada), 9.6% (Diversified) and 12.7% (Western US) of $\text{CVaR}_{0.95}$; the realized gaps (Table 1) never exceed 0.23%, more than an order of magnitude smaller (a factor of 30–55 across the three grids). The certificate is evaluated at the base per-cell cap $\bar{x} = 50$; under the variable-capacity regime 3c, where the ceiling rises to $x_{\max} = 75$, the same work-conservation argument enlarges it by only $\sqrt{x_{\max}/\bar{x}} \approx 1.22$, still well over an order of magnitude above the realized gaps. The bound holds *before any experiment is run*; what it buys is not tightness but structure, since the gap is governed by ε and the cross-region block norm and forced to zero in the no-robustification and no-cross-structure limits. We keep the a-priori form deliberately (substituting the realized $\|x^*\|_2$ would tighten it but make it post-hoc). The *tight* ceiling is measured, not bounded: the comonotone arm of Section 4.9 realizes the supremum over copulas, and across scenario seeds its gain (the comonotone ceiling Λ) is centred on zero with $|\Lambda| \lesssim 0.1\%$ of $\text{CVaR}_{0.95}$; the mean-leveling test of Section 4.6 bounds the same quantity independently. Both put Λ an order of magnitude below the schedule’s mean-driven value, so the mean field pins the schedule and the dependence model, elliptical or not, is immaterial.

Computational considerations. Each instance is small: the decision vector has $RT \leq 120$ entries, so the SOCP (2) solves in 0.19–0.26 s and the $S = 1000$ CVaR-SAA LP (11) in 0.9–1.3 s (CLARABEL/HiGHS, commodity laptop). The cost is in the *number* of instances: the full Phase 1 battery is on the order of 10^4 SOCP solves and Phase 2 adds 108 CVaR-SAA LPs, yet the whole study runs in under two hours on one core.

3.11 Reproducibility

All code is version-controlled (private repository, continuous-integration test suite, 202 unit tests, including a feasible-set equivalence test pinning the Phase 2 CVaR scheduler to the Phase 1 constraint set); experiment configurations are commit-locked before test-set evaluation; bootstrap and scenario seeds are fixed; and every number reported in Sections 4–5 traces to an archived summary table. Solvers: CLARABEL for the SOCP; HiGHS for the CVaR-SAA linear programs and deterministic LP baselines.

4 Results

4.1 The spatial assumption is valid (RQ1)

The premise holds. After removing each zone’s hour-of-day mean, cross-region correlations of carbon intensity in the two electrically coupled US/Canada cases are strong and survive de-seasonalization: in the US West, CISO–NEVP 0.78 and CISO–LDWP 0.72 (shared solar resource and weather across the Western Interconnection); in the Ontario–Eastern belt, CA–ON–NYISO 0.73 and MISO–PJM 0.70 (weather fronts traversing the Eastern Interconnection). The engineered portfolio is the low, *heterogeneous* end rather than a zero-correlation grid: its residual pairs span 0.00 (ERCOT–BPAT), 0.29 (CISO–ERCOT) and 0.37 (CISO–BPAT), mean $\bar{r} \approx 0.2$. This heterogeneous correlation is *diversifiable* (it drives the largest mean-leveling signal, Section 4.6), unlike the US West’s strong but *uniform*, common-mode correlation. Figure 3 shows the residual correlation matrices; overlay plots of standardized intensity (archived with the code) confirm visible co-movement in summer and winter weeks for the coupled cases. With the corroborating California/Nevada subset (0.65–0.78 within-state) and Iberia–France, the cases span the dependence spectrum from ≈ 0 to ≈ 0.8 , so whatever the scheduling result, it cannot be attributed to an absence of correlation in the data.




Figure 3: Residual (hour-of-day-removed) cross-region correlation of hourly carbon intensity, 2021–2025. Left: US West (CISO, BANC, LDWP, NEVP, AZPS). Right: Ontario–Eastern belt (CA-ON, NYISO, MISO, PJM). Cross-region correlation is strong and genuine in both, the assumption motivating spatial DRO is empirically valid.

4.2 The day-ahead scheduler and its savings (RQ1)

The scheduler migrates compute across regions against a day-ahead forecast. We report out-of-sample $\text{CVaR}_{0.95}$ on the *same* feasible set against a like-for-like comparator: carbon-aware scheduling with no inter-region transfer ($\Phi = 0$), each region scheduled by its own forecast. The transfer arm adds flows $f_{r \rightarrow s, t} \geq 0$ under a budget Φ . Unlike the Phase 1–2 falsification test, this transfer sweep and the emergency-severity crossover of Section 4.10 are exploratory extensions reported outside the pre-registered protocol.

Figure 4 reports the saving against Φ . Active transfer cuts out-of-sample $\text{CVaR}_{0.95}$ by 4.0–9.9% relative to the $\Phi = 0$ baseline across the three grids (Western 4.0%, Eastern 9.9%, Diversified 9.0%). The lever is the spatial *mean*: at each hour one region is cleanest on average, and migration ships work there. The saving is captured without any dependence model, consistent with the carbon-computing literature, which finds spatial shifting dominates temporal shifting [20, 23]. The out-of-sample reduction is largest at a moderate budget (on the US West a 4.4% peak near 10% of daily workload) and settles to the reported 4.04% plateau as the budget grows: at large budgets the deterministic schedule reallocates further toward the training-mean-cleanest region and slightly overfits out of sample, so we quote the conservative plateau as the headline rather than the peak. One caveat fixes the right reading of this number: the saving is a *gross spatial ceiling*. The transfer budget is set large enough not to bind and migration carries no per-unit, latency, or bandwidth cost, so 4.0–9.9% bounds the value of *perfect, costless* migration rather than netting out real interconnect economics; introducing a migration cost can only reduce it (limitation (i), Section 5.5).




Figure 4: The transfer lever. Out-of-sample $\text{CVaR}_{0.95}$ saving against the inter-region transfer budget Φ , relative to the carbon-aware no-transfer baseline ($\Phi = 0$). Active migration captures this spatial lever; the curve saturates as the feasible mean-based gain is exhausted. The robust layer is analyzed in Section 4.10.

4.3 Where the value comes from: the complexity–value frontier (RQ1)

Figure 5 places every model class on a single complexity–value frontier (carbon-blind, deterministic per-region, robust per-region, deterministic transfer, robust transfer). Deterministic transfer dominates; the passive covariance and copula layers add no height (Sections 4.4–4.9) and robust transfer pays only in the tail (Section 4.10). Three forces, developed in Section 5.2, make the transfer saving mean-driven and keep passive covariance from adding to it: the magnitude of the diurnal mean swing, common-mode (non-diversifiable) correlation, and a decoupled dirty tail.

4.4 The screening rule: passive covariance adds nothing (RQ2)

Table 1 reports the spatial gap (8) for the three pre-registered US/Canada cases: nine cells each (three constraint regimes $\times$ three α -levels), CV-selected ε^* , single test read on 2025. The gap never exceeds a few tenths of one percent of test $\text{CVaR}_{0.95}$ in either direction. Where the bootstrap calls a cell “detectable” the effect is genuine but economically negligible (on the order of a few hundred $\text{gCO}_2\text{eq/kWh}$ out of a CVaR of 1.5×10^6) and, decisively, its *sign* flips across adjacent α -levels within the same regime: an exploitable spatial structure would not reverse direction under a change of the flexible-load fraction, so these reversals are the signature of optimization at the boundary of materiality, not of a usable effect. (We confirmed the detectable cells are not numerical: re-solving




Figure 5: The complexity–value frontier. Cumulative out-of-sample emissions saved versus a carbon-blind scheduler, as each modelling layer is added. The value lives in deterministic transfer; passive dependence sophistication is flat; robust transfer is conditional on tail severity (Section 4.10). The transfer jump here is the same lever the headline states as a 4.0–9.9% reduction in tail-risk $\text{CVaR}_{0.95}$ over the like-for-like $\Phi = 0$ baseline, a stricter metric than the cumulative-versus-blind axis shown.

with CLARABEL and SCS reproduces the gap to four significant figures, so the sign instability is in the data, not the solver.) The California/Nevada panel agrees (gaps in $[-0.012, +0.058]\%$, DRO active throughout).

This is a null of the *strong* kind. In all three pre-registered cases cross-validation *chooses* a non-trivial ambiguity radius: $\varepsilon^* = 1$ in all nine cells for both Eastern US–Canada and Diversified, and in seven of nine for Western US (25 of the 27 cells overall). The DRO is therefore genuinely engaged (it is paying for robustness, not collapsing to the nominal problem) and the two arms still coincide. Had CV selected $\varepsilon^* = 0$ the null would be uninformative, with the arms matching by construction; here the robustification is demonstrably on and the cross-region covariance still buys nothing, which is what makes the null informative rather than vacuous. Figure 6 shows this directly: validation $\text{CVaR}_{0.95}$ dips to an interior minimum at a non-trivial ε^* before over-conservatism sets in, and the joint and shuffled curves coincide.

The Iberia–France panel is the instructive contrast and the project’s historical starting point. There the residual correlation is near zero, and CV does not even consistently engage the DRO: ε^* flickers between 0, 0.1, 1 and 10 across cells, collapsing to 0 in several. This is consistent with the thesis (with no cross-region structure, the machinery sees nothing worth robustifying) but makes the Iberian null the *weak* kind. We retain it only as low-correlation, real-grid external validity,




Figure 6: The DRO genuinely engages. Blocked 5-fold cross-validation $\text{CVaR}_{0.95}$ (relative to the nominal $\varepsilon = 0$ problem) against the Wasserstein radius ε , for the joint (solid) and shuffled (dashed) covariances. Cross-validation selects an interior ε^* (circled), not $\varepsilon = 0$, so the robustification is on; the two curves coincide, so the cross-region covariance changes nothing. Large ε is over-conservative.

and rest the central claim on the US/Canada cases where the DRO is unambiguously on. The engineered Diversified portfolio supplies the DRO-active low-correlation null ($\varepsilon^* = 1$ throughout, still no value) that Iberia cannot.

Three readings of Table 1 matter. First, in the adversarial diversification case (Diversified) the joint covariance actually *hurts* at low α (up to -0.23%): with no true cross-structure, the joint estimator fits noise that the block-diagonal arm is spared. Second, in the strongly correlated belt (Eastern US–Canada) the detectable cells flip sign between $\alpha = 0.30$ and $\alpha = 0.75$ within the same regime. Third, the US West (strongest correlation of all) produces no detectable cell beyond one of magnitude $+3 \times 10^{-6}\%$. Figure 7A plots all 27 cells against each case’s mean residual correlation: the cloud stays on zero across the whole range. **Validity of the spatial assumption does not imply value from it.**

The null is visible in the schedules themselves. Overlaying the optimal load for the joint and shuffled covariances on each region’s mean carbon field, the two schedules are indistinguishable to the eye, both placing compute in the cleaner hours subject to the deadline and ramp constraints; destroying the cross-region covariance changes essentially nothing.

4.5 The null survives a pre-registered robustness battery (RQ2)

Table 4 summarizes. Re-estimating the covariance with Ledoit–Wolf shrinkage, seasonal residuals, or AR(1) residuals never produces a spatial effect that passes the pre-registered agreement rule (positive and sign-stable across both residual estimators). The largest battery value anywhere, $+0.179\%$ (AR(1), Eastern US–Canada, $\alpha = 0.30$), appears *only* under AR(1) and not under the seasonal estimator, so the rule rejects it. Benjamini–Hochberg correction at $q = 0.05$ across all

Table 1: Spatial gap (% of test $\text{CVaR}_{0.95}$; positive = joint covariance better) for the three US/Canada cases. R3 = reference regime, R1 = +deadline, R2 = +variable carbon-coupled capacity. “det.” = bootstrap 95% CI excludes zero.

Regime	α	Diversified ($\bar{r} \approx 0.2$)		Eastern US–Canada ($\bar{r} \approx 0.45$)		Western US ($\bar{r} \approx 0.65$)	
		gap %	det.	gap %	det.	gap %	det.
R3	0.30	−0.233	✓	−0.022	✓	+0.043	
R3	0.50	−0.177	✓	−0.002		+0.032	
R3	0.75	+0.001		+0.034	✓	−0.005	
R1	0.30	−0.233	✓	−0.022	✓	+0.014	
R1	0.50	−0.177	✓	−0.002		+0.028	
R1	0.75	+0.001		+0.034	✓	+0.002	
R2	0.30	−0.211	✓	+0.038	✓	+0.008	
R2	0.50	−0.154	✓	+0.030	✓	+0.005	
R2	0.75	+0.062	✓	+0.009		+0.000	✓

144 real-data cells (4 grid-runs $\times$ 9 regime- α cells $\times$ 4 covariance estimators) leaves that same already-rejected cell as the largest surviving positive gap; nothing material survives both filters. A walk-forward refit (train 2021–2023, test 2024) reproduces the null out of time: maximum absolute gap 0.036% (Eastern US–Canada), 0.028% (Western US), 0.217% (Diversified, again negative); an 11-point log-denser ε grid changes no gap at the reported precision. A *constraint-tightness* sensitivity probes the most plausible route to spatial value: tripling the ramp limit’s tightness ($\Delta_r = 5$ vs 15 MW/h) strips the schedule’s freedom to track the mean field, yet the null is unmoved (maximum absolute gap 0.028% Western US, 0.044% Eastern US–Canada, 0.169% Diversified; all sign-flipping, $\varepsilon^* = 1$ retained). The covariance does not begin to pay even when the mean’s pull is mechanically curtailed. A *utilization* sweep makes the same point through the workload: packing the facility from 80% to 95% strips almost all scheduling freedom, yet the gap only *shrinks* (0.017% Western US, 0.048% Eastern US–Canada, 0.071% Diversified), and at a loose 50% it stays immaterial ($\leq 0.25\%$); $\varepsilon^* = 1$ throughout the $\{50, 80, 95\}\%$ range. Figure 8 collects the covariance-estimator and temporal battery: across all six arms and three grids, every spatial gap sits inside the no-value band.

Tail level: the null holds from $\text{CVaR}_{0.90}$ to $\text{CVaR}_{0.99}$. The primary read fixes the tail at $\beta = 0.95$. Because the residual co-movement lives in the *clean* tail (Section 4.7), the natural worry is that spatial value might surface only at a deeper risk level. Re-running the full pipeline (cross-validated ε , single 2025 read) at $\beta \in \{0.90, 0.95, 0.99\}$ rejects this (Table 2). The DRO stays engaged ($\varepsilon^* = 1$ almost everywhere) and the gap stays immaterial at every level. Probing the deepest tail widens the magnitudes slightly but never to materiality: the largest *positive* gap anywhere is +0.14% (Western US, $\beta = 0.99$), still a third of the 0.4% margin and not sign-stable, while in




Figure 7: The Phase 1 finding. **A:** spatial gap vs. mean residual cross-region correlation; value does not track correlation anywhere on the spectrum. **B:** mean-leveling test; in the covariance-only world (constant scheduling mean) the joint covariance pays off by up to +1.46%, proving the spatial signal is real but *masked* by the mean carbon field, not absent.

the adversarial Diversified grid the joint covariance becomes *more* negative at $\beta = 0.99$ (−0.36%, i.e. it hurts), exactly as the noise-fitting account predicts. The $\beta = 0.99$ estimates are necessarily noisier (the tail thins to a handful of days): the paired-bootstrap standard error rises to 0.15% for Western US (vs $\leq 0.04\%$ at $\beta = 0.90$), so that grid’s largest point estimate (+0.14%) is within one standard error of zero. The apparent widening at the deepest tail is sampling noise. This sweep is descriptive, *outside* the pre-registered 144-cell false-discovery-rate (FDR) family of Table 4; no sweep cell is both positive and materially large (every detectable cell is negative or below the 0.4% margin), so none would survive correction as a discovery.

Table 2: Spatial gap range (min, max over the nine cells, % of test CVaR_β) at three tail levels. The DRO is engaged ($\varepsilon^* = 1$ almost everywhere) throughout. No level yields a sign-stable positive gap reaching the 0.4% materiality margin.

Grid	$\beta = 0.90$	$\beta = 0.95$	$\beta = 0.99$
Western US	[−0.04, +0.01]	[−0.01, +0.04]	[0.00, +0.14]
Eastern US–Canada	[−0.02, +0.04]	[−0.02, +0.04]	[−0.01, +0.09]
Diversified	[−0.23, +0.08]	[−0.23, +0.06]	[−0.36, +0.02]

The null is not a conditioning artifact. Block-diagonalizing Σ also changes its conditioning, so the null might reflect the joint arm being harder to estimate rather than the cross-structure being valueless. The two arms are conditioned to the same order (ridge-regularized $\kappa \approx 1.1\text{--}1.6 \times 10^5$ for the joint covariance, $0.4\text{--}0.8 \times 10^5$ for the shuffled, the joint arm somewhat higher as expected for the larger model). The mean-leveling test of Section 4.6 is the decisive control: it reuses the




Figure 8: Robustness of the null. For each grid and each robustness arm (sample covariance, Ledoit–Wolf, seasonal and AR(1) residuals, walk-forward to 2024, and a $3\times$ tighter ramp), dots are the spatial gap for the nine (regime, α) cells and the tick marks the median. The shaded band is the $\pm 0.1\%$ no-value threshold. No arm produces a sign-stable, material gap; the few Diversified excursions are negative (the joint covariance mildly hurts).

same joint Σ at the *same* conditioning and, removing only the mean field, the covariance recovers up to $+1.46\%$. Estimation conditioning therefore cannot be what suppresses the spatial value; the mean field is.

Statistical power: the null is not an underpowered test. A null is only as strong as the test’s ability to reject it, so we quantify that ability directly. The paired day-bootstrap gives a standard error on the spatial gap of $SE = 0.005\text{--}0.039\%$ of $CVaR_{0.95}$ across the three primary grids (Table 3). The minimum spatial benefit detectable at 80% power (two-sided, $\alpha = 0.05$) is $MDE = (z_{0.975} + z_{0.80}) SE = 2.80 SE = 0.015\text{--}0.11\%$. The test therefore had the power to detect a spatial advantage *an order of magnitude below* a $\approx 1\%$ materiality threshold (the smallest gap a practitioner would act on; by the economic grounding of Section 5.4, $\approx 0.4\%$ of $CVaR_{0.95}$), and roughly the same order as the per-cell gaps themselves, yet no sign-stable positive gap of even this size survives. The null is a genuine absence of usable spatial value, not a test too blunt to see it.

Equivalence: the absence of a material effect is itself significant. Power bounds the effect the test *could* have missed; an equivalence test makes the complementary, affirmative claim that a material effect is *absent*. We apply the two-one-sided-tests (TOST) procedure [25, 26] *per cell*, using each cell’s own paired-bootstrap standard error (recovered from its 95% gap interval), against the economic materiality margin $\Delta = 0.4\%$ of $CVaR_{0.95}$ (Section 5.4; the $\approx 1\%$ practitioner threshold is looser still), testing $H_0: |\text{gap}| \geq \Delta$ against $H_1: |\text{gap}| < \Delta$. All 27 cells across the three grids are equivalent: every 90% confidence interval lies strictly inside $(-\Delta, +\Delta)$. The binding cell is the adversarial Diversified case (gap = -0.233% , cell $SE = 0.046\%$), which still clears both

one-sided tests, $z = (\Delta - |\text{gap}|)/\text{SE} = 3.7$ ($p = 1.3 \times 10^{-4}$); every other cell gives $z \geq 7$. We therefore do not merely fail to detect a spatial benefit but statistically reject one larger than the materiality threshold, in either direction, cell by cell, the strongest form a null can take.

The claim is threshold-sensitive, so we state where it breaks: the smallest margin each grid still clears (its largest $|\text{gap}| + 1.645 \text{ SE}$) is $\Delta^* = 0.05\%$ (Eastern US–Canada) and 0.10% (Western US), but 0.31% for the adversarial Diversified grid. Equivalence at the 0.4% economic margin thus holds with room to spare on the two interconnected grids and only narrowly on the engineered portfolio. Equivalently, the conclusion holds for any materiality margin $\Delta \geq 0.31\%$; the 0.4% figure is fixed a priori from the economic grounding of Section 5.4, not reverse-engineered to clear the test. That caveat does not threaten the thesis: the Diversified gap is *negative*, so the grid where equivalence is hardest to assert is the one where the joint covariance *hurts* rather than adds value.

Table 3: Statistical power of the spatial-gap test. SE is the paired-bootstrap standard error of the gap; $\text{MDE}_{80} = 2.80 \text{ SE}$ is the smallest true gap detectable at 80% power ($\alpha = 0.05$, two-sided). All are well below the $\approx 1\%$ materiality threshold.

Grid	SE (% CVaR _{0.95})	MDE ₈₀ (%)
Western US	0.036	0.10
Eastern US–Canada	0.005	0.015
Diversified	0.039	0.11

4.6 Why the covariance is masked: the mean-leveling test (RQ2)

The mean-leveling test isolates the cause (Table 5, Figure 7B). Under the `LEVEL` setting (regional time-averages equalized, hour-of-day shape kept) every gap is unchanged to three decimals, so *between-region* mean differences are not the mask. Under the `FLAT` setting (constant scheduling mean, the covariance-only world) the joint covariance suddenly pays: up to $+1.46\%$ in Diversified and $+0.41\%$ in Eastern US–Canada, all BH-significant, and CV no longer collapses both arms onto the same schedule. The spatial signal in the covariance is therefore *real and exploitable*, but in the real problem it is dominated by the within-day shape of the mean carbon field, which both arms share. The exception is Western US: even in the covariance-only world its gaps stay at zero (-0.036 to $+0.004\%$): its strong correlation is *common-mode*, all five zones move together, so destroying cross-dependence changes no scheduling decision. Common-mode correlation is non-diversifiable risk, with no hedge for a DRO to find.

The flat-setting result doubles as the test’s *positive control*: in a world constructed so the dependence layer *should* pay, the same protocol fires (up to $+1.46\%$), whereas on the real data it reads zero. That contrast is what licenses the main null as a true negative, a genuine absence of value, rather than a test too blunt to detect it.

Table 4: Robustness battery: maximum $|\text{gap}|$ (%) per covariance estimator, and temporal walk-forward (test year 2024). No cell passes the pre-registered agreement rule (positive and sign-stable under both seasonal and AR(1) residualization).

Case	Ledoit–Wolf	Seasonal	AR(1)	Walk-forward 2024
Diversified	0.233	0.355	0.035	0.217
Eastern US–Canada	0.037	0.034	0.179	0.036
Western US	0.043	0.066	0.017	0.028

Table 5: Mean-leveling test: spatial gap range (%) across the nine cells per case. `LEVEL` (equalize regional time-averages) leaves the null untouched; `FLAT` (constant scheduling mean = covariance-only world) reveals genuine spatial value except where correlation is common-mode (Western US).

Case	Real problem	<code>LEVEL</code>	<code>FLAT</code> (covariance-only)
Diversified	$-0.233 \dots + 0.062$	unchanged	$+0.39 \dots + 1.46$
Eastern US–Canada	$-0.022 \dots + 0.038$	unchanged	$+0.14 \dots + 0.41$
Western US	$-0.005 \dots + 0.043$	unchanged	$-0.04 \dots + 0.00$

4.7 Tail dependence: the structure is non-elliptical (RQ2)

Table 6 reports residual tail-dependence at $q = 0.95$ for representative pairs. Two findings. *First*, the upper tail is empirically at or below the Gaussian benchmark everywhere (excess -0.18 to $+0.04$): in the dirtiest hours, regions *de-couple* rather than spike together, leaving no upper-tail co-movement to exploit on the risk side. *Second*, the dependence is radially asymmetric, $\chi_L > \chi_U$: regions go *clean* together more strongly than dirty together, clearest in the solar-driven US West (CISO–LDWP $\chi_L = 0.48$ vs. $\chi_U = 0.25$; BANC–LDWP 0.40 vs. 0.17). Elliptical copulas force $\chi_L = \chi_U$, so a Mahalanobis–Wasserstein ball cannot represent this structure: for the part of the dependence that varies, the covariance is not merely masked (Section 4.6) but the *wrong object*. Figure 9 shows the asymmetry for the Ontario–Eastern belt.

4.8 The null is not an artifact of the coherent objective (RQ2)

Proposition 1’s mean-dominance argument leans on two structural facts: emissions are linear in the carbon vector, and CVaR is translation-invariant (cash-additive), so the deterministic mean separates from the dependence-bearing residual. A natural objection is that the null is then a property of the chosen risk measure rather than of carbon scheduling. We test this directly. Holding the data, the feasible set, and the out-of-sample metric fixed, we swap the coherent mean-standard-deviation DRO objective ($\min \langle \bar{\rho}, x \rangle + \varepsilon \sqrt{x^\top \Sigma x}$, the thesis) for a *non-coherent* mean-variance objective ($\min \langle \bar{\rho}, x \rangle + \lambda x^\top \Sigma x$), which is not cash-additive and hands the cross-region covariance an explicit quadratic term to act on, and re-run the shuffled-marginals falsification, sweeping the covariance




Figure 9: Upper-tail dependence $\chi_U(q)$ for the Ontario–Eastern belt region pairs (solid) against the Gaussian-copula benchmark at the matched correlation (dashed). The empirical curves sit at or below the benchmark as $q \rightarrow 1$: in the dirtiest hours the regions decouple, so there is no exploitable upper-tail co-movement for a richer ambiguity set to capture. The complementary radial asymmetry ($\chi_L > \chi_U$) is quantified in Table 6.

weight λ over four orders of magnitude (`risk_measure_swap` snapshot).

The swap does not rescue covariance; it indicts it (Table 7). Weighting the variance term up to $900\times$ the mean moves the schedule by up to 30%, so covariance genuinely enters the decision, yet the out-of-sample $\text{CVaR}_{0.95}$ gap between joint and shuffled covariance stays below 1.5% and sign-unstable, as under the coherent DRO (whose gap stays below 0.8%). At the weighting where the variance term balances the mean, the paired bootstrap interval excludes zero, but in the *adverse* direction on two of three grids: modelling the joint covariance *worsens* out-of-sample tail emissions by 1.5% (Western) and 0.8% (Diversified), and helps only trivially (+0.11%) on the Eastern belt. Even out-of-sample standard deviation, the quantity mean-variance actually optimises, shows no robust covariance benefit, and in-sample variance reduction exceeds out-of-sample throughout, the signature of overfitting. The null is therefore not an artifact of CVaR’s translation invariance: handed a non-coherent objective built to exploit it, the estimated cross-region dependence overfits and degrades the held-out tail. This strengthens the screening rule rather than bounding it.

Table 6: Residual tail dependence at $q = 0.95$ for selected pairs. “Excess” = empirical χ_U minus the Gaussian-copula value at the matched correlation ρ . Throughout, χ_U shows no exploitable upper-tail excess while $\chi_L > \chi_U$ signals non-elliptical (radially asymmetric) dependence.

Case	Pair	ρ	χ_U emp	χ_U Gauss	Excess	χ_L emp
Western US	CISO–LDWP	0.72	0.25	0.41	−0.16	0.48
Western US	BANC–LDWP	0.65	0.17	0.35	−0.18	0.40
Western US	CISO–NEVP	0.78	0.43	0.47	−0.05	0.39
Eastern US–Canada	CA–ON–NYISO	0.73	0.38	0.42	−0.04	0.31
Eastern US–Canada	MISO–PJM	0.70	0.43	0.39	+0.04	0.42
Diversified	CISO–BPAT	0.33	0.10	0.16	−0.06	0.26
Diversified	ERCOT–BPAT	0.00	0.04	0.05	−0.02	0.02

Table 7: Risk-measure swap. Re-running the shuffled-marginals test under a non-coherent mean-variance objective, at the weight where the variance term balances the mean. The schedule moves materially, but the out-of-sample $\text{CVaR}_{0.95}$ gap (modelling joint covariance minus shuffling it; positive means covariance helps) stays immaterial and is adverse on two grids. Paired bootstrap, 1000 resamples.

Grid	Schedule move	OOS $\text{CVaR}_{0.95}$ gap	95% CI
Western US	~30%	−1.50%	[−1.92, −1.18]
Eastern US–Canada	~20%	+0.11%	[+0.04, +0.19]
Diversified	~7%	−0.84%	[−1.02, −0.63]

4.9 Richer dependence models confirm the screening rule (RQ2)

If the covariance ball failed only because it is elliptical, a copula that encodes the empirical $\chi_L > \chi_U$ asymmetry should recover value. It does not. Re-running the falsification with Gaussian, Clayton, and the maximal (comonotone) copula on the same feasible set leaves the screening rule intact: across scenario seeds even the maximal coupling’s gain is centred on zero at the sampling-noise floor ($|\Lambda| \lesssim 0.1\%$ of $\text{CVaR}_{0.95}$). The binding constraint is the mean field rather than the shape of the dependence object, as Proposition 1 predicts.

4.10 The price of robustness: when does the crossover pay? (RQ3)

Passive sophistication adds nothing under nominal conditions; the question the title poses is when *robustness* pays. We make the spatial structure an active decision (inter-region flows) and add a two-stage robust commitment with migration-limited recourse, hedging day-ahead forecast error. By Proposition 1’s logic the mean dominates under nominal carbon, and indeed robust $\approx$ deterministic there: the DRO buys no mean reduction (a price-of-robustness result, not an emissions win), and the value of the stochastic solution is near zero at best. In the rolling online evaluation (Table 8) the

Table 8: Online robust vs. deterministic transfer, rolling 2025 evaluation with no injected emergency. Out-of-sample $\text{CVaR}_{0.95}$ reduction from robustifying (positive means robust is better); only the Diversified grid’s interval excludes zero, and there the robust controller is significantly worse.

Grid	$\text{CVaR}_{0.95}$ change	95% CI	Verdict
Western US	+0.28%	$[-0.65, +1.16]$	null
Eastern US–Canada	+0.04%	$[-0.10, +0.25]$	null
Diversified	−10.34%	$[-18.3, -4.4]$	robust worse

robust controller’s out-of-sample $\text{CVaR}_{0.95}$ is statistically indistinguishable from the deterministic one on the US West (+0.28%, 95% CI $[-0.65, +1.16]$) and Eastern (+0.04%, $[-0.10, +0.25]$) grids, but is significantly *worse* on the Diversified grid (−10.34%, $[-18.3, -4.4]$, a single representative seed whose four-seed mean is a milder but still negative −4.4%): hedging forecast error when no emergency is present costs tail performance where the regions are weakly coupled. This sharpens the decision rule: under nominal conditions robustness is at best free and can actively hurt, so it is worth buying only when emergencies past the crossover are expected.

The null holds in the deep tail. A robust controller is a tail hedge, so $\text{CVaR}_{0.95}$ might understate its value for a more tail-averse operator. Re-evaluating the same online controllers at progressively deeper risk levels, from $\text{CVaR}_{0.90}$ through $\text{CVaR}_{0.99}$ to the single worst day (the `dro_tail_sensitivity` snapshot), the gap stays immaterial throughout: none of the twelve grid-by-metric cells shows a significant robust improvement, and on the Diversified grid robustness stays worse. The mean-dominance ceiling holds beyond the headline tail level; day-ahead robustness does not earn its keep even for a maximally tail-averse operator.

Robustness earns its keep only in the tail. Modelling rare grid-emergency events (with probability 0.1 per day a region’s carbon scales by a severity M), the robust commitment’s advantage over the risk-neutral one grows with M . Across four independent emergency-scenario seeds the crossover survives on the strongly correlated US West grid only: the gain is −0.4% at $M = 1$, +0.6% (all four seeds) at $M = 2$, and +4.4% of $\text{CVaR}_{0.95}$ on average (s.d. 3.4%, three of four seeds) at $M = 4$ (Figure 10). On the Eastern and Diversified grids the gain hovers near zero with unstable sign and does not survive multi-seed testing, so any single-seed positive there is scenario-seed noise. The crossover is therefore real but grid-specific and only marginally significant (US West, one-sided), with $M^* \approx 3$ the materially relevant threshold.

Real grids stay below the crossover. The synthetic multiplier overstates real risk, and a data-grounded test confirms the crossover does not activate. On the *joint* (portfolio) axis the crossover lives on, the worst realized day reaches only 1.3–1.9× the portfolio mean across the three grids




Figure 10: The price-of-robustness crossover. Robust-minus-risk-neutral $\text{CVaR}_{0.95}$ advantage against synthetic single-region emergency severity M . The materially relevant crossover is $M^* \approx 3$ (grid-specific); the plotted curves are single representative seeds, and under four independent seeds the crossover is sign-stable only on the US West grid (+4.4% at $M = 4$). Realized joint severity stays at 1.3–1.9 $\times$ and the robust commitment does not pay on real top-5% emergency days, so robustness does not activate on observed data. Rule: deploy deterministic transfer; robustify only if you expect $M > M^*$.

(Western 1.34 $\times$, Eastern 1.29 $\times$, Diversified 1.89 $\times$; archived in `carbon_ceiling.csv`), well below $M^* \approx 3$. Per-region *relative* swings can be larger on very clean grids (BPAT 5.1 $\times$, Ontario 2.6 $\times$), but those sit on near-zero carbon bases and do not move the portfolio tail; Winter Storm Uri, the worst US grid event on record, reached only 1.3 $\times$. Decisively, when emergencies are drawn *from real data* (each emergency replaces a region’s day with a draw from its own empirical top-5% tail; `run_part3_real_emergency.py`), the robust commitment does *not* pay: its $\text{CVaR}_{0.95}$ gain over the risk-neutral schedule is negative or within noise on every grid and frequency (Western -0.3% , Diversified -0.9% to -2.3% , Eastern $+0.1\%$ to $+0.2\%$). Real emergencies are not severe enough, in joint absolute-emission terms, to reach the crossover, so deterministic transfer is dominant on observed data. We claim no universal impossibility, only that robustness does not activate across the zones and real emergencies we have.

5 Discussion

5.1 What the null means, and what it does not

The result is a *specification* finding, not a failure of carbon-aware scheduling. Carbon intensity *is* spatially correlated where the literature expects it to be (Section 4.1), and the DRO machinery *is* active (CV selects $\varepsilon^* = 1$ almost everywhere). What the experiment falsifies is the specific hypothesis that encoding that correlation in a covariance-based ambiguity set improves robust scheduling. The mean-leveling test result (Section 4.6) is what licenses the stronger claim: the covariance signal is genuine and, in a covariance-only world, worth up to 1.46% of tail emissions; it is dominated by the diurnal mean carbon field in the problem as posed. A reader could otherwise dismiss the null as “the data had no signal”; the mean-leveling test shows the signal is present but subordinate. The central contribution is a clean separation of the *validity* of a modelling assumption from its *value*.

5.2 Why the mean dominates: a mechanism

Three forces compound. (i) *Magnitude*. The within-day swing of mean carbon intensity (night-to-noon ratios of two-to-fourfold in solar-heavy grids) dwarfs the residual covariance, so the optimal schedule is driven almost entirely by *when* each region is clean on average; the second-order covariance term moves decisions only at the margin. (ii) *Common-mode correlation*. Where correlation is strongest (US West), it is positive and shared across all zones, so the risk is common rather than diversifiable. Diversification value requires offsetting fluctuations; positively co-moving regions offer no hedge, so destroying the cross-blocks (the shuffled arm) changes nothing. (iii) *Wrong tail*. Robust tail scheduling cares about joint *dirty* excursions, but the empirical upper tail is close to independent (χ_U at or below Gaussian), while the co-movement that does exist is in the *clean* tail ($\chi_L > \chi_U$), which is the regime a risk-averse scheduler need not protect against and a radial asymmetry an elliptical ball cannot encode in any case. The covariance is therefore both masked (dominated by the mean) and, for its varying part, mis-shaped (non-elliptical). The same mechanism surfaces operationally: when these grids are scheduled under tightening geographic and ramp constraints, the binding dimension is geographic flexibility rather than temporal smoothing, again locating the value in *where* each region is clean on average rather than in second-order co-movement.

5.3 Relation to prior work

The finding refines rather than contradicts the carbon-aware DRO literature. Per-region Wasserstein DRO for carbon-aware scheduling, robustifying each region’s carbon intensity in isolation, is established by Hall et al. [9]; their multi-datacenter formulation does not model the joint cross-region carbon distribution. They report robustness gains over sample-average approximation; the present model is the natural *multi-region* extension that treats carbon intensity as a correlated vector. Two quantitative observations place our null in that context. First, the robustification *is* engaged: cross-validation selects a non-trivial ambiguity radius ($\epsilon^* = 1$, Figure 6) rather than collapsing to the nominal problem, consistent with the motivation in [9]. Second, holding that per-region (in-isolation) robustification fixed, the cross-region covariance adds nothing: the spatial gap is null throughout. In our out-of-sample 2025 test even the per-region DRO improves on the nominal (mean-greedy) schedule by at most 0.18% of $\text{CVaR}_{0.95}$ across grids, itself a reflection of the mean-dominance we document, and the spatial extension adds nothing on top. The contribution is to map where the value is *not*: neither robustification depth nor spatial coupling improves on a mean-greedy per-region schedule for this problem. This spares practitioners modelling complexity that does not pay and gives theorists a precise target, namely non-elliptical tail structure that a Wasserstein ball over covariance cannot reach.

5.4 The decision rule: when does each layer pay?

The findings compose into a practitioner rule. **Layer 1 (deploy always):** day-ahead carbon-aware per-region scheduling. This is the baseline; temporal shifting captures the within-region diurnal carbon cycle. **Layer 2 (deploy if migration bandwidth exists):** deterministic inter-region transfer, the dominant spatial lever (4.0–9.9% over the $\Phi = 0$ baseline; Section 4.2). This channel needs accurate per-region mean forecasts and transfer-capacity planning, not dependence estimation; the saving is driven by the spatial *mean* (at each hour one region is cleanest on average) and is therefore deterministic. **Layer 3 (skip):** cross-region covariance or copula modelling. Both the Mahalanobis covariance ball and the richer Clayton copula add below 0.4% of $\text{CVaR}_{0.95}$ and are mildly counterproductive in the diversification-friendly case (the joint estimator fits noise). The signal is genuine (mean-leveling test recovers up to +1.46% in a covariance-only world) but masked by the within-day mean carbon field (Proposition 1). **Layer 4 (conditional):** robustify the transfer against day-ahead forecast error only if grid-emergency severity realistically exceeds $M^* \approx 3$. The robust two-stage commitment buys a small CVaR tail reduction at a mean premium (value of the stochastic solution near zero), a textbook price-of-robustness result. Realized joint severity stayed at 1.3–1.9 $\times$ the portfolio mean (well below M^*), and on real top-5% emergency days the robust commitment did not pay, so this layer’s promised value remained unrealized.

In absolute terms, for the Eastern US–Canada facility (≈ 160 MW aggregate, $\approx 460,000$ tCO₂/yr), the transfer lever is worth on the order of \$1.7M per year at \$80/tCO₂, while the cross-region-covariance channel is about \$37,000 per year and not reliably positive. The value worth pursuing lives in the transfer decision and the mean forecast, not the second moment.

What the protocol buys, beyond this problem. The value-pricing protocol (fix the optimizer, perturb one modelling object, pre-register, measure out-of-sample tail value) transfers to any correlated-uncertainty stochastic program: it lets a practitioner decide *empirically* whether a dependence, robustness, or richer-ambiguity layer earns its estimation and compute cost. Two cheap checks come first and can save the experiment outright: the mean-dominance bound (Proposition 1) flags, before any data is touched, when the mean field is large enough that no covariance layer can move the schedule materially, and the positive-control check confirms the test can detect value when value exists. Together they turn “is this layer worth modelling?” from a matter of faith into a checkable, pre-registered measurement.

5.5 Limitations

Six bound the claims. (i) *Representative workload and costless migration.* We evaluate one canonical splittable workload with stylised, uniform per-region demand, not real operator traces, and the headline transfer saving is computed *gross* of migration economics: no per-unit, latency, SLA, or network-bandwidth cost, and an unbinding transfer budget. It is therefore an upper bound on the value of migration, not a net operational figure; a positive migration cost can only reduce it. On the modelling side, a natural worry is that the spatial null is an artifact of the chosen objective (cost linear in carbon, and a translation-invariant CVaR); Section 4.8 tests this directly by swapping in a non-coherent mean-variance objective and finds the null robust. The constraint-tightness sensitivities (Section 4.5) address the most plausible feasible-set variations: a 3 $\times$ tighter ramp and a 50–95% utilization sweep both leave the null intact. A qualitatively different objective (e.g. hard per-job deadlines) remains untested. (ii) *Test horizon and non-stationarity.* The primary read is a single held-out year (2025); walk-forward to 2024 reproduces the null, but multi-year rolling evaluation would strengthen external validity. Grid carbon structure is also non-stationary: continued renewable build-out could raise cross-region correlation (shared solar) or volatility, so the “real grids stay below the crossover” reading is bounded by the period observed. (iii) *Dependence model.* The DRO uses a covariance (second-moment) ambiguity set by construction; Phase 2 (Appendix B) extends the test to non-elliptical (Clayton) and maximal (comonotone) copulas, but a full copula-ambiguity Wasserstein DRO in the Fan–Ji–Lejeune sense remains future work. (iv) *Geographic scope and the limit of generalization.* The region sets span

the Western Interconnection, the Eastern Interconnection, and Iberia–France across the dependence spectrum, but grids with different generation mixes (e.g. hydro-thermal systems with strong seasonal storage) remain untested. The result generalizes only to grids where the diurnal mean field dominates residual cross-region covariance: mean-dominance is an empirical regularity of carbon intensity (Section 5.2), not a universal law, and a grid whose residual covariance rivalled its mean swing could behave differently. (v) *Data licence and clocks*. Carbon intensity is Electricity Maps lifecycle estimates on a common local clock per region set; alternative methodologies or clock conventions could shift point estimates, though not by the order of magnitude separating the observed gaps from materiality. (vi) *Forecast model*. The day-ahead point forecast is deliberately a transparent hour-of-day climatological mean updated daily, not a learned predictor: anchoring the robust layer to a simple, reproducible forecast keeps the price-of-robustness result a property of hedging residual error, not of any one forecaster. A learned forecaster would only shrink the residual variance the DRO hedges, and with it the already-immaterial robustness premium; quantifying that is future work.

6 Conclusions

6.1 Answers to the research questions

RQ1: How much can the day-ahead migration scheduler save, and which lever dominates?

Active inter-region migration cuts out-of-sample $\text{CVaR}_{0.95}$ emissions by 4.0–9.9% over the carbon-aware no-transfer baseline ($\Phi = 0$) across three real US/Canada grids (Section 4.2). The dominant lever is the spatial *mean*: at each hour one region is cleanest on average, and migration ships work there, making the saving deterministic and independent of any dependence model. This aligns with the carbon-computing literature, which finds spatial shifting dominates temporal shifting [20, 23].

RQ2: When does adding passive modelling complexity improve tail emissions? Never, on the grids we tested. Across the dependence spectrum (the US West to the engineered diversified portfolio) the out-of-sample $\text{CVaR}_{0.95}$ spatial gap stays below 0.4% and is not sign-stable. The null survives Ledoit–Wolf shrinkage, seasonal and AR(1) residualization, Benjamini–Hochberg correction over 144 cells, walk-forward validation, the full tail range from $\text{CVaR}_{0.90}$ to $\text{CVaR}_{0.99}$, and a per-cell equivalence test (Sections 4.4–4.5). The DRO is genuinely engaged (CV selects $\varepsilon^* = 1$ in nearly every cell), so this is an *active* null: the robustification is on and the cross-region covariance still adds nothing. A mean-leveling test proves the covariance signal is real (+1.46% in a covariance-only world) but *masked* by the within-day mean carbon field (Section 4.6). A tail-dependence analysis shows the residual dependence is non-elliptical, upper-tail-independent, and radially asymmetric ($\chi_L > \chi_U$), hence structurally invisible to a covariance-based ambiguity

set (Section 4.7). Replacing the covariance ball with a Clayton copula built for that asymmetry still recovers no material value (Appendix B). Proposition 1 (mean-dominance) explains why: the translation invariance of CVaR makes the dependence model second-order to the mean field. The screening rule is that passive sophistication cannot pay where the diurnal mean field dominates cross-region dependence, the regime that holds on every grid studied here.

RQ3: When does robustifying against forecast error pay over deterministic transfer, and do real grids reach that regime? The robust two-stage commitment with migration-limited recourse buys a small CVaR tail reduction at a mean premium (the value of the stochastic solution is near zero), a classic price-of-robustness result: robustness earns value only in the tail. When grid-emergency severity is modelled as a rare (0.1/day) multiplicative scaling M of carbon intensity, the robust schedule materially outperforms the risk-neutral one only past a crossover $M^* \approx 3$ (Section 4.10). On the joint (portfolio) severity axis the crossover lives on, the worst realized day reached only 1.3–1.9 $\times$ the portfolio mean across the three grids, well below M^* ; per-region relative swings reach 5 $\times$ (BPAT) on near-zero carbon bases but do not move the portfolio tail. Decisively, on real top-5% emergency days the robust commitment did not pay (its CVaR_{0.95} gain was negative or within noise on every grid), so the robust layer’s promised value is characterized but unrealized on observed data. We claim no universal carbon-ceiling impossibility, only the tested one: across the zones and real emergencies we have, deterministic transfer dominated. Robustness remains a conditional option for operators with a concrete, defensible emergency model.

6.2 Contributions

This thesis contributes to carbon-aware scheduling and robust optimization: (1) a working day-ahead migration scheduler saving 4.0–9.9% over the $\Phi = 0$ baseline, with a layered decision rule for each modelling layer; (2) a pre-registered falsification methodology (shuffled-marginals, extended to copula-coupled resampling) that separates the *validity* of a spatial modelling assumption from its *value*, transferable to any correlated-uncertainty DRO; (3) a replicated, robustness-checked null on whether spatial dependence improves carbon-aware DRO scheduling, holding across multiple real grids and the dependence hierarchy from covariance ball to elliptical and lower-tail copulas; (4) a mechanistic explanation (the mean-dominance bound Proposition 1, demonstrated by the mean-leveling test) for the null, which supplies the screening rule; and (5) a reproducible, version-controlled, CI-tested pipeline (202 tests) with archived license-safe summary tables for every reported number.

6.3 Future work

Phase 2 settled the immediate question, whether a copula that sees the asymmetry recovers value (it does not), and sharpened the remaining agenda. A *copula-ambiguity Wasserstein DRO* in the Fan–Ji–Lejeune sense [5], built on a fitted vine [12, 13], would robustify the dependence structure itself rather than sample from one fitted copula; Proposition 1 predicts the mean field will still dominate, but the claim would then be fully model-free. The body establishes the transfer savings and the severity crossover (Sections 4.2, 4.10); the open problem is to robustify the recourse policy with affine decision rules and calibrated emergency models. We are not aware of prior work that combines Wasserstein-DRO robustification with active inter-region transfer, and making that combination pay under realistic forecast error is an open question. The transfer model also invites an *online, multistage* treatment with forecasts arriving sequentially and a Wasserstein ball centred on a learned forecast, and the mean-dominance bound generalizes into a problem-class condition for *when* cross-region dependence can improve a robust allocation. The null is therefore not a dead end but a sharper target: value, if any, comes from an active spatial decision rather than a better passive dependence model.

References

- [1] Y. Benjamini and Y. Hochberg, “Controlling the false discovery rate: a practical and powerful approach to multiple testing,” *J. R. Stat. Soc. B*, vol. 57, no. 1, pp. 289–300, 1995.
- [2] J. Blanchet, Y. Kang, and K. Murthy, “Robust Wasserstein profile inference and applications to machine learning,” *J. Appl. Probab.*, vol. 56, no. 3, pp. 830–857, 2019.
- [3] D. Bertsimas and M. Sim, “The price of robustness,” *Operations Research*, vol. 52, no. 1, pp. 35–53, 2004.
- [4] R. Bostandoost, W. A. Hanafy, et al., “Data-driven algorithm selection for carbon-aware scheduling,” in *HotCarbon*, 2024.
- [5] Z. Fan, R. Ji, and M. A. Lejeune, “Distributionally robust portfolio optimization under marginal and copula ambiguity,” *J. Optim. Theory Appl.*, 2024.
- [6] R. Gao and A. J. Kleywegt, “Distributionally robust stochastic optimization with dependence structure,” arXiv:1701.04200, 2017.
- [7] P. Embrechts, A. McNeil, and D. Straumann, “Correlation and dependence in risk management: properties and pitfalls,” in *Risk Management: Value at Risk and Beyond*, Cambridge Univ. Press, 2002, pp. 176–223.

- [8] P. Mohajerin Esfahani and D. Kuhn, “Data-driven distributionally robust optimization using the Wasserstein metric,” *Math. Program.*, vol. 171, pp. 115–166, 2018.
- [9] S. Hall, F. Micheli, G. Belgioioso, A. Radovanović, and F. Dörfler, “Carbon-Aware Computing for Data Centers with Probabilistic Performance Guarantees,” arXiv:2410.21510, 2024.
- [10] International Energy Agency, *Electricity 2024: Analysis and Forecast to 2026*. Paris: IEA, 2024.
- [11] H. Joe, *Dependence Modeling with Copulas*. CRC Press, 2014.
- [12] K. Aas, C. Czado, A. Frigessi, and H. Bakken, “Pair-copula constructions of multiple dependence,” *Insurance Math. Econ.*, vol. 44, no. 2, pp. 182–198, 2009.
- [13] J. Dißmann, E. C. Brechmann, C. Czado, and D. Kurowicka, “Selecting and estimating regular vine copulae and application to financial returns,” *Comput. Statist. Data Anal.*, vol. 59, pp. 52–69, 2013.
- [14] C. Czado, *Analyzing Dependent Data with Vine Copulas*. Cham: Springer, 2019.
- [15] M. Kaut, “A copula-based heuristic for scenario generation,” *Comput. Manag. Sci.*, vol. 11, no. 4, pp. 503–516, 2014.
- [16] P. Artzner, F. Delbaen, J.-M. Eber, and D. Heath, “Coherent measures of risk,” *Math. Finance*, vol. 9, no. 3, pp. 203–228, 1999.
- [17] A. Shapiro, D. Dentcheva, and A. Ruszczyński, *Lectures on Stochastic Programming: Modeling and Theory*. Philadelphia: SIAM, 2009.
- [18] D. Kuhn, P. Mohajerin Esfahani, V. A. Nguyen, and S. Shafieezadeh-Abadeh, “Wasserstein distributionally robust optimization: theory and applications in machine learning,” in *Operations Research & Management Science in the Age of Analytics*, INFORMS, 2019, pp. 130–166.
- [19] O. Ledoit and M. Wolf, “A well-conditioned estimator for large-dimensional covariance matrices,” *J. Multivar. Anal.*, vol. 88, no. 2, pp. 365–411, 2004.
- [20] A. Radovanović et al., “Carbon-aware computing for datacenters,” *IEEE Trans. Power Syst.*, vol. 38, no. 2, pp. 1270–1280, 2022.
- [21] R. T. Rockafellar and S. Uryasev, “Optimization of conditional value-at-risk,” *J. Risk*, vol. 2, no. 3, pp. 21–41, 2000.
- [22] A. Sklar, “Fonctions de répartition à n dimensions et leurs marges,” *Publ. Inst. Statist. Univ. Paris*, vol. 8, pp. 229–231, 1959.
- [23] P. Wiesner, I. Behnke, D. Scheinert, K. Gontarska, and L. Thamsen, “Let’s wait awhile: how temporal workload shifting can reduce carbon emissions in the cloud,” in *Proc. 22nd ACM/IFIP Middleware*, 2021, pp. 260–272.

- [24] R. Wijayawardana and A. A. Chien, “Scheduling cloud VMs on variable capacity datacenters,” in *Proc. ACM Symp. Cloud Computing (SoCC)*, 2025.
- [25] D. J. Schuirmann, “A comparison of the two one-sided tests procedure and the power approach for assessing the equivalence of average bioavailability,” *J. Pharmacokinet. Biopharm.*, vol. 15, no. 6, pp. 657–680, 1987.
- [26] D. Lakens, A. M. Scheel, and P. M. Isager, “Equivalence testing for psychological research: a tutorial,” *Adv. Methods Pract. Psychol. Sci.*, vol. 1, no. 2, pp. 259–269, 2018.

A Reproducibility Details

Scenario-count stability and the comonotone noise floor. Figure 11 re-evaluates the comonotone (upper-Fréchet) gap on Eastern US–Canada over five scenario seeds and a range of scenario counts S . The gain is sampling noise centred on zero: at the production $S = 1000$ the spread across seeds is $[-0.08, +0.10]\%$ with mean $+0.01\%$, and it narrows with S . This is the basis for treating the single-seed $+0.182\%$ comonotone gap of Section 4.9 as immaterial rather than a real effect, and it confirms $S = 1000$ is sufficient.




Figure 11: The comonotone gap is scenario-sampling noise. Mean (line) and range over five seeds (band) of the comonotone-vs-independence $\text{CVaR}_{0.95}$ gap on Eastern US–Canada, against the scenario count S . The band straddles zero and narrows with S ; the $\pm 0.1\%$ no-value band is shaded.

Repository and environment. All code lives in a private, version-controlled repository with a continuous-integration suite (202 unit tests in total; CI runs 186, excluding the 13 Electricity-Maps integration tests that require licensed data). Experiments run under Python with CVXPY; solvers are CLARABEL for the second-order cone program and HiGHS for deterministic LP baselines. Exact dependency versions are pinned in a tracked `requirements-lock.txt` (CLARABEL 0.11.1, HiGHS/highspy 1.14.0, CVXPY 1.9.1, NumPy 2.4.6, SciPy 1.17.1; Python 3.12.4), under which every reported number reproduces byte-identically. The licensed raw carbon-intensity data is never committed; only derived aggregate summary tables (CV-selected ε , test CVaR, spatial gaps, bootstrap CIs, BH p -values) are archived, so the non-redistribution academic licence is respected while every reported number remains traceable.

Pre-registration protocol. Each experiment follows `commit-lock` $\rightarrow$ `dry-run` $\rightarrow$ `single test read`. The full configuration (utilization 0.80; $\alpha \in \{0.30, 0.50, 0.75\}$; ε grid $\{0, 0.1, 1, 10, 100, 1000\}$; 5-fold blocked time-series CV; 1000 bootstrap resamples at a fixed seed; CVaR level 0.95; train 2021–2024, test 2025) is committed before any test-set evaluation, and a `-dry-run` gate runs cross-validation only. The 3c capacity bounds $(x_{\min}, x_{\max}) = (50, 75)$ were Goldilocks-calibrated on training data alone (bind fraction 0.28–0.37).

Reproducing the experiments. Every result regenerates with

```
python -m scripts.run_case_experiment -region-set <case> [flags]
```

where `<case>` is one of the internal region-set keys `us_west` (Western US), `taskc` (Eastern US–Canada), or `us_hetero` (Diversified), and flags select the estimator (`-shrinkage`, `-residualize`), mean-leveling test (`-ablate-mean`), walk-forward (`-test-year`), or grid density (`-eps-grid`). Snapshot files are named `<case>_regimes_<date>[_<variant>].csv`, and the snapshot README maps each to its source command. Table 9 maps each results subsection to its archived table.

Table 9: Mapping of reported results to archived summary tables.

Section	Result	Archived source
4.4	Spatial gaps (Table 1)	<code>{case}_regimes_2026-06-10.csv</code>
4.5	Robustness (Table 4)	<code>{case}_...{lw,seasonal,ar1,ty2024}.csv</code>
4.5	BH correction	<code>bh_correction.csv</code>
4.6	Mean-leveling test (Table 5)	<code>{case}_...ablate-{level,flat}.csv</code>
4.1,4.7	CA/Iberia panels	<code>taskA... es_pt_fr...csv</code>
4.2	Transfer value, curve	<code>transfer_value_curve..., part3_transfer_value...</code>
4.10	Crossover, realized severity	<code>part3_{emergency,real_emergency}..., carbon_ceiling...</code>
4.10	Online robust (Table 8)	<code>part4_online..., dro_tail_sensitivity...</code>

B Richer dependence models: confirming the screening rule

This appendix gives the full protocol behind the body’s screening-rule check (Section 4.9). The covariance test answers whether *elliptical* dependence helps; because the tail diagnostic flags non-elliptical structure ($\chi_L > \chi_U$, Table 6), a natural objection to the Phase 1 null is “you used the wrong dependence object.” We foreclose it by replacing the covariance ball with copula models that *can* represent the structure, and re-running the same falsification.

Copula-coupled resampling. We generalize the shuffled-marginals test from second moments to the full copula while preserving every region’s empirical marginal. Each region r keeps its pool of whole daily profiles $\{\rho_r^d \in \mathbb{R}^T\}$; a scenario is generated by drawing a copula sample $u \in [0, 1]^R$ and, for each region, selecting the training day whose daily-mean intensity has rank u_r (low rank = clean day), then taking that day’s full profile. The copula thus governs *only* the cross-region coupling, exactly as the shuffled covariance did. Three nested fitted models span the dependence hierarchy, and a fourth, the comonotone upper bound, is added as the ceiling: **(i)** the *independence* copula (regions decoupled, the Phase 1 shuffled arm); **(ii)** the *Gaussian* copula at the empirical rank-correlation (elliptical, what the covariance ball can see); **(iii)** an exchangeable *Clayton* copula, with lower-tail dependence $\lambda_L = 2^{-1/\theta}$ and zero upper-tail dependence [11], fitted by matching mean pairwise Kendall’s τ ($\theta = 2\tau/(1 - \tau)$), the object built for the $\chi_L > \chi_U$ finding; and **(iv)** the *comonotone* (upper-Fréchet) copula, the maximal coupling. Clayton is sampled by Marshall–Olkin frailty; no parametric marginal model is introduced. The exchangeable Clayton imposes a single θ and so cannot reproduce the pair-level heterogeneity of the tail diagnostic; a regular-vine copula with edge-specific families would. We do not fit one, for a principled reason: the comonotone arm realizes the supremum over *all* copulas, vines included, so it upper-bounds whatever any heterogeneous-dependence model could achieve. If the maximal-coupling arm recovers no material value, no vine can.

Copula CVaR scheduler. For each model we draw $S = 1000$ scenarios $\{\rho^s\}$ and solve the sample-average CVaR program in Rockafellar–Uryasev form [21],

$$\min_{x \in X, \tau, z \geq 0} \tau + \frac{1}{(1 - \beta)S} \sum_{s=1}^S z_s \quad \text{s.t.} \quad z_s \geq \langle \rho^s, x \rangle - \tau, \quad (11)$$

over the *same* feasible set X (4) and regimes as Phase 1, so the only thing that varies across arms is the dependence model. Each schedule is evaluated once on the 2025 panel by out-of-sample $\text{CVaR}_{0.95}$; the reported copula gaps are $\text{CVaR}(\text{indep}) - \text{CVaR}(\text{Gaussian})$, $\text{CVaR}(\text{indep}) - \text{CVaR}(\text{Clayton})$, and the asymmetry increment $\text{CVaR}(\text{Gaussian}) - \text{CVaR}(\text{Clayton})$, each with a paired day-bootstrap

CI. Algorithm 2 summarizes the procedure.

Algorithm 2 Copula-scenario CVaR scheduling (one region set)

- 1: **Input:** daily panels $\rho_{1:N}$ (train 2021–2024, test 2025); dependence models {independence, Gaussian, Clayton, comonotone}; regimes; α -levels; scenarios $S = 1000$, fixed seed
 - 2: **for** each dependence model C **do**
 - 3: fit C on training daily summaries (rank pseudo-observations); for Clayton, $\theta = 2\tau/(1 - \tau)$ from mean Kendall τ
 - 4: **for** each regime, each α **do**
 - 5: draw S copula samples $u^s \in [0, 1]^R$ ▷ indep./Gaussian/Clayton frailty/comonotone
 - 6: map each region’s u_r^s to the training day of that daily-summary rank; take its full profile
 $\Rightarrow$ scenario $\rho^s \in \mathbb{R}^{R \times T}$
 - 7: solve the CVaR-SAA program (11) over $\mathcal{X} \Rightarrow$ schedule x
 - 8: evaluate *once* on 2025 $\Rightarrow$ out-of-sample $\text{CVaR}_{0.95}$
 - 9: copula gaps vs. independence + paired day-bootstrap CIs; comonotone realizes $\sup_C$, the ceiling Λ
-

Result. Table 10 reports the copula gaps. Across the three grids no dependence model recovers material value: the Gaussian and Clayton gaps over independence stay below 0.2% of $\text{CVaR}_{0.95}$, and the asymmetry increment (Clayton over Gaussian, the object built for the $\chi_L > \chi_U$ structure) is smaller still. The comonotone ceiling Λ is scenario-sampling noise centred on zero ($|\Lambda| \lesssim 0.1\%$; Appendix A, Figure 11). The screening rule survives the richest passive dependence model available, exactly as Proposition 1 predicts: the binding constraint is the mean field, not the shape of the dependence object.

Table 10: Copula gaps (% of test $\text{CVaR}_{0.95}$): out-of-sample improvement of each dependence model over the independence (shuffled) arm, median over the nine (regime, α) cells. No model reaches the 0.4% materiality margin.

Case	Gaussian	Clayton	Clayton–Gaussian
Western US	0.066	0.038	0.089
Eastern US–Canada	0.158	0.014	0.182
Diversified	0.069	0.127	0.114

C Extended outlook: online and multistage robust transfer

The day-ahead migration scheduler, its savings, and the severity crossover are reported in the graded body (Sections 4.2, 4.10). This appendix records implementation and validation detail

for the transfer module and sketches the planned online/multistage extensions that lie beyond the capstone proper.

This appendix sketches where the line goes next, with *preliminary, proof-of-concept* experiments. They are prototype-grade, a stylised grid-emergency stress model, modest scenario counts, no pre-registration, and a rigorous treatment is the subject of a follow-on study; we include them only because they already complete the arc the capstone opened. The capstone showed that *passive* spatial modelling adds no value because the mean dominates. The natural question is whether making the spatial structure an *active decision* changes that. We add inter-region load flows $f_{r \rightarrow s,t} \geq 0$ so the executed load $y_{r,t} = x_{r,t} + \sum_s (f_{s \rightarrow r,t} - f_{r \rightarrow s,t})$ can migrate toward the cleaner region; the program stays an SOCP/LP.

Three findings emerge (Figure 12). **(1) Active transfer unlocks spatial value.** A transfer-budget sweep cuts out-of-sample CVaR_{0.95} by 4.0–9.9% across the three grids, the value the passive null left on the table, captured by exploiting the spatial *mean* (shipping work to the region cleanest on average). **(2) Robustness is worthless under normal carbon, even here.** A one-shot transfer DRO, a forecast-error (persistence) ambiguity set, and a two-stage robust commitment with costly recourse all leave robust $\approx$ deterministic, the mean-dominance of the main text extends into the active setting. **(3) But robustness pays under tail risk.** Modelling rare grid-emergency events (with probability 0.1 per day a region’s carbon is scaled by a severity M , as in a renewable drought or peaker dispatch) and re-solving the two-stage problem with migration-limited recourse traces a sharp *crossover*: at $M = 1$ the robust commitment is no better than the risk-neutral one (the main null), but as M grows it increasingly wins, by up to +4.4% of CVaR_{0.95} (four-seed mean, s.d. 3.4%) at $M = 4$ on the US West, the only grid where the crossover survives multi-seed testing. The robust commitment stays hedged while the risk-neutral one is burned when its favoured region spikes.

The synthesis is a clean phase diagram: distributional robustness is immaterial for carbon-aware scheduling under normal conditions (the mean dominates, whether the spatial structure is modelled passively or acted on actively), but becomes valuable precisely when rare tail events make the mean forecast unreliable. The value of robustness is thus a function of tail risk, the precise characterisation of which, together with a calibrated emergency model and a pre-registered evaluation, is the agenda for the follow-on work.

Implementation and validation. These results are not throwaway script output: the transfer model is implemented as a standalone, unit-tested module (`src/models/transfer_dro.py`), where the one-shot transfer DRO, the two-stage robust commitment, and the costly-recourse evaluation are each exercised by tests pinning the model’s invariants, conservation of work under inter-region flows ($\sum_{r,t} y_{r,t} = \sum_r W_r$), monotone emission reduction as the transfer budget grows, and exact recovery of the no-transfer schedule at zero budget. The flow variables $f_{r \rightarrow s,t}$ carry no




Figure 12: Transfer module validation. **A:** active transfer reduces out-of-sample $\text{CVaR}_{0.95}$ by 4.0–9.9% over the $\Phi = 0$ baseline. **B:** the severity crossover, the robust (CVaR-hedged) commitment’s advantage over the risk-neutral one as grid-emergency severity M grows; it turns materially positive only past $M^* \approx 3$. The same numbers are reported and interpreted in the body (Section 4.10).

self-loops and respect a transfer-capacity budget, and the two-stage program is solved as an LP per scenario with a shared CVaR epigraph. What remains *preliminary* is therefore the experimental protocol, the stylised emergency model, the modest scenario counts, and the absence of pre-registration, not the optimization itself: the crossover of Figure 12 rests on validated code. This is what makes promoting Part 3 to a rigorous study a matter of protocol and calibration, the steps below, rather than of rebuilding the machinery.

Directions beyond this capstone. Two extensions follow naturally and are left as future work. First, taking the transfer model *online*: a rolling-horizon robust controller in which forecasts arrive sequentially and demand and transmission are themselves uncertain, with a Wasserstein ball centred on a learned forecast and evaluated closed-loop on real operational traces. Second, *generalising* the mean-dominance bound into a problem-class condition for when cross-coordinate dependence can improve a robust allocation, of which carbon-aware scheduling is one instance among others (energy storage, EV fleets, spatial supply chains). Promoting the present Part 3 sketch to a rigorous, pre-registered study with a calibrated emergency model is the prerequisite for both. These directions are recorded in the project’s research roadmap.

D Glossary: plain-language terminology

The main technical terms used in this thesis, glossed in plain language; the precise definitions are in Section 3.

Carbon intensity: how dirty the electricity is at a given hour and place (grams of CO₂ per kWh); it swings with how much wind, solar, and hydro are on the grid.

Distributionally robust optimization (DRO): planning against the worst-case carbon distribution within a set of plausible distributions around the data, rather than trusting a single point forecast. Like packing for weather a little worse than the forecast.

Wasserstein distance / ball: a way to measure how far apart two probability distributions are (the cheapest cost of reshaping one into the other). The Wasserstein *ball* of radius ε is the set of all distributions within that distance of the data, every future close enough to what we observed to be worth hedging.

CVaR_{0.95} (Conditional Value-at-Risk): the average emissions on the worst 5% of days; a tail-risk measure, not the average day.

Second-order cone program (SOCP): the convex problem the scheduler reduces to, solved to global optimality in a fraction of a second.

Mahalanobis distance: a distance that weights directions by the carbon covariance, so it accounts for how regions move together. Like measuring distance on a stretched map where correlated regions sit closer.

Copula: the part of a joint distribution that captures only how regions move together, with each region's own behaviour stripped out.

Comonotone: the strongest possible copula, all regions rising and falling in perfect lockstep; used as a worst-case upper bound.

Tail dependence (χ_U, χ_L): whether regions hit their dirtiest (χ_U) or cleanest (χ_L) hours at the same time, a joint-extreme effect plain correlation can miss.

Residualization: removing each region's predictable hour-of-day (or day-to-day) average so the analysis works only on the genuine surprises, not the routine daily cycle.

Shuffled-marginals test: re-fitting the scheduler with the cross-region link destroyed but each region's own pattern intact, so the difference isolates the value of cross-region correlation.

Mean-leveling test: we deliberately flatten each region's average carbon level (so the only signal left is the cross-region co-movement) and re-run the scheduler. It shows whether that cross-

region structure is genuinely useful but normally hidden behind the much larger differences in average carbon.

Mean-dominance (Proposition 1): an a-priori bound showing the average carbon field caps how much any dependence model can change the schedule.

Out-of-sample / walk-forward: scoring a plan only on data it never trained on (out-of-sample); walk-forward is the rolling version, train on earlier years and test on a later untouched one, mimicking real deployment.

Equivalence test (TOST): a test that affirmatively shows an effect is smaller than a chosen materiality margin, turning “we found nothing” into “any effect this large is ruled out.”

Severity M : the multiple by which one region’s carbon spikes in a rare grid emergency ($M = 3$ is a tripling); the crossover M^* is where robustness starts to pay.

Value of the stochastic solution (VSS): how much planning against the full uncertainty distribution beats committing to a single point forecast; here it is near zero.

Transfer budget (Φ): how much compute may migrate between regions; $\Phi = 0$ is the honest no-transfer baseline, and raising it delivers the headline savings.

Annex A: Individual Contribution Statement

This Research Capstone was completed individually by **Marco Ortiz Togashi** under the supervision of Prof. Bissan Ghaddar. As sole author, I am responsible for all components of the work:

- **Research design.** Formulated the three research questions, designed the shuffled-marginals falsification test, and specified the pre-registration protocol that separates assumption validity from value.
- **Modelling.** Specified the Mahalanobis–Wasserstein DRO scheduling model and its SOCP reformulation, the constraint regimes (including the variable carbon-coupled capacity constraint adapted from the SoCC’25 formulation), and the mean-leveling test and tail-dependence diagnostics.
- **Data and implementation.** Assembled the multi-grid carbon-intensity and temperature panels, implemented the full experimental pipeline, the robustness battery, the Benjamini–Hochberg correction, and the figures, under version control with a continuous-integration test suite.
- **Analysis and writing.** Produced and interpreted all results, established the mechanistic explanation, and wrote the manuscript in full.

Generative AI tools were used as a coding and drafting assistant under my direction, as declared on the cover page; all research questions, modelling decisions, experimental designs, interpretations, and final text are my own. The inter-region migration mechanism itself follows established carbon-aware practice (Section 2); my individual contribution, developed under the supervisor's direction, is its distributionally robust treatment, the honest $\Phi = 0$ -anchored measurement of its spatial value, and the screening and price-of-robustness decision rules.

Marco Ortiz Togashi

Date