

A cultural atlas of vocal variation: yellow-naped amazons exhibit contact call dialects throughout their Mesoamerican range

Molly K. Genes,¹ Marcelo Araya-Salas,² Christine R. Dahlin,³ Timothy F. Wright^{1*}

¹Department of Biology, New Mexico State University, Las Cruces, New Mexico, USA

²Centro de Investigación en Neurociencias & Escuela de Biología, Universidad de Costa Rica, San Pedro, San José, Costa Rica

³Department of Biology, University of Pittsburgh at Johnstown, Johnstown, Pennsylvania, USA

*Correspondence:

Timothy Wright
wright@nmsu.edu

Keywords: contact call, cultural atlas, geographic variation, vocal dialect, vocal learning, yellow-naped amazon.

Abstract

Vocal dialects are a taxonomically widespread phenomenon which are typically only studied in a portion of a species' range. Thus, it is difficult to infer whether a geographic pattern of vocal dialects observed in one part of a species' range are typical across the range or whether local conditions influence their presence or absence. We examined the yellow-naped amazon, *Amazona auropalliata*, a parrot species with remarkable vocal learning capabilities. Although this species' native range spans across Mesoamerica, only Costa Rican populations have been evaluated long-term. Previous studies have shown that these populations have geographically and temporally stable vocal dialect patterns. Without data on populations outside of Costa Rica, it is impossible to know whether vocal dialects are present in northern range populations, and whether they show similar geographic structure to southern range populations. We recorded yellow-naped amazon contact calls at 47 different sites across the species' range between 2016 and 2019 and evaluated them for the presence of dialects. We visually classified 14 contact call types based on spectrographic similarity and used spectrographic cross-correlation, principal component analysis, and Mantel-based spatial autocorrelations to assess acoustic similarity; we also evaluated the robustness of our findings using simulated data. The results from our study show that the vocal patterns previously seen in Costa Rica are also present in northern range populations, supporting our hypothesis that this species has vocal dialects throughout its Mesoamerican range. Call types were regionally specific (e.g., vocal dialects occurred) across the range, and no call types were repeated across multiple regions. We did, however, observe distinctive structural characteristics that are found in multiple call types, suggesting that different call types stem from a common origin. Alternatively, similarity in the acoustic features of call types may also be a result of physiological and anatomical features that are common to all members of the species. Vocal dialects in this species are likely maintained through a tendency toward philopatry and matching call types to enhance social identification.

Deleted: s

Deleted: that are

Deleted: exhibit

Deleted: if

Deleted: or not

Deleted: exhibit

Deleted: and

Deleted: d

Deleted: it should be considered that

Deleted: across

48 1 Introduction

50 Vocal learning, defined as the ability to replicate a sound produced by a conspecific or other
52 species based on social experience (Bradbury & Balsby, 2016; Tyack, 2020), can generate high
54 levels of variation in acoustic structure. Cultural transmission of vocal traits occurs when naïve
56 individuals in a population match their vocal behavior to a template observed from conspecifics,
58 such as parents or contemporaries (Whitehead, 2010; Whiten, 2019). Imperfect learning of these
60 traits can lead to changes in the behaviors of a population, a phenomenon known as cultural drift
(Podos and Warren, 2007). When acted upon in conjunction with geographic or reproductive
isolation, cultural drift in the context of learned vocalizations can produce vocal dialects, a
distinct mosaic pattern of geographic variation in acoustic signals (Wright et al., 2008; Martins et
al., 2018). Research on vocal dialects in wild populations is usually limited to a portion of a
species' range (but see Kroodsma et al., 1999). Thus, it is generally unclear whether vocal dialect
formation is intrinsic to a species or dependent on local conditions.

Since vocal dialects were first documented in the white-crowned sparrow, *Zonotrichia*
62 *leucophrys* (Marler & Tamura, 1964), they have been identified in a much wider array of taxa,
including whales (Deecke et al., 2000), hummingbirds (Wiley, 1971), seals (Casey et al., 2018),
64 bats (Boughman and Wilkinson, 1998), and parrots (Wright, 1996). Despite this wide taxonomic
distribution, songbirds have remained the focal point of research regarding both vocal learning
66 (Beecher and Brenowitz, 2005; MacDougall-Shackleton, 2009) and vocal dialects (West and
King, 1985; Maney et al., 2003; Pipek et al., 2018; Loo and Cain, 2021). Parrots are well-known
68 for their vocal mimicry skills in captivity and there is increasing evidence that under natural
conditions this open-ended learning results in a number of different patterns of variation,
70 including vocal dialects, clinal variation, and individually-specific vocal signatures (Wright and
Dahlin, 2018; Smith-Vidaurre et al., 2021, [Smeele et al., 2023](#)).

72 Yellow-naped amazons are large, social parrots that occur within lowland, seasonally dry
tropical forests along both the Pacific and Caribbean coasts of Mesoamerica, ranging from
74 southern Mexico to northern Costa Rica (Wright et al., 2018). They exhibit fission-fusion flock
dynamics, communally roosting in the evenings and departing in smaller groups around sunrise
76 to forage (Wright, 1999). The first evidence of vocal dialects in a parrot was documented in
Costa Rican populations of this species (Wright, 1996). In 2005, Wright *et al.* (2008) recorded
78 contact calls [in 2005](#) at the same sites as previously recorded in Costa Rica and found that the
acoustic variation discovered in 1996 was temporally and geographically stable. During both the
80 1996 and 2005 surveys, some birds were recorded using multiple dialects, known as
bilingualism, at some sites along dialect boundaries (Wright, 1996; Wright et al., 2008). Genetic
82 analysis showed that yellow-naped amazon vocal dialects are not genetically distinct (Wright and
Wilkinson, 2001), suggesting that this species preferentially learns local call types (Wright et al.,
84 2008), a conclusion supported by translocation experiments (Salinas-Melgoza and Wright,
2012). In addition, playback experiments showed that members of this species are less
86 responsive to playbacks from different dialects (Wright & Doring 2001). These data indicate that
yellow-naped amazon dialects may be maintained [over time](#) because using the proper dialect
88 provides social benefits to individuals (Wright and Dahlin 2018).

At present, there is little data to indicate whether yellow-naped amazon vocal dialects occur only
90 in Costa Rica and neighboring populations or extend throughout the species' range. Furthermore,

Deleted: a

92 it remains unclear whether patterns of geographic structure exist above the dialect level in this
 94 species. We hypothesized that vocal dialect patterns observed in the southern portion of the
 range would also be present in the northern portion. To address this question, we recorded
 contact calls from yellow-naped amazon populations in Costa Rica, Nicaragua, Honduras,
 96 Guatemala, and Mexico from 2016-2019. We expected to see a mosaic pattern of geographic
 variation in the contact calls, where **by** calls within populations are more similar in their acoustic
 98 structure than between populations (Wright and Dahlin, 2018). Alternatively, variation in
 yellow-naped amazon contact calls could be clinal. In this case, we **would expect** to see the
 100 acoustic structure of calls progressively diverge from one end of the range to the other (Lee et
 al., 2019). We used spectrographic cross-correlation, PCA, and Mantel-based spatial
 102 autocorrelations to assess these two alternatives. In addition, we compared the fit of models
 predicting vocal variation based on our visual dialect classification to a model based on
 104 geographic distance alone, and evaluated the robustness of the conclusions using data simulating
 either clinal or dialectal variation.

Deleted: expected

106 2 Materials and methods

106 2.1 Data collection

108 We recorded the contact calls of unmarked and unsexed yellow-naped amazons occurring in and
 around the same roost sites at which population sizes were surveyed by Dupin *et al.* (2020).
 110 Contact calls are the most frequently emitted acoustic signal in this species, particularly in and
 around the roost (Wright, 1996). We recorded calls during the morning as individuals left night
 112 roosts or in the **evening** as they arrived. We recorded calls from both flying and perched birds as
 this species is known to use contact calls in both contexts (Wright, 1996). We aimed to record a
 114 minimum of 10 calls each from six different birds at each site. Calls were recorded using a
 Sennheiser ME67 shotgun microphone and a Marantz PMD660 solid state digital recorder at a
 116 frequency of 22.05 kHz, with an amplitude resolution of 16 bits. Individuals were differentiated
 in recordings via detailed dictation by the recordist. This project was approved by New Mexico
 118 State University's Institutional Animal Care and Use Committee (IACUC protocols 2016-023
 and 2018-009).

Deleted: morning

120 2.2 Sampling sites

We recorded contact calls from yellow-naped amazons during June and July of 2016, 2018, and
 122 2019; both months fall just after the species' breeding season (Matuzak and Brightsmith, 2007,
 Dahlin et al., 2018). Sites were **chosen**, based on previous survey history, local anecdotes,
 124 **information from** local organizations working with yellow-naped amazons, and eBird reports.
 We attempted to record contact calls at as many locations as logistically possible where yellow-
 126 naped amazon populations were known to exist at the time of sampling, or were historically
 present (Dahlin et al., 2018). In 2016, we recorded calls at 22 sites in Costa Rica and 9 in
 128 Nicaragua, in 2018 and 2019 we recorded at 10 sites in southern Mexico, as well as 2 sites in
 Guatemala during 2019, and 4 sites in the Bay Islands, Honduras in 2019; one site in the Bay
 130 Islands was a private location which remains undisclosed at the owner's request. In total, we
 recorded contact calls at 47 sites across the yellow-naped amazon range (Figure 1). We were
 132 unable to record calls from a reportedly small population on the Pacific coast of El Salvador, and

Deleted: that

Deleted: determined

larger populations reported to occur along the Caribbean coast of Honduras and Nicaragua due to logistical challenges.

2.3 Data processing

Contact calls were processed digitally and visualized using the sound analysis program Raven Pro version 1.5. Calls from each field site were selected for analysis after a visual assessment of the original sound files and identification of one or more local types. Contact calls in yellow-naped amazons are typically short (0.2 - 0.5 seconds), monosyllabic, have a dominant frequency around 2 kHz, and are repeated in long strings by birds while perched and in flight; thus, calls across the range were identified as contact calls when they fit these patterns. After all contact calls within each file were identified, the highest quality calls were selected by manually annotating the start and end times on the spectrograms using Raven.

2.4 Visual assessment of call types

Annotations from Raven were imported into R for further analysis using the function `imp_raven` from the package `Rraven` (Araya-Salas, 2020). We visually assessed variation in call structure by creating a spectrogram catalogue of all calls using the `catalog` function from the `warbleR` package (Araya-Salas & Smith-Vidaurre, 2017) (see Data Availability). To do this, we used a window length of 512, overlap of 85%, and a bandpass filter with low and high frequencies of 0.5 kHz and 2.5 kHz, respectively. We chose these limits to filter out low and high frequency background noise such as traffic, insects, and other bird species.

Each contact call was categorized as being of low, medium, or high signal quality by a single observer (MG). Low quality calls were those that had high levels of noise interference, overlap with another call, and/or had no visible harmonic bands in the signal; calls of medium quality had some visible bands but lacked higher-frequency harmonics and might contain some noise interference. High quality calls had multiple, dark **harmonic** bands and no noise interference. Each spectrogram was evaluated for the presence of overlap with other calls, and whether a 2.5 kHz upper frequency limit excluded any overlapping signal or background noise. All low-quality calls were removed from the dataset along with calls that overlapped with another signal. Call selections that included noise even after the bandpass filtering were examined individually to determine whether that noise would interfere with the analysis. If noise did not overlap substantially with the call, it was retained for further analysis.

After our visual assessment, we evaluated the number of remaining calls per bird at each site, and the number of birds at each site. Birds with **fewer** than 3 calls and sites with **fewer** than 3 birds were removed from the dataset (**Supplemental Table 1**). We adjusted the start and end times of the remaining calls using the `seltailor` function from the `warbleR` package in R (Araya-Salas & Smith-Vidaurre, 2017) (see Data Availability). Each call was then categorized based on its acoustic structure; calls with noticeably different acoustic structures were categorized as **distinct call types**. This method **does require subjective judgement by observers**, but has proven to be useful as a tool to assess the overall degree of difference between **yellow-naped amazon** call types, as described by Wright and colleagues in their previous studies on yellow-naped amazons (Wright, 1996; Wright et al., 2008). All calls were grouped by country for subsequent analyses to aid in visual assessment of results.

Deleted: less

Deleted: less

Deleted: a

Deleted: is subjective

182 2.5 Statistical analyses

184 We used spectrographic cross-correlation (SPCC) to estimate pairwise dissimilarity across all
 186 calls using the cross_correlation function from the package warbleR in R (Araya-Salas & Smith-
 188 Vidaurre, 2017). Window length was set at 512, overlap was set at 85%, and low and high
 190 frequency bounds were set at 0.5 kHz and 2.5 kHz, respectively. Pairwise cross-correlation
 192 values were converted into vectors representing variation in acoustic structure using classic
 metric multidimensional scaling (hereafter, MDS) and Kruskal non-metric MDS plots. Kernel
 plots were then produced to show the distribution of each call type using concentric contour
 lines. The smallest circles indicate high concentrations of calls in one area of acoustic space. All
 code used for SPCC and MDS plots was modeled on code by Smith-Vidaurre *et al.* (2021) (see
 Data Availability).

We used principal component analysis on spectrographic features as an alternative approach to
 measure similarity between call types across the range. We accomplished this by first using the
 spectro_analysis function from the warbleR package in R (Araya-Salas & Smith-Vidaurre, 2021)
 to measure 26 acoustic parameters on each call, and then the prcomp function from the R stats
 package (R Core Team, 2021) to apply principal component analysis (PCA) to reduce
 dimensionality. We plotted the first 2 principal components for each region to compare to the
 MDS bidimensional plots (above). To test for equal variances, we used the function leveneTest()
 from the car package (Fox and Weisberg, 2019), and to evaluate the normality of our dataset we
 used the function shapiro.test() from the package MASS (Venables and Ripley, 2002). We
 evaluated differences among call types by performing Welch's ANOVA using oneway.test()
 from the R stats package (R Core Team, 2021) on principle components 1 and 2 from our PCA
 output. To determine which call type comparisons showed significant differences, we conducted
 a Tukey post-hoc analysis on our ANOVA results using the posthocTGH() from the rosetta
 package (version 0.3.6) (Peters and Verboon, 2023).

We explored the relationship between geographic distance and acoustic similarity of contact calls
 using a custom routine based on the Mantel correlogram with the function mantel.correlog from
 the package vegan (version 2.5-7) (Oksanen et al., 2020). Mantel correlograms show the change
 in dissimilarity (in this case acoustic dissimilarity) as a function of spatial distance. Hence, they
 can be useful to assess patterns of variation in yellow-naped amazon calls across the species'
 geographic range. Dissimilarity is typically measured over a fixed set of even spatial distances
 across a specified geographic range, but results can vary markedly for different spatial
 resolutions. Therefore, we calculated several Mantel correlograms with different spatial
 resolutions (6 distances from 25 to 250 km) and estimated the mean and standard deviations
 across the correlograms for each distance value after interpolation.

We tested whether distance or dialect assignment had a significant effect on call structure using a
 multiple regression model for distance matrices (Lichstein, 2007) in the function MRM from the
 R package ecodist (Goslee and Urban, 2007). This method uses a permutation approach to
 estimate statistical significance of the regression coefficients to account for the lack of
 independency of observations. A dissimilarity matrix based on the pairwise cross-correlation
 matrix (1 - spcc matrix) was used as a response variable while a pairwise distance matrix and a
 dialect membership binary matrix, were used as a predictor. The latter, created with the function
 binary_triangular_matrix in the R package PhenotypeSpace (Araya-Salas and Odom 2022), uses

Deleted: , with closer lines representing a higher density of calls...

Deleted: In addition to creating plots for each country, we also generated cumulative plots over the entire range for each analysis to assess whether variation appeared to be random or clinal.

Deleted: l

Deleted: W

Deleted: to assess for equal variances

Deleted: the

Deleted: to evaluate the normality of our dataset

Deleted: ed

Deleted: to evaluate differences among call types

Deleted: we conducted a Games-Howell post-hoc analysis on our ANOVA results using the posthocTGH() from the rosetta package...

Deleted: measures

Deleted: ed

Deleted: to statistically evaluate whether distance or dialect assignment had a significant effect on call structure in yellow-naped amazons

246 0s and 1s to represent whether two calls belong to the same dialect (0) or not (1), and therefore
can be used to assess the variation in call structure explained by dialects.

248 Finally, we use simulations to evaluate whether our statistical approach could tease apart
250 dialectal from clinal variation in call acoustic structure. For this, we created synthetic data sets
with the same number of locations and calls per location as in the actual data set. Locations were
252 equally spaced across the x axis (separated by one unit) in a bidimensional space. The position of
each observation within each location was set randomly by adding random variation from a
normal distribution (centered at 0 with standard deviation of 0.1) on both the x and y axis. For
254 each pattern of acoustic variation we simulated a vector representing a hypothetical acoustic
feature. The clinal geographic pattern was simulated by simply adding random variation (from a
normal distribution centered at 0 with standard deviation of 0.2) to the x axis value of each
256 observation. The dialectal geographic pattern was simulated by randomly assigning an integer
number to each location (i.e. the same value to all observations within a location) and then
258 adding random variation to each observation (from a normal distribution centered at 0 with
standard deviation of 0.2). We also simulated a random variation pattern in which the acoustic
260 feature value was assigned randomly regardless of locality of geographic distance to other
observations. The three continuous simulated vectors (clinal, dialectal and random variation) as
262 well as the geographic positions were converted into pairwise distance matrices using the
function dist in R (R Core Team, 2021). Distances were zero-centered and transformed to unit
264 variance to enable effect size comparisons between different models. As with the real data set,
dialect membership was represented by a binary matrix that uses 0s and 1s to represent whether
266 two calls belong to the same dialect (0) or not (1), created by the function
binary_triangular_matrix from the package PhenotypeSpace (Araya-Salas and Odom 2022). We
268 also evaluated whether geographic distance and dialect membership (predictors) and the three
acoustic feature variables (responses) were associated using multiple regression for distance
270 matrices (Lichstein, 2007). Single response models were run for each acoustic feature variable.
272 The simulation was replicated 100 times and the results were averaged across replicates.

3 Results

274 3.1 Contact calls and quality assessment

We surveyed for yellow-naped amazons at 72 locations in Mesoamerica where they are known to
276 exist or have occurred historically. We observed this species at 56 of those sites and recorded 3
or more individuals at 47 sites (Supplemental Table 1). In 2016 our team recorded contact calls
278 at 22 locations in Costa Rica and 9 in Nicaragua. We recorded calls at 10 sites in Mexico
between 2018 and 2019. In 2019 we recorded contact calls at 2 sites in Guatemala and 4 in the
280 Bay Islands of Honduras. We identified and selected 2,566 yellow-naped amazon contact calls
from our field recordings and removed 116 of those during quality assessment due to the
282 presence of noise overlap or low signal quality. In total, 24 birds were removed from the
remaining dataset due to retention of fewer than three contact calls. Three sites from Costa Rica
284 (Ensenada, Pelon Bajura, and Palo Verde) were removed as a result of having fewer than 3 birds
(Supplemental Table 1). None of the other countries surveyed had sites removed during quality
286 assessment.

Deleted:

Deleted:

Deleted:

Deleted: M

Deleted: were also used to evaluate the association between geographic distance and dialect membership (predictors) and the three acoustic feature variables (responses)

Deleted:

We retained 2,338 yellow-naped amazon contact calls from our original dataset for analysis. We classified 21.5% of calls as medium quality, and 78.5% of calls as high quality. Contact calls from Mexico represented 23.3% of contact calls in the final dataset; Guatemala 1.5%; the Bay Islands, Honduras 9.3%; Nicaragua 23%; and Costa Rica 43% of all calls.

3.2 Call types across the range

Our visual classification of yellow-naped amazon contact calls yielded 14 different call types across the species range (Figure 2). We identified 4 types in Mexico and 2 types in Guatemala. In the Bay Islands Honduras, we [found](#) 2 distinct types. Previously, Wright *et al.* reported 3 call types in Costa Rica during a 1994 survey and the same 3 types and a new sub-type during a second survey in 2005 (Wright, 1996; Wright et al., 2008); our visual analysis of contact calls collected from these same regions during 2016 indicated that birds in Costa Rica [gave](#) 2 different call types, both of which were reported in the previous surveys. The third call type previously reported on the northern border of Costa Rica in 1994 and 2005 was not recorded there in 2016 but was seen in populations in the south of Nicaragua, along with 3 other call types [across Nicaragua](#). Sixteen percent of the 243 birds in our dataset [were bilingual](#). Overall, these observations were infrequent, and only 19 bilingual birds were retained in our dataset for analyses. These individuals were primarily concentrated in Costa Rica (N = 16), although bilingual birds were also recorded in Mexico (N = 2) and Guatemala (N = 1). It should be noted that we did observe and record bilingual birds in Nicaragua; however, none of those birds were preserved in our dataset due to low quality recordings or an insufficient number of recordings. Our visual assessment suggested that the mosaic patterns of vocal variation consistent with dialects was present throughout the range, rather than a clinal variation of call types. Dialect types in this species appear to result from discrete modifications to a basic call structure (Figure 3) that result in superficially similar call types appearing in different parts of the species range.

Deleted: recorded

Deleted: i

Deleted: exhibited bilingualism

Deleted: c

3.3 Spectrographic cross-correlations

SPCC analyses for each country supported some, but not all of our call classifications for the yellow-naped amazon contact call types identified across the range (Figure 4). When all identified types from across the range were plotted in acoustic space, it was difficult to detect separate distributions as many overlapped to a significant extent. When we examined acoustic overlap on a country-level basis, we noticed that the types sampled in the northern ([Mexico and Guatemala](#)) and eastern ([Bay Islands](#)) portions of the range showed more acoustic separation amongst themselves than did call types in the southern ([Nicaragua and Costa Rica](#)) portion of the range, where dialects were first described in this species ([see Supplemental Figure 1 for regional PCA plots](#)).

3.4 Principal component analysis of spectral measures

We used PCA on spectral measures [from all calls](#) as an alternative measure of validation for our visually classified contact call types. The first 5 loadings from our analysis accounted for 77.4% of cumulative variation (Figure 4 [and Supplemental Table 2](#)). PC1 represented 26.7% of explained variation and primarily accounted for differences in frequency variables such as mean dominant frequency, and mean frequency (in kHz). PC2 explained 17.6% of variation and

represented differences in temporal variables such as signal duration and duration of the call in the 75th quantile (Supplemental Table 2).

Plots of PC1 and PC2 for call types in each country generally reflected the patterns exhibited in the corresponding SPCC plots (Figure 4). When we created a PC plot for all call types identified across the range, there was little visual separation among call types (data not shown). An ANOVA on PC1 and PC2 for all call types combined did indicate significant differences between call types for both PC1 ($F = 294.3$, $df = 299.41$, $p < 0.001$) and PC2 ($F = 53.292$, $df = 300.36$, $p < 0.001$). Our Tukey post-hoc analysis on the ANOVA values from PC1 revealed that 74% of pairwise call type comparisons showed significant differences at the $p < 0.001$ level. Forty-one percent of pairwise call type comparisons from the Tukey post-hoc analysis on ANOVA values from PC2 showed significant differences at the $p < 0.001$ level (Supplemental Table 3).

3.5 Spatial patterns of acoustic similarity

The Mantel-based spatial autocorrelogram illustrated two distinct patterns in our data. First, within 250 km yellow-naped amazon contact calls generally exhibit high similarity (Figure 5). Second, after 250km, acoustic similarity decreases markedly and remains at a low level for the remaining geographic distance range.

The multiple regression of distance matrices revealed a significant association between acoustic dissimilarity and dialect membership ($p \leq 0.0001$) but no association with geographic distance ($p = 0.281$). Call dissimilarity was higher between calls from different dialects relative to calls from the same dialect. The effect size of dialect membership (0.65) was three orders of magnitude higher than the absolute effect size of geographic distance (-0.003).

The simulated data revealed that our statistical approach is capable of discerning between the two hypothesized patterns of geographic variation. When modeling acoustic similarity in clinal variation, the effect of geographic distance on acoustic similarity was consistently higher (effect size mean = 1.000; $sd = 0.001$) than that of dialect membership (effect size mean = -0.024; $sd = 0.005$). The opposite pattern was observed when predicting acoustic similarity in data simulating dialectal variation: the effect size of dialect membership was consistently higher (mean = 1.440; $sd = 0.257$) than that of geographic distance (mean = 0.020; $sd = 0.178$). In both simulations 95% of the replicates produced significant effects for both predictors. Finally, for the random variation simulation, geographic distance and dialect membership both produce equally low effect sizes (mean = -0.001; $sd = 0.009$ and mean = 0.001; $sd = 0.016$; respectively). Effects were significant in fewer than 95% of these replicated models.

4 Discussion

Historically, variation in the contact calls of yellow-naped amazons in Costa Rica has been described as dialectal, with a mosaic pattern of variation in which calls within a locale are similar but change dramatically in acoustic structure at dialect boundaries (Wright, 1996). Based on visual categorization of sound spectrograms, we defined 14 different call types confirming that the pattern of dialects previously seen in Costa Rica is present throughout the Mesoamerican range of this species. Currently wild yellow-naped amazons in many parts of their range are

Deleted:

Deleted: 2.2⁻¹⁶

Deleted: 2.2⁻¹⁶

Deleted: were

Deleted: ly

Deleted: t

Deleted: were

Deleted: ly

Deleted: t

Deleted: The significance value for the post-hoc tests was set to $p < 0.001$.

Deleted: 2

Deleted: After this distance

Deleted: to

Deleted: al

Deleted: that

Deleted: less

Deleted: Our

Deleted: classification

Deleted: of

Deleted: observed across the species' range confirm

Deleted: this

isolated into discrete regional populations due to loss of habitat and declining populations (Dupin et al., 2020). In each region we sampled, we were able to visually detect more than one acoustically discrete call type with a distinct geographic distribution. Our SPCC and PCA analyses validated many, but not all, of our visually identified acoustic types. Importantly, multiple regressions of distance matrices from our data gave stronger support for dialect variation than clinal variation, with effect sizes of similar magnitude to the same analysis run on data simulating dialect variation. In general, our data indicate that conditions which lead to dialect formation are intrinsic to the social structure and communication system of yellow-naped amazons, which appears to depend heavily on vocal learning.

4.1 Global patterns of vocal variation

In 1996, Wright identified 3 distinct contact call types in Costa Rican populations (Wright, 1996), and later confirmed the geographic and temporal stability of those same call types (Wright et al., 2005). Our visual assessment of over 2000 contact calls recorded from 44 sites across the range of the yellow-naped amazon revealed the presence of 14 distinct call types, including those identified in the initial description of dialects in Costa Rica. Results from our SPCC analyses and PCA support many of our visual classifications; however, they also indicate that several call types were acoustically similar. Overall, the SPCC and PCA plots showed more separation in the northern and eastern portions of the range (e.g. the Apex and Island call types from the Bay Islands). The southern portion of the range had more call type overlap (e.g. South and North call types in Costa Rica). This pattern may result from the repetition of similar acoustic structures in different parts of the range, or a lack of resolution in our acoustic analysis approaches. While SPCC is widely-used method of assessing acoustic similarity (Cortopassi and Bradbury, 2000; Sharp and Hatchwell, 2006; Araya-Salas et al., 2019) and has been used in several previous studies of acoustic variation in parrots (Wright, 1996; Wright et al., 2008), the limits of its capabilities should be considered. The cross-corr function from the package warbleR in R generates correlation coefficients by sliding one spectrogram over another and comparing amplitudes for each frequency and time point across each call (Clark et al., 1987; Araya-Salas and Smith-Vidaurre, 2017; Sawant et al., 2022). Thus, when call types use the same bandwidths and have very similar fundamental frequencies, the cross correlation will indicate that the calls are acoustically similar, even if there are visually detectable temporal and frequency modulation differences in the overall structure (Supplemental Figure 2). Although SPCC is considered to be somewhat robust to noise, field recordings may contain considerable acoustic interference from insects, other birds, vehicles, and farm animals. We implemented a bandpass filter to remove high and low frequency noise and discarded those recordings with substantial overlapping noise within the frequency range as the calls themselves but retained calls with lower levels of masking noise.

Several other parrot species have been found to exhibit vocal dialects, including the Australian ringneck parrot, *Barnardius zonarius* (Baker, 2000); galah, *Eolophus roseicapillus* (Baker, 2003); Australian palm cockatoo, *Probosciger aterrimus* (Keighley et al., 2017); Cuban parrot, *Amazona leucocephala* (Reynolds and Hayes, 2010); and invasive populations of the monk parakeet, *Myiopsitta monachus* (Buhrman-Deever et al., 2007; but see Smith-Vidaurre et al., 2021) for evidence of different patterns in native populations. There is little understanding of the process by which parrot species form vocal dialects, but several hypotheses have been suggested to explain their function (Wright and Dahlin, 2018). Podos and Warren (2007) postulated that

Deleted: e

Deleted: e

Deleted:

Deleted: northern

Deleted: half

Deleted: , like with the

Deleted: , like in

Deleted: are

Deleted: are similar in the same bandwidths being measured, the cross ...

Deleted: in addition to yellow-naped amazons

Deleted: (

Deleted:) (Buhrman-Deever et al., 2007; Smith-Vidaurre et al., 2021)

Deleted: as to

Deleted: why

Deleted: exhibit

Deleted: formed

464 female preference for local call types is a driver of acoustic divergence. In this case, female
 466 sexual preference for males with a local trait would result in some level of genetic divergence
 468 between populations (Wilkins et al., 2013). A study by Wright and Wilkinson (2001) reported no
 470 genetic correlation with dialects in yellow-naped amazon contact calls, stating instead that
 individuals exhibited high dispersal across dialect boundaries. Parrots may also exhibit vocal
 dialect as a badge or signal of group affiliation and familiarity (Sewall et al., 2016).
 Alternatively, geographic variation in dialects could be a consequence of cultural drift, when
 individuals learning calls make copying errors, and then are themselves copied accurately (Podos
 and Warren, 2007).

472 Overall, yellow-naped amazon vocal dialects are most consistent with a process of cultural drift.
 In general, we noticed that yellow-naped amazon contact calls exhibit some similar
 474 spectrographic structures in different call types across the range, such as the presence of gaps
 476 within a call, frequency shifts, and differences in duration. We observed that neighboring call
 types exhibit subtle differences in acoustic structure as opposed to whole-structure change. In
 conjunction, these facts suggest that call types have a common origin in a 'proto-dialect' call
 478 type (Figure 3), and that subsequent variation is a result of cultural drift, possibly in isolated
 480 populations. It is also possible that call types share similar acoustic features due to anatomical
and physiological characteristics of the vocal control system that are common across all
 482 members of the species, and constrain vocal system modulations that produce sounds. Long-term
 studies by Wright and colleagues on yellow-naped amazons in Costa Rica lend credence to the
 cultural drift hypothesis (Wright et al., 2008; C. Dahlin et al. in prep). Additionally, there is
 support for the social identification hypothesis when call types are examined on a smaller scale.
 Wright (2000) found marked differences in neighboring dialects. If dialects were solely a result
 of cultural drift, neighboring call types would show a strong tendency to be more similar and
 show clinal variation across the species' range. The absence of strong clinal patterns suggest that
 488 there is some selection for neighboring populations to produce acoustically distinct call types,
 consistent with the hypothesis that calls function for group identification (Buhrman-Deever et al.,
 490 2007).

Several birds in our dataset exhibited more than one call type. In most cases of bilingualism there
 492 appeared to be a dominant call type used with high frequency by an individual, and a secondary
 type that a bird would emit infrequently. Wright reported the occasional occurrence of bilingual
 494 birds in Costa Rica during his 1994 survey, and Wright *et al.* observed them infrequently in
 Costa Rica during their 2005 survey (Wright, 1996; Wright et al., 2005). Bilingualism may exist
 496 in these populations to facilitate individual movement between social groups during fission and
 fusion. If, during foraging, individuals encounter conspecifics that give a different call type,
 498 learning the additional call type might increase foraging success and enhance protection provided
 by group membership. It is also possible that bilingualism occurs primarily in dispersing
 500 individuals that have learned one dialect in their natal roost and then dispersed to a roost with a
different dialect, and learned the new dialect while retaining the original one (Salinas-Melgoza
 502 and Wright, 2012). In this case, bilingualism would have no functional advantage.

4.2 Regional patterns and implications for cultural evolution

504 Yellow-naped amazon populations currently occur in regional clusters across their range with
 few or no birds observed in between, due to rapid population declines associated with habitat

Deleted: as seen

Deleted: shows evidence

Deleted: like

Deleted: imperfect learning and

Deleted: should be also considered that it is

Deleted: have

Deleted: because

Deleted: of

Deleted: Previous temporal

Deleted:

Deleted: *et al*

Deleted: .

Deleted: also

Deleted: Previously,

Deleted: that

Deleted: exhibited marked differences

Deleted: an epiphenomenon of dispersal is occurring, where
 individuals

Deleted: ,

Commented [TW1]: Add citation
 Salinas-Melgoza, A., & Wright, T. F. (2012). Evidence
 for Vocal Learning and Limited Dispersal as Dual
 Mechanisms for Dialect Maintenance in a Parrot. *PLoS*
ONE, 7(11), e48667. doi:
 10.1371/journal.pone.0048667

Deleted: that

Deleted: birds

Deleted:

528 loss and fragmentation (Dupin et al., 2020). This habitat fragmentation in the birds' geographic
530 range may serve to enhance the mosaic pattern of call distribution. However, there is no evidence
532 that the current call types have arisen due to this recent isolation, and many of these population
534 clusters contain more than one call type. It seems more likely that call types predate this
536 isolation, as some of the earliest recordings from this area show distinct regional call types (see
538 Wright and Wilkinson 2001). These regional call types may be maintained by a tendency
540 towards philopatry and selection towards matching regional types to enhance group
identification, as suggested by playback responses of dialect variants of another vocalization
type, pair duets (Wright and Dorin, 2001). In southern Nicaragua and Costa Rica, where yellow-
naped amazon populations are more contiguous, many call types are restricted to a certain
geographic area, while others are currently more widespread. In the northern-most country of
Mexico, yellow-naped amazons are much more patchily distributed, which may explain why call
types exhibit almost no acoustic overlap (Sewall et al., 2016).

542 Though we were able to sample populations across the yellow-naped amazons' range, there are
544 several regions we were unable to survey due to time, manpower, and safety limitations. These
546 regions include the country of El Salvador, the southeastern Caribbean region of Honduras, and
548 the northwestern Caribbean region of Nicaragua. Although we were not able to record birds in
550 these areas, our data are likely to be an accurate representation of the pattern of vocal variation
552 across the range as we were able to sample populations at the northern- and southern-most
bounds and on both the Pacific and Caribbean slopes of Mesoamerica. A study by Herrera and
colleagues in El Salvador suggest there are an estimated 250 yellow-naped amazons in the
country (2020). It is also likely that some healthy populations remain in the large, unsampled
regions of Nicaragua and Honduras where forests are relatively intact and human populations are
smaller. Future yellow-naped amazon dialect studies should focus on evaluating the temporal
and geographic stability of call types across the range, including those unsampled populations.

4.3 Summary and future directions

554 Our cultural atlas of vocal variation in contact calls across the range of the yellow naped amazon
556 supports the hypothesis that vocal dialects are present across the range of this species. Contact
558 call types exhibited separation by geographic region with only rare cases of bilingual birds, a
560 trend seen in original studies on Costa Rican populations. These results suggest that the factors
562 that give rise to and maintain vocal dialects are inherent to the social structure and
564 communication system of this species. Our observation of repeated structural similarities in
geographically distinct acoustic variants suggests that calls may have diversified from a shared
basic structure through the occurrence of learning errors and their subsequent cultural
transmission in populations. Recent isolation of populations may reinforce these tendencies.
Future studies on yellow-naped amazons could examine functional responses to dialect variants
of contact calls, potentially in reintroduced populations that lack local dialect types.

5 Conflict of Interest

566 The authors declare that the research was conducted in absence of any commercial or financial
relationships that could be construed as a potential conflict of interest.

6 Author Contributions

Deleted: isolation

Deleted: ,

Deleted: h

Deleted: reason

Deleted: to believe

Deleted: Rather,

Deleted: i

Deleted: we believe that our

Deleted: is still

Deleted:

Molly K. Genes (néé Dupin) formulated the questions, collected data, analyzed the data, and wrote the paper; Marcelo Araya-Salas analyzed the data; Christine R. Dahlin collected data and supervised the research; Timothy F. Wright planned the study, collected data, supervised the research and data analysis, and assisted in writing the paper.

7 Funding

This research was funded by the World Parrot Trust (grant date April 2018), and New Mexico State University's College of Arts and Sciences.

8 Acknowledgements

We are very grateful to the colleagues that aided in the collection field data between 2016 and 2019: Alyssa Trimeloni, Dr. Grace Smith-Vidaurre, Dominique Hellmich, Lorena Cabada-Gomez, Carlos Campos, Dr. James Gilardi, Sam Williams and the Macaw Recovery Network, Eric Anderson, Edith Belen Jimenez Diaz, Cristina Contreras Meda, Candelario Giron Montes, Martín Lezama Lopez, and Cara Dunbar. We would also like to extend a thank you to the following organizations which facilitated in data collection via permission to record on their land: Area de Conservación Guanacaste, La Biosfera de la Encrucijada, and Finca Patrocinio and One Earth Organization. Also, thank you to Theodore Muka, Sophia Lasche, and Adam Hopper for providing feedback and editing.

9 Contribution to the field

Vocal learning research in wild animals has gained momentum over the last several decades, yet the focus of most studies remains on songbird species. Many parrot species are known to be open-ended vocal learners and can acquire new vocalizations well into adulthood. Our study on the yellow-naped amazon calls across its [Mesoamerican](#) range illustrate that the mosaic pattern of discrete call types (e.g., vocal dialects) previously described in one part of the range, are characteristic of all surveyed populations. This suggests that the processes that give rise to these dialects are a general characteristic of the species. Additionally, our analyses show that the call types we identified across the range share a basic structure that has likely been modified in different populations via learning errors and cultural drift. More generally, wild parrots have abundant potential to reveal more about vocal learning processes and how this form of behavioral plasticity contributes to communication and social interactions.

10 References

- Araya-Salas, M., & K. Odom. 2022. PhenotypeSpace: and R package to quantify and compare phenotypic trait spaces. R package version 0.1.0.
- Araya-Salas, M., and Smith-Vidaurre, G. (2017). warbleR: an R package to streamline analysis of animal acoustic signals. *Methods Ecol Evol.* 8:2. doi: 10.1111/2041-210X.12624.
- Araya-Salas, M., Smith-Vidaurre, G., Mennill, D. J., González-Gómez, P. L., Cahill, J., and Wright, T. F. (2019). Social group signatures in hummingbird displays provide evidence of co-occurrence of vocal and visual learning. *Proc R Soc B.* 286:1903. doi: 10.1098/rspb.2019.0666.
- Baker, M. C. (2000). Cultural diversification in the flight call of the Ringneck Parrot in Western Australia. *Condor.* 102:4. doi: 10.2307/1370319.

Baker, M. C. (2003). Local similarity and geographic differences in a contact call of the Galah
 (Cacatua roseicapilla assimilis) in Western Australia. *Emu*. 103:3. doi: 10.1071/MU02035.
 Beecher, M. D., and Brenowitz, E. A. (2005). Functional aspects of song learning in songbirds.
Trends Ecol Evol. 20:3. doi: 10.1016/j.tree.2005.01.004.
 Boughman, J. W., and Wilkinson, G. S. (1998). Greater spear-nosed bats discriminate group
 mates by vocalizations. *Anim Behav*. 55:6. doi: 10.1006/anbe.1997.0721.
 Bradbury, J. W., and Balsby, T. J. S. (2016). The functions of vocal learning in parrots. *Behav*
Ecol Sociobiol. 70:3. doi: 10.1007/s00265-016-2068-4.
 Buhrman-Deever, S. C., Rappaport, A. R., and Bradbury, J. W. (2007). Geographic variation in
 contact calls of feral North American populations of the Monk Parakeet. *Condor*. 109:2. doi:
 10.1093/condor/109.2.389.
 Casey, C., Reichmuth, C., Costa, D. P., and Boeuf, B. Le (2018). The rise and fall of dialects in
 northern elephant seals. *Proc R Soc B*. 285:1892. doi: 10.1098/RSPB.2018.2176.
 Clark, C. W., Marler, P., and Beeman, K. (1987). Quantitative analysis of animal vocal
 phonology: an application to swamp sparrow song. *Ethology*. 76:2. doi: 10.1111/j.1439-
 0310.1987.tb00676.x.
 Cortopassi, K. A., and Bradbury, J. W. (2000). The comparison of harmonically rich sounds
 using spectrographic cross-correlation and principal coordinates analysis. *Bioacoustics*.
 11:2. doi: 10.1080/09524622.2000.9753454.
 Dahlin, C. R., Blake, C., Rising, J., and Wright, T. F. (2018). Long-term monitoring of Yellow-
 naped Amazons (Amazona auropalliata) in Costa Rica: breeding biology, duetting, and the
 negative impact of poaching. *J Field Ornithol*. 89:1. doi: 10.1111/jofo.12240.
 Deecke, V. B., Ford, J. K. B., and Spong, P. (2000). Dialect change in resident killer whales:
 Implications for vocal learning and cultural transmission. *Anim Behav*. 60:5. doi:
 10.1006/anbe.2000.1454.
 Dupin, M. K., Dahlin, C. R., and Wright, T. F. (2020). Range-wide population assessment of the
 endangered yellow-naped amazon (Amazona auropalliata). *Diversity*. 12:10. doi:
 10.3390/d12100377.
 Fox, J., Weisberg, S. (2019). *An R Companion to Applied Regression*, Third edition. Thousand
 Oaks, CA: Sage.
 Goslee, S. C., and Urban, D. L. (2007). The ecodist package for dissimilarity-based analysis of
 ecological data. *J Stat Softw*. 22. doi: 10.18637/jss.v022.i07.
 Herrera, N., Lara, K., and Funes, C. (2020). Estado poblacional de la Lora Nuca Amarilla
 (Amazona auropalliata) en El Salvador. *Zeledonia*. 24:1. Retrieved from
<https://www.zeledonia.com>
 Keighley, M. V., Langmore, N. E., Zdenek, C. N., and Heinsohn, R. (2017). Geographic
 variation in the vocalizations of Australian palm cockatoos (Probosciger aterrimus).
Bioacoustics. 26:1. doi: 10.1080/09524622.2016.1201778.
 Kroodsma, D. E., Byers, B. E., Halkin, S. L., Hill, C., Minis, D., Bolsinger, J. R., Dawson, J. A.,
 Donelan, E., Farrington, J., Gill, F. B. and Houlihan, P. (1999). Geographic variation in
 black-capped chickadee songs and singing behavior. *Auk*. 116:2. doi: 10.2307/4089373
 Lee, J. H., Podos, J., and Sung, H. C. (2019). Distinct patterns of geographic variation for
 different song components in Daurian Redstarts Phoenicurus aureus. *Bird Study*. 66:1. doi:
 10.1080/00063657.2019.1614144.
 Lichstein, J. W. (2007). Multiple regression on distance matrices: a multivariate spatial analysis
 tool. *Plant Ecol*. 188. doi: 10.1007/s11258-006-9126-3.

666 Loo, Y. Y., and Cain, K. E. (2021). A call to expand avian vocal development research. *Front*
Ecol Evol. 9. doi: 10.3389/fevo.2021.757972.
 668 MacDougall-Shackleton, S. A. (2009). The importance of development: What songbirds can
 teach us. *Can J Exp Psychol.* 63:1. doi: 10.1037/a0015414.
 670 Maney, D. L., MacDougall-Shackleton, E. A., MacDougall-Shackleton, S. A., Ball, G. F., and
 Hahn, T. P. (2003). Immediate early gene response to hearing song correlates with receptive
 672 behavior and depends on dialect in a female songbird. *J Comp Physiol A.* 189. doi:
 10.1007/s00359-003-0441-z.
 674 Martins, B. A., Rodrigues, G. S. R., and de Araújo, C. B. (2018). Vocal dialects and their
 implications for bird reintroductions. *Perspect Ecol Conserv.* 16:2. doi:
 10.1016/j.pecon.2018.03.005.
 676 Matuzak, G. D., and Brightsmith, D. J. (2007). Roosting of Yellow-naped Parrots in Costa Rica:
 estimating the size and recruitment of threatened populations. *J Field Ornithol.* 78:2. doi:
 678 10.1111/j.1557-9263.2007.00099.x.
 680 Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., et al. (2020).
 Vegan: Community Ecology Package. R package version 2.5-7. [https://CRAN.R-](https://CRAN.R-project.org/package=vegan)
 682 [project.org/package=vegan](https://CRAN.R-project.org/package=vegan)
 package version 0.3.6. <https://r-packages.gitlab.io/rosetta/>
 684 Pipek, P., Petrusková, T., Petrušek, A., Diblíková, L., Eaton, M. A., and Pyšek, P. (2018).
 Dialects of an invasive songbird are preserved in its invaded but not native source range.
 686 *Ecography.* 41:2. doi: 10.1111/ecog.02779.
 Podos, J., and Warren, P. S. (2007). The evolution of geographic variation in birdsong. *Adv Stud*
 688 *Behav.* 37. doi: 10.1016/S0065-3454(07)37009-5.
 R Core Team, R. (2021). *R: A Language and Environment for Statistical Computing*. Vienna,
 690 Austria: R Foundation for Statistical Computing.
 Reynolds, M. B. J., and Hayes, W. K. (2010). Geographic variation in the flight call of the Cuban
 692 Parrot (*Amazona leucocephala*) and its taxonomic relevance. *J Caribbean Ornithol.* 23:1.
 Retrieved from <https://jco.birdscaribbean.org/index.php/jco>
 694 Salinas-Melgoza, A., and Wright, T. F. (2012). Evidence for vocal learning and limited dispersal
 as dual mechanisms for dialect maintenance in a parrot. *PLoS One.* 7:11. doi:
 696 10.1371/journal.pone.0048667.
 Sawant, S., Arvind, C., Joshi, V., and Robin, V. V. (2022). Spectrogram cross-correlation can be
 698 used to measure the complexity of bird vocalizations. *Methods Ecol Evol.* 13:2. doi:
 10.1111/2041-210X.13765.
 700 Sewall, K. B., Young, A. M., and Wright, T. F. (2016). Social calls provide novel insights into
 the evolution of vocal learning. *Anim Behav.* 120. doi: 10.1016/j.anbehav.2016.07.031.
 702 Sharp, S. P., and Hatchwell, B. J. (2006). Development of family specific contact calls in the
 Long-tailed Tit *Aegithalos caudatus*. *Ibis.* 148:4. doi: 10.1111/j.1474-919X.2006.00568.x.
 704 [Smeele, S. Q., Senar, J. C., Aplin, L. M., & McElreath, M. B. \(2023\). Evidence for vocal](#)
 706 [signatures and voice-prints in a wild parrot. Roy. Soc. Open Sci., 10:230835. doi:](#)
[doi:10.1098/rsos.230835](#)
 708 Smith-Vidaurre, G., Araya-Salas, M., and Wright, T. F. (2021). Individual signatures outweigh
 social group identity in contact calls of a communally nesting parrot. *Behav Ecol.* 31:2. doi:
 10.1093/BEHECO/ARZ202.

- 710 Tyack, P. L. (2020). A taxonomy for vocal learning. *Philo T R Soc B*. 375:1789. doi:
10.1098/rstb.2018.0406.
- 712 Venables, W. N., and Ripley, B. D. (2002). *Modern Applied Statistics with S*, Fourth edition.
New York: Springer.
- 714 West, M. J., and King, A. P. (1985). Studying dialects in songbirds: Finding the common ground.
Behav Brain Sci. 8:1. doi: 10.1017/S0140525X0001997X.
- 716 Whitehead, H. (2010). Conserving and managing animals that learn socially and share cultures.
Learn Behav. 38:3. doi: 10.3758/LB.38.3.329.
- 718 Whiten, A. (2019). Cultural Evolution in Animals. *Annu Rev Ecol Evol S*. 50. doi:
10.1146/annurev-ecolsys-110218-025040.
- 720 Wiley, R. H. (1971). Song Groups in a Singing Assembly of Little Hermits. *Condor*. 73:1. doi:
10.2307/1366121.
- 722 Wilkins, M. R., Seddon, N., and Safran, R. J. (2013). Evolutionary divergence in acoustic
signals: Causes and consequences. *Trends Ecol Evol*. 28:3. doi: 10.1016/j.tree.2012.10.002.
- 724 Wright, T. (1999). Vocal Communication In Wild Populations of the Yellow-naped Amazon
(*Amazona auropalliata*). *AFA Watchbird*. 26:1, 7-9.
- 726 Wright, T. F. (1996). Regional dialects in the contact call of a parrot. *Proc R Soc B*. 263:1372.
doi: 10.1098/rspb.1996.0128.
- 728 Wright, T. F., and Dahlin, C. R. (2018). Vocal dialects in parrots: Patterns and processes of
cultural evolution. *Emu*. 118:1. doi: 10.1080/01584197.2017.1379356.
- 730 Wright, T. F., Dahlin, C. R., and Salinas-Melgoza, A. (2008). Stability and change in vocal
dialects of the yellow-naped amazon. *Anim Behav*. 76:3. doi:
10.1016/j.anbehav.2008.03.025.
- 732 Wright, T. F., and Dorin, M. (2001). Pair duets in the yellow-naped amazon (Psittaciformes:
Amazona auropalliata): responses to playbacks of different dialects. *Ethology*. 107:2. doi:
10.1046/j.1439-0310.2001.00632.x.
- 736 Wright, T. F., Lewis, T., Lezama-Lopez, M., Smith-Vidaurre, G., and Dahlin, C. (2018). Yellow-
naped amazon (*Amazona auropalliata*) populations are markedly low and rapidly declining
in Costa Rica and Nicaragua. *Bird Conserv Int*. 29:2. doi: 10.1017/S0959270918000114.
- 738 Wright, T. F., Rodriguez, A. M., and Fleischer, R. C. (2005). Vocal dialects, sex-biased
dispersal, and microsatellite population structure in the parrot *Amazona auropalliata*. *Mol*
Ecol. 14:4. doi: 10.1111/j.1365-294X.2005.02466.x.
- 740 Wright, T. F., and Wilkinson, G. S. (2001). Population genetic structure and vocal dialects in an
amazon parrot. *Proc R Soc B*. 268:1467. doi: 10.1098/rspb.2000.1403.
- 742
- 744

10 Data Availability Statement

- 746 The original contributions presented in the study are publicly available. All relevant code for
analyses can be found at the following GitHub repository: [https://github.com/mgenes/cultural-](https://github.com/mgenes/cultural-atlas-of-vocal-variation)
748 [atlas-of-vocal-variation](https://github.com/mgenes/cultural-atlas-of-vocal-variation). Data for analyses can be found at
<https://doi.org/10.5061/dryad.9cnp5hqq7>.

750

11 Figure Captions

752 Figure 1. (a) A map of all sites where 3 or more birds were recorded during 2018 and 2019. Sites
754 from the study conducted by Wright in 2016 were also included (Wright et al., 2019). The color
756 and shape of each point corresponds to the year the site was sampled. Purple circles represent
758 sites with fewer than 3 birds that were removed from the dataset prior to analysis. The
undisclosed site located in Honduras is not shown. (b) A species range polygon for the yellow-
naped amazon provided by BirdLife (BirdLife International and Handbook of the Birds of the
World, 2019).

760 Figure 2. Spectrograms of each identified call type with the country of origin listed above and
the call type name listed below the image. All images are shown on a 0.6 second timescale at 3.5
kHz frequency.

762 Figure 3. An image depicting typical structural variants of yellow-naped amazon contact calls
and hypothesized routes by which one type might evolve into another through cultural drift.

764 Figure 4. A multi-panel image with four columns showing SPCC plots (A), PCA plots (B), a
766 geographic map (C), and a legend (D). Rows represent the sampled countries, arranged in order
from north to south. The geographic map for Honduras depicts the call type for the undisclosed
location using an asterisk (*). Black circles represent a single site.

768 Figure 5. A Mantel-based spatial autocorrelogram depicting the acoustic similarity of calls
770 against increasing geographic distance. Grey shading indicates the 95% confidence interval
around the estimated correlation coefficient.

Deleted: sampled
Deleted: Wright et al.
Deleted:
Deleted: conducted in 2016 are also included.
Deleted:

Deleted:
Deleted: