

More on approximate methods for large datasets

Mauricio A. Álvarez, PhD

Curso de entrenamiento ArcelorMittal

A comment on notation

- Before, we used $\mathbf{K}_{f,f}$ to refer to the matrix $\mathbf{K}(\mathbf{X}, \mathbf{X})$. In this slides, we will also use \mathbf{K}_{nn} to refer to the same matrix.

Contents

Subset of regressors

Nyström method

Subset of datapoints

Bayesian committee machine

Methods based on inducing variables

Deterministic training conditional (DTC) approximation

Fully Independent Training Conditional (FITC) Approximation

Partially Independent Training Conditional (PITC) Approximation

DTC, FITC and PITC for multiple output GPs

Mean GP predictor (I)

- We can get the GP regression predictive equations if we start by defining

$$f(\mathbf{x}) = \sum_{i=1}^n \alpha_i k(\mathbf{x}, \mathbf{x}_i),$$

where $\alpha = [\alpha_1, \dots, \alpha_n]^\top$ and

$$\alpha \sim \mathcal{N}(\mathbf{0}, \mathbf{K}^{-1}).$$

- We have used $\mathbf{K} = \mathbf{K}(\mathbf{X}, \mathbf{X})$. We will also use \mathbf{K}_{nn} to refer to the kernel matrix $\mathbf{K}(\mathbf{X}, \mathbf{X})$.
- For the Gaussian regression case and with training data $\mathcal{D} = \{\mathbf{X}, \mathbf{y}\}$, the likelihood function is

$$p(\mathbf{y}|\mathbf{X}, \alpha, \sigma_n^2) = \mathcal{N}(\mathbf{y}|\mathbf{K}\alpha, \sigma_n^2\mathbf{I}).$$

Mean GP predictor (II)

- By using the prior on α , $p(\alpha)$, and the likelihood $p(\mathbf{y}|\mathbf{X}, \alpha, \sigma_n^2)$, we get the following posterior for α

$$p(\alpha|\mathbf{y}, \mathbf{X}, \sigma_n^2) = \mathcal{N}(\alpha|\Sigma\mathbf{K}^\top\sigma_n^{-2}\mathbf{y}, \Sigma)$$

where $\Sigma = (\mathbf{K} + \sigma_n^{-2}\mathbf{K}^\top\mathbf{K})^{-1}$.

- \mathbf{K} is a symmetric matrix and

$$\Sigma = (\mathbf{K} + \sigma_n^{-2}\mathbf{K}^\top\mathbf{K})^{-1} = [\sigma_n^{-2}\mathbf{K}(\sigma_n^2\mathbf{I} + \mathbf{K})]^{-1} = (\sigma_n^2\mathbf{I} + \mathbf{K})^{-1}\mathbf{K}^{-1}\sigma_n^2.$$

- The posterior mean in $p(\alpha|\mathbf{y}, \mathbf{X}, \sigma_n^2)$ is simply

$$\bar{\alpha} = \Sigma\mathbf{K}^\top\sigma_n^{-2}\mathbf{y} = (\sigma_n^2\mathbf{I} + \mathbf{K})^{-1}\mathbf{K}^{-1}\sigma_n^2\mathbf{K}\sigma_n^{-2}\mathbf{y} = (\sigma_n^2\mathbf{I} + \mathbf{K})^{-1}\mathbf{y}.$$

Mean GP predictor (III)

- We can now compute the predictive distribution for $p(f(\mathbf{x}_*)|\mathbf{X}, \mathbf{y})$ by marginalising α using the posterior distribution $p(\alpha|\mathbf{y}, \mathbf{X}, \sigma_n^2)$.
- According to what we saw before, $f(\mathbf{x}_*) = \sum_{i=1}^n \alpha_i k(\mathbf{x}_*, \mathbf{x}_i) = \mathbf{k}^\top(\mathbf{x}_*)\alpha$, where $\mathbf{k}^\top(\mathbf{x}_*) = [k(\mathbf{x}_*, \mathbf{x}_1), \dots, k(\mathbf{x}_*, \mathbf{x}_n)]$.
- The mean predictive is then given as

$$\bar{f}(\mathbf{x}_*) = \mathbb{E}(f(\mathbf{x}_*)) = \mathbf{k}^\top(\mathbf{x}_*)\mathbb{E}(\alpha) = \mathbf{k}^\top(\mathbf{x}_*)(\sigma_n^2\mathbf{I} + \mathbf{K})^{-1}\mathbf{y},$$

which corresponds to the expression for the mean prediction in GP regression.

- What if instead of having $f(\mathbf{x}) = \sum_{i=1}^n \alpha_i k(\mathbf{x}, \mathbf{x}_i)$, with $\alpha \sim \mathcal{N}(\mathbf{0}, \mathbf{K}^{-1})$, we use $f(\mathbf{x}) = \sum_{i=1}^m \alpha_i k(\mathbf{x}, \mathbf{x}_i)$ with $\alpha \sim \mathcal{N}(\mathbf{0}, \mathbf{K}_{mm}^{-1})$?

What is \mathbf{K}_{mm} ?

- Several methods, including Subset of Regressors, consider selecting a subset I of the n datapoints.
- The set I has size $m < n$.
- The remaining $n - m$ datapoints form the set R .
- I is the subset of included datapoints whereas R is the set of remaining datapoints.
- The matrix \mathbf{K} can be partitioned as

$$\mathbf{K} = \begin{bmatrix} \mathbf{K}_{mm} & \mathbf{K}_{m(n-m)} \\ \mathbf{K}_{(n-m)m} & \mathbf{K}_{(n-m)(n-m)} \end{bmatrix}$$

- A key difference with the inducing variable methods is that the set I is part of the n datapoints, whereas \mathbf{u} can be any points.

Subset of regressors (I)

- We can consider a subset of regressors $m < n$ such that

$$f_{\text{SR}}(\mathbf{x}) = \sum_{i=1}^m \alpha_i k(\mathbf{x}, \mathbf{x}_i), \quad \alpha \sim \mathcal{N}(\mathbf{0}, \mathbf{K}_{mm}^{-1}).$$

- Following the same procedure than before, the likelihood function is given as

$$p(\mathbf{y}|\mathbf{X}, \alpha, \sigma_n^2) = \mathcal{N}(\mathbf{y}|\mathbf{K}_{nm}\alpha_m, \sigma_n^2\mathbf{I}).$$

- With the prior over α_m and the likelihood $p(\mathbf{y}|\mathbf{X}, \alpha, \sigma_n^2)$, the posterior distribution over α_m follows as

$$p(\alpha_m|\mathbf{y}, \mathbf{X}, \sigma_n^2) = \mathcal{N}(\alpha_m|\Sigma_{mm}\mathbf{K}_{mn}\sigma_n^{-2}\mathbf{y}, \Sigma)$$

where $\Sigma_{mm} = (\mathbf{K}_{mm} + \sigma_n^{-2}\mathbf{K}_{mn}\mathbf{K}_{nm})^{-1} = (\sigma_n^2\mathbf{K}_{mm} + \mathbf{K}_{mn}\mathbf{K}_{nm})^{-1}\sigma_n^2$.

Subset of regressors (II)

- Predictions are made using $f_{\text{SR}}(\mathbf{x}_*) = \mathbf{k}_m^\top(\mathbf{x}_*)\alpha_m$, where

$$\mathbf{k}_m^\top(\mathbf{x}_*) = [k(\mathbf{x}_*, \mathbf{x}_1), \dots, k(\mathbf{x}_*, \mathbf{x}_m)].$$

- Using the posterior distribution $p(\alpha_m | \mathbf{y}, \mathbf{X}, \sigma_n^2)$ to marginalise α_m from $f_{\text{SR}}(\mathbf{x}_*)$, the predictive distribution for $f_{\text{SR}}(\mathbf{x}_*)$ has moments

$$\begin{aligned}\bar{f}_{\text{SR}} &= \mathbf{k}_m^\top(\mathbf{x}_*)(\sigma_n^2 \mathbf{K}_{mm} + \mathbf{K}_{mn} \mathbf{K}_{nm})^{-1} \mathbf{K}_{mn} \mathbf{y} \\ \mathbb{V}[f_{\text{SR}}] &= \sigma_n^2 \mathbf{k}_m^\top(\mathbf{x}_*)(\sigma_n^2 \mathbf{K}_{mm} + \mathbf{K}_{mn} \mathbf{K}_{nm})^{-1} \mathbf{k}_m(\mathbf{x}_*).\end{aligned}$$

- Computational complexity is $\mathcal{O}(nm^2)$.

SR marginal likelihood

- Using

$$\mathbf{f}_{\text{SR}}(\mathbf{x}) = \sum_{i=1}^m \alpha_i k(\mathbf{x}, \mathbf{x}_i), \quad \boldsymbol{\alpha} \sim \mathcal{N}(\mathbf{0}, \mathbf{K}_{mm}^{-1}),$$

the marginal distribution for $p(\mathbf{f}_{\text{SR}}) = \mathcal{N}(\mathbf{f}_{\text{SR}} | \mathbf{0}, \mathbf{K}_{nm} \mathbf{K}_{mm}^{-1} \mathbf{K}_{mn})$.

- Let $\tilde{\mathbf{K}} = \mathbf{K}_{nm} \mathbf{K}_{mm}^{-1} \mathbf{K}_{mn}$.

- The log-marginal likelihood under this model follows as

$$\log p_{\text{SR}}(\mathbf{y} | \mathbf{X}) = -\frac{1}{2} \log |\tilde{\mathbf{K}} + \sigma_n^2 \mathbf{I}| - \frac{1}{2} \mathbf{y}^\top (\tilde{\mathbf{K}} + \sigma_n^2 \mathbf{I})^{-1} \mathbf{y} - \frac{n}{2} \log(2\pi).$$

How to choose l ?

- ❑ Randomly from \mathbf{X} .
- ❑ Run clustering over $\{\mathbf{x}_i\}_{i=1}^n$ and obtain m points that are closest to the m centres.

Contents

Subset of regressors

Nyström method

Subset of datapoints

Bayesian committee machine

Methods based on inducing variables

Deterministic training conditional (DTC) approximation

Fully Independent Training Conditional (FITC) Approximation

Partially Independent Training Conditional (PITC) Approximation

DTC, FITC and PITC for multiple output GPs

Approximation of the eigenfunctions of $k(\mathbf{x}, \mathbf{x}')$

- The Nyström method approximates the i eigenfunction of a kernel function $k(\mathbf{x}, \mathbf{x}')$ using

$$\phi_i(\mathbf{x}) \simeq \frac{\sqrt{n}}{\lambda_i^{\text{mat}}} \mathbf{k}^\top(\mathbf{x}) \mathbf{u}_i,$$

where $\mathbf{k}^\top(\mathbf{x}) = [k(\mathbf{x}_1, \mathbf{x}), \dots, k(\mathbf{x}_n, \mathbf{x})]$ and λ_i^{mat} and \mathbf{u}_i are obtained from solving the matrix eigenproblem

$$\mathbf{K} \mathbf{u}_i = \lambda_i^{\text{mat}} \mathbf{u}_i.$$

- The eigenvectors are normalised $\mathbf{u}_i^\top \mathbf{u}_i = 1$.

Approximation of the eigenvectors of \mathbf{K}

- We compute the eigenvalues/vectors for \mathbf{K}_{mm} , denoted as $\{\lambda_i^{(m)}\}_{i=1}^m$ and $\{\mathbf{u}_i^{(m)}\}_{i=1}^m$.
- We use these to compute the eigenvalues/vectors for \mathbf{K} ,

$$\begin{aligned}\tilde{\lambda}_i^{(n)} &\triangleq \frac{n}{m} \lambda_i^{(m)}, & i = 1, \dots, m \\ \tilde{\mathbf{u}}_i^{(n)} &\triangleq \sqrt{\frac{m}{n}} \frac{1}{\lambda_i^{(m)}} \mathbf{K}_{nm} \mathbf{u}_i^{(m)}, & i = 1, \dots, m\end{aligned}$$

- We approximate \mathbf{K} using

$$\mathbf{K} \approx \tilde{\mathbf{K}} = \sum_{i=1}^p \tilde{\lambda}_i^{(n)} \tilde{\mathbf{u}}_i^{(n)} \left(\tilde{\mathbf{u}}_i^{(n)} \right)^\top.$$

- Setting $p = m$ then leads to

$$\tilde{\mathbf{K}} = \sum_{i=1}^m \tilde{\lambda}_i^{(n)} \tilde{\mathbf{u}}_i^{(n)} \left(\tilde{\mathbf{u}}_i^{(n)} \right)^\top = \mathbf{K}_{nm} \mathbf{K}_{mm}^{-1} \mathbf{K}_{mn}.$$

K by $\tilde{\mathbf{K}}$

- The Nyström method replaces \mathbf{K} by $\tilde{\mathbf{K}} = \mathbf{K}_{nm}\mathbf{K}_{mm}^{-1}\mathbf{K}_{mn}$ in the mean and variance prediction equations of GP regression.
- The original GP predictive distribution $p(f(\mathbf{x}_*)|\mathbf{X}, \mathbf{y})$ has moments

$$\begin{aligned}\bar{f}(\mathbf{x}_*) &= \mathbf{k}^\top(\mathbf{x}_*) [\mathbf{K} + \sigma_n^2 \mathbf{I}]^{-1} \mathbf{y} \\ \mathbb{V}(f(\mathbf{x}_*)) &= k(\mathbf{x}_*, \mathbf{x}_*) - \mathbf{k}^\top(\mathbf{x}_*) [\mathbf{K} + \sigma_n^2 \mathbf{I}]^{-1} \mathbf{k}(\mathbf{x}_*).\end{aligned}$$

- For the Nyström method, the predictive distribution $p(f(\mathbf{x}_*)|\mathbf{X}, \mathbf{y})$ has moments

$$\begin{aligned}\bar{f}_N(\mathbf{x}_*) &= \mathbf{k}^\top(\mathbf{x}_*) [\tilde{\mathbf{K}} + \sigma_n^2 \mathbf{I}]^{-1} \mathbf{y} \\ \mathbb{V}(f_N(\mathbf{x}_*)) &= k(\mathbf{x}_*, \mathbf{x}_*) - \mathbf{k}^\top(\mathbf{x}_*) [\tilde{\mathbf{K}} + \sigma_n^2 \mathbf{I}]^{-1} \mathbf{k}(\mathbf{x}_*).\end{aligned}$$

Contents

Subset of regressors

Nyström method

Subset of datapoints

Bayesian committee machine

Methods based on inducing variables

Deterministic training conditional (DTC) approximation

Fully Independent Training Conditional (FITC) Approximation

Partially Independent Training Conditional (PITC) Approximation

DTC, FITC and PITC for multiple output GPs

Subset of datapoints

- A simple approximation to the full-sample GP predictor is to keep the GP predictor on a smaller subset of size m of the data.
- We can use a greedy algorithm to select which points are taken into the active set I .

Greedy approximation

- The algorithm starts with the active set I being empty, and the set R containing the indices of all training examples.
- On each iteration one index is selected from R and added to I .
- This is achieved by evaluating some criterion Δ and selecting the data point that optimizes this criterion.
- It can be too expensive to evaluate Δ on all points in R , so some working set $J \subset R$ can be chosen instead, usually at random from R .

Algorithm: greedy approximation

```
input:  $m$ , desired size of active set
2: Initialization  $I = \emptyset$ ,  $R = \{1, \dots, n\}$ 
   for  $j := 1 \dots m$  do
4:   Create working set  $J \subseteq R$ 
     Compute  $\Delta_j$  for all  $j \in J$ 
6:    $i = \operatorname{argmax}_{j \in J} \Delta_j$ 
     Update model to include data from example  $i$ 
8:    $I \leftarrow I \cup \{i\}$ ,  $R \leftarrow R \setminus \{i\}$ 
   end for
10: return:  $I$ 
```

Algorithm 8.1: General framework for greedy subset selection. Δ_j is the criterion function evaluated on data point j .

Selection criteria

- The *informative vector machine* (IVM) (Lawrence et al., 2003) efficiently computes the *differential entropy score*

$$\Delta_j \triangleq H[p(f_j)] - H[p^{\text{new}}(f_j)] ,$$

where $H[p(f_j)]$ is the entropy of the Gaussian process at $j \in R$ without including observation j and $H[p^{\text{new}}(f_j)]$ is the entropy at $j \in R$ when including the observation j .

- The *information gain* criterion $\text{KL}(p^{\text{new}}(f_j) \parallel p(f_j))$ can also be used as a selection criterion (Seeger, 2003).

Contents

Subset of regressors

Nyström method

Subset of datapoints

Bayesian committee machine

Methods based on inducing variables

Deterministic training conditional (DTC) approximation

Fully Independent Training Conditional (FITC) Approximation

Partially Independent Training Conditional (PITC) Approximation

DTC, FITC and PITC for multiple output GPs

Split the data into p parts

- Let \mathbf{f}_* be the vector of function values at the test locations.
- The Bayesian committee machine (BCM) splits the dataset into p parts, $\mathcal{D}_1, \dots, \mathcal{D}_p$, where $\mathcal{D}_i = \{\mathbf{X}_i, \mathbf{y}_i\}$.
- It assumes that

$$p(\mathbf{y}_1, \dots, \mathbf{y}_p \mid \mathbf{f}_*, \mathbf{X}) \simeq \prod_{i=1}^p p(\mathbf{y}_i \mid \mathbf{f}_*, \mathbf{X}_i).$$

- The above approximation leads to

$$q(\mathbf{f}_* \mid \mathcal{D}_1, \dots, \mathcal{D}_p) \propto p(\mathbf{f}_*) \prod_{i=1}^p p(\mathbf{y}_i \mid \mathbf{f}_*, \mathbf{X}_i) = c \frac{\prod_{i=1}^p p(\mathbf{f}_* \mid \mathcal{D}_i)}{p^{p-1}(\mathbf{f}_*)},$$

where we have used $p(\mathbf{y}_i \mid \mathbf{f}_*, \mathbf{X}_i) \propto p(\mathbf{f}_* \mid \mathcal{D}_i)/p(\mathbf{f}_*)$ and c is a constant.

Predictive distribution for the BCM (I)

- The numerator and denominator in the expression before only involve Gaussian distributions.
- We can use the technique of *completing the square* to compute the mean and covariance for $q(\mathbf{f}_* | \mathcal{D}_1, \dots, \mathcal{D}_p) = q(\mathbf{f}_* | \mathcal{D})$.

Predictive distribution for the BCM (II)

- It can be shown that the predictive mean and predictive covariance for $q(\mathbf{f}_* | \mathcal{D})$ are given as

$$\mathbb{E}_q[\mathbf{f}_* | \mathcal{D}] = [\text{cov}_q(\mathbf{f}_* | \mathcal{D})] \sum_{i=1}^p [\text{cov}(\mathbf{f}_* | \mathcal{D}_i)]^{-1} \mathbb{E}[\mathbf{f}_* | \mathcal{D}_i]$$

$$[\text{cov}_q(\mathbf{f}_* | \mathcal{D})]^{-1} = -(p-1)\mathbf{K}_{**}^{-1} + \sum_{i=1}^p [\text{cov}(\mathbf{f}_* | \mathcal{D}_i)]^{-1},$$

where \mathbf{K}_{**} corresponds to the covariance matrix at the test points.

- $\mathbb{E}[\mathbf{f}_* | \mathcal{D}_i]$ and $\text{cov}(\mathbf{f}_* | \mathcal{D}_i)$ are computed using the expressions for the predictive distribution in GP regression.

Contents

Subset of regressors

Nyström method

Subset of datapoints

Bayesian committee machine

Methods based on inducing variables

Deterministic training conditional (DTC) approximation

Fully Independent Training Conditional (FITC) Approximation


Partially Independent Training Conditional (PITC) Approximation

DTC, FITC and PITC for multiple output GPs

Inducing variable methods

- There is a set of methods that explicitly introduce additional variables into the GP prior.
- Such additional variables $\mathbf{u} = [u_1, \dots, u_m]^\top$ are known as *inducing variables*.
- These latent variables are values of the GP evaluated at a set of inputs \mathbf{Z} .
- The main idea of these methods is to exploit *conditional independencies* between \mathbf{f} and \mathbf{f}_* based on \mathbf{u} .
- The role of \mathbf{u} is to *induce* dependencies between training and test cases.

Pictorial representation



A comment on notation

- Depending on context, $\mathbf{K}_{\mathbf{u},\mathbf{u}}$ might also be called \mathbf{K}_{mm}

Approximate the likelihood or the prior

- We can introduce these methods by using the inducing variables to exploit conditional dependencies in the likelihood or in the prior.
- In what follows, we will use the unifying view by Quiñonero-Candela and Rasmussen (2005) where the conditional dependencies are in the prior.
- This view assumes the exact likelihood function for GP regression $p(\mathbf{y} \mid \mathbf{f}) = \mathcal{N}(\mathbf{y} \mid \mathbf{f}, \sigma_{\text{noise}}^2 \mathbf{I})$.

Exact prior

- The exact GP prior $p(\mathbf{f}_*, \mathbf{f})$ can be recovered by marginalising \mathbf{u} from $p(\mathbf{f}_*, \mathbf{f}, \mathbf{u})$,

$$p(\mathbf{f}_*, \mathbf{f}) = \int p(\mathbf{f}_*, \mathbf{f}, \mathbf{u}) d\mathbf{u} = \int p(\mathbf{f}_*, \mathbf{f} \mid \mathbf{u}) p(\mathbf{u}) d\mathbf{u}$$

where $p(\mathbf{u}) = \mathcal{N}(\mathbf{u} \mid \mathbf{0}, \mathbf{K}_{\mathbf{u},\mathbf{u}})$.

- The exact prior conditionals $p(\mathbf{f} \mid \mathbf{u})$ and $p(\mathbf{f}_* \mid \mathbf{u})$ are given as

$$\begin{aligned} p(\mathbf{f} \mid \mathbf{u}) &= \mathcal{N}(\mathbf{f} \mid \mathbf{K}_{\mathbf{f},\mathbf{u}} \mathbf{K}_{\mathbf{u},\mathbf{u}}^{-1} \mathbf{u}, \mathbf{K}_{\mathbf{f},\mathbf{f}} - \mathbf{Q}_{\mathbf{f},\mathbf{f}}) \\ p(\mathbf{f}_* \mid \mathbf{u}) &= \mathcal{N}(\mathbf{f}_* \mid \mathbf{K}_{*,\mathbf{u}} \mathbf{K}_{\mathbf{u},\mathbf{u}}^{-1} \mathbf{u}, \mathbf{K}_{*,*} - \mathbf{Q}_{*,*}), \end{aligned}$$

with $\mathbf{Q}_{\mathbf{a},\mathbf{b}} \triangleq \mathbf{K}_{\mathbf{a},\mathbf{u}} \mathbf{K}_{\mathbf{u},\mathbf{u}}^{-1} \mathbf{K}_{\mathbf{u},\mathbf{b}}$.

Approximate prior

- The joint prior $p(\mathbf{f}_*, \mathbf{f})$ can be approximated by $q(\mathbf{f}_*, \mathbf{f})$ using

$$p(\mathbf{f}_*, \mathbf{f}) \simeq q(\mathbf{f}_*, \mathbf{f}) = \int q(\mathbf{f}_* | \mathbf{u}) q(\mathbf{f} | \mathbf{u}) p(\mathbf{u}) d\mathbf{u},$$

where $p(\mathbf{u}) = \mathcal{N}(\mathbf{u} | \mathbf{0}, \mathbf{K}_{\mathbf{u}, \mathbf{u}})$.

- Depending on how we approximate the conditional priors $q(\mathbf{f}_* | \mathbf{u})$ and $q(\mathbf{f} | \mathbf{u})$, we get different type of approximations.

Contents

Subset of regressors

Nyström method

Subset of datapoints

Bayesian committee machine

Methods based on inducing variables

Deterministic training conditional (DTC) approximation

Fully Independent Training Conditional (FITC) Approximation

Partially Independent Training Conditional (PITC) Approximation

DTC, FITC and PITC for multiple output GPs

DTC: priors

- DTC has been introduced as a likelihood approximation method based on inducing inputs, under the names of Projected Latent Variables (PLV) or Projected Process Approximation (PPA).
- In the DTC approximation, the training conditional distribution is deterministic and the test conditional distribution is exact, this is

$$\begin{aligned}q_{\text{DTC}}(\mathbf{f} \mid \mathbf{u}) &= \mathcal{N}(\mathbf{f} \mid \mathbf{K}_{\mathbf{f},\mathbf{u}}\mathbf{K}_{\mathbf{u},\mathbf{u}}^{-1}\mathbf{u}, \mathbf{0}) \\q_{\text{DTC}}(\mathbf{f}_* \mid \mathbf{u}) &= p(\mathbf{f}_* \mid \mathbf{u}) \\&= \mathcal{N}(\mathbf{f}_* \mid \mathbf{K}_{*,\mathbf{u}}\mathbf{K}_{\mathbf{u},\mathbf{u}}^{-1}\mathbf{u}, \mathbf{K}_{*,*} - \mathbf{Q}_{*,*}).\end{aligned}$$

- The joint prior implied by DTC follows as

$$q_{\text{DTC}}(\mathbf{f}, \mathbf{f}_*) = \mathcal{N}\left(\begin{bmatrix} \mathbf{f} \\ \mathbf{f}_* \end{bmatrix} \middle| \mathbf{0}, \begin{bmatrix} \mathbf{Q}_{\mathbf{f},\mathbf{f}} & \mathbf{Q}_{\mathbf{f},*} \\ \mathbf{Q}_{*,\mathbf{f}} & \mathbf{K}_{*,*} \end{bmatrix}\right)$$

DTC: predictive distribution

- Using the Gaussian likelihood model, $p(\mathbf{y} | \mathbf{f}) = \mathcal{N}(\mathbf{y} | \mathbf{f}, \sigma_{\text{noise}}^2 \mathbf{I})$, the predictive distribution follows as

$$q_{\text{DTC}}(\mathbf{f}_* | \mathbf{y}) = \mathcal{N}(\mathbf{f}_* | \boldsymbol{\mu}_{*,\text{DTC}}, \boldsymbol{\Sigma}_{*,\text{DTC}}),$$

where

$$\boldsymbol{\mu}_{*,\text{DTC}} = \mathbf{Q}_{*,\mathbf{f}} (\mathbf{Q}_{\mathbf{f},\mathbf{f}} + \sigma_{\text{noise}}^2 \mathbf{I})^{-1} \mathbf{y} = \sigma^{-2} \mathbf{K}_{*,\mathbf{u}} \boldsymbol{\Sigma} \mathbf{K}_{\mathbf{u},\mathbf{f}} \mathbf{y},$$

$$\boldsymbol{\Sigma}_{*,\text{DTC}} = \mathbf{K}_{*,*} - \mathbf{Q}_{*,\mathbf{f}} (\mathbf{Q}_{\mathbf{f},\mathbf{f}} + \sigma_{\text{noise}}^2 \mathbf{I})^{-1} \mathbf{Q}_{\mathbf{f},*} = \mathbf{K}_{*,*} - \mathbf{Q}_{*,*} + \mathbf{K}_{*,\mathbf{u}} \boldsymbol{\Sigma} \mathbf{K}_{*,\mathbf{u}}^{\top},$$

with $\boldsymbol{\Sigma} = \left(\sigma_{\text{noise}}^{-2} \mathbf{K}_{\mathbf{u},\mathbf{f}} \mathbf{K}_{\mathbf{f},\mathbf{u}} + \mathbf{K}_{\mathbf{u},\mathbf{u}} \right)^{-1}$.

- Comparing against the subset of regressors, the predictive means are the same in both methods. The covariance for the SoR was equal to $\mathbf{K}_{*,\mathbf{u}} \boldsymbol{\Sigma} \mathbf{K}_{*,\mathbf{u}}^{\top}$.
- Because $\mathbf{K}_{*,*} - \mathbf{Q}_{*,*}$ is always positive, the predictive variance in DTC is larger than the SoR's predictive variance.;

Contents

Subset of regressors

Nyström method

Subset of datapoints

Bayesian committee machine

Methods based on inducing variables

Deterministic training conditional (DTC) approximation

Fully Independent Training Conditional (FITC) Approximation

Partially Independent Training Conditional (PITC) Approximation


DTC, FITC and PITC for multiple output GPs

FITC: priors

- FITC has been introduced as a likelihood approximation method based on inducing inputs, under the name Sparse Gaussian Processes using Pseudo-inputs (SGPP) by Snelson and Ghahramani (2005).
- In the FITC approximation, the training conditional distribution includes a variance term and the test conditional distribution is exact, this is

$$q_{\text{FITC}}(\mathbf{f} \mid \mathbf{u}) = \prod_{i=1}^n p(f_i \mid \mathbf{u}) = \mathcal{N}(\mathbf{f} \mid \mathbf{K}_{\mathbf{f},\mathbf{u}} \mathbf{K}_{\mathbf{u},\mathbf{u}}^{-1} \mathbf{u}, \text{diag}[\mathbf{K}_{\mathbf{f},\mathbf{f}} - \mathbf{Q}_{\mathbf{f},\mathbf{f}}])$$
$$q_{\text{FITC}}(\mathbf{f}_* \mid \mathbf{u}) = p(\mathbf{f}_* \mid \mathbf{u}) = \mathcal{N}(\mathbf{f}_* \mid \mathbf{K}_{*,\mathbf{u}} \mathbf{K}_{\mathbf{u},\mathbf{u}}^{-1} \mathbf{u}, \mathbf{K}_{*,*} - \mathbf{Q}_{*,*}).$$

Pictorial representation



FITC vs DTC

- The joint prior implied by FITC follows as

$$q_{\text{FITC}}(\mathbf{f}, \mathbf{f}_*) = \mathcal{N} \left(\begin{bmatrix} \mathbf{f} \\ \mathbf{f}_* \end{bmatrix} \middle| \mathbf{0}, \begin{bmatrix} \mathbf{Q}_{\mathbf{f},\mathbf{f}} - \text{diag}[\mathbf{Q}_{\mathbf{f},\mathbf{f}} - \mathbf{K}_{\mathbf{f},\mathbf{f}}] & \mathbf{Q}_{\mathbf{f},*} \\ \mathbf{Q}_{*,\mathbf{f}} & \mathbf{K}_{*,*} \end{bmatrix} \right)$$

- From before, the joint prior implied by DTC was

$$q_{\text{DTC}}(\mathbf{f}, \mathbf{f}_*) = \mathcal{N} \left(\begin{bmatrix} \mathbf{f} \\ \mathbf{f}_* \end{bmatrix} \middle| \mathbf{0}, \begin{bmatrix} \mathbf{Q}_{\mathbf{f},\mathbf{f}} & \mathbf{Q}_{\mathbf{f},*} \\ \mathbf{Q}_{*,\mathbf{f}} & \mathbf{K}_{*,*} \end{bmatrix} \right)$$

- Compared to DTC, FITC uses the exact covariance function in the main diagonal.

FITC: predictive distribution

- Using the Gaussian likelihood model, $p(\mathbf{y} | \mathbf{f}) = \mathcal{N}(\mathbf{y} | \mathbf{f}, \sigma_{\text{noise}}^2 \mathbf{I})$, the predictive distribution follows as

$$q_{\text{FITC}}(\mathbf{f}_* | \mathbf{y}) = \mathcal{N}(\mathbf{f}_* | \boldsymbol{\mu}_{*,\text{FITC}}, \boldsymbol{\Sigma}_{*,\text{FITC}}),$$

where

$$\boldsymbol{\mu}_{*,\text{FITC}} = \mathbf{Q}_{*,\mathbf{f}} (\mathbf{Q}_{\mathbf{f},\mathbf{f}} + \Lambda)^{-1} \mathbf{y} = \mathbf{K}_{*,\mathbf{u}} \boldsymbol{\Sigma} \mathbf{K}_{\mathbf{u},\mathbf{f}} \Lambda^{-1} \mathbf{y},$$

$$\boldsymbol{\Sigma}_{*,\text{FITC}} = \mathbf{K}_{*,*} - \mathbf{Q}_{*,\mathbf{f}} (\mathbf{Q}_{\mathbf{f},\mathbf{f}} + \Lambda)^{-1} \mathbf{Q}_{\mathbf{f},*} = \mathbf{K}_{*,*} - \mathbf{Q}_{*,*} + \mathbf{K}_{*,\mathbf{u}} \boldsymbol{\Sigma} \mathbf{K}_{*,\mathbf{u}}^{\top},$$

with $\boldsymbol{\Sigma} = (\mathbf{K}_{\mathbf{u},\mathbf{f}} \Lambda^{-1} \mathbf{K}_{\mathbf{f},\mathbf{u}} + \mathbf{K}_{\mathbf{u},\mathbf{u}})^{-1}$ and $\Lambda = \text{diag}[\mathbf{K}_{\mathbf{f},\mathbf{f}} - \mathbf{Q}_{\mathbf{f},\mathbf{f}} + \sigma_{\text{noise}}^2 \mathbf{I}]$.

Contents

Subset of regressors

Nyström method

Subset of datapoints

Bayesian committee machine

Methods based on inducing variables

Deterministic training conditional (DTC) approximation

Fully Independent Training Conditional (FITC) Approximation

Partially Independent Training Conditional (PITC) Approximation

DTC, FITC and PITC for multiple output GPs


PITC: priors

- In the PITC approximation, the training conditional distribution has a block-diagonal covariance and the test conditional distribution is exact, this is

$$q_{\text{PITC}}(\mathbf{f} \mid \mathbf{u}) = \mathcal{N}(\mathbf{f} \mid \mathbf{K}_{\mathbf{f},\mathbf{u}}\mathbf{K}_{\mathbf{u},\mathbf{u}}^{-1}\mathbf{u}, \text{blockdiag}[\mathbf{K}_{\mathbf{f},\mathbf{f}} - \mathbf{Q}_{\mathbf{f},\mathbf{f}}])$$
$$q_{\text{PITC}}(\mathbf{f}_* \mid \mathbf{u}) = p(\mathbf{f}_* \mid \mathbf{u}) = \mathcal{N}(\mathbf{f}_* \mid \mathbf{K}_{*,\mathbf{u}}\mathbf{K}_{\mathbf{u},\mathbf{u}}^{-1}\mathbf{u}, \mathbf{K}_{*,*} - \mathbf{Q}_{*,*}),$$

where $\text{blockdiag}[A]$ is a block diagonal matrix. The blocks have not been specified.

Pictorial representation



PITC: prior

- The joint prior implied by PITC follows as

$$q_{\text{PITC}}(\mathbf{f}, \mathbf{f}_*) = \mathcal{N} \left(\begin{bmatrix} \mathbf{f} \\ \mathbf{f}_* \end{bmatrix} \middle| \mathbf{0}, \begin{bmatrix} \mathbf{Q}_{\mathbf{f},\mathbf{f}} - \text{blockdiag}[\mathbf{Q}_{\mathbf{f},\mathbf{f}} - \mathbf{K}_{\mathbf{f},\mathbf{f}}] & \mathbf{Q}_{\mathbf{f},*} \\ \mathbf{Q}_{*,\mathbf{f}} & \mathbf{K}_{*,*} \end{bmatrix} \right)$$

PITC: predictive distribution

- Using the Gaussian likelihood model, $p(\mathbf{y} | \mathbf{f}) = \mathcal{N}(\mathbf{y} | \mathbf{f}, \sigma_{\text{noise}}^2 \mathbf{I})$, the predictive distribution follows as

$$q_{\text{PITC}}(\mathbf{f}_* | \mathbf{y}) = \mathcal{N}(\mathbf{f}_* | \boldsymbol{\mu}_{*,\text{PITC}}, \boldsymbol{\Sigma}_{*,\text{PITC}}),$$

where

$$\begin{aligned}\boldsymbol{\mu}_{*,\text{PITC}} &= \mathbf{Q}_{*,\mathbf{f}} (\mathbf{Q}_{\mathbf{f},\mathbf{f}} + \boldsymbol{\Lambda})^{-1} \mathbf{y} = \mathbf{K}_{*,\mathbf{u}} \boldsymbol{\Sigma} \mathbf{K}_{\mathbf{u},\mathbf{f}} \boldsymbol{\Lambda}^{-1} \mathbf{y}, \\ \boldsymbol{\Sigma}_{*,\text{PITC}} &= \mathbf{K}_{*,*} - \mathbf{Q}_{*,\mathbf{f}} (\mathbf{Q}_{\mathbf{f},\mathbf{f}} + \boldsymbol{\Lambda})^{-1} \mathbf{Q}_{\mathbf{f},*} = \mathbf{K}_{*,*} - \mathbf{Q}_{*,*} + \mathbf{K}_{*,\mathbf{u}} \boldsymbol{\Sigma} \mathbf{K}_{*,\mathbf{u}}^{\top},\end{aligned}$$

with $\boldsymbol{\Sigma} = (\mathbf{K}_{\mathbf{u},\mathbf{f}} \boldsymbol{\Lambda}^{-1} \mathbf{K}_{\mathbf{f},\mathbf{u}} + \mathbf{K}_{\mathbf{u},\mathbf{u}})^{-1}$ and
 $\boldsymbol{\Lambda} = \text{blockdiag}[\mathbf{K}_{\mathbf{f},\mathbf{f}} - \mathbf{Q}_{\mathbf{f},\mathbf{f}} + \sigma_{\text{noise}}^2 \mathbf{I}]$.

Contents

Subset of regressors

Nyström method

Subset of datapoints

Bayesian committee machine

Methods based on inducing variables

Deterministic training conditional (DTC) approximation

Fully Independent Training Conditional (FITC) Approximation

Partially Independent Training Conditional (PITC) Approximation

DTC, FITC and PITC for multiple output GPs

Marginal likelihood of the full Gaussian process.

- The marginal likelihood of the model is given by

$$p(\mathbf{y}|\mathbf{X}, \phi) = \mathcal{N}(\mathbf{0}, \mathbf{K}_{\mathbf{f},\mathbf{f}} + \Sigma)$$

where $\mathbf{y} = [\mathbf{y}_1^\top, \dots, \mathbf{y}_D^\top]^\top$ is the set of output functions, $\mathbf{K}_{\mathbf{f},\mathbf{f}}$ covariance matrix with blocks $\text{cov}[f_d, f_{d'}]$, Σ matrix of noise variances, ϕ is the set of parameters of the covariance matrix and $\mathbf{X} = \{\mathbf{x}_1, \dots, \mathbf{x}_N\}$ is the set of input vectors.

- Learning from the log-likelihood involves the inverse of $\mathbf{K}_{\mathbf{f},\mathbf{f}} + \Sigma$, which grows with complexity $\mathcal{O}(N^3 D^3)$

Predictive distribution of the full Gaussian process.

- Predictive distribution at \mathbf{X}_*

$$p(\mathbf{y}_* | \mathbf{y}, \mathbf{X}, \mathbf{X}_*, \phi) = \mathcal{N}(\boldsymbol{\mu}_*, \boldsymbol{\Lambda}_*)$$

with


$$\boldsymbol{\mu}_* = \mathbf{K}_{\mathbf{f}_*, \mathbf{f}}(\mathbf{K}_{\mathbf{f}, \mathbf{f}} + \boldsymbol{\Sigma})^{-1} \mathbf{y}$$

$$\boldsymbol{\Lambda}_* = \mathbf{K}_{\mathbf{f}_*, \mathbf{f}_*} - \mathbf{K}_{\mathbf{f}_*, \mathbf{f}}(\mathbf{K}_{\mathbf{f}, \mathbf{f}} + \boldsymbol{\Sigma})^{-1} \mathbf{K}_{\mathbf{f}, \mathbf{f}_*} + \boldsymbol{\Sigma}$$

- Prediction is $\mathcal{O}(ND)$ for the mean and $\mathcal{O}(N^2 D^2)$ for the variance.

Conditional prior distribution.


Sample from $p(u)$



$$f_d(\mathbf{x}) = \int_{\mathcal{X}} G_d(\mathbf{x} - \mathbf{z}) u(\mathbf{z}) d\mathbf{z}$$


Conditional prior distribution.

Sample from $p(u)$



$$f_d(\mathbf{x}) = \int_{\mathcal{X}} G_d(\mathbf{x} - \mathbf{z}) u(\mathbf{z}) d\mathbf{z}$$


Discretize u



$$f_d(\mathbf{x}) \approx \sum_{\forall k} G_d(\mathbf{x} - \mathbf{z}_k) u(\mathbf{z}_k)$$


Conditional prior distribution.

Sample from $p(u)$




$$f_d(\mathbf{x}) = \int_{\mathcal{X}} G_d(\mathbf{x} - \mathbf{z}) u(\mathbf{z}) d\mathbf{z}$$

Discretize u



$$f_d(\mathbf{x}) \approx \sum_{\forall k} G_d(\mathbf{x} - \mathbf{z}_k) u(\mathbf{z}_k)$$

Sample from $p(u|\mathbf{u})$



$$f_d(\mathbf{x}) \approx \int_{\mathcal{X}} G_d(\mathbf{x} - \mathbf{z}) \mathbb{E}[u(\mathbf{z})|\mathbf{u}] d\mathbf{z}$$

The conditional independence assumption (I)

- This form for $f_d(\mathbf{x})$ leads to the following likelihood

$$p(\mathbf{f}|\mathbf{u}, \mathbf{Z}) = \mathcal{N}(\mathbf{f}|\mathbf{K}_{\mathbf{f},\mathbf{u}}\mathbf{K}_{\mathbf{u},\mathbf{u}}^{-1}\mathbf{u}, \mathbf{K}_{\mathbf{f},\mathbf{f}} - \mathbf{K}_{\mathbf{f},\mathbf{u}}\mathbf{K}_{\mathbf{u},\mathbf{u}}^{-1}\mathbf{K}_{\mathbf{u},\mathbf{f}}),$$

where

- \mathbf{u} discrete sample from the latent function
- \mathbf{Z} set of input vectors corresponding to \mathbf{u}
- $\mathbf{K}_{\mathbf{u},\mathbf{u}}$ cross-covariance matrix between latent functions
- $\mathbf{K}_{\mathbf{f},\mathbf{u}} = \mathbf{K}_{\mathbf{u},\mathbf{f}}^\top$ cross-covariance matrix between latent and output functions

- Even though we conditioned on \mathbf{u} , we still have dependencies between outputs due to the uncertainty in $p(u|\mathbf{u})$.

The conditional independence assumption (II)

Our key assumption is that the outputs will be independent even if we have only observed \mathbf{u} rather than the whole function u .

$\mathbf{K}_{f_1 f_1} - \mathbf{K}_{f_1 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_1}$	$\mathbf{K}_{f_1 f_2} - \mathbf{K}_{f_1 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_2}$	$\mathbf{K}_{f_1 f_3} - \mathbf{K}_{f_1 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_3}$
$\mathbf{K}_{f_2 f_1} - \mathbf{K}_{f_2 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_1}$	$\mathbf{K}_{f_2 f_2} - \mathbf{K}_{f_2 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_2}$	$\mathbf{K}_{f_2 f_3} - \mathbf{K}_{f_2 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_3}$
$\mathbf{K}_{f_3 f_1} - \mathbf{K}_{f_3 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_1}$	$\mathbf{K}_{f_3 f_2} - \mathbf{K}_{f_3 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_2}$	$\mathbf{K}_{f_3 f_3} - \mathbf{K}_{f_3 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_3}$

The conditional independence assumption (II)

Our key assumption is that the outputs will be independent even if we have only observed \mathbf{u} rather than the whole function u .

$\mathbf{K}_{f_1 f_1} - \mathbf{K}_{f_1 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{u f_1}$	0	0
0	$\mathbf{K}_{f_2 f_2} - \mathbf{K}_{f_2 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{u f_2}$	0
0	0	$\mathbf{K}_{f_3 f_3} - \mathbf{K}_{f_3 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{u f_3}$

Better approximations can be obtained when $E[u|\mathbf{u}]$ approximates u .

Comparison of marginal likelihoods

Integrating out \mathbf{u} , the marginal likelihood is given as

$$p(\mathbf{y}|\mathbf{Z}, \mathbf{X}, \theta) = \mathcal{N}(\mathbf{y}|\mathbf{0}, \mathbf{K}_{\mathbf{f},\mathbf{u}}\mathbf{K}_{\mathbf{u},\mathbf{u}}^{-1}\mathbf{K}_{\mathbf{u},\mathbf{f}} + \text{blockdiag}[\mathbf{K}_{\mathbf{f},\mathbf{f}} - \mathbf{K}_{\mathbf{f},\mathbf{u}}\mathbf{K}_{\mathbf{u},\mathbf{u}}^{-1}\mathbf{K}_{\mathbf{u},\mathbf{f}}] + \Sigma).$$

Comparison of marginal likelihoods

Integrating out \mathbf{u} , the marginal likelihood is given as

$$p(\mathbf{y}|\mathbf{Z}, \mathbf{X}, \theta) = \mathcal{N}(\mathbf{y}|\mathbf{0}, \mathbf{K}_{f,u}\mathbf{K}_{u,u}^{-1}\mathbf{K}_{u,f} + \text{blockdiag}[\mathbf{K}_{f,f} - \mathbf{K}_{f,u}\mathbf{K}_{u,u}^{-1}\mathbf{K}_{u,f}] + \Sigma).$$

$\mathbf{K}_{f_1 f_1}$	$\mathbf{K}_{f_1 f_2}$	$\mathbf{K}_{f_1 f_3}$
$\mathbf{K}_{f_2 f_1}$	$\mathbf{K}_{f_2 f_2}$	$\mathbf{K}_{f_2 f_3}$
$\mathbf{K}_{f_3 f_1}$	$\mathbf{K}_{f_3 f_2}$	$\mathbf{K}_{f_3 f_3}$

 \approx

$\mathbf{K}_{f_1 f_1}$	$\mathbf{K}_{f_1 u}\mathbf{K}_{u u}^{-1}\mathbf{K}_{u f_2}$	$\mathbf{K}_{f_1 u}\mathbf{K}_{u u}^{-1}\mathbf{K}_{u f_3}$
$\mathbf{K}_{f_2 u}\mathbf{K}_{u u}^{-1}\mathbf{K}_{u f_1}$	$\mathbf{K}_{f_2 f_2}$	$\mathbf{K}_{f_2 u}\mathbf{K}_{u u}^{-1}\mathbf{K}_{u f_3}$
$\mathbf{K}_{f_3 u}\mathbf{K}_{u u}^{-1}\mathbf{K}_{u f_1}$	$\mathbf{K}_{f_3 u}\mathbf{K}_{u u}^{-1}\mathbf{K}_{u f_2}$	$\mathbf{K}_{f_3 f_3}$

Comparison of marginal likelihoods

Integrating out \mathbf{u} , the marginal likelihood is given as

$$p(\mathbf{y}|\mathbf{Z}, \mathbf{X}, \theta) = \mathcal{N}(\mathbf{y}|\mathbf{0}, \mathbf{K}_{f,u}\mathbf{K}_{u,u}^{-1}\mathbf{K}_{u,f} + \text{blockdiag}[\mathbf{K}_{f,f} - \mathbf{K}_{f,u}\mathbf{K}_{u,u}^{-1}\mathbf{K}_{u,f}] + \Sigma).$$

$\mathbf{K}_{f_1 f_1}$	$\mathbf{K}_{f_1 f_2}$	$\mathbf{K}_{f_1 f_3}$
$\mathbf{K}_{f_2 f_1}$	$\mathbf{K}_{f_2 f_2}$	$\mathbf{K}_{f_2 f_3}$
$\mathbf{K}_{f_3 f_1}$	$\mathbf{K}_{f_3 f_2}$	$\mathbf{K}_{f_3 f_3}$

 \approx

$\mathbf{K}_{f_1 f_1}$	$\mathbf{K}_{f_1 u}\mathbf{K}_{u u}^{-1}\mathbf{K}_{u f_2}$	$\mathbf{K}_{f_1 u}\mathbf{K}_{u u}^{-1}\mathbf{K}_{u f_3}$
$\mathbf{K}_{f_2 u}\mathbf{K}_{u u}^{-1}\mathbf{K}_{u f_1}$	$\mathbf{K}_{f_2 f_2}$	$\mathbf{K}_{f_2 u}\mathbf{K}_{u u}^{-1}\mathbf{K}_{u f_3}$
$\mathbf{K}_{f_3 u}\mathbf{K}_{u u}^{-1}\mathbf{K}_{u f_1}$	$\mathbf{K}_{f_3 u}\mathbf{K}_{u u}^{-1}\mathbf{K}_{u f_2}$	$\mathbf{K}_{f_3 f_3}$

$\mathbf{K}_{f_1 f_1}$	$\mathbf{K}_{f_1 f_2}$	$\mathbf{K}_{f_1 f_3}$
$\mathbf{K}_{f_2 f_1}$	$\mathbf{K}_{f_2 f_2}$	$\mathbf{K}_{f_2 f_3}$
$\mathbf{K}_{f_3 f_1}$	$\mathbf{K}_{f_3 f_2}$	$\mathbf{K}_{f_3 f_3}$

 \approx

\mathbf{G}

 \mathbf{X}

\mathbf{G}^T

Discrete case $[\mathbf{G}]_{i,k} = G_d(\mathbf{x}_i - \mathbf{z}_k)$

Predictive distribution for the sparse approximation

Predictive distribution

$$p(\mathbf{y}_* | \mathbf{y}, \mathbf{X}, \mathbf{X}_*, \mathbf{Z}, \theta) = \mathcal{N}(\tilde{\boldsymbol{\mu}}_*, \tilde{\boldsymbol{\Lambda}}_*) , \text{ with}$$

$$\tilde{\boldsymbol{\mu}}_* = \mathbf{K}_{\mathbf{f}_*, \mathbf{u}} \mathbf{A}^{-1} \mathbf{K}_{\mathbf{u}, \mathbf{f}} (\mathbf{D} + \boldsymbol{\Sigma})^{-1} \mathbf{y}$$

$$\tilde{\boldsymbol{\Lambda}}_* = \mathbf{D}_* + \mathbf{K}_{\mathbf{f}_*, \mathbf{u}} \mathbf{A}^{-1} \mathbf{K}_{\mathbf{u}, \mathbf{f}_*} + \boldsymbol{\Sigma}$$

$$\mathbf{A} = \mathbf{K}_{\mathbf{u}, \mathbf{u}} + \mathbf{K}_{\mathbf{u}, \mathbf{f}} (\mathbf{D} + \boldsymbol{\Sigma})^{-1} \mathbf{K}_{\mathbf{f}, \mathbf{u}}$$

$$\mathbf{D}_* = \text{blockdiag} [\mathbf{K}_{\mathbf{f}_*, \mathbf{f}_*} - \mathbf{K}_{\mathbf{f}_*, \mathbf{u}} \mathbf{K}_{\mathbf{u}, \mathbf{u}}^{-1} \mathbf{K}_{\mathbf{u}, \mathbf{f}_*}]$$

Remarks


- For learning the computational demand is in the calculation of the block-diagonal term which grows as $\mathcal{O}(N^3 D) + \mathcal{O}(NDM^2)$ (with $Q = 1$). Storage is $\mathcal{O}(N^2 D) + \mathcal{O}(NDM)$.
- For inference, the computation of the mean grows as $\mathcal{O}(DM)$ and the computation of the variance as $\mathcal{O}(DM^2)$, after some pre-computations and for one test point.
- The functional form of the approximation is almost identical to that of the Partially Independent Training Conditional (PITC) approximation Quiñero-Candela and Rasmussen (2005).

Additional conditional independencies

- The N^3 term in the computational complexity and the N^2 term in storage in PITC are still expensive for larger data sets.
- An additional assumption is independence over the data points.


Additional conditional independencies

- The N^3 term in the computational complexity and the N^2 term in storage in PITC are still expensive for larger data sets.
- An additional assumption is independence over the data points.



Additional conditional independencies

- The N^3 term in the computational complexity and the N^2 term in storage in PITC are still expensive for larger data sets.
- An additional assumption is independence over the data points.



Comparison of marginal likelihoods

The marginal likelihood is given as

$$p(\mathbf{y}|\mathbf{Z}, \mathbf{X}, \theta) = \mathcal{N}(\mathbf{0}, \mathbf{K}_{f,u} \mathbf{K}_{u,u}^{-1} \mathbf{K}_{u,f} + \text{diag}[\mathbf{K}_{f,f} - \mathbf{K}_{f,u} \mathbf{K}_{u,u}^{-1} \mathbf{K}_{u,f}] + \Sigma) .$$

Comparison of marginal likelihoods

The marginal likelihood is given as

$$p(\mathbf{y}|\mathbf{Z}, \mathbf{X}, \theta) = \mathcal{N}(\mathbf{0}, \mathbf{K}_{f,u} \mathbf{K}_{u,u}^{-1} \mathbf{K}_{u,f} + \text{diag}[\mathbf{K}_{f,f} - \mathbf{K}_{f,u} \mathbf{K}_{u,u}^{-1} \mathbf{K}_{u,f}] + \Sigma).$$

$\mathbf{K}_{f_1 f_1}$	$\mathbf{K}_{f_1 f_2}$	$\mathbf{K}_{f_1 f_3}$
$\mathbf{K}_{f_2 f_1}$	$\mathbf{K}_{f_2 f_2}$	$\mathbf{K}_{f_2 f_3}$
$\mathbf{K}_{f_3 f_1}$	$\mathbf{K}_{f_3 f_2}$	$\mathbf{K}_{f_3 f_3}$

 \approx

$\mathbf{K}_{f_1 f_1}$	$\mathbf{K}_{f_1 u} \mathbf{K}_{u u}^{-1} \mathbf{K}_{u f_2}$	$\mathbf{K}_{f_1 u} \mathbf{K}_{u u}^{-1} \mathbf{K}_{u f_3}$
$\mathbf{K}_{f_2 u} \mathbf{K}_{u u}^{-1} \mathbf{K}_{u f_1}$	$\mathbf{K}_{f_2 f_2}$	$\mathbf{K}_{f_2 u} \mathbf{K}_{u u}^{-1} \mathbf{K}_{u f_3}$
$\mathbf{K}_{f_3 u} \mathbf{K}_{u u}^{-1} \mathbf{K}_{u f_1}$	$\mathbf{K}_{f_3 u} \mathbf{K}_{u u}^{-1} \mathbf{K}_{u f_2}$	$\mathbf{K}_{f_3 f_3}$

Comparison of marginal likelihoods

The marginal likelihood is given as

$$p(\mathbf{y}|\mathbf{Z}, \mathbf{X}, \theta) = \mathcal{N}(\mathbf{0}, \mathbf{K}_{f,u} \mathbf{K}_{u,u}^{-1} \mathbf{K}_{u,f} + \text{diag}[\mathbf{K}_{f,f} - \mathbf{K}_{f,u} \mathbf{K}_{u,u}^{-1} \mathbf{K}_{u,f}] + \Sigma)$$

$\mathbf{K}_{f_1 f_1}$	$\mathbf{K}_{f_1 f_2}$	$\mathbf{K}_{f_1 f_3}$	\approx	$\mathbf{K}_{f_1 f_1}$	$\mathbf{K}_{f_1 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_2}$	$\mathbf{K}_{f_1 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_3}$
$\mathbf{K}_{f_2 f_1}$	$\mathbf{K}_{f_2 f_2}$	$\mathbf{K}_{f_2 f_3}$		$\mathbf{K}_{f_2 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_1}$	$\mathbf{K}_{f_2 f_2}$	$\mathbf{K}_{f_2 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_3}$
$\mathbf{K}_{f_3 f_1}$	$\mathbf{K}_{f_3 f_2}$	$\mathbf{K}_{f_3 f_3}$		$\mathbf{K}_{f_3 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_1}$	$\mathbf{K}_{f_3 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_2}$	$\mathbf{K}_{f_3 f_3}$

$\mathbf{K}_{f_1 f_1}$	$\mathbf{K}_{f_1 f_2}$	$\mathbf{K}_{f_1 f_3}$	\approx	$\mathbf{Q}_{f_1 f_1}$	$\mathbf{K}_{f_1 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_2}$	$\mathbf{K}_{f_1 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_3}$
$\mathbf{K}_{f_2 f_1}$	$\mathbf{K}_{f_2 f_2}$	$\mathbf{K}_{f_2 f_3}$		$\mathbf{K}_{f_2 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_1}$	$\mathbf{Q}_{f_2 f_2}$	$\mathbf{K}_{f_2 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_3}$
$\mathbf{K}_{f_3 f_1}$	$\mathbf{K}_{f_3 f_2}$	$\mathbf{K}_{f_3 f_3}$		$\mathbf{K}_{f_3 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_1}$	$\mathbf{K}_{f_3 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_2}$	$\mathbf{Q}_{f_3 f_3}$

Comparison of marginal likelihoods

The marginal likelihood is given as

$$p(\mathbf{y}|\mathbf{Z}, \mathbf{X}, \theta) = \mathcal{N}(\mathbf{0}, \mathbf{K}_{f,u} \mathbf{K}_{u,u}^{-1} \mathbf{K}_{u,f} + \text{diag}[\mathbf{K}_{f,f} - \mathbf{K}_{f,u} \mathbf{K}_{u,u}^{-1} \mathbf{K}_{u,f}] + \Sigma)$$

$\mathbf{K}_{f_1 f_1}$	$\mathbf{K}_{f_1 f_2}$	$\mathbf{K}_{f_1 f_3}$	\approx	$\mathbf{K}_{f_1 f_1}$	$\mathbf{K}_{f_1 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_2}$	$\mathbf{K}_{f_1 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_3}$
$\mathbf{K}_{f_2 f_1}$	$\mathbf{K}_{f_2 f_2}$	$\mathbf{K}_{f_2 f_3}$		$\mathbf{K}_{f_2 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_1}$	$\mathbf{K}_{f_2 f_2}$	$\mathbf{K}_{f_2 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_3}$
$\mathbf{K}_{f_3 f_1}$	$\mathbf{K}_{f_3 f_2}$	$\mathbf{K}_{f_3 f_3}$		$\mathbf{K}_{f_3 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_1}$	$\mathbf{K}_{f_3 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_2}$	$\mathbf{K}_{f_3 f_3}$

$\mathbf{K}_{f_1 f_1}$	$\mathbf{K}_{f_1 f_2}$	$\mathbf{K}_{f_1 f_3}$	\approx	$\mathbf{Q}_{f_1 f_1}$	$\mathbf{K}_{f_1 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_2}$	$\mathbf{K}_{f_1 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_3}$
$\mathbf{K}_{f_2 f_1}$	$\mathbf{K}_{f_2 f_2}$	$\mathbf{K}_{f_2 f_3}$		$\mathbf{K}_{f_2 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_1}$	$\mathbf{Q}_{f_2 f_2}$	$\mathbf{K}_{f_2 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_3}$
$\mathbf{K}_{f_3 f_1}$	$\mathbf{K}_{f_3 f_2}$	$\mathbf{K}_{f_3 f_3}$		$\mathbf{K}_{f_3 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_1}$	$\mathbf{K}_{f_3 u} \mathbf{K}_{uu}^{-1} \mathbf{K}_{uf_2}$	$\mathbf{Q}_{f_3 f_3}$

Comparison of marginal likelihoods

The marginal likelihood is given as

$$p(\mathbf{y}|\mathbf{Z}, \mathbf{X}, \theta) = \mathcal{N}(\mathbf{0}, \mathbf{K}_{f,u} \mathbf{K}_{u,u}^{-1} \mathbf{K}_{u,f} + \text{diag}[\mathbf{K}_{f,f} - \mathbf{K}_{f,u} \mathbf{K}_{u,u}^{-1} \mathbf{K}_{u,f}] + \Sigma)$$

$\mathbf{K}_{f_1 f_1}(\mathbf{x}_1, \mathbf{x}_1)$	$(\mathbf{K}_{f_1 u} \mathbf{K}_{u u}^{-1} \mathbf{K}_{u f_1})(\mathbf{x}_1, \mathbf{x}_2)$	$(\mathbf{K}_{f_1 u} \mathbf{K}_{u u}^{-1} \mathbf{K}_{u f_1})(\mathbf{x}_1, \mathbf{x}_3)$
$(\mathbf{K}_{f_1 u} \mathbf{K}_{u u}^{-1} \mathbf{K}_{u f_1})(\mathbf{x}_2, \mathbf{x}_1)$	$\mathbf{K}_{f_1 f_1}(\mathbf{x}_2, \mathbf{x}_2)$	$(\mathbf{K}_{f_1 u} \mathbf{K}_{u u}^{-1} \mathbf{K}_{u f_1})(\mathbf{x}_2, \mathbf{x}_3)$
$(\mathbf{K}_{f_1 u} \mathbf{K}_{u u}^{-1} \mathbf{K}_{u f_1})(\mathbf{x}_3, \mathbf{x}_1)$	$(\mathbf{K}_{f_1 u} \mathbf{K}_{u u}^{-1} \mathbf{K}_{u f_1})(\mathbf{x}_3, \mathbf{x}_2)$	$\mathbf{K}_{f_1 f_1}(\mathbf{x}_3, \mathbf{x}_3)$



$\mathbf{K}_{f_1 f_1}$	$\mathbf{K}_{f_1 f_2}$	$\mathbf{K}_{f_1 f_3}$
$\mathbf{K}_{f_2 f_1}$	$\mathbf{K}_{f_2 f_2}$	$\mathbf{K}_{f_2 f_3}$
$\mathbf{K}_{f_3 f_1}$	$\mathbf{K}_{f_3 f_2}$	$\mathbf{K}_{f_3 f_3}$

\approx

$\mathbf{Q}_{f_1 f_1}$	$\mathbf{K}_{f_1 u} \mathbf{K}_{u u}^{-1} \mathbf{K}_{u f_2}$	$\mathbf{K}_{f_1 u} \mathbf{K}_{u u}^{-1} \mathbf{K}_{u f_3}$
$\mathbf{K}_{f_2 u} \mathbf{K}_{u u}^{-1} \mathbf{K}_{u f_1}$	$\mathbf{Q}_{f_2 f_2}$	$\mathbf{K}_{f_2 u} \mathbf{K}_{u u}^{-1} \mathbf{K}_{u f_3}$
$\mathbf{K}_{f_3 u} \mathbf{K}_{u u}^{-1} \mathbf{K}_{u f_1}$	$\mathbf{K}_{f_3 u} \mathbf{K}_{u u}^{-1} \mathbf{K}_{u f_2}$	$\mathbf{Q}_{f_3 f_3}$

Comparison of marginal likelihoods

The marginal likelihood is given as

$$p(\mathbf{y}|\mathbf{Z}, \mathbf{X}, \theta) = \mathcal{N}(\mathbf{0}, \mathbf{K}_{f,u} \mathbf{K}_{u,u}^{-1} \mathbf{K}_{u,f} + \text{diag}[\mathbf{K}_{f,f} - \mathbf{K}_{f,u} \mathbf{K}_{u,u}^{-1} \mathbf{K}_{u,f}] + \Sigma) .$$

$\mathbf{K}_{f_1 f_1}(\mathbf{x}_1, \mathbf{x}_1)$	$(\mathbf{K}_{f_1 u} \mathbf{K}_{u u}^{-1} \mathbf{K}_{u f_1})(\mathbf{x}_1, \mathbf{x}_2)$	$(\mathbf{K}_{f_1 u} \mathbf{K}_{u u}^{-1} \mathbf{K}_{u f_1})(\mathbf{x}_1, \mathbf{x}_3)$
$(\mathbf{K}_{f_1 u} \mathbf{K}_{u u}^{-1} \mathbf{K}_{u f_1})(\mathbf{x}_2, \mathbf{x}_1)$	$\mathbf{K}_{f_1 f_1}(\mathbf{x}_2, \mathbf{x}_2)$	$(\mathbf{K}_{f_1 u} \mathbf{K}_{u u}^{-1} \mathbf{K}_{u f_1})(\mathbf{x}_2, \mathbf{x}_3)$
$(\mathbf{K}_{f_1 u} \mathbf{K}_{u u}^{-1} \mathbf{K}_{u f_1})(\mathbf{x}_3, \mathbf{x}_1)$	$(\mathbf{K}_{f_1 u} \mathbf{K}_{u u}^{-1} \mathbf{K}_{u f_1})(\mathbf{x}_3, \mathbf{x}_2)$	$\mathbf{K}_{f_1 f_1}(\mathbf{x}_3, \mathbf{x}_3)$

$$\mathbf{Q}_{f_1, f_1}$$

Computational requirements

- The computational demand is now equal to $\mathcal{O}(NDM^2)$. Storage is $\mathcal{O}(NDM)$.
- For inference, the computation of the mean grows as $\mathcal{O}(DM)$ and the computation of the variance as $\mathcal{O}(DM^2)$, after some pre-computations and for one test point.
- Similar to the Fully Independent Training Conditional (FITC) approximation Quiñonero-Candela and Rasmussen (2005); Snelson and Ghahramani (2005).

Deterministic approximation

- We could also assume that given the latent functions the outputs are deterministic.
- The marginal likelihood is given as

$$p(\mathbf{y}|\mathbf{Z}, \mathbf{X}, \theta) = \mathcal{N}(\mathbf{0}, \mathbf{K}_{f,u} \mathbf{K}_{u,u}^{-1} \mathbf{K}_{u,f} + \Sigma) .$$

- Computation complexity is the same as FITC.
- Deterministic training conditional approximation (DTC).

References I

- Neil Lawrence, Matthias Seeger, and Ralf Herbrich. Fast sparse gaussian process methods: The informative vector machine. In S. Becker, S. Thrun, and K. Obermayer, editors, *Advances in Neural Information Processing Systems*, volume 15, pages 625–632. MIT Press, 2003.
- Joaquin Quiñero-Candela and Carl Edward Rasmussen. A unifying view of sparse approximate Gaussian process regression. *Journal of Machine Learning Research*, 6:1939–1959, 2005.
- Matthias Seeger. *Bayesian Gaussian Process Models: PAC-Bayesian Generalisation Error Bounds and Sparse Approximations*. PhD thesis, School of Informatics, University of Edinburgh, 2003.
- Edward Snelson and Zoubin Ghahramani. Sparse Gaussian processes using pseudo-inputs. In *NIPS 2005*, 2005.