

MA-2012-14 to 26

AI24BTECH11017 - Maanya

14) Let $R = \mathbb{Z} \times \mathbb{Z}$ and $I = \mathbb{Z} \times \mathbb{Z} \setminus \{0\}$. Then which of the following statements is correct?

- a) I is a maximal ideal but not a prime ideal of R .
- b) I is a prime ideal but not a maximal ideal of R .
- c) I is both maximal ideal as well as a prime ideal of R .
- d) I is neither a maximal ideal nor a prime ideal of R .

15) The function $u(r, \theta)$ satisfying the Laplace equation

$$\frac{\partial^2 u}{\partial r^2} + \frac{1}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} = 0, \quad e < r < e^2$$

subject to the conditions $u(e, \theta) = 1$, $u(e^2, \theta) = 0$ is

- a) $\ln(e/r)$
- b) $\ln(e/r^2)$
- c) $\ln(e^2/r)$
- d) $\sum_{n=1}^{\infty} \left(\frac{r-e^2}{e-e^2} \right) \sin n\theta$

16) The functional

$$\int_0^1 (y^2 + (y' + 2y')^2 + kyy' + y^2) dx, \quad y(0) = 0, y(1) = 1, y'(0) = 2, y'(1) = 3$$

is path independent if k equals

- a) 1
- b) 2
- c) 3
- d) 4

17) If a transformation $y = uv$ transforms the given differential equation

$$f(x)y'' - 4f'(x)y' + g(x)y = 0$$

into the equation of the form $v'' + h(x)v = 0$, then u must be

- a) $1/f^2$
- b) xf
- c) $1/2f$
- d) f^2

18) The expression

$$\frac{1}{D_x^2 - D_y^2} \sin(x - y)$$

is equal to

- a) $-\frac{x}{2} \cos(x - y)$
- b) $-\frac{x}{2} \sin(x - y) + \cos(x - y)$
- c) $\frac{x}{2} \cos(x - y) + \sin(x - y)$
- d) $\frac{3x}{2} \sin(x - y)$

19) The function $\phi(x)$ satisfying the integral equation

$$\int_0^x e^{-t^2} \phi(\xi) d\xi = \frac{x^2}{2}$$

is

- a) $\frac{x^2}{2}$
- b) $x + \frac{x^2}{2}$
- c) $x - \frac{x^2}{2}$
- d) $1 + \frac{x^2}{2}$

20) Given the data:

x	1	2	3	4	5
y	-1	2	-3	4	-5

If the derivative of $y(x)$ is approximated as:

$$y'(x_i) = \frac{1}{h} \left(\Delta y_i + \frac{1}{2} \Delta^2 y_i - \frac{1}{4} \Delta^3 y_i \right),$$

then the value of $y'(2)$ is

- a) 4
- b) 8
- c) 12
- d) 16

21) If

$$A = \begin{pmatrix} 1 & 0 & 0 \\ 50 & 1 & 0 \\ 50 & 0 & 1 \end{pmatrix},$$

then A^{50} is

- a) $\begin{pmatrix} 1 & 0 & 0 \\ 50 & 1 & 0 \\ 50 & 0 & 1 \end{pmatrix}$
- b) $\begin{pmatrix} 1 & 0 & 0 \\ 48 & 0 & 0 \\ 48 & 0 & 1 \end{pmatrix}$
- c) $\begin{pmatrix} 1 & 0 & 0 \\ 25 & 1 & 0 \\ 25 & 0 & 1 \end{pmatrix}$
- d) $\begin{pmatrix} 1 & 0 & 0 \\ 24 & 1 & 0 \\ 24 & 0 & 1 \end{pmatrix}$

22) If

$$y = \sum_{n=0}^{\infty} c_n x^n$$

is assumed to be a solution of the differential equation

$$x^2 y'' - xy' - 3(1+x^2)y = 0,$$

then the values of r are

- a) 1 and 3

- b) -1 and 3
- c) 1 and -3
- d) -1 and -3

23) Let the linear transformation $T : \mathbb{R}^2 \rightarrow \mathbb{R}^3$ be defined by

$$T(x_1, x_2) = (x_1, x_1 + x_2, x_2).$$

Then the nullity of T is:

- a) 0
- b) 1
- c) 2
- d) 3

24) The approximate eigenvalue of the matrix

$$A = \begin{bmatrix} -15 & 4 & 3 \\ 10 & -12 & 6 \\ 20 & -4 & 2 \end{bmatrix}$$

obtained after two iterations of the Power method, with the initial vector $[1 \ 1 \ 1]^T$, is:

- a) 7.768
- b) 9.468
- c) 10.548
- d) 19.468

25) The root of the equation $xe^x = 1$ between 0 and 1, obtained by using two iterations of the bisection method, is:

- a) 0.25
- b) 0.50
- c) 0.75
- d) 0.65

Q.26 to Q.55 carry two marks each.

26) Let

$$\oint_C \left(\frac{1}{(z-2)^2} + \frac{(a-2)^2}{z} + 4 \right) dz = 4\pi i,$$

where the closed curve C is the triangle having vertices at:

$$i, \quad \frac{1}{\sqrt{2}}, \quad \text{and} \quad \frac{-1}{\sqrt{2}} i,$$

the integral being taken in the anti-clockwise direction. Then one value of a is:

- a) $1 + i$
- b) $2 + i$
- c) $3 + i$
- d) $4 + i$