## Matrix Theory Assignment 1

## AI24BTECH11017-Maanya Sri

- 16) Using mathematical induction, prove that  $\tan^{-1}\left(\frac{1}{3}\right) + \tan^{-1}\left(\frac{1}{7}\right) + \dots + \tan^{-1}\left\{\frac{1}{(n^2+n+1)}\right\} =$
- $\tan^{-1}\left\{\frac{n}{(n+2)}\right\}_{k=1}^{k}$  (1993-5 Marks) 17) Prove that  $\sum_{k=1}^{n} (-3)^{r-1} {}^{3n}C_{2n-1} = 0$ , where  $k = \frac{(3n)}{2}$  and n is an even positive integer. (1993-5 Marks)
- 18) If x is not an integral multiple of  $2\pi$  use mathematical induction to prove that :  $\cos x + \cos 2x + \dots + \cos nx = \cos \frac{n+1}{2} x \sin \frac{nx}{2} \csc \frac{x}{2}$
- 19) Let *n* be a positive integer and  $(1 + x + x^2)^n = a_0 + a_1 x + \dots + a_{2n} x^{2n}$ . Show that  $a_0^2 - a_1^2 + a_2^2 + a_2^2 + a_2^2 = a_n$
- 20) Using mathematical induction prove that for every integer  $n \ge 1, (3^{2n} 1)$  is divisible by  $2^{n+2}$  but not by  $2^{n+3}$ . (1996-3 Marks)
- 21) Let  $0 < A_i < \pi$  for i = 1, 2, ..., n. Use mathematical induction to prove that  $\sin A_1 +$  $\sin A_2... + \sin A_n \le n \sin \left(\frac{A_1 + A_2 + .... + A_n}{n}\right)$  where  $\ge 1$  is a natural number .{You may use the fact that  $p \sin x + (1 - p) \sin y \le \sin [px + (1 - p)y]$ , where  $0 \le p \le 1$  and  $0 \le x, y \le \pi.$ (1997-5 Marks)
- 22) Let p be a prime number and m a positive integer. By mathematical induction on m, or otherwise, prove that whenever r is an integer such that p does not divide r, p divides  ${}^{mp}C_r$  [Hint: You may use the fact that  $(1+x)^{(m+1)p} = (1+x)^p (1+x)^{mp}$ ] (1998-8 Marks)
- 23) Let *n* be any positive integer. Prove that  $\sum_{k=0}^{m} \frac{\binom{2^{n-k}}{k}}{\binom{2^{n-1}}{2^{n-2k+1}}} \cdot \frac{2^{n-4k+1}}{2^{n-2k+1}} 2^{n-2k} = \frac{\binom{n}{m}}{\binom{2^{n-2m}}{n-m}} 2^{n-2m}$  for each non-be gatuve integer  $m \le n$ . (Here  $\binom{p}{q} = {}^pC_q$ ).
- 24) For any positive integer  $m, n(\text{with } n \geq m)$ , let  $\binom{n}{m} = {}^{n}C_{m}$ . Prove that  $\binom{n}{m} + \binom{n-1}{m} + \binom{n-1}{m}$  $\binom{n-2}{m} + \dots + \binom{m}{m} = \binom{n+1}{m+2}$ . Hence or otherwise, prove that

$$\binom{n}{m} + 2\binom{n-1}{m} + 3\binom{n-2}{m} + \dots + (n-m+1)\binom{m}{m} = \binom{n+2}{m+2}$$
 (24.1)

- 25) For every positive integer n, prove that  $\sqrt{(4n+1)} < \sqrt{n} + \sqrt{n+1} < \sqrt{4n+2}$ . Hence or otherwise, prove that  $\left[\sqrt{n} + \sqrt{(n+1)}\right] = \left[\sqrt{4n+1}\right]$ , where [x] denotes the greatest integer not exceeding x.
- 26) Let a, b, c be positive real numbers such that  $b^2 4ac > 0$  and let  $\alpha_1 = c$ . Prove by induction that  $\alpha_{n+1} = \frac{a\alpha_n^2}{(b^2 - 2a(\alpha_1 + \alpha_2 + ...\alpha_n))}$  is well-defined and  $\alpha_{n+1} < \frac{\alpha_n}{2}$  for all n = 1,2,... (Here, 'well-defined' means that the denominator in the expression for  $\alpha_{n+1}$  is not zero.) (2001-5 Marks)
- 27) Use mathematical induction to show that  $(25)^{n+1} 24n + 5735$  is divisible by  $(24)^2$ for all n = 1, 2, ....(2002-5 Marks)

28) Prove that

$$2^{k} \binom{n}{0} \binom{n}{k} - 2^{k-1} \binom{n}{2} \binom{n}{1} \binom{n-1}{k-1} + 2^{k-2} \binom{n-2}{k-2} - \dots \cdot (-1)^{k} \binom{n}{k} \binom{n-k}{0} = \binom{n}{k}$$
 (28.1)

(2003-2 Marks)

29) A coin has probability p of showing head when tossed. It is tossed n times.Let  $p_n$  denote the probability that no two (or more) consequtive heads occur. Prove that  $p_1 = 1, p_2 = 1 - p^2$  and  $p_n = (1 - p) \cdot p_{n-1} + p(1 - p)p_{n-2}$  for all  $n \ge 3$ . Prove by induction on n, that  $p_n = A\alpha^n + B\beta^n$  for all  $n \ge 1$ , where  $\alpha$  and  $\beta$  are the roots of quadratic equation  $x^2 - (1 - p)x - p(1 - p) = 0$  and  $A = \frac{p^2 + \beta - 1}{\alpha\beta - \alpha^2}$ ,  $B = \frac{p^2 + \alpha - 1}{\alpha\beta - \beta^2}$ . (2000-5 Marks)