

Computer-Assisted Theorem Proving: 7CCSACTP

Exercise Sheet 2

Important Note: Please download Isabelle from the link provided in the slides and bring a laptop to the large group tutorial. Performing the proofs in Isabelle will be an integral part of those tutorials.

Before you start, please open a new Isabelle/HOL theory file called `Week_1.thy` and add the following to its beginning.

```
theory Week_2
imports Main
begin
```

Exercise 2.1 Addition is Associative

Consider the function $(+)$ defined in Isabelle/HOL for natural numbers.

Prove the following theorem, first pen-and-paper, and then formally in Isabelle.

lemma *add_associative*: $((a::nat) + b) + c = a + (b + c)$

Hints: 1. Perform the proof by induction, using substitutions *subst*, backward reasoning using the method *rule*, and forward reasoning using *OF*.

2. Think about which variable to perform induction on. Addition on natural numbers is defined by recursion on its first argument: look at *thm add_0_left add_Suc*.

3. You will need to identify multiple lemmas about how *add* works. Do not prove those lemmas. Use *sorry* as the proof. This is the top-down approach.

4. There is a proof which uses only the following lemmas: *add_0_left*, *add_Suc*, *nat.inject*, *iffD2*

Exercise 2.2 Multiplication is Monotone

Consider the multiplication function $(*)$ defined in Isabelle. Prove the following property of it, i.e. that it is monotonically increasing in both arguments. You should prove it both, pen-and-paper and in Isabelle.

lemma *mult_mono*:

fixes $a\ b\ c\ d :: nat$ — Note the alternative way of fixing the types of the constants
assumes $a \leq b\ c \leq d$
shows $a * c \leq b * d$

Your pen-and-paper proof should indicate

1. what lemmas are you assuming,
2. on what variable are you performing induction,
3. what are assumptions and the proof goals in the base case and the step case, and what is the induction hypothesis, and
4. how are you instantiating the induction hypothesis, i.e. how do you show that its assumptions hold and which quantified variables are instantiated with which constants.

Hints: 1. The proof should be by induction

2. The following lemmas suffice to prove the goal with only substitution, forward reasoning using *OF*, backward reasoning using *rule*, and induction:

le_0_eq, *iffD1*, *mult_zero_right*, *le_refl*, *not_le*, *Suc_leI*, *le_antisym*, *le_SucI*, *mult_le_mono1*, *mult_le_mono2*, *le_trans*

3. In the step case, perform a case analysis with *proof (cases c ≤ d)*. In the second case, together with the assumption $c \leq \text{Suc } d$, you can derive $c = \text{Suc } d$.

Exercise 2.3 Induction Hypothesis Generalisation

Subtraction on natural numbers is truncated: $n - m = 0$ whenever $n \leq m$. Prove that, as long as $m \leq n$, subtracting m and adding it back gives n again.

lemma *sub_add*: $(m :: nat) \leq n \implies (n - m) + m = n$

(a) First try to prove the lemma by plain induction on n , i.e. with *proof(induction n)*. Write down the goal of the step case, its assumption and the induction hypothesis. Explain, pen-and-paper, why the induction hypothesis is not strong enough to prove the step case. Consider the case $m = \text{Suc } k$.

(b) Now prove the lemma, first pen-and-paper and then formally in Isabelle, by generalising the induction hypothesis over m , i.e. with *proof(induction n arbitrary: m)*. Your pen-and-paper proof should state the generalised induction hypothesis and say with which term you instantiate its universally quantified variable in the step case.

Hints: 1. In the step case, perform a case analysis on m with *proof(cases m)*: either $m = 0$ or $m = \text{Suc } k$ for some k . You can name the case assumption, e.g. *case m_eq: (Suc k)*.

2. Use *thm Suc.IH* in the step case to compare the induction hypothesis you get with and without *arbitrary: m*. The induction hypothesis has an assumption, which you can discharge with *OF*.

3. The following lemmas suffice: *le_0_eq*, *iffD1*, *diff_zero*, *add_0_left*, *add_0_right*, *Suc_le_mono*, *diff_Suc_Suc*, *add_Suc_right*.

(c) Challenge (optional) Prove the same lemma, pen-and-paper and in Isabelle, by induction on m instead. Which variable must you generalise now, and why?

lemma *sub_add_on_m*: $(m::nat) \leq n \implies (n - m) + m = n$

Hint: In the case $n = 0$, the assumption $Suc\ m \leq 0$ is impossible: look at *thm Suc_neq_Zero*.