Tables

Bo Ma[1*]

[1*]Auckland University of Technology.

Tables for: *JDCNet Uses Confidence Gated Privileged Modality Distillation for Cost Preserving X Ray Inference*, in the order in which they appear in the manuscript (main text, then appendix).

Table 1. Methodological comparison of JDCNet with related privileged-modality and confidence-aware distillation methods. "Fixed-cost deploy." denotes whether the deployed model is audited to have the same parameters and MACs as a single-modality baseline; "Negative-channel reporting" denotes whether the study reports alternative transfer mechanisms that were tested and did not help.

Method	Modality pairing	Same-pt.?	Fixed-cost deploy.?	Gating / selection mechanism	Target granularity	Negative-channel reporting
JDCNet (ours)	CT (train only) → X-ray	Yes	Yes, X-ray-only	Hard per-sample confidence mask	Argmax label or untempered two-class distribution	Yes
Chen et al.	Multimodal magnetic resonance imaging (MRI; train) → unimodal	Yes	Not audited	Down-weighting of incorrect teacher outputs	Segmentation maps	No
Cahan et al.	Chest X-ray ↔ CT pulmonary angiography	Yes	Not audited	Latent-diffusion prior	Continuous latent representation	No
DANTE	Cross-modal (disentangled)	Not specified	Not audited	Shared/ specific disentanglement	Continuous representation	No
K-MaT	Multi-anatomy teacher mixture	Not CT–X-ray	Not audited	Instance routing across teachers	Continuous	No
KRD	Generic (non-medical)	N/A	N/A	Continuous reliability weight	Softened distribution	No
MST-Distill	Generic mixture-of-teacher	N/A	N/A	Continuous instance routing	Softened distribution	No
BDKD	Generic calibration-balanced KD	N/A	N/A	Continuous divergence weighting	Softened distribution	No

Table 2. Summary of the 510-patient paired comparisons (five folds × three seeds). Unless stated otherwise, ΔBA is against the batch-512 supervised baseline. p: paired t-test over the 15 fold/seed cells (patient-level bootstrap for the row so marked); p_{Holm}: Holm-adjusted across the 16 JDCNet configurations. The two JDCNet rows marked "selected" were chosen post hoc from 16 configurations (Table 4).

Comparison	ΔBA	95% CI	+/0/−	p	p_{Holm}
Does the CT view carry class signal? (CT-only reference)					
CT mid-slice	+0.045	[+0.019, +0.070]	11/0/4	0.005	–
CT 3-slice	+0.051	[+0.025, +0.079]	12/0/3	0.003	–
Does JDCNet move the X-ray student? (original sweep)					
3-slice, soft-KL, τ=0.70, λ=1.0 (selected)	+0.0345	[+0.0112, +0.0571]	10/0/5	0.014	0.22
Mid, hard, τ=0.80, λ=1.5 (selected)	+0.0329	[+0.0074, +0.0584]	10/1/4	0.028	0.42
Average of all 16 configurations (unselected)	+0.020	[+0.003, +0.039]	13/0/2	0.060	–
Average of the 8 initial configurations (unselected)	+0.016	[−0.001, +0.035]	8/0/7	0.11	–
Pre-registered supplementary experiment (batch 128, 16 supervised restarts per cell)					
Supervised batch 128 (r=0) minus supervised batch 512	−0.005	[−0.032, +0.022]	8/0/7	0.75	–
JDCNet best of 16 minus supervised best of 16, selected on validation (primary)	+0.037	[+0.016, +0.058]	11/1/3	0.005	–
Same comparison, patient-level bootstrap	+0.037	[+0.007, +0.068]	–	0.017	–
Average of all 16 configurations minus mean of 16 restarts	+0.019	[+0.006, +0.031]	12/0/3	0.016	–
Do alternative channels move the X-ray student?					
Gated logit KD, mid-slice	−0.022	[−0.055, +0.010]	5/0/10	0.23	–
Gated logit KD, DRR	−0.064	[−0.095, −0.034]	3/0/12	0.002	–
Plain logit KD, 3-slice	−0.036	[−0.062, −0.013]	4/0/11	0.016	–
Contrastive alignment, best of 4	+0.008	[−0.021, +0.038]	7/1/7	0.61	–

Table 3. Data sources and their roles. Only the 510-patient paired BIMCV cohort supports the main comparisons.

Cohort / source	Size	Pairing	Role in paper	Main caveat
Primary BIMCV paired cohort	510 patients ($113^{+}/397^{-}$)	Same-patient CT–X-ray	JDCNet sweep, CT-only references, logit-KD and contrastive comparators, batch-matched supervised restarts	Single public source; classes from two releases; binary task.
Balanced BIMCV subsets	226–228 patients (1:1); 216 with a rendered CT teacher view	Same-patient CT–X-ray	CT-view comparison, BiomedCLIP control, earlier reruns (appendix)	Smaller, class-balanced; different split.
BIMCV rerun split	466 patients with a rendered CT teacher view	Same-patient CT–X-ray	Earlier gate-temperature rerun (appendix)	Paired comparison on the validation partition only.
Historical Cohen paired subset	19 patients / 26 X-rays	Same-patient CT–X-ray	Early feasibility and instability diagnosis	Too small for ranking.
External MIDRC radiographs	126 images ($63^{+}/63^{-}$)	X-ray only	Frozen-student stress test of the 510-patient models	Different source, detector mix, labels, and population.

Table 4. All 16 JDCNet configurations on the 510-patient BIMCV cohort (ResNet-18 student; five folds × seeds 42–44). Stage: I, initial grid; A, extension A (λ=1.5); B, extension B (soft-KL), in the order they were run. +/0/−: fold/seed cells with positive, zero, and negative ΔBA against the batch-512 supervised baseline. CI: recorded percentile bootstrap. p: paired t-test over the 15 cells; p_{H}: Holm-adjusted over the 16 configurations. Δ_{128}: mean difference from the average of the 16 batch-128 supervised restarts of the same cell. Rule: meets the planned decision rule (mean $\geq +0.03$ and lower CI bound > 0).

Teacher	τ	λ	Target	Stage	+/0/−	ΔBA	95% CI	p	p_{H}	Δ_{128}	Rule
3-slice	0.70	1.0	soft-KL	B	10/0/5	+0.0345	[+0.0112, +0.0571]	0.014	0.22	+0.033	Yes
Mid	0.80	1.5	hard	A	10/1/4	+0.0329	[+0.0074, +0.0584]	0.028	0.42	+0.032	Yes
Mid	0.70	1.0	hard	I	10/0/5	+0.0298	[−0.0002, +0.0597]	0.079	1.00	+0.029	–
Mid	0.70	1.5	hard	A	9/0/6	+0.0296	[+0.0026, +0.0577]	0.061	0.86	+0.029	–
Mid	0.70	1.0	soft-KL	B	10/0/5	+0.0264	[+0.0017, +0.0547]	0.082	1.00	+0.025	–
3-slice	0.80	1.0	soft-KL	B	10/0/5	+0.0258	[−0.0029, +0.0546]	0.11	1.00	+0.025	–
3-slice	0.80	1.0	hard	I	10/0/5	+0.0247	[+0.0012, +0.0504]	0.082	1.00	+0.024	–
3-slice	0.70	0.5	hard	I	10/0/5	+0.0239	[−0.0032, +0.0501]	0.11	1.00	+0.023	–
Mid	0.80	1.0	soft-KL	B	9/0/6	+0.0231	[−0.0083, +0.0544]	0.19	1.00	+0.022	–
3-slice	0.80	1.5	hard	A	9/0/6	+0.0181	[−0.0105, +0.0464]	0.25	1.00	+0.017	–
3-slice	0.80	0.5	hard	I	9/0/6	+0.0151	[−0.0185, +0.0479]	0.40	1.00	+0.014	–
Mid	0.80	0.5	hard	I	10/0/5	+0.0144	[−0.0063, +0.0358]	0.22	1.00	+0.013	–
Mid	0.80	1.0	hard	I	10/0/5	+0.0127	[−0.0177, +0.0423]	0.44	1.00	+0.012	–
3-slice	0.70	1.0	hard	I	7/0/8	+0.0066	[−0.0182, +0.0330]	0.63	1.00	+0.005	–
Mid	0.70	0.5	hard	I	7/0/8	+0.0005	[−0.0247, +0.0264]	0.97	1.00	−0.001	–
3-slice	0.70	1.5	hard	A	9/0/6	−0.0031	[−0.0315, +0.0253]	0.84	1.00	−0.004	–

Table 5. Absolute test-partition performance on the 510-patient cohort, mean (SD) over 15 fold/seed cells. Macro-F1 is the unweighted mean of the per-class F1 scores; sensitivity refers to the COVID-19 class.

Method	BA	AUC	Macro-F1	Sensitivity	Specificity
Supervised X-ray, batch 512	0.625 (0.050)	0.674 (0.049)	0.568	0.629 (0.139)	0.620 (0.128)
Supervised X-ray, batch 128 (r=0)	0.620 (0.066)	0.668 (0.053)	0.558	0.646 (0.130)	0.594 (0.127)
JDCNet 3-slice soft-KL	0.659 (0.041)	0.713 (0.027)	0.618	0.607 (0.153)	0.711 (0.125)
JDCNet mid hard	0.658 (0.038)	0.692 (0.045)	0.607	0.642 (0.138)	0.674 (0.132)
Plain logit KD, mid	0.614 (0.071)	0.676 (0.072)	0.565	0.550 (0.214)	0.679 (0.228)
Gated logit KD, mid	0.603 (0.079)	0.660 (0.068)	0.528	0.558 (0.290)	0.647 (0.302)
Gated logit KD, DRR	0.561 (0.059)	0.613 (0.077)	0.429	0.689 (0.287)	0.433 (0.324)
CT-only reference, mid	0.670 (0.051)	0.747 (0.050)	0.653	0.531 (0.155)	0.809 (0.097)
CT-only reference, 3-slice	0.676 (0.045)	0.754 (0.032)	0.662	0.534 (0.136)	0.818 (0.096)
Contrastive alignment, best	0.633 (0.047)	0.661 (0.052)	0.577	0.642 (0.090)	0.623 (0.092)

Table 6. CT-teacher reliability by partition (mean over the 15 fold/seed teachers of each view). Cov.: fraction of samples with teacher confidence above τ; Acc. ret./rej.: teacher accuracy on samples with confidence above/below τ=0.80; ECE: expected calibration error (15 bins); T_{val}: temperature fitted on the validation partition by negative log-likelihood. The gate acts on the training partition.

View	Partition	Acc.	ECE	Cov. τ=0.70	Cov. τ=0.80	Acc. ret./rej.	T_{val}
Mid-slice	train	0.943	0.032	0.89	0.84	0.965 / 0.700	1.72
	val	0.787	0.169	0.83	0.76	0.828 / 0.689	1.72
	test	0.747	0.189	0.82	0.75	0.772 / 0.568	1.72
3-slice	train	0.962	0.021	0.95	0.92	0.981 / 0.727	1.71
	val	0.789	0.174	0.89	0.82	0.813 / 0.674	1.71
	test	0.755	0.194	0.90	0.82	0.792 / 0.551	1.71
Mean proj.	train	0.881	0.057	0.84	0.74	0.920 / 0.688	2.52
	val	0.764	0.176	0.79	0.68	0.807 / 0.687	2.52
	test	0.723	0.190	0.79	0.69	0.765 / 0.523	2.52
DRR	train	0.865	0.060	0.84	0.77	0.937 / 0.590	1.15
	val	0.746	0.194	0.80	0.71	0.732 / 0.681	1.15
	test	0.699	0.202	0.79	0.71	0.768 / 0.524	1.15

Table 7. Anonymized case-level outputs of a mid-slice CT teacher and the confidence gate ($\tau_{\text{gate}} = 0.80$; fold 0, seed 42). Case labels do not encode patient identifiers, and no pixels, filenames, or identifiers are reproduced. The source collection is the BIMCV-COVID19 portal, https://bimcv.cipf.es/bimcv-projects/bimcv-covid19. The four examples are illustrative, not a performance estimate.

Case	Teacher confidence	Teacher prediction	Ground truth	Gate outcome
Case A	0.9992	COVID-19	COVID-19	Accepted
Case B	0.9997	Non-COVID	COVID-19	Accepted
Case C	0.7619	COVID-19	COVID-19	Rejected
Case D	0.7707	Non-COVID	COVID-19	Rejected

Table 8. Supporting evidence layers and what each contributes.

Evidence layer	Main observation	Role
19-patient paired cohort	Seed-level swings between chance and perfect scores	Feasibility only
Balanced-subset gated-logit scan	Best configuration positive, CI crosses zero	Instability diagnosis
510-patient logit-KD runs	Negative or neutral across four teacher views	Comparator evidence
226-patient BiomedCLIP control	No detectable difference from ResNet-18 ($+0.002$, CI ± 0.05)	Capacity control
Pre-registered restarts	No detectable batch-size effect; nested JDCNet advantage $+0.037$	Selection and budget control
External MIDRC test	All students at chance	Generalization warning

Table 9. Inference cost at 224×224 input in 32-bit floating point (FP32), batch size 1, model forward pass only. MACs count convolution and linear layers. Latency is the median (interquartile range) of 1,000 timed passes after 50 warm-ups on a laptop with an Intel Core i7-13700H CPU (one thread) and an NVIDIA GeForce RTX 4050 Laptop GPU (PyTorch 2.5.1); a repeated run differed by up to a factor of two on the GPU because of power management, so timings are illustrative. The supervised, JDCNet, and logit-KD students share one deployed graph and therefore one row.

Deployed model	Params / MACs	CPU latency	GPU latency	GPU peak mem.
ResNet-18 X-ray classifier (supervised, JDCNet, logit KD)	11.18 M / 1.81 G	55.4 (5.5) ms	2.63 (0.45) ms	62 MB
Module-augmented student (historical pilot)	0.57 M / 3.02 G	99.8 (11.1) ms	1.93 (0.29) ms	34 MB

Table 10. External stress test of the 510-patient models on 126 MIDRC radiographs (63/63). Values are means over the 15 fold/seed checkpoints of each method; 95% CIs reflect image-level sampling. Ens. AUC: AUC of the checkpoint-averaged score. Pos. rate: fraction of images predicted COVID-19.

Students (checkpoints)	BA [95% CI]	AUC [95% CI]	Ens. AUC	Pos. rate
Supervised, batch 512 (15)	0.514 [0.427, 0.599]	0.505 [0.404, 0.606]	0.517	0.68
Supervised, batch 128 (15)	0.498 [0.412, 0.584]	0.489 [0.387, 0.590]	0.477	0.57
JDCNet 3-slice soft-KL (15)	0.492 [0.406, 0.578]	0.498 [0.397, 0.599]	0.485	0.67
JDCNet mid hard (15)	0.503 [0.417, 0.589]	0.511 [0.409, 0.612]	0.507	0.60

Table 11. Benefits and limitations of JDCNet as supported by the present evidence.

Dimension	Benefit or evidence	Limitation or risk
Deployed model	CT is used only in training; the deployed graph is the supervised ResNet-18 classifier, so per-image inference cost is unchanged.	Same-patient CT is needed for training; one extra teacher forward pass per training sample.
Within-cohort performance	Nested, batch-matched comparison: +0.037 BA (patient-level CI $[+0.007, +0.068]$); 15 of 16 configurations positive.	One cohort; no single configuration significant after adjustment; validation partition used for two selections.
Operating point	Higher specificity and AUC, toward the CT teacher's profile.	Sensitivity unchanged or lower; retrospective metrics only.
Generalization	–	All models at chance on an external radiograph set; single source; binary task.
Clinical translation	X-ray-only inputs avoid CT at inference.	No workflow, outcome, calibration, or regulatory evaluation.

Table A1. Tests computed directly from the archived per fold/seed records of the 16 JDC-Net configurations (510-patient cohort; n=15 paired cells; differences against the batch-512 supervised baseline). Stage: I, initial; A, extension A; B, extension B. p_t: paired t-test; p_W: Wilcoxon signed-rank; p_b: bootstrap (shift method, 10,000 resamples); p_s: exact sign test; p_{BH}: Benjamini–Hochberg-adjusted p_t. ΔSens, ΔSpec, ΔAUC: mean paired differences in sensitivity, specificity, and AUC.

View	τ	λ	Target	Stage	ΔBA	p_t	p_W	p_b	p_s	p_{BH}	ΔSens	ΔSpec	ΔAUC
3-slice	0.70	1.0	soft	B	+0.0345	0.014	0.022	0.003	0.30	0.22	−0.022	+0.091	+0.039
Mid	0.80	1.5	hard	A	+0.0329	0.028	0.035	0.010	0.18	0.22	+0.013	+0.053	+0.018
Mid	0.70	1.0	hard	I	+0.0298	0.079	0.095	0.050	0.30	0.22	−0.001	+0.061	+0.033
Mid	0.70	1.5	hard	A	+0.0296	0.061	0.14	0.032	0.61	0.22	−0.005	+0.064	+0.006
Mid	0.70	1.0	soft	B	+0.0264	0.082	0.14	0.051	0.30	0.22	+0.016	+0.036	+0.022
3-slice	0.80	1.0	soft	B	+0.0258	0.11	0.14	0.082	0.30	0.23	−0.021	+0.073	+0.026
3-slice	0.80	1.0	hard	I	+0.0247	0.082	0.17	0.050	0.30	0.22	−0.037	+0.086	+0.027
3-slice	0.70	0.5	hard	I	+0.0239	0.11	0.14	0.087	0.30	0.23	−0.005	+0.053	+0.018
Mid	0.80	1.0	soft	B	+0.0231	0.19	0.23	0.15	0.61	0.33	−0.044	+0.091	+0.018
3-slice	0.80	1.5	hard	A	+0.0181	0.25	0.25	0.22	0.61	0.36	−0.018	+0.054	+0.038
3-slice	0.80	0.5	hard	I	+0.0151	0.40	0.30	0.37	0.61	0.54	+0.016	+0.014	+0.018
Mid	0.80	0.5	hard	I	+0.0144	0.22	0.23	0.18	0.30	0.34	+0.016	+0.013	+0.014
Mid	0.80	1.0	hard	I	+0.0127	0.44	0.49	0.41	0.30	0.54	−0.019	+0.045	+0.016
3-slice	0.70	1.0	hard	I	+0.0066	0.63	0.76	0.63	1.00	0.72	−0.052	+0.065	+0.012
Mid	0.70	0.5	hard	I	+0.0005	0.97	0.98	0.97	1.00	0.97	−0.061	+0.062	+0.006
3-slice	0.70	1.5	hard	A	−0.0031	0.84	0.93	0.83	0.61	0.90	−0.077	+0.071	0.000
Average of the 8 initial					+0.0160	0.11	0.28	0.078	1.00	–	–	–	–
Average of all 16					+0.0197	0.060	0.026	0.032	0.007	–	–	–	–

Table A2. Comparator interpretation. Unless marked otherwise, rows are paired differences against the supervised X-ray baseline on the 510-patient cohort.

Family	Representative result	Reading
CT-only reference	Mid $+0.045$ $[+0.019, +0.070]$; 3-slice $+0.051$ $[+0.025, +0.079]$	CT carries signal
JDCNet	15/16 configurations positive; nested, batch-matched comparison $+0.037$ $[+0.016, +0.058]$	Small within-cohort gain
Gated logit KD	Mid -0.022 $[-0.055, +0.010]$; 3-slice -0.026; mean proj. -0.014; DRR -0.064 $[-0.095, -0.034]$	No gain
Plain logit KD	3-slice -0.036 $[-0.062, -0.013]$; other views -0.016 to -0.001, CIs include zero	No gain
Contrastive alignment	Best of 4: $+0.008$ $[-0.021, +0.038]$	No gain
BiomedCLIP (226 subset)	$+0.002$ $[-0.048, +0.050]$ vs. ResNet-18 on the same subset	No gain
Gated logit KD (earlier balanced-subset scan)	Best of eight configurations $+0.034$, CI crosses zero	Not reproduced at 510

Table A3. Contrastive alignment (InfoNCE pretraining followed by supervised fine-tuning) on the 510-patient cohort ($n = 15$ paired fold/seed cells per row).

Teacher view	τ_{NCE}	BA mean [95% CI]	ΔBA [95% CI]	$+/0/-$
3-slice stack	0.20	0.633 [0.610, 0.656]	$+0.008$ $[-0.021, +0.038]$	7/1/7
3-slice stack	0.07	0.627 [0.594, 0.657]	$+0.003$ $[-0.023, +0.030]$	7/0/8
Mid-slice	0.20	0.620 [0.594, 0.643]	-0.005 $[-0.030, +0.020]$	5/0/10
Mid-slice	0.07	0.616 [0.590, 0.642]	-0.008 $[-0.038, +0.027]$	5/0/10

Table A4. CT view comparison on the class-balanced 226-patient BIMCV subset (five folds × seeds 42–44, n=15 cells per row). ΔBA is the paired difference against the supervised X-ray baseline trained on the same subset.

Teacher view	Method	ΔBA [95% CI]	+/0/−
Mid-slice	CT-only ref.	$+0.011$ $[-0.024, +0.046]$	8/0/7
	Plain KD	-0.022 $[-0.062, +0.021]$	6/0/9
	Gated KD	-0.014 $[-0.053, +0.023]$	8/0/7
3-slice stack	CT-only ref.	$+0.019$ $[-0.013, +0.058]$	8/3/5
	Plain KD	-0.030 $[-0.083, +0.016]$	7/0/8
	Gated KD	-0.045 $[-0.092, +0.000]$	5/0/10
Mean projection	CT-only ref.	$+0.045$ $[+0.008, +0.081]$	9/1/5
	Plain KD	-0.031 $[-0.082, +0.019]$	5/1/9
	Gated KD	-0.015 $[-0.058, +0.027]$	6/2/7
DRR	CT-only ref.	$+0.023$ $[-0.023, +0.066]$	9/0/6
	Plain KD	-0.043 $[-0.095, +0.014]$	4/1/10
	Gated KD	-0.005 $[-0.048, +0.039]$	8/0/7

Table A5. Batch-matched gate-temperature rerun on a 466-patient split (single seed, n=5 folds per row; validation-partition BA). τ': raw-confidence threshold equivalent to gating at τ with temperature T_g. ΔBA is the paired difference against the batch-matched supervised baseline (validation BA 0.664 for the 3-slice row set and 0.604 for the mid-slice row set, which use different student manifests); the last column is the change relative to T_g=1.0.

Configuration	T_g	τ'	BA	ΔBA [95% CI]	vs. T_g=1.0
3-slice soft-KL, τ=0.70	0.5	0.60	0.711	+0.047 [+0.028, +0.064]	+0.025
3-slice soft-KL, τ=0.70	1.0	0.70	0.686	+0.022 [−0.011, +0.068]	0.000
3-slice soft-KL, τ=0.70	2.0	0.84	0.691	+0.027 [−0.001, +0.061]	+0.005
Mid hard, τ=0.80	0.5	0.67	0.680	+0.076 [+0.035, +0.117]	+0.007
Mid hard, τ=0.80	1.0	0.80	0.673	+0.069 [+0.034, +0.106]	0.000
Mid hard, τ=0.80	2.0	0.94	0.659	+0.054 [+0.024, +0.088]	−0.014

Table A6. Pre-registered supplementary comparisons on the 510-patient splits (batch-matched supervised restarts). Cell-level: paired over 15 fold/seed cells (bootstrap CI, paired t-test). Patient-level: paired bootstrap over 510 patients with out-of-fold predictions pooled per seed (10,000 resamples).

Comparison	Cell-level ΔBA [95% CI]	p	Patient-level ΔBA [95% CI]	p
Supervised batch 128 (r=0) − batch 512	−0.005 [−0.032, +0.022]	0.75	−0.004 [−0.038, +0.029]	0.82
Supervised best of 16 − supervised r=0	+0.013 [−0.011, +0.038]	0.33	–	–
JDCNet best of 16 − supervised best of 16 (primary)	+0.037 [+0.016, +0.058]	0.005	+0.037 [+0.007, +0.068]	0.017
JDCNet best of 8 − supervised best of 8	+0.029 [+0.007, +0.049]	0.022	–	–
JDCNet best of 16 − supervised batch 512	+0.046 [+0.021, +0.072]	0.004	+0.046 [+0.010, +0.082]	0.012
3-slice soft-KL − supervised r=0	+0.039 [+0.002, +0.074]	0.063	+0.039 [+0.003, +0.077]	0.036
Mid hard − supervised r=0	+0.038 [+0.009, +0.069]	0.035	+0.038 [+0.001, +0.073]	0.045
Average of 16 configurations − mean of 16 restarts	+0.019 [+0.006, +0.031]	0.016	–	–

Table A7. Pre-specified batch-matched comparison of every JDCNet configuration against the first batch-128 supervised restart (r=0) of the same cell (510-patient cohort; rows in the order of Table A1, where the exploration stage of each configuration is listed). Cell level: mean paired difference over the 15 fold/seed cells, bootstrap CI, and paired t-test p_t. Patient level: paired bootstrap over the 510 patients with out-of-fold predictions pooled per seed (10,000 resamples) and two-sided bootstrap p_b. p_{H}: Holm-adjusted over the 16 configurations.

				Cell level					Patient level		
View	τ	λ	Target	+/0/−	ΔBA	95% CI	p_t	p_{H}	95% CI	p_b	p_{H}
3-slice	0.70	1.0	soft	11/0/4	+0.039	[+0.002, +0.074]	0.063	0.95	[+0.003, +0.077]	0.036	0.54
Mid	0.80	1.5	hard	9/0/6	+0.038	[+0.009, +0.069]	0.035	0.56	[+0.001, +0.073]	0.045	0.63
Mid	0.70	1.0	hard	9/0/6	+0.034	[+0.000, +0.071]	0.086	1.00	[−0.001, +0.071]	0.061	0.72
Mid	0.70	1.5	hard	9/1/5	+0.034	[+0.004, +0.069]	0.064	0.95	[+0.005, +0.064]	0.021	0.33
Mid	0.70	1.0	soft	9/0/6	+0.031	[−0.000, +0.064]	0.091	1.00	[−0.000, +0.062]	0.050	0.66
3-slice	0.80	1.0	soft	10/0/5	+0.030	[−0.005, +0.065]	0.12	1.00	[−0.002, +0.063]	0.065	0.72
3-slice	0.80	1.0	hard	9/1/5	+0.029	[−0.006, +0.064]	0.14	1.00	[−0.002, +0.062]	0.069	0.72
3-slice	0.70	0.5	hard	10/0/5	+0.029	[+0.000, +0.057]	0.077	1.00	[−0.001, +0.060]	0.060	0.72
Mid	0.80	1.0	soft	10/0/5	+0.028	[−0.011, +0.065]	0.19	1.00	[−0.008, +0.065]	0.13	1.00
3-slice	0.80	1.5	hard	9/0/6	+0.023	[−0.016, +0.064]	0.30	1.00	[−0.017, +0.061]	0.26	1.00
3-slice	0.80	0.5	hard	11/0/4	+0.020	[−0.003, +0.041]	0.11	1.00	[−0.012, +0.052]	0.20	1.00
Mid	0.80	0.5	hard	10/0/5	+0.019	[−0.011, +0.051]	0.26	1.00	[−0.011, +0.051]	0.21	1.00
Mid	0.80	1.0	hard	9/0/6	+0.017	[−0.016, +0.049]	0.33	1.00	[−0.015, +0.050]	0.29	1.00
3-slice	0.70	1.0	hard	9/1/5	+0.011	[−0.012, +0.035]	0.39	1.00	[−0.024, +0.046]	0.52	1.00
Mid	0.70	0.5	hard	9/0/6	+0.005	[−0.026, +0.034]	0.75	1.00	[−0.030, +0.040]	0.77	1.00
3-slice	0.70	1.5	hard	8/0/7	+0.002	[−0.035, +0.039]	0.94	1.00	[−0.036, +0.037]	0.95	1.00
Average of all 16				10/0/5	+0.024	[−0.000, +0.051]	0.092	–	–	–	–