

Exposure Category by Gender: Followup Analysis

Yash Jaggi

Methods

To determine the relationship between gender and the main substance ingested, we stratified exposure counts by gender (Table 1). There was no overall systematic difference in exposures between genders.

	F	M	Total	p-value
Exposure Category				0.2
Alcohol	3 (5.1%)	4 (7.7%)	7 (6.3%)	
Analgesic	11 (19%)	7 (13%)	18 (16%)	
Antidepressants	12 (20%)	5 (9.6%)	17 (15%)	
CO, As, CN	4 (6.8%)	7 (13%)	11 (9.9%)	
Combination	7 (12%)	10 (19%)	17 (15%)	
Sedatives	4 (6.8%)	3 (5.8%)	7 (6.3%)	
Street Drugs	6 (10%)	11 (21%)	17 (15%)	
Unknown	12 (20%)	5 (9.6%)	17 (15%)	
Total	59 (100%)	52 (100%)	111 (100%)	

Table 1: Exposure Category by Gender. Cells show count of cases, percentage of total by gender in parentheses.

	Gender		Total	p-value ¹
	F	M		
Exposure Category				0.2
Alcohol	3	4	7	
Analgesic	11	7	18	
Antidepressants	12	5	17	
CO, As, CN	4	7	11	
Combination	7	10	17	
Sedatives	4	3	7	
Street Drugs	6	11	17	
Unknown	12	5	17	
Total	59	52	111	

¹Fisher's exact test