

# GEODESICS AND VALUES OF QUADRATIC FORMS

GREG MCSHANE

## 1. INTRODUCTION

**Theorem 1.1.** *Let  $p$  be a prime then the equation*

$$x^2 = -1$$

*admits a solution in  $\mathbb{F}_p$  iff  $p = 2$  or  $p - 1$  is a multiple of 4.*

**Theorem 1.2** (Fermat). *Let  $p$  be a prime then the equation*

$$x^2 + y^2 = p$$

*has a solution in integers iff  $p = 2$  or  $p - 1$  is a multiple of 4.*

There are many proofs of these theorems but the approach initiated by Heath-Brown in [9] has inspired many admirers if not imitators see for example the very nice account of Elsholtz [6]. In some senses this manuscript is a companion to Elsholtz's where instead of looking at the number theory as combinatorics we work in an explicitly geometric context. As such we refer the reader to Elsholtz for historical perspective and the like.

**1.1. Involutions.** The essential ingredients in the Heath-Brown paper are: a finite set  $X$  equipped with a pair of involutions

- any fixed point of the one of the involutions, should it exist, is a solution of the equation.
- the other involution has a unique fixed point which is easy to compute.

The existence of the unique fixed point of the second involution allows one to conclude that the  $X$  has an odd number of elements and so that any involution has a fixed point.

**1.2. Arcs on a punctured surface.** The starting point for this work was a series of observations concerning  $\lambda$ -lengths of simple arcs on a once punctured torus equipped with a hyperbolic structure for example  $\mathbb{H}/\Gamma'$  where  $\Gamma' < \mathrm{SL}(2, \mathbb{Z})$  is the commutator subgroup. An important feature of a punctured surface is that it admits *uniform cusp regions* that is there is a neighborhood of each puncture, or more properly cusp, isometric to

$$\{z \in \mathbb{H}, \mathrm{Re} z \geq 1\} / \langle z \mapsto z + 2 \rangle$$

Such a torus can be obtained from a pair of ideal triangles by "gluing" and the sides of the ideal triangles form a triple of complete simple

geodesics on the surface. The corners of the triangles glue up to a neighborhood of the puncture. We define an *arc* to be any complete geodesic on a punctured surface with both of its ends terminating at cusps. The three sides of the ideal triangle(s) above form a triple of disjoint simple arcs. Each of these arcs has infinite length but if we only consider the portion outside the uniform cusp region then its length is finite. The  $\lambda$ -length of the arc is exponential of half the length of this finite portion. Whilst this definition works well for simple arcs, since the portion of the arc outside the uniform cusp region is connected, more care is needed if the arc is not simple.

## 2. KLEIN FOUR GROUP AND THE BURNSIDE LEMMA

We give a proof of Theorem 1.1 using the Burnside Lemma. Recall that if  $G$  is a group acting on a finite set  $X$  then the Burnside Lemma says

$$(1) \quad |G||X/G| = \sum_g |X^g|$$

where, as usual,  $X^g$  denotes the set of fixed points of the element  $g$  and  $X/G$  the orbit space.

Let  $p \neq 2$ ,  $X = \mathbb{F}_p^*$  and  $G$  be the group generated by the two involutions

$$\begin{aligned} x &\mapsto -x \\ x &\mapsto 1/x. \end{aligned}$$

The group  $G$  has exactly four elements namely:

- the trivial element which has  $p - 1$  fixed points
- $x \mapsto -x$  which has no fixed points
- $x \mapsto 1/x$  has exactly two fixed points namely 1 and  $-1$ .
- $g : x \mapsto -1/x$  is the remaining element and the theorem is equivalent to the existence of a fixed point for it.

Note that since  $\mathbb{F}_p$  is a field  $|X^g| = \#\{x^2 = -1, x \in \mathbb{F}_p^*\}$  is either 0 or 2. Now for our choice of  $X$  and  $G$  equation (1) yields

$$(2) \quad 4|X/G| = (p - 1) + 2 + |X^g|.$$

The LHS is always divisible by 4 so the RHS is too and it follows from this that

$$|X^g| = \begin{cases} 0 & (p - 1) = 2 \pmod{4} \\ 2 & (p - 1) = 0 \pmod{4} \end{cases}$$

This proves Theorem 1.1.

**2.1. Extending.** Thus we have shown that  $-1$  is a quadratic residue modulo  $p$  if  $p$  is of the form  $4k + 1$ . It is natural to consider the other questions considered by Fermat: namely for which values of  $p$  are  $-2$  and  $-3$  residues?

In fact  $-2$  is a residue if  $p$  is 2 or of the form  $8k + 1$  or  $8k + 3$ . Showing this in the spirit of Heath-Brown requires one to consider a group generated by the involutions

$$\begin{aligned} x &\mapsto -x \\ x &\mapsto 2/x. \end{aligned}$$

One immediately sees that things are more complicated as the second involution has fixed points if and only if 2 is a residue whereas  $x \mapsto 1/x$  always had exactly two fixed points. Thus there are two cases:

- $p = 8k + 1$  and both 2 and  $-2$  are residues
- $p = 8k + 3$  and  $-2$  is a residue but 2 is not.

To prove this second assertion one must show that  $x \mapsto 2/x$  has no fixed point so that, by Burnside,  $x \mapsto -2/x$  has two fixed points both of which are square roots of  $-2$ . Thus one must show that the only solution of the associated diophantine equation

$$np = x^2 - 2y^2,$$

is the trivial solution  $n = x = y = 0$ . Now using the fact that  $x^2 - 2y^2$  is the norm of  $x + y\sqrt{2} \in \mathbb{Z}[\sqrt{2}]$  which is a euclidean ring for this norm, one reduces to considering just the solutions of

$$p = x^2 - 2y^2.$$

Finally, one concludes by showing that if  $x, y$  are integers then  $x^2 - 2y^2$  never takes the value 3 mod 8.

**2.2. The case  $p = 11$ .** The first real case of interest in understanding  $x \mapsto -2/x$  is that of  $\mathbb{F}_{11}^*$ . Evidently,  $11 = 3^2 + 2 \times 1^2$  so that  $\bar{3}$  and  $-\bar{3} = \bar{8}$  are the fixed points of the involution.

The reduction homomorphism  $x \mapsto \bar{x}$  allows one to identify the elements of  $\mathbb{F}_p$  with the equivalence classes that constitute the quotient  $\mathbb{Z}/p\mathbb{Z}$ . It is usual to choose the integers  $0, 1, 2, \dots, p-1$  as representatives for the latter, however, we shall find it convenient to work with another set of representatives which are the even integers  $0, 2, 4, \dots, 2p-2$ . Using the euclidean algorithm to compute  $\bar{x}^{-1} \in \mathbb{F}_{11}$  we have the following table:

|                  |    |    |    |    |    |    |    |    |    |    |
|------------------|----|----|----|----|----|----|----|----|----|----|
| $x$              | 12 | 2  | 14 | 4  | 16 | 6  | 18 | 8  | 20 | 10 |
| $\bar{x}$        | 1  | 2  | 3  | 4  | 5  | 6  | 7  | 8  | 9  | 10 |
| $\bar{x}^{-1}$   | 12 | 6  | 4  | 14 | 20 | 2  | 8  | 18 | 16 | 10 |
| $-2\bar{x}^{-1}$ | 20 | 10 | 14 | 16 | 4  | 18 | 6  | 8  | 12 | 2  |

One notes that there are two fixed points of  $-\bar{x} \mapsto -2\bar{x}^{-1}$  namely  $\frac{1}{14} = \bar{3}$  and  $\bar{8}$ .

### 3. COUNTING SUMS OF SQUARES

The transformation  $z \mapsto z + 1$  generates an infinite cyclic group acting on  $\mathbb{H}$ . The standard fundamental domain for this group is an infinite strip, which we will refer to as the *fundamental strip*, consisting of all the  $z \in \mathbb{C}$  such that the real part is between 0 and 1.

**Lemma 3.1.** *Let  $n \geq 2$  be an integer. The number of ways of writing  $n$  as a sum of squares*

$$n = c^2 + d^2$$

*with  $c, d$  coprime integers is equal to the number of points of  $\Gamma.\{i\}$ , the  $\mathrm{SL}(2, \mathbb{Z})$  orbit of  $i$ , in the fundamental strip at height  $\frac{1}{n}$ .*

Note that we are counting  $c^2 + d^2$  and  $d^2 + c^2$  as *different* representations of  $n$ .

*Proof.* Suppose there is such a point which we denote  $w$  verifying the hypotheses in particular

$$w = \frac{ai + b}{ci + d}, \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma,$$

then

$$\mathrm{Im} w = \mathrm{Im} \frac{ai + b}{ci + d} = \frac{\mathrm{Im} i}{c^2 + d^2}.$$

So  $n = c^2 + d^2$ .

Conversely if  $c, d$  are coprime integers then there exists  $a, b$  such that

$$ad - bc = 1 \Rightarrow \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma.$$

By applying a suitable iterate of the parabolic transformation  $z \mapsto z + 1$ , one can choose  $w$  such that  $0 \leq \mathrm{Re} w < 1$ . □

Suppose that  $n$  can be written as a sum of squares  $c^2 + d^2$  and  $w$  is the corresponding point in the fundamental strip then we can associate a Poincaré geodesic to  $w$  in a natural way, we simply take the vertical line that passes through  $w$ . This geodesic joins two points in the ideal boundary of  $\mathbb{H}$  namely  $\infty$  and  $\frac{ac+bd}{n} \in \mathbb{R}$ . Although not strictly necessary we choose to associate a geometric quantity to this geodesic - Penner's  $\lambda$ -length.

#### 4. INVERSIONS

We denote by  $\mathbb{H}$  the Poincaré upper half plane and  $\partial\mathbb{H}$  its ideal boundary ie  $\mathbb{R} \cup \{\infty\}$ . Recall that an *inversion* is an orientation reversing isometry of  $\mathbb{H} \cup \partial\mathbb{H}$ . A Poincaré geodesic is either a vertical line or a semicircle orthogonal to  $\mathbb{R}$ . In both cases it is uniquely determined by its endpoints in the ideal boundary. To each Poincaré geodesic is associated a unique inversion which fixes it pointwise. The inversion  $\phi_h : z \mapsto -\bar{z}$  fixes 0 and  $\infty$  and so the arc joining them. The group of isometries acts transitively on pairs of distinct points  $a, b \in \partial\mathbb{H}$  and so there is an inversion that fixes the geodesic joining  $a, b$  which is in fact conjugate to  $\phi_h$ . The inversion fixing 1,  $-1$  is easily seen to be  $\phi_v : z \mapsto \frac{1}{\bar{z}}$ .

Note that if  $a, b$  are coprime integers then:

- The image of  $\frac{a}{b}$  under  $\phi_h$  is  $-\frac{a}{b}$  and the  $\lambda$ -length of the geodesic joining them is  $2ab$ .
- The image of  $\frac{a}{b}$  under  $\phi_v$  is  $\frac{b}{a}$  and the  $\lambda$ -length of the geodesic joining them is  $|a^2 - b^2|$ .

**Lemma 4.1.** *Let  $p > 2$  be a prime then:*

- *There are exactly two arcs of  $\lambda$ -length  $p$  invariant under  $\phi_v$ ;*
- *There is no arc of  $\lambda$ -length  $p$  invariant under  $\phi_h$ .*

#### 5. CONGRUENCE SUBGROUPS

The Hecke congruence subgroup  $\Gamma_0(N)$  of level  $N$  is the subgroup of  $\Gamma = \text{SL}(2, \mathbb{Z})$  is a normal. It is a subgroup of  $\Gamma_0(N)$ :

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} 1 & * \\ 0 & 1 \end{pmatrix} \pmod{N}.$$

For  $N = 2$  this is generated by just two elements namely:

$$P = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix} \text{ and } Q = \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix}.$$

The product  $P^{-1}Q$  is an element of order 2:

$$P^{-1}Q = \begin{pmatrix} -1 & -1 \\ 2 & 1 \end{pmatrix}.$$

So the quotient  $\mathbb{H}/\Gamma_0(2)$  is a non-compact orbifold with two cusps and a single cone point.

The action of this group on  $\mathbb{Q} \cup \{\infty\}$  is not transitive and there are two orbits. Now  $\Gamma_0(2) < \Gamma(2)$  so each of these orbits is a union of  $\Gamma(2)$ -orbits. Since  $\Gamma(2)$  preserves the parity of the numerator and denominator of a fraction there are exactly three  $\Gamma(2)$  orbits corresponding to  $\frac{0}{1} = 0$ ,  $\frac{1}{1} = 1$ ,  $\frac{1}{0} = \infty$ . Now since  $P$  maps 0 to one:

$$\Gamma_0(2)\{0\} = \Gamma(2)\{0\} \cup \Gamma(2)\{1\}.$$




FIGURE 1. On the left a fundamental domain for  $\Gamma_0^t(2)$  with side pairings. On the right the quotient surface  $\mathbb{H}/\Gamma_0^t(2)$ , the dark region is a cusp region.

In the previous section we considered the action of the involution  $x \mapsto -2/x$  on  $\mathbb{F}_p^*$ . It is natural to study the action of the corresponding involution of  $\mathbb{H}$  that is  $z \mapsto -2/z$  but unfortunately this does not normalise  $\Gamma_0(2)$ . However it does normalise the *anti Hecke congruence-group*:

$$\Gamma_0^t(2) := \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} 1 & 0 \\ * & 1 \end{pmatrix} \pmod{N} \right\} < \Gamma(2).$$

The Hecke congruence group and the anti Hecke group are isomorphic and in fact conjugate in  $\Gamma$  by  $z \mapsto -1/z$ . We can determine the orbits of this group on  $\mathbb{Q}$  using this conjugation and we have:

$$\Gamma_0^t(2)\{\infty\} = \Gamma(2)\{\infty\} \cup \Gamma(2)\{1\}.$$

**5.1. Cusp regions on  $\mathbb{H}/\Gamma_0^t(2)$ .** We denote by  $F$  the set  $\{z, \text{Im } z > 1\}$  this is a *horoball* in  $\mathbb{H}$  centered at  $\infty$ . The image of  $F$  under the  $\text{SL}(2, \mathbb{Z})$  action consists of  $F$  and infinitely many disjoint circles, the so-called *Ford circles*, each tangent to the real line at some rational  $m/n$ . We adopt the convention that  $F$  is also a Ford circle of infinite radius.

The following is well known and is easily checked:

**Lemma 5.1.** *The Ford circle tangent to the real line at  $m/n$  has Euclidean diameter  $1/n^2$ .*

The image of a Ford circle on  $\mathbb{H}/\Gamma(2)$  is a *cusp region* around one of the three cusps  $0, 1, \infty$  of  $\mathbb{H}/\Gamma(2)$ . The map  $z \mapsto z/z + 1$  normalises  $\Gamma(2)$  and so induces an automorphism, in fact an involution of  $\mathbb{H}/\Gamma(2)$  and the quotient is naturally identified with  $\mathbb{H}/\Gamma_0^t(2)$ . Thus  $\mathbb{H}/\Gamma_0^t(2)$  inherits a system of cusp regions from  $\mathbb{H}/\Gamma_0(2)$ . The involution  $z \mapsto$




FIGURE 2. Ford circles with tangent points and curvatures. Recall that the curvature of a euclidean circle is the reciprocal of its radius.

$-2/z$  normalises  $\Gamma_0^t(2)$  so induces an automorphism of  $\mathbb{H}/\Gamma_0^t(2)$  which fixes the points labelled  $1+i$  and  $i\sqrt{2}$ , swaps the cusps labelled  $\infty = \frac{1}{0}$  and  $0 = \frac{0}{1}$  but which does not swap the cusp regions inherited from  $\mathbb{H}/\Gamma(2)$ . We remedy this by choosing a pair of cusp regions which are tangent at the fixed point of the automorphism.

**5.2. Arcs on  $\mathbb{H}/\Gamma_0^t(2)$ .** The surface has two cusps and so there are two kind of arc

- arcs that join distinct cusps  $0/1$  and  $1/0$
- arcs that have both ends at the same either cusps  $0/1$  or  $1/0$ .

**Lemma 5.2.** *Arcs of the first kind, that is those which join distinct cusps  $0/1$  and  $1/0$ , have the same  $\lambda$ -length for the inherited cusp regions and our modified cusp regions.*

Since the automorphism swaps cusps only arcs of the first kind can be invariant for it. Now any arc of the first kind lifts to a vertical line ending at some rational  $\frac{m}{n} \in \mathbb{Q}$ . It follows that for each  $p$  prime there are exactly  $p-1$  arcs of the first kind, namely the projections to  $\mathbb{H}/\Gamma_0^t(2)$  of the Poincaré geodesics  $\infty, \frac{2k}{p}$  with  $k = 1, 2, \dots, p-1$  and each of these has  $\lambda$ -length  $p$  for our choice of cusp regions.

## REFERENCES

- [1] M. Aigner *Markov's Theorem and 100 Years of the Uniqueness Conjecture*, Springer (2013)
- [2] Aigner M., Ziegler G.M. *Representing numbers as sums of two squares*. In: Proofs from the book. Springer, Berlin, Heidelberg. (2010)
- [3] A. Baragar, *On the Unicity Conjecture for Markoff Numbers* Canadian Mathematical Bulletin , Volume 39 , Issue 1 , 01 March 1996 , pp. 3 - 9
- [4] J. O. Button, *The uniqueness of the prime Markoff numbers*, J. London Math. Soc. (2) 58 (1998), 9–17.
- [5] Dolan, S. (2021). 105.38 A very simple proof of the two-squares theorem. The Mathematical Gazette, 105(564), 511-511. doi:10.1017/mag.2021.120
- [6] Elsholtz C.A *Combinatorial Approach to Sums of Two Squares and Related Problems*. In: Chudnovsky D., Chudnovsky G. (eds) Additive Number Theory. Springer, New York, NY. (2010)
- [7] Lester R Ford, *Automorphic Functions*

- [8] Generalov, A.I. A combinatorial proof of Euler-Fermat's theorem on the representation of the primes  $p=8k+3$  by the quadratic form  $x^2 + 2y^2$ . J Math Sci 140, 690–691 (2007). <https://doi.org/10.1007/s10958-007-0008-6>
- [9] Heath-Brown, Roger. *Fermat's two squares theorem*. Invariant (1984)
- [10] Neil Herriot, Communication with Jim Propp <https://faculty.uml.edu/jpropp/reach/Herriot/ptolemywriteup.html>
- [11] Jackson, Terence H.. "A Short Proof That Every Prime  $p = 3 \pmod{8}$  Is of the Form  $x^2 + 2y^2$ ." The American Mathematical Monthly 107 (2000): 447 - 447.
- [12] M.L. Lang, S.P Tan, *A simple proof of the Markoff conjecture for prime powers* Geometriae Dedicata volume 129, pages15–22 (2007)
- [13] G. McShane, *Simple geodesics and a series constant over Teichmuller space* Invent. Math. (1998)
- [14] Northshield, Sam. *A Short Proof of Fermat's Two-square Theorem*. The American Mathematical Monthly. 127. 638-638. (2020).
- [15] R. C. Penner, *The decorated Teichmueller space of punctured surfaces*, Communications in Mathematical Physics 113 (1987), 299–339.
- [16] James Propp, The combinatorics of frieze patterns and Markoff numbers, in Integers, Volume 20 (2020) <http://math.colgate.edu/~integers/u12/u12.pdf>
- [17] J-P. Serre, *A Course in Arithmetic*, Graduate Texts in Mathematics, Springer-Verlag New York 1973
- [18] D. Zagier, *A one-sentence proof that every prime  $p = 1 \pmod{4}$  is a sum of two squares*, American Mathematical Monthly, 97 (2): 144

INSTITUT FOURIER 100 RUE DES MATHS, BP 74, 38402 ST MARTIN D'HÈRES  
CEDEX, FRANCE

*Email address:* mcshane at univ-grenoble-alpes.fr