

Mathematics 1

1-2

Units

Review the Units section of the
NCEES Handbook

Professional Publications, Inc.

FERC

Mathematics 1

1-3

Fundamental Constants

[illegible]PPI • www.ppi2pass.com

Professional Publications, Inc.

FERC

1-4

Conversion)

[illegible]PPI • www.ppi2pass.com

1-5

Mathematics

PPI • www.ppi30000.comPPI • www.ppi30000.com

Professional Publications, Inc.


FERC

Mathematics 1

1-6a

Straight Line

Figure 4.1 Straight Line




$$Ax + By + C = 0 \quad 4.1$$

$$y = mx + b \quad 4.2$$

$$y - y_1 = m(x - x_1) \quad 4.3$$

$$m = \frac{y_2 - y_1}{x_2 - x_1} \quad 4.4$$

Figure 4.2 Two Lines Intersecting in Two-Dimensional Space



$$m_1 = \frac{-1}{m_2} \quad 4.6$$

$$d = \sqrt{(y_2 - y_1)^2 + (x_2 - x_1)^2} \quad 4.7$$

Professional Publications, Inc.

FERC

Mathematics 1

1-6b

Straight Line

Example (EFPRB):

MATHEMATICS-30

The equation $y = a_1 + a_2x$ is an algebraic expression for which of the following?

- (A) a cosine expansion series
- (B) projectile motion
- (C) a circle in polar form
- (D) a straight line

$y = mx + b$ is the slope-intercept form of the equation of a straight line. Thus, $y = a_1 + a_2x$ describes a straight line.

The answer is (D).

MATHEMATICS-31

Find the slope of the line defined by $y - x = 5$.

- (A) $5 + x$
- (B) $-1/2$
- (C) $1/4$
- (D) 1

The slope-intercept form of the equation of a straight line is $y = mx + b$, where m is the slope and b is the y -intercept.

$$y - x = 5$$

$$y = x + 5$$

The coefficient of x , m , is

$$m = 1$$

The answer is (D).

Professional Publications, Inc.

FERC

Mathematics 1

1-6c

Straight Line

Example (EFPRB):

MATHEMATICS-32Find the equation of a line with slope = 2 and y -intercept = -3.

- (A) $y = -3x + 2$
 (B) $y = 2x - 3$
 (C) $y = \frac{2}{3}x + 1$
 (D) $y = 2x + 3$

The slope-intercept form of the given equation is

$$y = 2x - 3$$

The answer is (B).

Professional Publications, Inc.

FERC

Mathematics 1

1-6d

Straight Line

Example (EFPRB):

MATHEMATICS-33

Find the equation of the line that passes through the points (0, 0) and (2, -2).

- (A) $y = x$ (B) $y = -2x + 2$ (C) $y = -2x$ (D) $y = -x$

Since the line passes through the origin, the y -intercept is 0. Thus, the equation simplifies to $y = mx$. Substituting for the known points,

$$y = \left(\frac{-2 - 0}{2 - 0} \right) x$$

$$= -x$$

The answer is (D).

Professional Publications, Inc.

FERC

Mathematics 1

1-6e

Straight Line

Example (EFPRB):

MATHEMATICS-29

In finding the distance, d , between two points, which equation is the appropriate one to use?

(A) $d = \sqrt{(x_1 - x_2)^2 - (y_2 - y_1)^2}$

(B) $d = \sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2}$

(C) $d = \sqrt{(x_1^2 - x_2^2) + (y_1^2 - y_2^2)}$

(D) $d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$

The distance formula is defined as follows.

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$$

The answer is (D).

Professional Publications, Inc.

FERC

Mathematics 1

1-7a

Quadratic Equations

$$x_1, x_2 = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \quad 4.8$$

Example (FERM):

Problem 5

What are the solutions to the following equation?

$$x^2 - x - 12 = 0$$

(A) $x_1 = 1; x_2 = 12$

(B) $x_1 = 4; x_2 = -3$

(C) $x_1 = -1; x_2 = 4$

(D) $x_1 = 6; x_2 = -2$

Solution

There are two ways to solve the equation. The first method is to factor the equation.

$$x^2 - x - 12 = (x + 3)(x - 4) = 0$$

$$x = -3 \text{ or } x = 4$$

The second method is to use the quadratic equation.

$$\begin{aligned} x_1, x_2 &= \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} \\ &= \frac{-(-1) \pm \sqrt{(-1)^2 - (4)(1)(-12)}}{(2)(1)} \\ &= \frac{1 \pm 7}{2} \end{aligned}$$

$$x_1 = 4$$

$$x_2 = -3$$

Answer is B.

Professional Publications, Inc.

FERC

Mathematics 1

1-7b

Quadratic Equations

Example (EFPRB):

MATHEMATICS-6

What is the solution of the equation $50x^2 + 5(x - 2)^2 = -1$, where x is a real-valued variable?

- (A) -6.12 and -3.88 (B) -0.52 and 0.700
 (C) 7.55 (D) no solution

For real-valued x , the left-hand side of the equation must always be greater than or equal to zero, since all terms containing x are squared. There is no solution to this equation for real values of x .

The answer is (D).

Professional Publications, Inc.

FERC

Mathematics 1

1-7c

Quadratic Equations

Example (EFPRB):

MATHEMATICS-7

What are the roots of the cubic equation $x^3 - 8x - 3 = 0$?

- (A) $x = -7.90, -3, -0.38$
 (B) $x = -3, -2, 2$
 (C) $x = -3, -0.38, 2$
 (D) $x = -2.62, -0.38, 3$

By inspection, $+3$ is a root, and $(x - 3)$ is a factor. Factor out $(x - 3)$.

$$\frac{x^3 - 8x - 3}{x - 3} = x^2 + 3x + 1$$

Use the quadratic equation to solve $x^2 + 3x + 1 = 0$.

$$x = 3, \frac{-3 \pm \sqrt{9 - 4}}{2} \\ = -2.62, -0.38, 3$$

The answer is (D).

Professional Publications, Inc.


FERC

Mathematics 1

1-8a

Conic Sections

Figure 4.3 Conic Sections Produced by Cutting Planes



Professional Publications, Inc.


FERC

Mathematics 1

1-8b


Conic Sections

Figure 4.4 Parabola



$$(y - k)^2 = 2p(x - h) \quad 4.12$$

Figure 4.5 Ellipse



$$\frac{(x - h)^2}{a^2} + \frac{(y - k)^2}{b^2} = 1 \quad 4.14$$

Professional Publications, Inc.


FERC

Mathematics 1

1-8c


Conic Sections

Figure 4.6 Hyperbola



$$\frac{(x-h)^2}{a^2} - \frac{(y-k)^2}{b^2} = 1 \quad 4.18$$

Figure 4.7 Circle



$$(x-h)^2 + (y-k)^2 = r^2 \quad 4.22$$

Professional Publications, Inc.

FERC

Mathematics 1

1-8d

Conic Sections

Example (EFPRB):

MATHEMATICS-9What is the radius of the circle defined by $x^2 + y^2 - 4x + 8y = 7$?

- (A) $\sqrt{3}$ (B) $2\sqrt{5}$ (C) $3\sqrt{3}$ (D) $4\sqrt{3}$

Since the general equation for a circle is $(x-a)^2 + (y-b)^2 = r^2$, rearrange the equation given to fit the general equation.

$$x^2 - 4x + y^2 + 8y = 7$$

Complete the binomial forms.

$$x^2 - 4x + 4 + y^2 + 8y + 16 = 7 + 4 + 16$$

$$(x-2)^2 + (y+4)^2 = 27$$

$$r^2 = 27$$

$$r = 3\sqrt{3}$$

The answer is (C).

Professional Publications, Inc.


FERC

Mathematics 1

1-9

Right Triangles

Figure 4.9 Right Triangle



$$\sin \theta = \frac{y}{r} \quad 4.30$$

$$\cos \theta = \frac{x}{r} \quad 4.31$$

$$\tan \theta = \frac{y}{x} \quad 4.32$$

$$\cot \theta = \frac{x}{y} \quad 4.33$$

$$\csc \theta = \frac{r}{y} \quad 4.34$$

$$\sec \theta = \frac{r}{x} \quad 4.35$$

Professional Publications, Inc.

FERC

Mathematics 1

1-10a

Trigonometric Identities

$$\csc \theta = \frac{1}{\sin \theta} \quad 4.36$$

$$\sec \theta = \frac{1}{\cos \theta} \quad 4.37$$

$$\cot \theta = \frac{1}{\tan \theta} \quad 4.38$$

$$\tan \theta = \frac{\sin \theta}{\cos \theta} \quad 4.39$$

$$\cot \theta = \frac{1}{\tan \theta} = \frac{\cos \theta}{\sin \theta} \quad 4.40$$

$$\sin^2 \theta + \cos^2 \theta = 1 \quad 4.41$$

$$\tan^2 \theta + 1 = \sec^2 \theta \quad 4.42$$

$$\cot^2 \theta + 1 = \csc^2 \theta \quad 4.43$$

$$\sin 2\alpha = 2 \sin \alpha \cos \alpha \quad 4.44$$

$$\begin{aligned} \cos 2\alpha &= \cos^2 \alpha - \sin^2 \alpha \\ &= 1 - 2 \sin^2 \alpha \end{aligned}$$

$$= 2 \cos^2 \alpha - 1 \quad 4.45$$

$$\tan 2\alpha = \frac{2 \tan \alpha}{1 - \tan^2 \alpha} \quad 4.46$$

$$\cot 2\alpha = \frac{\cot^2 \alpha - 1}{2 \cot \alpha} \quad 4.47$$

Professional Publications, Inc.

FERC

Mathematics 1

1-10b

Trigonometric Identities

• Two-angle formulas

$$\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta \quad 4.48$$

$$\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta \quad 4.49$$

$$\tan(\alpha + \beta) = \frac{\tan \alpha + \tan \beta}{1 - \tan \alpha \tan \beta} \quad 4.50$$

$$\cot(\alpha + \beta) = \frac{\cot \alpha \cot \beta - 1}{\cot \alpha + \cot \beta} \quad 4.51$$

$$\sin(\alpha - \beta) = \sin \alpha \cos \beta - \cos \alpha \sin \beta \quad 4.52$$

$$\cos(\alpha - \beta) = \cos \alpha \cos \beta + \sin \alpha \sin \beta \quad 4.53$$

$$\tan(\alpha - \beta) = \frac{\tan \alpha - \tan \beta}{1 + \tan \alpha \tan \beta} \quad 4.54$$

$$\cot(\alpha - \beta) = \frac{\cot \alpha \cot \beta + 1}{\cot \beta - \cot \alpha} \quad 4.55$$

• Half-angle formulas

$$\sin \frac{\alpha}{2} = \pm \sqrt{\frac{1 - \cos \alpha}{2}} \quad 4.56$$

$$\cos \frac{\alpha}{2} = \pm \sqrt{\frac{1 + \cos \alpha}{2}} \quad 4.57$$

$$\tan \frac{\alpha}{2} = \pm \sqrt{\frac{1 - \cos \alpha}{1 + \cos \alpha}} \quad 4.58$$

$$\cot \frac{\alpha}{2} = \pm \sqrt{\frac{1 + \cos \alpha}{1 - \cos \alpha}} \quad 4.59$$

Professional Publications, Inc.

FERC

Mathematics 1

1-10c

Trigonometric Identities

• Miscellaneous formulas

$$\sin \alpha \sin \beta = \frac{1}{2}(\cos(\alpha - \beta) - \cos(\alpha + \beta)) \quad 4.60$$

$$\cos \alpha \cos \beta = \frac{1}{2}(\cos(\alpha - \beta) + \cos(\alpha + \beta)) \quad 4.61$$

$$\sin \alpha \cos \beta = \frac{1}{2}(\sin(\alpha + \beta) + \sin(\alpha - \beta)) \quad 4.62$$

$$\sin \alpha + \sin \beta = 2 \sin \left(\frac{\alpha + \beta}{2} \right) \cos \left(\frac{\alpha - \beta}{2} \right) \quad 4.63$$

$$\sin \alpha - \sin \beta = 2 \cos \left(\frac{\alpha + \beta}{2} \right) \sin \left(\frac{\alpha - \beta}{2} \right) \quad 4.64$$

$$\cos \alpha + \cos \beta = 2 \cos \left(\frac{\alpha + \beta}{2} \right) \cos \left(\frac{\alpha - \beta}{2} \right) \quad 4.65$$

$$\cos \alpha - \cos \beta = -2 \sin \left(\frac{\alpha + \beta}{2} \right) \sin \left(\frac{\alpha - \beta}{2} \right) \quad 4.66$$

Professional Publications, Inc.


FERC

Mathematics 1

1-11a

General Triangles

Figure 4.11 General Triangle



$$\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} \quad 4.67$$

$$a^2 = b^2 + c^2 - 2bc \cos A \quad 4.68$$

$$b^2 = a^2 + c^2 - 2ac \cos B \quad 4.69$$

$$c^2 = a^2 + b^2 - 2ab \cos C \quad 4.70$$

Professional Publications, Inc.

FERC

Mathematics 1

1-11b

General Triangles

Example (EFPRB):

MATHEMATICS-19If $\sin \alpha = x$, what is $\sec \alpha$?

- (A) $\sqrt{1-x^2}$ (B) $\frac{x}{\sqrt{1-x^2}}$ (C) $\frac{1}{\sqrt{1-x^2}}$ (D) $\frac{x}{\sqrt{1+x^2}}$

Since $\sin \alpha$ is the side facing angle α divided by the hypotenuse, the hypotenuse = 1. Therefore,

$$\begin{aligned} \text{side adjacent to angle } \alpha &= \sqrt{1-x^2} \\ \sec \alpha &= \frac{\text{hypotenuse}}{\text{side adjacent to angle } \alpha} = \frac{1}{\sqrt{1-x^2}} \end{aligned}$$

The answer is (C).

Professional Publications, Inc.

FERC


Mathematics 1

1-11c

General Triangles

Example (EFPRB):

MATHEMATICS-21

If the sine of angle A is given as K , what is the tangent of angle A?

(A) $\frac{hK}{o}$

(B) $\frac{aK}{h}$

(C) $\frac{ha}{K}$

(D) $\frac{hK}{a}$

$$\begin{aligned}\sin A &= \frac{o}{h} \\ &= K \\ \tan A &= \frac{o}{a} \\ &= \left(\frac{h}{a}\right) \left(\frac{o}{h}\right) \\ &= \frac{hK}{a}\end{aligned}$$

The answer is (D).

Professional Publications, Inc.

FERC

Mathematics 1

1-11d

General Triangles

Example (EFPRB):

MATHEMATICS-22

Which is true regarding the signs of the natural functions for angles between 90° and 180° ?

- (A) The tangent is positive.
- (B) The cotangent is positive.
- (C) The cosine is negative.
- (D) The sine is negative.

In the second quadrant, the natural functions and their signs are as follows.

sin	positive
cos	negative
tan	negative
cot	negative
sec	negative
csc	positive

The answer is (C).

Professional Publications, Inc.

FERC

Mathematics 1

1-11e

General Triangles

Example (EFPRB):

MATHEMATICS-23

What is the inverse natural function of the cosecant?

- (A) secant (B) sine (C) cosine (D) tangent

In a right triangle, the cosecant is the hypotenuse divided by the opposite side. The sine is the opposite side divided by the hypotenuse.

$$\sin \theta = \frac{1}{\csc \theta}$$

The answer is (B).

Professional Publications, Inc.

FERC

Mathematics 1

1-11f

General Triangles

Example (EFPRB):

MATHEMATICS-24

What is the sum of the squares of the sine and cosine of an angle?

- (A) 0 (B) 1 (C) $\sqrt{3}$ (D) 2

For any angle,

$$\cos^2 x + \sin^2 x = 1$$

The answer is (B).

Professional Publications, Inc.

FERC

Mathematics 1

1-11g

General Triangles

Example (EFPRB):

MATHEMATICS-25

What is an equivalent expression for $\sin 2x$?

- (A) $\frac{1}{2} \sin x \cos x$ (B) $2 \sin x \cos \frac{1}{2}x$ (C) $-2 \sin x \cos x$ (D) $\frac{2 \sin x}{\sec x}$

The double angle formula for the sine function is

$$\begin{aligned}\sin 2x &= 2 \sin x \cos x \\ &= \frac{2 \sin x}{\sec x}\end{aligned}$$

The answer is (D).

Professional Publications, Inc.

FERC

Mathematics 1

1-12a

Mensuration of Areas and Volumes

Figure 4.12 Parabola





Figure 4.13 Ellipse





Figure 4.14 Circular Segment



$$A = \frac{1}{2}r^2(\phi - \sin \phi) \quad [\phi \text{ must be in radians}]$$

$$\phi = \frac{s}{r} = 2 \arccos\left(\frac{r-d}{r}\right)$$

Professional Publications, Inc.

FERC

Mathematics 1

1-12b

Mensuration of Areas and Volumes

Figure 4.15 Circular Sector





Figure 4.16 Parallelogram



Professional Publications, Inc.

FERC

Mathematics 1

1-12c

Mensuration of Areas and Volumes

Figure 4.17 Regular Polygon (n equal sides)





Figure 4.18 Sphere



Professional Publications, Inc.

FERC

Mathematics 1

1-12d

Mensuration of Areas and Volumes

Figure 4.19 Right Circular Cone





Figure 4.20 Right Circular Cylinder



Professional Publications, Inc.


FERC

Mathematics 1

1-12e

Mensuration of Areas and Volumes

Figure 4.21 Paraboloid of Revolution



Professional Publications, Inc.

FERC