# Foundations of Sensitivity Analysis

From Local Sensitivity to Global Uncertainty

Leon M. Arriola James M. Hyman

October 16, 2026

*Preliminary version v1_25_0, for review only*

Draft

© 2026 Leon M. Arriola and James M. Hyman.
All rights reserved.

**Conflict of interest.** The authors declare no competing financial interests or personal relationships that could have influenced the work reported in this book.

**Funding.** The development of this material was supported in part by the Tulane University Department of Mathematics and by the Mathematical and Theoretical Biology Institute (MTBI) at Arizona State University. The companion code was developed with support from Tulane University.

Draft

# Contents

**Preface** **xi**

**Notation and Conventions** **xvii**

**I The Sensitivity Question** **1**

**1 Introduction and Overview** **3**
1.1 The Central Question . . . . . 3
1.2 Two Directions: Forward and Adjoint Sensitivity . . . . 7
1.2.1 Forward Sensitivity Analysis: "What if?" . 8
1.2.2 Adjoint Sensitivity Analysis: "How come?" . 10
1.3 Global Sensitivity: When Local Is Not Enough . . . . . 10
1.4 Sensitivity Analysis and Uncertainty Quantification . . 11
1.5 Four Ideas That Run Through the Book . . . . . 12
1.6 A Note on Parameterization . . . . . 14
1.7 Structure of the Book . . . . . 15
1.8 Exercises . . . . . 17
Notes . . . . . 19

**2 Measuring Local Sensitivity** **21**
2.1 Ratios as Precursors to Derivatives . . . . . 21
2.2 Absolute, Relative, and Regularized Change . . . . . 22
2.2.1 Absolute Change . . . . . 22
2.2.2 Relative Change . . . . . 23
2.2.3 Regularized Change . . . . . 23
2.3 The Sensitivity Index . . . . . 24
2.3.1 The Sign Rule . . . . . 24
2.3.2 Formal Definition . . . . . 25
2.3.3 Interpreting the Magnitude . . . . . 27
2.4 Worked Examples . . . . . 27
2.4.1 Volume of a Cylinder . . . . . 27

Draft

2.4.2 Arc Length of a Line . . . 28
2.4.3 Area Under a Line . . . 29
2.4.4 Root of a Nonlinear Equation . . . 29
2.4.5 A Linear Prediction Model (Statistical Connection) . . . 30
2.5 The Sensitivity Index for the SIR Model . . . 30
2.5.1 Model Derivation from First Principles . . . 30
2.5.2 The Basic Reproduction Number . . . 32
2.5.3 Reparameterizing the SIR Model . . . 32
2.5.4 Analytic Computation . . . 34
2.6 Choosing Interpretable POIs . . . 34
2.7 The Tornado Plot . . . 36
2.8 A Preview: Second-Order Terms . . . 38
2.9 Computational Laboratory: Sensitivity Indices for the SIR Model . . . 38
2.10 Exercises . . . 40
Notes . . . 43

**II Computing Sensitivity in Practice 45**

**3 Computing Sensitivity in Practice 47**
3.1 The Sensitivity Jacobian . . . 47
3.2 Finite Difference Formulas . . . 48
3.2.1 Forward Difference . . . 48
3.2.2 Centered Difference . . . 49
3.2.3 Fourth-Order Formula (Rarely Used in Practice) . . . 49
3.2.4 Practical Workflow: Checking the Step with an Error Indicator . . . 49
3.2.5 Cost Comparison . . . 50
3.3 Step Size Selection: The U-Curve . . . 51
3.3.1 Two Sources of Error . . . 51
3.3.2 The U-Curve and the Optimal Step Size . . . 52
3.4 Second-Order Terms . . . 54
3.4.1 Pure Second-Order Derivatives . . . 55
3.4.2 Mixed Second-Order Derivatives . . . 55
3.5 Correlated POIs: The Most Common Mistake . . . 56
3.5.1 Diagnosing Correlation . . . 56
3.5.2 The Impact of Ignoring Correlation . . . 57
3.5.3 Four Practical Responses . . . 58

Draft

3.6 Black-Box Models and the Pseudocode Template . . . . 59
3.7 Computational Laboratory: Finite Differences . . . . . 61
3.8 Exercises . . . . . 63
Notes . . . . . 65

**III Analytic Forward Sensitivity Analysis 67**

**4 Analytic Forward Sensitivity Analysis 69**
4.1 The FSE Derivation Pattern . . . . . 69
4.1.1 The Pattern in Three Steps . . . . . 70
4.1.2 Application to Three Model Types . . . . . 70
4.2 Algebraic Systems and Equilibria . . . . . 71
4.2.1 A Single Nonlinear Equation . . . . . 71
4.2.2 Linear Systems: The FSE for $A\vec{u} = \vec{b}$ . . . . 72
4.2.3 The Competitive Lotka–Volterra Model . . . 72
4.2.4 The Car Rental Problem: O$\Delta$E Sensitivity . 74
4.3 Forward Sensitivity for ODE Systems . . . . . 75
4.3.1 Derivation of the ODE FSE . . . . . 75
4.3.2 Time-Dependent Sensitivity Indices . . . . . 76
4.3.3 The SIR Model: Full FSE Treatment . . . . 76
4.4 The Proliferation Problem and Its Consequences . . . . 77
4.5 Choosing a Method . . . . . 80
4.6 Computational Laboratory: Analytic FSA of the SIR Model . . . . . 81
4.7 Exercises . . . . . 82
Notes . . . . . 84

**IV The Adjoint Approach 87**

**5 The Adjoint in Linear Algebra 89**
5.1 A Preview: The Same Answer, Two Solves Instead of Fifty-One . . . . . 89
5.2 The Strategy, Stated Explicitly . . . . . 91
5.3 The Commutative Diagram . . . . . 93
5.4 Cost Comparison and the Decision Rule . . . . . 94
5.4.1 When to Use the Adjoint . . . . . 94
5.4.2 Cost Comparison Plot . . . . . 95
5.5 The Competitive Population: Full Adjoint Treatment . 96
5.5.1 FSE Approach: Three Parameters, Four Solves 96

Draft

5.5.2 Adjoint Approach: Three Solves for Two Response Functionals . . . 96
5.5.3 Verification . . . 97
5.6 Reverse Mode and Backpropagation . . . 97
5.6.1 The Computation Graph . . . 97
5.6.2 Forward Mode: One Pass Per Input . . . 97
5.6.3 Reverse Mode: Two Passes for All Inputs . . . 98
5.6.4 The Backpropagation Connection . . . 98
5.7 Bridge to Chapter 6 . . . 100
5.8 Computational Laboratory: Forward vs. Adjoint . . . 101
5.9 Exercises . . . 102
Notes . . . 105

**6 The Adjoint for Dynamical Systems 107**
6.1 The Timeline Figure . . . 108
6.2 Why Backward? Intuition Before Derivation . . . 108
6.3 The Lagrange Multiplier Derivation . . . 109
6.3.1 A Warm-Up: Exponential Decay . . . 109
6.3.2 The Five-Step Derivation . . . 110
6.3.3 Setup . . . 110
6.3.4 The Vector Case . . . 115
6.4 The SIR Model: Full Adjoint Treatment . . . 116
6.4.1 Setup and Response Functional . . . 117
6.4.2 The Jacobian . . . 117
6.4.3 Adjoint ODEs . . . 117
6.4.4 Sensitivity Formulas . . . 118
6.4.5 The Adjoint Variable and Its Interpretation 118
6.5 Discrete vs. Continuous Adjoint . . . 121
6.6 Computational Laboratory: The SIR Adjoint in Practice 122
6.7 Exercises . . . 124
Notes . . . 127

**7 Adjoint Sensitivity in Practice 129**
7.1 The Annuity Function: A Worked Comparison . . . 129
7.1.1 The Problem . . . 129
7.1.2 Forward Mode: Three Passes . . . 130
7.1.3 Reverse Mode: Two Passes . . . 130
7.1.4 Symbolic Differentiation . . . 131
7.1.5 Exact Operation Counts . . . 131
7.2 Algorithmic Differentiation . . . 134
7.2.1 Forward Mode . . . 134

Draft

7.2.2 Reverse Mode . . . 134
7.2.3 Available Tools . . . 135
7.3 The Master Decision Table . . . 136
7.4 Practical Guidance and Common Pitfalls . . . 138
7.4.1 Discretize-Then-Adjoint vs. Adjoint-Then-Discretize 138
7.4.2 Memory Requirements for the Adjoint . . . 138
7.4.3 Adaptive Step-Size Solvers . . . 139
7.4.4 Discontinuities Break AD . . . 139
7.5 Computational Laboratory: Four Methods, One Model . 140
7.6 Exercises . . . 142
Notes . . . 144

**V Limitations and Extensions 147**

**8 Caveat Emptor, Limitations of Local Sensitivity Analysis 149**
8.1 Method-Question Mismatch . . . 150
8.2 Bifurcations and SA Failure . . . 151
8.2.1 The Quadratic Example . . . 151
8.2.2 The Epidemic Threshold . . . 153
8.3 Ill-Conditioning . . . 157
8.3.1 The Condition Number . . . 157
8.3.2 SA and Ill-Conditioning . . . 158
8.4 Separatrices and Trajectory Sensitivity . . . 159
8.4.1 What Is Differentiated Decides the Failure . 160
8.5 When Global SA Is Needed . . . 161
8.5.1 Large Parameter Uncertainty . . . 161
8.5.2 Highly Nonlinear Models . . . 162
8.5.3 Correlated Parameters . . . 162
8.5.4 The Portfolio Example . . . 162
8.6 When SA Guides Data Collection . . . 164
8.7 Computational Laboratory: Bifurcation and Ill-Conditioning 166
8.8 Exercises . . . 167
Notes . . . 169

**9 Global Sensitivity Analysis and Sobol Indices 171**
9.1 Why Local SA Is Not Enough: The Sigma-Normalized Bridge . . . 172
9.1.1 The Portfolio Example . . . 172
9.1.2 Two SIR Models . . . 173
9.1.3 Time-Varying Sensitivity . . . 173
9.2 Sampling Methods . . . 174

Draft

9.2.1 Random Sampling . . . 174
9.2.2 Latin Hypercube Sampling . . . 175
9.2.3 Low-Discrepancy Sampling and Scrambled Sobol' Sequences . . . 175
9.3 Regression-Based SA: LHS/PRCC . . . 177
9.3.1 The PRCC Calculation . . . 178
9.3.2 When PRCC is Appropriate . . . 178
9.4 Sobol Decomposition and Variance-Based SA . . . 179
9.4.1 The Intuition: Conditional Expectation . . . 179
9.4.2 The ANOVA Decomposition . . . 179
9.4.3 Sobol Indices . . . 180
9.5 Computing Sobol Indices: The Saltelli Algorithm . . . 182
9.5.1 The Saltelli Construction . . . 182
9.5.2 The Jansen Estimators . . . 183
9.5.3 Convergence . . . 184
9.6 Morris Screening . . . 187
9.7 Sobol Indices for Correlated POIs . . . 190
9.8 Beyond Variance: Moment-Independent Sensitivity . . . 191
9.9 Comparison of the Main SA Methods . . . 193
9.10 Verifying an SA Result You Did Not Compute . . . 195
9.10.1 An audit in six questions . . . 195
9.10.2 A worked audit . . . 196
9.10.3 The referee's version . . . 197
9.11 Bridge to the Broader UQ Landscape . . . 198
9.12 Computational Laboratory: Complete Global SA Pipeline for SIR . . . 199
9.13 Exercises . . . 200
Notes . . . 204

**Appendices** **209**

**A Inner Products, Norms, and Function Spaces** **209**
A.1 Inner Products and Norms . . . 209
A.2 The $L^2$ Inner Product . . . 210
A.3 Integration by Parts as Transposition . . . 210
A.4 Orthogonality and the ANOVA Decomposition . . . 210
A.5 Exercises . . . 211

**B The Adjoint Formalism** **213**
B.1 Linear Operators and Their Adjoints . . . 213
B.2 The Lagrange Identity . . . 214

Draft

B.3 Solvability Conditions . . . . . 214
B.4 Connection to Chapters 5 and 6 . . . . . 215
B.5 Exercises . . . . . 215

**C Eigenvalue Sensitivity and Singular Value Decomposition 217**
C.1 Sensitivity of Eigenvalues . . . . . 217
C.1.1 Symmetric Matrices . . . . . 217
C.1.2 Non-Symmetric Matrices . . . . . 217
C.2 Singular Value Decomposition . . . . . 219
C.2.1 SVD and the Condition Number . . . . . 219
C.2.2 SVD and SA . . . . . 219
C.2.3 SVD and the Pseudoinverse . . . . . 219
C.3 Exercises . . . . . 220

**D Periodic Solutions and Floquet Theory 223**
D.1 Floquet Theory . . . . . 223
D.2 Sensitivity of the Period . . . . . 224
D.3 The van der Pol Example . . . . . 224
D.4 Sensitivity of Floquet Multipliers . . . . . 225
D.5 Exercises . . . . . 226

**E The Adjoint in Optimization 227**
E.1 Linear Programming Duality . . . . . 227
E.2 Sensitivity of the Optimal Value . . . . . 228
E.3 Quadratic Programming . . . . . 228
E.4 Connection to Optimal Control . . . . . 229
E.5 Exercises . . . . . 229

**Bibliography 233**

**Glossary 239**

**Index 245**

Draft

Draft

# Preface

This book develops sensitivity analysis (SA) as a mathematical discipline, not merely as a collection of computational recipes. SA asks the foundational question of quantitative modeling: which parameters most strongly determine the model's predictions?

Four ideas run through every chapter:

1. The sensitivity index is a normalized derivative.
2. The adjoint is the transpose, in whatever space you work.
3. Local SA is a linearization, with all the limitations that implies.
4. A sensitivity index describes a model, not the world.

Students who understand these four ideas (not just their definitions but their implications and connections) will be equipped to design, implement, and critique SA studies in any application domain.

The SIR epidemic model serves as the running example throughout. Every method is demonstrated on this one model, so students can track increasing mathematical sophistication against a fixed biological context. All computational examples use MATLAB; each chapter after the first includes a structured laboratory and a reusable code template. Companion Colab notebooks, one per chapter, recompute examples from the text in Python and assert selected results against fresh computation. The MATLAB reference library (eighteen functions across four modules, with an automated test suite) is available alongside them. A solutions manual covers every exercise and is written for self-study as much as for use as an instructor's key. The notebooks, the library, and the manual are hosted at:

`https://github.com/machyman/arriola2026sa`

The adjoint method (Chapters 5–7) is the conceptual centerpiece of the book. It is the same idea encountered three times in three different mathematical spaces: the matrix transpose in linear algebra, the reversed computation graph in algorithmic differentiation, and a backward ODE in dynamical systems. Chapter 8, *Caveat Emptor*, asks when sensitivity anal-

ysis can mislead, and is the chapter students most frequently cite as the one that changed how they think about mathematical models.

**Relationship to existing literature.** Cacuci (1981) established the adjoint formalism for nonlinear systems; Giles and Pierce (2000) developed it for aeronautical design; Cao et al. (2002) gave the computational treatment for ODE systems; and Griewank and Walther (2008) treat the computation-graph view comprehensively. The present book differs from each of these in its pedagogical aim: it is one of the few texts, and to our knowledge the first at this level, to introduce all three forms of the adjoint (matrix, graph, and ODE) as instances of a single unifying idea, and that teaches students when *not* to trust the results they compute.

The last of those is where this book departs furthest from its neighbors. Chapter 9 closes with a short protocol for auditing a sensitivity analysis someone else performed. Six questions establish which decision the result informs, whether the index reported answers that question, and whether the method's assumptions were met. We know of no comparable treatment in the existing texts, and it is the part of the book we would point a referee to first.

The closest neighbor is Smith's *Uncertainty Quantification* (2024), which SIAM offers as a one-semester sensitivity analysis course text, and the overlap is larger than the titles suggest. The difference is level and route rather than territory. Smith assumes a graduate background and reaches sensitivity analysis through the full uncertainty-quantification apparatus. This book is pitched at a more elementary audience: it assumes calculus, linear algebra, and ODEs, and reaches sensitivity analysis through one worked epidemic model. It can be read first, or in parallel with the sensitivity analysis chapters there.

**A note on register.** Results are developed in the running text rather than stated as numbered theorems. The book contains two theorems and one proposition, and the sparseness is deliberate: at this level the derivation is what carries the understanding, and a formal statement placed beside it tends to be read instead of the argument rather than alongside it. Where a result needs to be quotable it is displayed and labeled.

## To the Instructor

The book supports a one-semester course for advanced undergraduates and beginning graduate students. Students need calculus, linear algebra, and ODEs; no measure theory or functional analysis is assumed. A minimal one-semester course can omit the computational AD material in Chapter 7 and all appendices without loss of continuity. Chapters 1–4 and 8–9 form

a self-contained global-SA course for students who will not use the adjoint in their research. Exercises tagged [Bloom L5] and [L6] are designed for semester projects and take-home examinations; those tagged [L2]–[L4] are suitable for weekly problem sets. Every chapter but the first closes with a Computational Laboratory. Chapter 1 has none by design: it orients the reader before any method has been introduced, so there is nothing yet to compute. Chapter 9 is twice the length of a typical chapter and is best given two weeks; the others take one.

One further word on assessment. Students now have capable computational assistants, and the exercises in this book are not equally exposed to them. An assistant will produce a competent answer to most [L2] and [L3] exercises, which are recall and routine application, so those are best used for practice rather than for grading, or set under supervision. The [L5] and [L6] project exercises are far less exposed, because they require judgments about a specific model that an assistant cannot make on the student's behalf. The audit exercises introduced in this edition, marked as such in their statements, are deliberately resistant for a different reason: each presents a plausible finished analysis and asks what is wrong with it. Producing a confident answer is easy, and recognizing that a confident answer addresses the wrong question is the skill being assessed. Worked solutions to every exercise are public, in the solutions manual posted with the notebooks, so an exercise used for grading is best varied from its printed form.

**Prerequisites.** Students should have completed: (1) calculus through multivariable (partial derivatives, chain rule, change of variables); (2) linear algebra through eigenvalues and the matrix transpose; (3) a first course in ordinary differential equations (existence and uniqueness, systems of ODEs, basic phase-plane analysis); and (4) introductory MATLAB or a comparable computing environment (the first laboratory session introduces every MATLAB construct the book uses, so prior MATLAB experience is helpful but not required). No probability theory, measure theory, or functional analysis is assumed.

**For students.** You will almost certainly have a computational assistant available while you work through this book, and you should use it. It will differentiate an expression, set up a sampling scheme, or write a Sobol estimator faster and more reliably than you can, and there is nothing to be gained by pretending otherwise.

What it cannot do is tell you which sensitivity index answers your question. That choice depends on what you intend to do with the answer, on which simplifications your model already contains, and on whether the assumptions behind a method hold for the system in front of you. None of that is visible in the expression you hand over. An assistant asked which pa-

rameter matters most will return a ranking, and it will return one whether or not the question was well posed. Table 9.4 in Chapter 9 catalogs the answers that follow from asking the wrong version of that question, and it was written from the published literature rather than from anything a machine produced. The errors are ours first.

The book therefore asks you to develop a habit that is useful well beyond this subject: treat any result you did not derive yourself as provisional until you have checked what question it answers and what it assumes. Section 9.10 sets out how to do that, and several exercises give you finished analyses to audit rather than problems to solve.

**In memory of Leon M. Arriola (1957–2026).**

Leon Arriola and I began working on sensitivity analysis together more than twenty-five years ago. What started as a shared mathematical project became, over time, one of the sustaining collaborations of my life. Leon brought deep insight, patience, humor, and generosity to the work. He cared deeply about helping others learn, and every page of this book reflects his commitment to making hard ideas genuinely accessible.

Leon passed away in April 2026, before he could see the manuscript reach its final form. Completing this book, and offering it to the mathematical community in his name, is one way I know to carry forward what he taught me, and what he gave so freely to his students and colleagues over a lifetime of teaching.

*James M. Hyman, Tulane University*
*New Orleans, Louisiana*
*April 2026*

**Acknowledgments.** I thank the students in our sensitivity analysis courses at the University of Wisconsin-Whitewater, Tulane University, and the Mathematical and Theoretical Biology Institute (MTBI) at Arizona State University, whose questions and confusion shaped every page of this book.

I am grateful to the colleagues who generously read drafts and offered suggestions that improved the manuscript. I particularly thank Ralph C. Smith (North Carolina State University) for his careful reading and detailed comments.

**Declaration on the use of AI tools.** Generative AI tools were used in preparing this book: in drafting and revising narrative text, in writing and testing the companion code, in checking algebraic and numerical results against independent computation, and in literature search. This assistance was subject to the following verification. The companion notebooks, which readers can run for themselves, recompute examples from each chapter and

assert values against fresh computation rather than merely printing them. A ledger checked against the source before each release accounts for every decimal value and every computed count that the chapters and appendices quote: each is reproduced by a notebook check, verified as data the text supplies or as printed arithmetic, or classified with its reason recorded. Values quoted from a figure are reproduced instead by the generator of that figure. Every figure generated from computation is produced by a generator included in the source distribution, and those generators are byte-reproducible. Every bibliographic entry has been checked against a primary source, and the checks are recorded in the `annote` field of each entry in the `.bib` file; that process corrected four incorrect DOIs and errors of volume, number, journal or author list in thirteen further entries. This work was carried out after Leon Arriola's death, and James Hyman alone is accountable for it. The authors assume responsibility for all content.

The companion code was developed with support from the Tulane University Department of Mathematics.

**Dedication.**

*To Leon, who taught us to ask which parameters matter*
*— and to our students, past and future,*
*who will carry the question forward.*

Draft

Draft

# Notation and Conventions

This book uses one running model and a compact, consistent notation. The tables below collect the symbols that recur across chapters, so a reader who meets a symbol mid-derivation can find its meaning without hunting for its first appearance. A symbol defined and used only within a single example is not listed here.

Three typographic conventions hold throughout. Vectors are set with an arrow, $\vec{u}$; matrices are uppercase upright or italic by context, $A$, $J_F$; and scalars are italic, $c$, $\beta$. A subscript on a rate or time constant names the compartment or transition it governs ($\gamma_R$, $\tau_m$), never a running index. A sensitivity index is written $\mathcal{S}$ with the parameter as subscript and the response as superscript: $\mathcal{S}_\beta^J$ is the sensitivity of the response $J$ to the parameter $\beta$.

The objects the framework is stated in terms of recur throughout and are collected here.

| Symbol | Meaning |
|---|---|
| $\vec{u}(t)$ | the model state vector; for the SIR model, $(S, I, R)$ |
| $p$ | a generic parameter (parameter of interest, POI) |
| $q$, $Q$, $Y$ | a generic response (quantity of interest, QOI); $Y$ in the global-SA chapter |
| $T$ | terminal time of the adjoint / optimal-control problem |
| $A$ | a generic matrix (eigenvalue and SVD sections) |
| $D$, $D_i$ | total and partitioned output variance in the Sobol decomposition |

## The sensitivity index

The central object is the normalized (relative) sensitivity index. For a scalar response $Q$ and a parameter $p$,

$$\mathcal{S}_p^Q \;=\; \frac{p}{Q}\,\frac{\partial Q}{\partial p},$$

the fractional change in $Q$ per fractional change in $p$. It is dimensionless, and $\mathcal{S}_p^Q = 1$ means unit elasticity: a one-percent change in $p$ produces a

one-percent change in $Q$. The book writes *parameter of interest* (POI) for the input under study and *quantity of interest* (QOI) for the response.

| Symbol | Meaning |
|---|---|
| $\mathcal{S}_p^Q$ | normalized sensitivity index of response $Q$ to parameter $p$ |
| $S_i$ | first-order Sobol index of parameter $i$ |
| $S_{T_i}$ | total-effect Sobol index of parameter $i$ |
| $S_i^\sigma$ | $\sigma$-normalized (uncertainty-weighted) index of parameter $i$ |
| $\mu_i^*$ | Morris mean of absolute elementary effects for parameter $i$ |
| $\delta_i$ | Borgonovo (moment-independent) delta index |
| $\eta^2$ | correlation ratio / first-order effect measure |

The distinctions among $S_i$, $S_{T_i}$, $S_p^\sigma$, and $\mu_i^*$, which question each answers, and which error each guards against, are set out in Table 9.4 in Chapter 9. That table, not this one, is the place to resolve which index a given task requires.

## The adjoint and operator symbols

| Symbol | Meaning |
|---|---|
| $\vec{\lambda}$ | adjoint variable (costate); components $\lambda_S, \lambda_I, \lambda_R$ |
| $J_F$ | Jacobian of the model right-hand side $F$ |
| $F_p$ | partial derivative of $F$ with respect to a parameter $p$ |
| $g_u$ | partial derivative of the integrand $g$ with respect to the state |
| $L,\ L^\dagger$ | a linear operator and its adjoint |
| $\kappa$ | condition number ($\kappa_2 = \sigma_1/\sigma_n$) |
| $\Phi(t)$ | fundamental matrix (Appendix D) |
| $\Phi_j$ | normalized Shapley effect of parameter $j$ (Chapter 9) |
| $N$ | sample size, as in the Saltelli cost $N(m+2)$ (Chapter 9) |

## The SIR model and its parameters

Every method in the book is demonstrated on one susceptible–infectious–recovered model. Its parameters are the following.

| Symbol | Meaning | Nominal |
|---|---|---|
| $S, I, R$ | susceptible, infectious, recovered counts | – |
| $N$ | total population, $S + I + R = N$ | – |
| $c$ | contacts per individual per day | 5 |
| $\beta$ | transmission probability per contact | 0.06 |
| $\tau_R$ | mean infectious period (days) | 7 |
| $\gamma_R$ | recovery rate, $\gamma_R = 1/\tau_R$ | – |
| $\tau_m$ | mean time in the population (days) | 10,000 |
| $\nu_m$ | demographic rate, $\nu_m = 1/\tau_m$ | – |
| $\mathcal{R}_0$ | basic reproduction number, $c\beta\tau_R$ | 2.1 |
| $\lambda(t)$ | force of infection, $c\beta I(t)/N$ | – |

## A note on $\lambda$

One Greek letter carries more than one meaning in this book, by established convention in the fields it borrows from, and the reader who knows which is which will not be misled.

In the epidemic model, $\lambda(t) = c\beta I(t)/N$ is the *force of infection*, the instantaneous rate at which a susceptible individual becomes infected. This is the universal notation of mathematical epidemiology, and the book keeps it. It is written as $\lambda$ only in the modeling chapters (Chapters 2 and 3), where no adjoint is present.

In the sensitivity chapters, $\vec{\lambda}$ is the *adjoint variable*, or costate. It is a vector, carrying one component per state equation ($\lambda_S$, $\lambda_I$, $\lambda_R$), and the book writes it with an arrow wherever it is a vector and with a subscript for a named component. Chapter 5 first uses it at the nodes of a computation graph, as $\lambda_i = \partial J/\partial v_i$ (Section 5.6), and previews the adjoint ODE for $\lambda(t)$. Chapter 6 introduces that ODE first for a single-state equation, where the adjoint is genuinely a scalar $\lambda(t)$; from the vector case onward it carries the arrow. The two meanings of $\lambda$ never appear in the same equation. When an adjoint chapter needs the force of infection, it writes the product $c\beta I(t)/N$ in full rather than $\lambda(t)$.

Two further uses are confined to the appendices and are standard there. In Appendix C, $\lambda_k$ is an *eigenvalue*. In Appendix E, $\vec{\lambda}$ is the vector of *Karush–Kuhn–Tucker multipliers* of a constrained optimization problem. That the multiplier and the adjoint share a symbol is not a coincidence: Appendix E shows they are the same object, the sensitivity of an optimum to a constraint, seen from two directions.

## A note on $\sigma$

Like $\lambda$, the letter $\sigma$ carries more than one standard meaning, and position and subscript make clear which is intended. As a *standard deviation*, $\sigma_i$ is the uncertainty in parameter $i$ and appears in the $\sigma$-normalized index $S_i^{\sigma}$ (Chapter 9). As *biting rates*, $\sigma_v$ and $\sigma_h$ are model parameters in the vector-borne example (Chapter 3). As *singular values*, $\sigma_1 \geq \sigma_2 \geq \cdots \geq \sigma_n$ are the diagonal entries of the SVD and define the condition number $\kappa_2 = \sigma_1/\sigma_n$ (Chapter 8, Appendix C). And $\sigma(\cdot)$ with a function argument is the *sigmoid* used to restore differentiability at a switch (Chapter 7). These uses do not overlap within any one expression.

## A note on $N$

The capital $N$ is the third symbol in this book to carry two standard meanings, and the two are separated by chapter rather than by decoration. In the modeling chapters, $N$ is the *total population*, $S + I + R = N$, which is the universal notation of mathematical epidemiology and appears in the force of infection $c\beta I(t)/N$.

In Chapter 9, $N$ is the *sample size*: the number of parameter vectors drawn from the joint distribution, and the base sample size in the Saltelli cost $N(m + 2)$. This is equally standard in the sensitivity literature, where $N(m+2)$ is the customary way to state the price of a Sobol' analysis, and the book keeps it so that a reader moving between this text and that literature meets the same expression.

The two never appear together. Chapter 9 carries no population $N$: its epidemic examples are written in terms of $\mathcal{R}_0 = c\beta/\gamma_R$ and of sample estimates, not of the force of infection. A reader who keeps the chapter in mind will not be misled, and a subscripted $N$ always denotes a population subgroup, as in the vector and host populations $N_v$ and $N_h$ of the malaria example (Chapter 3).

Draft

# Part I

# The Sensitivity Question

Draft

Draft

# Chapter 1
# Introduction and Overview

*Sensitivity analysis begins when a model is forced to say what matters.*

An infectious disease has just entered a new country. The public health response team has built a compartmental model with fourteen parameters: transmission rates, recovery rates, contact patterns, immunity duration, disease-induced mortality, and more. The team has funding to deploy exactly two interventions and time to measure exactly three parameters with high precision before a policy decision must be made.

The question due by tomorrow morning is this: *which three parameters are worth measuring, and which two interventions will most effectively reduce the spread?*

This is the problem that sensitivity analysis is designed to answer. Before the mathematics, we need a language for asking the question precisely.

## 1.1 ▪ The Central Question

Every mathematical model takes inputs and produces outputs. The inputs include parameters whose exact values may be uncertain or whose values might be changed through intervention. The outputs include whatever the model is designed to predict. Sensitivity analysis asks: *which inputs have the most effect on the outputs we care about?*

To answer this precisely, we adopt two terms used throughout the book.

A *Parameter of Interest* (POI) is any input to the model whose variation is worth studying. In practice, a parameter becomes a POI for one of three reasons, and the reason matters for how you interpret the sensitivity index.

1. *Epistemic uncertainty* (uncertainty from lack of knowledge). The parameter is fixed by nature but imperfectly known: different studies

report different estimates, the data are sparse, or the measurement process is imprecise. Epistemic uncertainty is *reducible*: better data, improved experiments, or longer observation windows can sharpen the estimate. Sensitivity analysis identifies which epistemically uncertain parameters most affect the output, directing limited data-collection resources toward the measurements that would reduce output uncertainty the most. In the disease example, the transmission probability per contact is epistemically uncertain: studies estimate it from contact-tracing data, but estimates vary across populations and settings.

2. *Aleatory uncertainty* (inherent stochastic variability). The parameter genuinely varies from individual to individual, place to place, or time to time, and this variation is *irreducible*: it persists even with unlimited data. More observations characterize the distribution more precisely but do not eliminate the randomness. In the disease example, the contact rate varies stochastically across the population: individuals differ in how many contacts they make per day. Sensitivity analysis with aleatory uncertainty typically asks how much of the output variance is attributable to each stochastic source, motivating the global SA methods of Chapter 9.

3. *Control variable* (a lever the investigator can move). The parameter is known and could, in principle, be changed by a policy decision, intervention, or design choice. Sensitivity analysis in this context answers the question: which intervention has the most leverage over the outcome I care about? In the disease example, the contact rate can be reduced by social distancing policies, and the transmission probability can be reduced by mask requirements or vaccination. The sensitivity index tells the policy maker which reduction, contacts or transmission, produces the larger reduction in $\mathcal{R}_0$, or else, as happens for $\mathcal{R}_0$ in Chapter 2, that the two are equally leveraged and the choice has to be made on feasibility and cost instead.

Many parameters fall into more than one category simultaneously: the contact rate is both uncertain (its current value is imprecisely known) and controllable (it can be targeted by policy). In practice, the interpretation of a sensitivity index should always be read in light of which role the POI is playing. A large sensitivity index for an epistemically uncertain parameter is a call to collect better data; for an aleatory parameter, it is a call to characterize the distribution more carefully; for a control variable, it is a call to prioritize that intervention.

In the disease example above, the transmission probability per contact is primarily a POI because it is epistemically uncertain (different studies report different values) and because it can be targeted by an intervention (vaccines reduce it). The contact rate is a POI for the same two reasons: it varies across populations (epistemic and aleatory uncertainty) and can be reduced through social distancing policies (control variable).

A *Quantity of Interest* (QOI) is any model output that the investigator cares about predicting or controlling. The peak number of infectious individuals, the total number of deaths over a six-month period, and the basic reproduction number $\mathcal{R}_0$ are all plausible QOIs for an epidemic model. The choice of QOI is not neutral: different choices lead to different rankings of parameter importance.

The relationship between POIs and QOIs is the subject of this book. Specifically, we want to know how much each POI affects each QOI, so that we can rank parameters by importance, design interventions efficiently, and allocate measurement resources wisely.

Draft

*For Readers with a Statistics Background*

The POI/QOI vocabulary will feel familiar: a POI resembles a predictor variable and a QOI resembles a response variable from regression. The sensitivity index introduced in Chapter 2 is the same quantity known as an *elasticity* in econometrics (the percentage change in output per percentage change in input); a dimensionless measure of proportional change. In Chapter 8 the sigma-normalized sensitivity index will turn out to be identical to the standardized regression coefficient reported by most statistical software. In Chapter 9 the first-order Sobol index is the proportion of output variance explained by one parameter acting alone, the same quantity as $\eta^2$ (eta-squared) in a classical ANOVA (the ratio of between-group variance to total variance), and the same idea behind permutation feature importance and Shapley-value attribution (a game-theoretic method that allocates a model's output among its inputs) in machine learning.

**For readers from machine learning:** Forward sensitivity analysis is the local, interpretable cousin of computing a gradient; it measures how the output changes when one input is perturbed. *Forward-mode* automatic differentiation (AD) does this one parameter at a time, propagating dual numbers through the computation; it is the analogue of the FSE in Chapter 4. `torch.autograd` and similar frameworks, however, use *reverse-mode* AD (backpropagation) by default, which computes gradients with respect to all inputs in a single backward pass. This is the adjoint method of Chapters 5–7, applied to a computation graph rather than an ODE. The adjoint method (Chapters 5–7) is, mathematically, the same operation as backpropagation in a neural network: the chain rule applied in reverse order, computing the sensitivity of one output with respect to all inputs in a single backward pass. If you have trained a neural network, you have already used the adjoint method under a different name.
What this book adds beyond regression and ML is threefold: how to compute these quantities for models defined by equations rather than data; when the linearization underlying all of these quantities breaks down and global variance-based methods are required; and the mathematical unification showing that gradient descent, backpropagation, and sensitivity analysis are facets of the same idea.

One of the most persistent misconceptions in mathematical modeling is the assumption that all parameters matter equally until proven otherwise. They do not. In virtually every model studied in this book, a small number

of POIs account for most of the variation in the QOIs, while the remaining parameters have negligible effect. Identifying that small number is one of the primary goals of sensitivity analysis, and the answer is rarely obvious from the model equations alone.

*Self-check.* In the outbreak scenario above, classify each of the following as a POI, a QOI, or neither: the transmission probability per contact; the peak number of infectious individuals; the day the first case was reported; the mean number of days a patient is infectious.

*Answer:* POI: transmission probability (uncertain; can be reduced by vaccines), mean infectious period (uncertain; can be reduced by treatment). QOI: peak number of infectious individuals (model output we care about). Neither: day of first case (a historical fact, not a model parameter or prediction). Note that the mean infectious period could also be a QOI in a different study; this classification depends on the question being asked.

Figure 1.1 illustrates the basic structure. The forward problem takes nominal POI values as input and produces the QOI as output. Sensitivity analysis adds one more step: it asks how the QOI changes as the POIs vary.





**Figure 1.1.** *The forward problem takes nominal values for the input* POIs *and produces the associated output* QOI*. Sensitivity analysis quantifies how changes in the* POIs *propagate to the* QOI *by computing the normalized partial derivative* $\mathcal{S}_{p_i}^{q_j}$ *for each* POI–QOI *pair.*

## 1.2 ▪ Two Directions: Forward and Adjoint Sensitivity

There are two complementary ways to ask the sensitivity question, and they differ in a fundamental way.

Figure 1.2 captures the distinction geometrically. In panel (a), a perturbation to one input POI, $p_j$, spreads forward through the model and changes the output QOI. This is *forward sensitivity analysis*. Panel (b) reverses the direction: a specified variation in the QOI is traced backward to every POI that could have caused it. This is *adjoint sensitivity analysis*.

This book counts the cost of a method in solves, and the baseline solve at the nominal values always counts. Perturbing each of $m$ POIs once then

costs $m + 1$ solves. For one QOI the adjoint costs two: the baseline forward solve and one backward adjoint solve. For $r$ QOIs it costs $r + 1$: the same forward solve plus one adjoint solve per QOI.





**Figure 1.2.** *(a) Forward sensitivity analysis answers "What if?" questions. A perturbation to one input* POI $p_j$ *propagates forward through the model and produces a change in the output* QOI *. Because the model must be solved once at the baseline and once for each* POI*, forward sensitivity requires $m + 1$ solves for $m$* POIs*. (b) Adjoint sensitivity analysis answers "How come?" questions. A specified variation $\delta q_j$ in the output* QOI *enters the model from the right and is propagated* backward *through it. A single adjoint solve simultaneously recovers the sensitivity of $q_j$ with respect to every* POI $p_1, \ldots, p_m$*. The adjoint is developed in Chapters 5 and 6; for now, the key contrast is computational: forward SA costs $m + 1$ solves, adjoint SA costs* 2*.*

The "What if?" and "How come?" labels are not merely rhetorical. They correspond to genuinely different computational strategies and genuinely different questions.

### 1.2.1 ▪ Forward Sensitivity Analysis: "What if?"

Consider a disease model for an emerging outbreak. The transmission probability $\beta$ is estimated from early case data, but the estimate carries significant uncertainty. A public health officer asks: *if our estimate of $\beta$ is wrong by 10%, how much does our predicted peak infection count change?*

This is a forward question. We are tracing the effect of a change in an input POI (the transmission probability) forward to an output QOI (the peak infection count). Forward sensitivity analysis answers this question by computing a sensitivity index: a number that tells us, for a small percentage change in a POI, what percentage change to expect in the QOI.

*Connection to gradient computation.* Forward SA is the interpretable, application-oriented cousin of computing a gradient. When a machine-learning framework computes $\partial \mathcal{L}/\partial \theta$ for one parameter $\theta$ using forward-mode automatic differentiation, or when you perturb one input and measure the resulting output change (an ablation study), you are performing forward sensitivity analysis. The sensitivity index normalizes that gradient into a dimensionless, unit-free quantity that can be compared across parameters measured on different scales. Figure 1.3 contrasts the two directions on a single small computation graph: one forward pass per input, against one backward pass per output.





**Figure 1.3.** *Computation graph for $f(x_1, x_2) = (x_1 + x_2) \cdot x_2$, with intermediate node $v = x_1 + x_2$, illustrating the two modes of sensitivity analysis. (a) Forward SA propagates tangent vectors $(\dot{x}_1, \dot{x}_2)$ from each input to the output; one forward pass per input (POI). This is identical to forward-mode automatic differentiation. (b) Adjoint SA propagates the adjoint $\bar{f} = 1$ backward from the output to all inputs simultaneously, one backward pass per output (QOI). Edge weights are local partial derivatives: $\bar{v} = \bar{f} \cdot (\partial f/\partial v) = x_2$; the direct $f \to x_2$ edge carries $\bar{f} \cdot v = x_1 + x_2$. Final gradients: $\bar{x}_1 = \bar{v} = x_2$; $\bar{x}_2 = \bar{v} + \bar{f} \cdot v = x_2 + (x_1 + x_2) = x_1 + 2x_2$. This is identical to backpropagation in a neural network. The cost advantage: forward mode costs $m$ passes (one per input, each also evaluating $f$); adjoint mode costs two passes, one forward and one backward, regardless of the number of inputs.*

A sensitivity index of $+0.8$ would mean that a 10% increase in $\beta$ produces approximately an 8% increase in the peak infection count. A sensitivity index of $-1.0$ for the recovery rate would mean that a 10% increase in the recovery rate produces a 10% decrease in the peak. The sign tells you the direction; the magnitude tells you the amplification. Chapter 2 develops this definition carefully and shows how to compute these indices for a wide range of models.

*Self-check.* A forward sensitivity analysis gives $S^{\mathcal{R}_0}_{\tau_R} = +1.0$ (the notation is defined formally in Chapter 2), where $\tau_R$ is the mean infectious period (in days) and $\mathcal{R}_0$ is the basic reproduction number. If the mean infectious period increases by 5%, approximately what happens to $\mathcal{R}_0$? Is this good

news or bad news for disease control?

*Answer:* $\mathcal{R}_0$ increases by $5\% \times 1.0 = 5\%$. The change is exact here, because $\mathcal{R}_0 = c\beta\tau_R$ is proportional to $\tau_R$. This is bad news: a larger $\mathcal{R}_0$ means more transmission. The positive sign tells us that POI and QOI move in the same direction, so longer infectious periods lead to more disease spread.

### 1.2.2 ▪ Adjoint Sensitivity Analysis: "How come?"

Now consider a different problem. City planners want to site a new industrial facility near a residential area. The facility emits pollutants, and regulators have set a standard: the average pollutant concentration at a specified monitoring station must not exceed a given threshold. A transport model predicts whether any candidate site meets this standard.

The planner's question is different in character from the previous one. Rather than asking "if this parameter changes, what happens to the output?", they want to know: *for a given allowed change in the output (the pollutant concentration), how much variation in the inputs (wind speed, emission rate, atmospheric stability) is permissible?*

This is an adjoint question. It runs the sensitivity analysis in reverse: from a specified variation in the QOI back to the POIs that could have caused it. For models with many POIs and only a few QOIs, adjoint sensitivity analysis is dramatically more efficient than computing forward sensitivities for every parameter individually. Chapters 5 and 6 explain why this is so and how to exploit it.

*Self-check.* An environmental regulator measures pollutant concentration at a monitoring station and finds it exceeds the legal threshold. She wants to determine which weather assumptions in the transport model are most likely responsible for the exceedance. Is this a forward or adjoint question? Explain in one sentence.

*Answer:* Adjoint. The regulator starts with a specified change in the QOI (exceedance of the threshold) and traces backward to identify which input POIs (weather assumptions) most plausibly caused it. A forward analysis would instead ask how changing a specific weather assumption would affect the predicted concentration, which runs in the opposite direction.

## 1.3 ▪ Global Sensitivity: When Local Is Not Enough

Both forward and adjoint sensitivity analysis are *local*: they measure the effect of small perturbations to the POIs near a fixed nominal value. This is powerful, but it assumes that the model behaves approximately linearly in the vicinity of that nominal value.

Many real models are not linear, and many POIs carry substantial uncertainty that cannot be called "small." In these cases, local sensitivity indices

can be misleading. If the model is nonlinear in a POI, a linear prediction built from its local index can badly misjudge the effect of varying that POI across its full range of plausible values.

*Example* 1.1 (A linear prediction over a large range). Consider $q(p) = p^2$ and suppose the nominal value is $p^* = 0.1$. The local sensitivity index is $S_p^q = (p/q)\,(dq/dp) = (p/p^2)(2p) = 2$. But if $p$ is uncertain and could range from 0.1 to 1.0, the QOI ranges from 0.01 to 1.0, a hundredfold change. The linear prediction from $p^* = 0.1$ is $q(1) \approx 0.01 + 0.2 \times 0.9 = 0.19$, short of the true value by a factor of more than five. The index itself is not at fault. It equals 2 at every $p$, and read as the slope of the log-log plot of $q$ against $p$ (2.6), it gives the hundredfold change exactly: $(1.0/0.1)^2 = 100$. What fails is the straight-line extrapolation from one point. When the index also changes across the range, no single local value describes the whole range. Global SA, introduced in Chapter 9, is designed to capture exactly this kind of large-range nonlinear behavior.

Chapter 9 introduces *global sensitivity analysis*, which asks how the output varies when the POIs are varied across their entire plausible ranges simultaneously. The primary tool there, the Sobol index, measures how much of the total output variance is attributable to each POI. Global SA does not replace local SA; it complements it, answering questions that local SA cannot.

Readers from statistics and machine learning will recognize this question: it is the same one answered by $\eta^2$ in a classical ANOVA (proportion of variance explained by each factor), by permutation feature importance (how much does output variance increase when one feature is shuffled?), and by Shapley-value attribution methods such as SHAP. Chapter 9 makes these connections precise and shows exactly when each method is appropriate.

## 1.4 ▪ Sensitivity Analysis and Uncertainty Quantification

Sensitivity analysis and uncertainty quantification are related but distinct enterprises.

Sensitivity analysis, as developed in this book, begins as a *deterministic* enterprise: we assign nominal values to the POIs and compute how the QOI changes when those values are perturbed. The POIs are treated as numbers with specific values, not as random variables. The sensitivity index is a derivative: a precise mathematical object that exists when the model is differentiable. This local deterministic form is the focus of Chapters 2–8. Chapter 9 extends sensitivity analysis to the global setting, where the POIs are treated as random variables drawn from specified distributions, and the goal shifts from linearized local derivatives to variance-based attributions

of output uncertainty.

Uncertainty quantification treats the POIs as random variables with associated probability distributions. The goal is to determine the distribution of the QOI given the distributions of the POIs. This is a statistical enterprise, and its primary tools (Monte Carlo methods, Bayesian inference, polynomial chaos) are developed in depth in Smith's *Uncertainty Quantification: Theory, Implementation, and Applications* [59], which complements the present book. Smith's Chapters 8 and 9 develop local and global sensitivity analysis at the graduate level, extending the material in this book with functional-analytic and measure-theoretic foundations. Smith's Chapter 7 treats parameter identifiability and influence using the sensitivity Jacobian introduced computationally in our Chapter 3 and developed analytically in Chapter 4, connecting it to the surrogate-model literature. The SA foundations developed here follow Arriola and Hyman [2, 3]. For the broader development of SA as a formal discipline, see Saltelli et al. [51], Razavi et al. [46], and the annotated historical timeline in Tarantola et al. [62].

The connection between SA and uncertainty quantification is real and important. Global SA, covered in Chapter 9, bridges the gap: Sobol indices are defined in terms of variances, which requires probabilistic inputs, but they can be estimated by methods that build directly on the ideas in this book.

## 1.5 ▪ Four Ideas That Run Through the Book

Every chapter of this book revisits one or more of four fundamental ideas. We name them here so that students can recognize them when they reappear, each time in a new setting.

*Unifying Idea 1: The sensitivity index is a normalized derivative.*

At its core, SA answers a single question: if this input changes by a small percentage, by what percentage does the output change? That ratio is the sensitivity index. It is not a mysterious new object: it is the derivative, normalized so that its magnitude is independent of units and scales.

This idea appears first in Chapter 2, where we define the sensitivity index formally. It reappears in Chapter 3 as a finite-difference approximation and in Chapters 4–6 as an analytic derivative. In Chapter 9 it appears inside the variance-based formulas, whenever the response is close to linear over the input range. Every local sensitivity index in this book is an instance of this one idea. Chapter 9 also uses global measures that are built from other quantities and contain no local index. Smith [59] develops normalized derivative-based global sensitivity indices in §9.1 and §9.3, connecting the local framework of this chapter to the global variance- and derivative-based

methods covered in our Chapter 9.

**Unifying Idea 2: The adjoint is the transpose in whatever space you are working in.**

The adjoint sensitivity method, which makes it possible to compute the sensitivities of one QOI with respect to hundreds of POIs at the cost of solving just two problems instead of hundreds, rests on a single mathematical operation: transposition. For a linear system $A\vec{u} = \vec{b}$, the adjoint involves the transposed matrix $A^T$. For a computation graph (the structure underlying every computer program), the adjoint is the chain rule applied in reverse order, which is the backpropagation operation familiar from machine learning. For an ordinary differential equation, the adjoint is a second ODE running backward in time.

These three forms look different on the surface. They are the same idea.

Chapter 5 introduces the adjoint through the matrix transpose, which is the simplest and most familiar form. Chapter 6 extends it to differential equations. At each transition, the book explicitly identifies the connection back to this unifying idea.

**Unifying Idea 3: Local SA is a linearization.**

Every sensitivity index in Chapters 2–8 is based on a derivative, and a derivative is a linear approximation to a generally nonlinear function. This approximation is valid when the perturbation is small and the model is smooth. It fails near bifurcation points, where the model changes qualitative behavior (Section 8.4.1 sorts the ways it fails there). It fails when the POI uncertainty is large. It fails when the model is strongly nonlinear in the POI of interest.

Chapter 8 (*Caveat Emptor*) documents the conditions under which local SA breaks down. Chapter 9 provides the global alternative. Both chapters are applications of this one principle: local SA is a linearization, and linearizations have limits.

**Unifying Idea 4: A sensitivity index describes a model, not the world.**

Every index in this book is computed from a model's equations. It reports how that model responds, and it is silent on whether the model's simplifications are faithful to the system the model stands for. Holding those two claims apart is the most easily overlooked discipline in applied SA, and the most consequential when a result informs a decision.

Box put the underlying point in a sentence: "Essentially, all models are wrong, but some are useful" [8]. The useful ones are wrong in ways that do not matter for the question being asked. Sensitivity analysis is how you find out which of a model's wrongnesses matter, which parts of the simplification

are load-bearing. Picasso said something close about his own trade in 1923: "Art is a lie that makes us realize truth, at least the truth that is given us to understand."[1] A model is a deliberate lie about a system, told so that something true about the system becomes visible. The indices in this book measure the lie.

Two failure modes follow, and they point in opposite directions. A POI can dominate a model's output and matter little in the field, if the model's structure hands it more leverage than the system does. And a factor that is absent from the model returns a sensitivity of exactly zero, which is not evidence that the factor is unimportant, only that it was not modeled. A closed epidemic model with no births or deaths returns exactly zero sensitivity to the mean residence time in the population, because that parameter never appears on the right-hand side. Nothing whatever has been established about demography. The demographic model used from Chapter 4 onward, and in the Chapter 6 laboratory, does carry that parameter, and the contrast is worth holding onto. A mean residence time of $10^4$ days set against a 90-day outbreak means almost nobody enters or leaves the population while the epidemic runs, and the computed sensitivity is $-7.9 \times 10^{-4}$, three orders of magnitude below the others. Small for a reason and zero by construction look alike in a table, and only one of them says anything about demography.

This distinction is hardest to hold when the result was produced by someone, or something, other than you. Whoever computed the index did not make the modeling decisions, and so cannot tell you which of the model's simplifications are load-bearing for your question. What arrives is a number and a ranking, with the assumptions that generated them left behind. A borrowed result is therefore a claim about a model whose construction you may not have seen, and reading it as a claim about the world compounds the two gaps rather than one. Section 9.10 returns to this once the global methods are in place, since that is where the temptation is strongest.

Saltelli, Tarantola and Campolongo argued on these grounds that sensitivity analysis belongs inside the modeling cycle rather than appended to it [54]. The reminders placed through the chapters that follow are there to keep the distinction visible at the points where it is easiest to lose.

## 1.6 ▪ A Note on Parameterization

One practical decision that profoundly affects the usefulness of SA results deserves mention before Chapter 2.

[1] "Picasso Speaks," interview with Marius de Zayas, *The Arts*, New York, May 1923, p. 315.

The sensitivity index depends on how a POI is defined. In a compartmental epidemic model, the recovery process can be parameterized either by the *recovery rate* $\gamma_R$ (units: per day) or by the *mean infectious period* $\tau_R = 1/\gamma_R$ (units: days). These two parameterizations are mathematically equivalent, each determining the other. But their sensitivity indices are not equal: they have the same magnitude and opposite signs.

Which one should you use? The answer is: whichever one you can discuss in ordinary language without confusion. People who work with epidemic models (clinicians, public health officials, and policy makers) think naturally in terms of days. A clinician understands immediately what it means to say "the mean infectious period is 7 days." They do not think in recovery rates. Reporting $\mathcal{S}_{\gamma_R}^{\mathcal{R}_0} = -1.0$ requires the reader to invert $\gamma_R$ mentally before the result makes intuitive sense. Reporting $\mathcal{S}_{\tau_R}^{\mathcal{R}_0} = +1.0$ is immediately clear: a longer infectious period increases transmission.

Throughout this book, we choose POIs that can be understood in everyday terms: durations rather than rates, probabilities rather than their complements, counts rather than densities when the distinction matters. This is not just a matter of aesthetics. Sensitivity indices that require mental inversion to interpret are sensitivity indices that will be misread.

Chapter 2 formalizes this principle and shows how to convert between parameterizations. Readers who find that their model's natural parameters are rates will learn how to restate their results in terms of the corresponding durations before reporting them.

## 1.7 ▪ Structure of the Book

The book is organized in five parts. Each part builds on the previous, and the SIR epidemic model [26, 64] serves as the recurring example throughout, so that students can track a single application as the mathematical tools become progressively more powerful. The SIR model divides a population into three compartments: Susceptible ($S$), Infectious ($I$), and Recovered ($R$); it is derived from first principles in Section 2.5.

**Part I: The Sensitivity Question** (Chapters 1–2) establishes the central problem and introduces the sensitivity index as the primary tool. Chapter 2 develops the formal definition, the sign interpretation, the tornado plot as a display format, and the principle of interpretable parameterization. By the end of Part I, students can compute and interpret sensitivity indices for any differentiable function of several variables.

**Part II: Computing Sensitivity in Practice** (Chapter 3) gives students an immediately usable computational method for any model, including models that cannot be differentiated analytically. Finite-difference ap-

proximations, step-size selection, the sensitivity Jacobian, and the treatment of correlated POIs are developed here. Students who work through Chapter 3 can perform SA on any model they can run, including inherited code and black-box simulation software.

**Part III: Analytic Forward Sensitivity Analysis** (Chapter 4) shows how to differentiate model equations directly to obtain exact sensitivity information at lower computational cost. The forward sensitivity equation is derived for algebraic systems, difference equations, and ordinary differential equations. The chapter ends by quantifying the computational bottleneck that motivates the adjoint approach.

**Part IV: The Adjoint Approach** (Chapters 5–7) introduces adjoint sensitivity analysis through three progressively more abstract settings: linear systems in Chapter 5, dynamical systems in Chapter 6, and computational tools including algorithmic differentiation in Chapter 7. The central theme is Unifying Idea 2: the adjoint is the matrix transpose in whatever mathematical space the problem inhabits.

**Part V: Limitations and Extensions** (Chapters 8–9) addresses the boundaries of local SA. Chapter 8 documents the conditions under which sensitivity indices give misleading results: bifurcation points, ill-conditioned systems, and method-question mismatch. Chapter 9 introduces global SA through variance decomposition, Sobol indices, and partial rank correlation coefficients. It closes with connections to distribution-based uncertainty quantification, developed at length in Smith's *Uncertainty Quantification* [59].

**Computing environment.** Every chapter after this one includes a structured computational laboratory that implements the chapter's core idea on the SIR epidemic model. Students who complete these laboratories will leave the course with a portable toolkit for performing SA on any model they encounter in research or practice. Companion Colab notebooks recompute examples from each chapter in Python and assert results against fresh computation, and a ledger accounts for every decimal value and computed count that the chapters and appendices quote. The MATLAB reference library (eighteen functions across four modules, with an automated test suite) is available alongside them. All companion materials are hosted at:

`https://github.com/machyman/arriola2026sa`

Each function in that library will match a concept or algorithm in the text. `sir_nominal.m` will define the SIR parameters used in every chapter; `sensitivity_jacobian.m` the centered-difference Jacobian from Chapter 3; `sobol_jansen.m` the Jansen estimators from Chapter 9. The repository is

organized to mirror the book's structure, so the code for any chapter is found without searching.

**Notation.** Throughout the book, the sensitivity index of a QOI $q$ with respect to a POI $p$ is written $\mathcal{S}_p^q$. POIs are always chosen to be interpretable in everyday terms: durations, probabilities, counts. The relationship to the mathematical parameters in the model equations is made explicit wherever the two differ.

***Scope of This Book: Scalar QOIs***

Every sensitivity index in this book is computed for a *scalar* QOI, a single number such as the peak infection count, the total disease burden, or the value of $\mathcal{R}_0$. Real applications sometimes require sensitivity of an entire output *trajectory*: how does the full time series $I(t)$ change when $\beta$ changes? Extending SA to functional (trajectory-valued) QOIs requires either reducing the trajectory to a scalar (which loses information) or computing a time-dependent SI at each time point (as in Chapter 4's Figure 4.1). A rigorous treatment using the adjoint sensitivity analysis procedure (ASAP) for both algebraic and differential systems is developed in Smith [59] Ch. 8 §§8.1–8.6, and is a natural next step after mastering the scalar case treated here.

## 1.8 ▪ Exercises

Each exercise carries a Bloom's Taxonomy tag indicating cognitive level: *L2* (understand), *L3* (apply), *L4* (analyze), *L5* (evaluate), *L6* (create and synthesize). Higher-level exercises require constructing an argument or designing a procedure, not just applying a formula.

**Exercise 1.1** *(*[Bloom L2]*).* Explain in your own words the difference between a POI and a QOI. For a model describing the spread of antibiotic-resistant bacteria in a hospital, give two examples of each. Then explain, in one sentence, what sensitivity analysis would tell you about this model that simply running the model would not.

**Exercise 1.2** *(*[Bloom L2]*).* What is the difference between forward and adjoint sensitivity analysis? Describe each in terms of the "What if?" and "How come?" framing. For each of the following questions, classify it as a *forward question* or an *adjoint question* based on the type of question being asked (not on which algorithm would be most efficient to use computationally; efficiency is the subject of Chapters 5 and 6).

(a) If the mean latent period increases by 10%, how does the peak infection count change?

(b) The peak infection count is higher than expected. Which parameter uncertainties are most likely to have caused this?

(c) Among all parameters in the model, which one has the largest effect on the basic reproduction number?

**Exercise 1.3** *([Bloom L4]).* For each of the following modeling scenarios, state whether forward SA, adjoint SA, or global SA is most appropriate. Justify each choice in one or two sentences.

(a) A climate model has 500 tunable parameters. The investigator wants to know which 10 parameters most affect the mean surface temperature over a 100-year run. Each model run takes 8 hours.

(b) A pollution transport model has 6 input parameters. The regulator wants to know: if the pollution at a specific monitoring station exceeds the legal threshold, which input assumptions most likely caused this?

(c) A pharmacokinetic model has 4 parameters, each of which is known only to within a factor of 2. The investigator wants to know the range of predicted drug concentrations consistent with this uncertainty.

**Exercise 1.4** *([Bloom L5]).* You are advising a team modeling the transmission of dengue fever. The model has 9 parameters. The team has funding to measure exactly 3 parameters precisely in the field and to implement exactly 2 vector-control interventions. Write a one-page plan describing how you would use SA to guide these decisions. Your plan should specify: what QOI you would focus on and why; how you would determine which 3 parameters to measure; what additional information (if any) you would need beyond the SA results to recommend interventions; and one important limitation of your approach that the team should be aware of.

**Exercise 1.5** *([Bloom L6]).* A published report on a disease model states: "Sensitivity analysis was performed on all 22 model parameters simultaneously, demonstrating that the model is robust to parameter uncertainty." The analysis was conducted by computing local sensitivity indices at the best-fit parameter values.

Identify at least two substantive problems with this description or its conclusions. For each problem you identify, explain specifically what the authors should have done differently.

Draft

## Notes

The sensitivity index as a normalized derivative appears throughout applied mathematics under different names. Relative sensitivity, elasticity (in economics), and logarithmic derivative all refer to the same quantity $p\,\partial q/\partial p \cdot q^{-1}$. The term *sensitivity index* as used here follows Arriola and Hyman [2, 3], who established the notation and sign conventions adopted in this book.

The SIR compartmental model originates with Kermack and McKendrick [26], whose 1927 paper remains a landmark of mathematical epidemiology. Anderson and May [1] developed the framework extensively, including the derivation of $\mathcal{R}_0 = c\beta/\gamma_R$ and its role as the threshold parameter for epidemic invasion. The basic reproduction number $\mathcal{R}_0$ and its precise definition for structured population models is treated rigorously by Diekmann, Heesterbeek, and Metz [15] and by van den Driessche and Watmough [63].

For a comprehensive survey of sensitivity analysis methodology, the books by Saltelli et al. [52, 51] are authoritative. The recent review by Razavi et al. [46] assesses the state of the field and identifies open research problems. Tarantola et al. [62] provide an annotated timeline of the major methodological developments. Smith [59] develops uncertainty quantification as a complementary topic, with particular strength on Bayesian and surrogate-model approaches.

Draft

Draft

# Chapter 2
# Measuring Local Sensitivity

*A sensitivity index is a normalized derivative with a practical job.*

*Calibration tells you which parameter values fit the data.*
*Sensitivity analysis tells you which parameters matter to the answer.*

Two studies of the same epidemic model arrive at the following conclusions.

Study A reports: "Increasing the recovery rate by 1% decreases $\mathcal{R}_0$ by 1%."

Study B reports: "Increasing the mean infectious period by 1% increases $\mathcal{R}_0$ by 1%."

Both statements are mathematically correct. The recovery rate and the mean infectious period are reciprocals of each other, so the two studies analyzed the same model, used the same mathematics, and obtained the same numerical result. Yet they convey opposite signs and, more importantly, very different intuitions. Study A requires the reader to mentally invert a parameter before the public health implication becomes clear. Study B is immediately interpretable: patients who are infectious longer spread the disease more.

This chapter introduces the sensitivity index, the central quantitative tool of this book, and develops the conventions that make sensitivity analysis results readable rather than merely correct.

## 2.1 ▪ Ratios as Precursors to Derivatives

Comparison by ratio is one of the most fundamental quantitative habits in applied science. In finance, the price-to-earnings ratio tells an investor how many dollars they pay for one dollar of profit. In aeronautical engineering, the lift-to-drag ratio determines whether a wing design is viable. In

Draft

experimental physics, the signal-to-noise ratio separates measurement from artifact. None of these ratios requires calculus. Each one gives a dimensionless number that can be compared across different systems and used to make decisions.

The sensitivity index is a ratio of this same type: it measures how much the output changes, per unit change in the input, normalized so that the result is dimensionless and comparable across parameters with different units and scales.

To see why normalization matters, consider the price-to-earnings ratio $P/E$. Suppose a stock trades at \$50 per share and has earned \$2.50 over the past year. The $P/E$ ratio is $50/2.50 = 20$: you pay \$20 for each dollar of annual earnings. Now suppose earnings increase by \$0.50 to \$3.00. A raw comparison of the two earnings figures gives $\$3.00 - \$2.50 = \$0.50$, which tells you nothing useful without knowing the original value. The normalized comparison is $(3.00 - 2.50)/2.50 = 20\%$, a 20% increase in earnings, regardless of whether the numbers are in dollars, euros, or any other currency.

A derivative, in the same spirit, measures the rate of change of the output per unit change in the input. The sensitivity index normalizes this derivative so that it answers a specifically useful question: if the input changes by a small percentage, by what percentage does the output change?

*Self-check.* A second stock trades at \$45 per share and has earned \$3.00 per year. Calculate the $P/E$ ratio for each stock. Which stock is a better value by this measure, and why?

> *Answer:* Stock 1: $P/E = 50/2.50 = 20$. Stock 2: $P/E = 45/3.00 = 15$. Stock 2 has a lower $P/E$ ratio, meaning you pay less per dollar of earnings, a better value by this measure. The ratio provides a dimensionless comparison that is fair across the two stocks despite their different prices and earnings.

## 2.2 ▪ Absolute, Relative, and Regularized Change

Before defining the sensitivity index, we need precise language for describing how quantities change. Three measures of change appear throughout the book, each appropriate in different circumstances.

### 2.2.1 ▪ Absolute Change

Let $u$ denote the solution to a model, and suppose a perturbation $\delta u$ is applied. The *absolute change* is simply the difference between the perturbed and original solutions:

$$\Delta u_{\text{Abs}} := \delta u = \hat{u} - u, \tag{2.1}$$

where $\hat{u} = u + \delta u$ is the perturbed solution. A 1-inch change in the length of a string has absolute change 1 inch, regardless of the original length.

### 2.2.2 ▪ Relative Change

The 1-inch change means something very different if the string is 6 inches long versus 1 mile long. The *relative change* accounts for this by normalizing against the original value:

$$\Delta u_{\text{Rel}} := \frac{\delta u}{u} = \frac{\hat{u} - u}{u}. \tag{2.2}$$

For the 6-inch string, the relative change is $1/6 \approx 16.7\%$. For the 1-mile string (63,360 inches), it is $1/63{,}360 \approx 0.0016\%$. The same absolute change has a very different significance depending on context, and relative change captures this.

### 2.2.3 ▪ Regularized Change

Relative change has one practical problem: if $u \approx 0$, the denominator in (2.2) is near zero and the ratio becomes unreliable. In dynamical models, the solution can pass through or near zero during its evolution: the infectious compartment $I(t)$ in an epidemic approaches zero as the outbreak resolves, for instance. The *regularized change* handles this case by adding a small correction to the denominator:

$$\Delta u_{\text{Reg}} := \frac{\delta u}{u_{\text{ref}} + u}. \tag{2.3}$$

Here $u_{\text{ref}} > 0$ is a reference scale carrying the same units as $u$, chosen from the problem: a typical or nominal magnitude of the state, or the smallest value we care to resolve. The scale cannot be dropped. Writing $1 + u$ in its place adds a dimensionless 1 to a dimensional quantity, and the resulting index would change whenever the units change. When $u \ll u_{\text{ref}}$, the regularized change approximates absolute change measured in units of $u_{\text{ref}}$; when $u \gg u_{\text{ref}}$, it approximates relative change $\delta u/u$. It interpolates smoothly between the two regimes, and the crossover sits where the choice of $u_{\text{ref}}$ puts it, which is why that choice should be stated whenever the index is reported. (For models where $u$ can be negative, use $u_{\text{ref}} + |u|$; all SIR state variables satisfy $u \geq 0$.)

Looking more carefully at regularized change reveals a pattern that recurs throughout applied mathematics: when a formula fails at a special value, adding a small stabilizing term to the denominator restores well-definedness without significantly changing the result away from the problematic point. The same idea appears in numerical linear algebra (Tikhonov

regularization of ill-conditioned systems) and in Bayesian statistics (pseudo-counts in categorical models). The regularized sensitivity index in Chapter 3 is one more instance.

*Self-check.* The infectious population in a disease model drops from $I = 0.04$ to $\hat{I} = 0.03$. Calculate the absolute, relative, and regularized change. Which measure gives the most informative description of this transition, and why?

*Answer:* Absolute: $\delta I = 0.03 - 0.04 = -0.01$. Relative: $-0.01/0.04 = -0.25$, a 25% decrease. Regularized: taking $u_{\text{ref}} = 1$ as the reference scale, since the compartments here are fractions of the whole population and 1 is that population, $-0.01/(1 + 0.04) \approx -0.0096$, close to absolute. The reference scale is what makes the denominator dimensionally sound; it is not a bare 1 added to a fraction.

Since $I$ is small relative to 1, relative change gives the most interpretable result: a 25% decrease in the infectious population is meaningful regardless of scale. For $I \ll 1$, absolute and regularized change are both small in magnitude and harder to interpret without the normalization that relative change provides.

## 2.3 ▪ The Sensitivity Index

We now define the central tool of local sensitivity analysis.

### 2.3.1 ▪ The Sign Rule

Before any formal definition, the most important thing to know about the sensitivity index is what its sign means.

**The sign rule:** A positive sensitivity index means that the QOI and POI move in the same direction: increase the POI and the QOI increases. A negative sensitivity index means they move in opposite directions: increase the POI and the QOI decreases. The magnitude tells you by how much: a sensitivity index of 2 means that a 1% change in the POI produces approximately a 2% change in the QOI.

**Important:** the sign belongs to the *parameterization*, not to the underlying biology. The recovery rate $\gamma_R$ and the mean infectious period $\tau_R = 1/\gamma_R$ describe the same biological process, yet $\mathcal{S}^{\mathcal{R}_0}_{\gamma_R} < 0$ while $\mathcal{S}^{\mathcal{R}_0}_{\tau_R} > 0$. When two papers report opposite signs for what appears to be the same parameter, check whether they are using rate versus duration parameterizations before concluding there is a contradiction. Section 2.5 illustrates this with the SIR model.

Students sometimes interpret a negative sensitivity index as indicating an error in the calculation. It is not. A negative index is often the best possible news: if the sensitivity of $\mathcal{R}_0$ with respect to the mean infectious

period is positive, that means longer illness leads to more transmission, and the corresponding statement with the recovery rate would be negative: faster recovery reduces transmission. Both facts are correct; the sign simply tells you which direction the influence runs. With this intuition established, the formal definition follows naturally.

### 2.3.2 ▪ Formal Definition

**Definition 2.1** (Sensitivity Index)**.** *Let $q$ denote a* QOI *that depends differentiably on a* POI *$p$. The* normalized sensitivity index *of $q$ with respect to $p$ is*

$$\mathcal{S}_p^q := \lim_{\delta p \to 0} \frac{\left(\frac{\delta q}{q}\right)}{\left(\frac{\delta p}{p}\right)} = \frac{p}{q}\frac{\partial q}{\partial p}, \qquad q, p \neq 0. \tag{2.4}$$

*This quantity is called an* elasticity *in economics and econometrics, where it measures the proportional responsiveness of one variable to another. The sensitivity index is its counterpart in mathematical modeling.*

*Remark* 2.2 (Near-zero QOI). When $q = 0$ at the nominal parameter values, the normalized SI is undefined (division by zero). In this case, use the absolute sensitivity $\partial q/\partial p$ directly, or the regularized form $S^q_{p,\mathrm{reg}} = \frac{p}{q_{\mathrm{ref}}+|q|}\frac{\partial q}{\partial p}$, where $q_{\mathrm{ref}} > 0$ carries the units of $q$ and is chosen from the problem as in (2.3). Throughout this book we assume $q \neq 0$ and $p \neq 0$ unless stated otherwise; this holds for all QOIs and POIs in the SIR examples.

*Practical switching rule.* In dynamical models, a QOI may approach zero even when it is nonzero at the nominal point, for example, $I(t)$ becomes small near $t = T$ in epidemic models with a finite observation window. Use the normalized SI when $|q^*| \geq 0.01 \times \max_t |q(t)|$; switch to the absolute sensitivity $\partial q/\partial p$ when $|q^*|$ is smaller than this threshold. If the timing of a near-zero crossing itself matters (for example, the epidemic end date), redefine the QOI to be the crossing time rather than the compartment value.

Three equivalent forms of this definition are in common use. All three give the same number; which one is most convenient depends on the problem at hand.

*Ratio form* (Definition 2.1): the ratio of relative changes in the QOI and the POI. This form is closest to the P/E ratio analogy from Section 2.1.

*Derivative form*: multiply the partial derivative by the ratio of nominal values:

$$\mathcal{S}_p^q = \frac{p}{q}\frac{\partial q}{\partial p}. \tag{2.5}$$

This is the form used most often in computation. The factor $p/q$ does the normalization work: it converts $\partial q/\partial p$, which has units of $[Q]/[P]$ and depends on the choice of units, into a dimensionless percent-for-percent ratio. Concretely, if $\mathcal{S}_p^q = 2$, then a 1% increase in $p$ produces approximately a 2% increase in $q$, regardless of whether $p$ is measured in days, kilograms, or dollars.

*Logarithmic form*: for positive POIs and QOIs,

$$\mathcal{S}_p^q = \frac{\partial \ln q}{\partial \ln p}. \tag{2.6}$$

This form reveals that the sensitivity index is the slope of the log-log plot of $q$ versus $p$, a fact that becomes useful when comparing SI values across very different scales.

The three forms are equivalent whenever $q$ and $p$ are positive and $q$ is differentiable with respect to $p$.

*Remark* 2.3. When $p$ or $q$ may be negative, but neither is zero, use the derivative form (2.5) directly; the logarithmic form (2.6) requires $p > 0$ and $q > 0$. Remark 2.2 treats a zero $q$. In compartmental epidemic models this holds wherever the compartments and rates are strictly positive, as at a standard endemic equilibrium; it can fail where a compartment or rate vanishes, such as a disease-free equilibrium or a switched-off intervention, and there the logarithmic form does not apply.

*Unifying Idea 1.* The sensitivity index is the central object of this book. It appears here as a ratio of relative changes. In Chapters 3 and 4 it reappears as a finite-difference approximation and as the solution of a sensitivity equation. In Chapters 5 and 6, it appears as the output of the adjoint calculation. Every one of these is the same normalized derivative, asked in the same way: *if the input changes by x%, by what percentage does the output change?* In Chapter 9, it reappears inside the variance-based formulas whenever the response is close to linear over the input range.

**Why normalization is not optional.** Without the $p/q$ factor, comparing raw derivatives across parameters is not meaningful. In the SIR model, $\partial \mathcal{R}_0/\partial c$ has units of days (since $c$ has units day$^{-1}$), while $\partial \mathcal{R}_0/\partial \beta$ has different units entirely. Asking which raw derivative is larger is like asking whether 1 pound is larger than 2 inches: the numbers may have any magnitude, but the comparison is incommensurable because the units differ. The normalized SI eliminates this problem. Both $\mathcal{S}_c^{\mathcal{R}_0}$ and $\mathcal{S}_\beta^{\mathcal{R}_0}$ equal $+1$; dimensionless, unit-free, so ranking them is legitimate and the tornado plot is an honest comparison. This is why the tornado plot sorts by $|\mathcal{S}_p^q|$ rather than by $|\partial q/\partial p|$: only the normalized index gives a ranking that is independent of how parameters happen to be measured.

*Self-check.* Use each of the three forms to compute the sensitivity index of the area $A = \pi r^2$ with respect to the radius $r$. Verify that all three give the same answer.

*Answer: Ratio form:* $\delta A/A = 2\pi r\,\delta r/(\pi r^2) = 2\delta r/r$, so $\mathcal{S}_r^A = (\delta A/A)/(\delta r/r) = 2$.
*Derivative form:* $\partial A/\partial r = 2\pi r$, so $\mathcal{S}_r^A = (r/\pi r^2)(2\pi r) = 2$.
*Logarithmic form:* $\ln A = \ln \pi + 2\ln r$, so $\partial \ln A/\partial \ln r = 2$.
All three give $\mathcal{S}_r^A = 2$: a 1% increase in radius produces a 2% increase in area. The circle's area is twice as sensitive to radius as to a parameter that appears linearly.

### 2.3.3 ▪ Interpreting the Magnitude

The sensitivity index answers a specific quantitative question: if the POI changes by $x\%$, the QOI changes by approximately $x \cdot \mathcal{S}_p^q\%$. For small $x$, this estimate is accurate to first order. For larger $x$, second-order corrections become relevant, and Chapter 3 discusses how to assess whether those corrections matter.

As a practical interpretive guide:

- $|\mathcal{S}_p^q| > 1$: the QOI is more sensitive than the POI (the change in QOI exceeds the change in POI, percentage-for-percentage).
- $|\mathcal{S}_p^q| = 1$: the QOI changes in exact proportion to the POI.
- $|\mathcal{S}_p^q| < 1$: the QOI is less sensitive than the POI (the change in QOI is smaller than the change in POI).
- $|\mathcal{S}_p^q| \approx 0$: the QOI is essentially insensitive to this POI near the nominal value.

## 2.4 ▪ Worked Examples

The following examples build in complexity, beginning with purely geometric functions and ending with the epidemic model that serves as the book's primary running example.

### 2.4.1 ▪ Volume of a Cylinder

The volume of a cylinder with radius $r$ and height $h$ is $V = \pi r^2 h$. We want to know which dimension (radius or height) has the greater effect on the volume.

Using the derivative form (2.5):

$$\mathcal{S}_r^V = \frac{r}{V}\frac{\partial V}{\partial r} = \frac{r}{\pi r^2 h} \cdot 2\pi r h = 2. \tag{2.7}$$

$$\mathcal{S}_h^V = \frac{h}{V}\frac{\partial V}{\partial h} = \frac{h}{\pi r^2 h} \cdot \pi r^2 = 1. \tag{2.8}$$

The volume is twice as sensitive to radius as to height. A 1% increase in radius produces a 2% increase in volume; the same percentage increase in height produces only a 1% increase. This follows from the fact that $r$ appears squared in the volume formula while $h$ appears linearly.

For a designer building a cylindrical tank with a fixed material budget, this result has a direct implication: small manufacturing errors in the radius matter twice as much as the same-sized errors in the height.

*Self-check.* For a cone with volume $V = \frac{1}{3}\pi r^2 h$, compute $\mathcal{S}_r^V$ and $\mathcal{S}_h^V$. Compare with the cylinder results. Explain why the answer is the same despite the different formula.

*Answer:* $\mathcal{S}_r^V = (r/V)(\partial V/\partial r) = (r/V)\cdot(2\pi rh/3) = 2$. $\mathcal{S}_h^V = 1$. Identical to the cylinder.
The reason: the sensitivity index depends on the *exponent* of each variable in the formula. Both $V_{\text{cone}} = \frac{1}{3}\pi r^2 h$ and $V_{\text{cylinder}} = \pi r^2 h$ have $r$ to the power 2 and $h$ to the power 1. The constant $\frac{1}{3}$ cancels in the logarithmic form $\partial \ln V/\partial \ln r = 2$. This is a general principle: for a monomial $q = c\,p_1^{a_1}p_2^{a_2}\cdots$, the sensitivity index with respect to $p_i$ is simply $a_i$.

### 2.4.2 ▪ Arc Length of a Line

Let $u(t) = mt + b$ be a line with slope $m$ and intercept $b$, defined on $[0, a]$. The arc length is

$$J(m, b) = a\sqrt{1 + m^2}.$$

Computing the sensitivity indices:

$$\mathcal{S}_m^J = \frac{m}{J}\frac{\partial J}{\partial m} = \frac{m}{a\sqrt{1+m^2}}\cdot\frac{am}{\sqrt{1+m^2}} = \frac{m^2}{1+m^2}. \tag{2.9}$$

$$\mathcal{S}_b^J = \frac{b}{J}\frac{\partial J}{\partial b} = 0, \tag{2.10}$$

since $J$ does not depend on $b$.

The arc length is completely insensitive to the intercept. Geometrically, this is clear: shifting the line vertically does not change its length. Analytically, $b$ does not appear in the formula for $J$, so its derivative is zero. A zero sensitivity index is not an error; it tells us that one POI has no local effect on the QOI, which is itself an informative finding.

*Self-check.* For the line $u(t) = 3t + 2$ on $[0, 4]$, compute $\mathcal{S}_m^J$ numerically. Verify: if $m$ increases from 3 to 3.03 (a 1% increase), does $J$ change by approximately $\mathcal{S}_m^J$%?

*Answer:* $\mathcal{S}_m^J = m^2/(1+m^2) = 9/(1+9) = 0.9$.
At $m = 3$: $J = 4\sqrt{10} \approx 12.649$. At $m = 3.03$: $J = 4\sqrt{1 + 3.03^2} = 4\sqrt{10.1809} \approx 12.763$. Change: $(12.763 - 12.649)/12.649 \approx 0.90\% \approx 0.9 \times 1\%$. ✓

### 2.4.3 ▪ Area Under a Line

For the same line $u(t) = mt + b$ on $[0, a]$, the area is

$$J(m, b) = \int_0^a (mt + b)\, dt = \tfrac{1}{2}ma^2 + ba.$$

Both parameters now affect the area:

$$\mathcal{S}_m^J = \frac{m}{\frac{1}{2}ma^2 + ba} \cdot \tfrac{1}{2}a^2 = \frac{am}{am + 2b}, \tag{2.11}$$

$$\mathcal{S}_b^J = \frac{b}{\frac{1}{2}ma^2 + ba} \cdot a = \frac{2b}{am + 2b}. \tag{2.12}$$

Note that $\mathcal{S}_m^J + \mathcal{S}_b^J = 1$, so the two indices share a unit total for this particular linear functional. This is not a general property: sensitivity indices do not sum to any fixed value for arbitrary models or QOIs. Which parameter matters more depends on the values: $\mathcal{S}_m^J > \mathcal{S}_b^J$ when $am > 2b$, and the inequality reverses when $am < 2b$.

This example illustrates a general phenomenon: when two POIs both influence the same QOI, their relative importance depends on the nominal parameter values, not just on the model structure. SA must always be reported at specific nominal values.

### 2.4.4 ▪ Root of a Nonlinear Equation

Consider the nonlinear equation

$$f(u; \alpha) := \sin(\alpha u) - u = 0.$$

For $\alpha = 2$, the equation has three roots; for $\alpha > 1$, there is at least one nonzero root. We want to know how the positive root $u^*(\alpha)$ changes with $\alpha$.

Differentiating $f(u; \alpha) = 0$ implicitly with respect to $\alpha$:

$$u\cos(\alpha u) + \alpha\cos(\alpha u)\,\frac{\partial u}{\partial \alpha} - \frac{\partial u}{\partial \alpha} = 0.$$

Solving for $\partial u/\partial \alpha$:

$$\frac{\partial u}{\partial \alpha} = \frac{u\cos(\alpha u)}{1 - \alpha\cos(\alpha u)}.$$

At $\alpha = 2$, the positive root is $u^* \approx 0.9477$, giving

$$\mathcal{S}_\alpha^{u^*} = \frac{\alpha}{u^*} \cdot \frac{u^*\cos(\alpha u^*)}{1 - \alpha\cos(\alpha u^*)} \approx -0.39.$$

A 1% increase in $\alpha$ decreases the positive root by approximately 0.39%. This result requires no numerical experiment once the derivative is known; the sensitivity index provides the estimate directly.

### 2.4.5 ▪ A Linear Prediction Model (Statistical Connection)

Sensitivity indices apply equally to statistical models. Consider a linear prediction model for hospital readmission risk:

$$\hat{P} = \beta_0 + \beta_1 x_1 + \beta_2 x_2, \tag{2.13}$$

where $x_1$ = patient age (years) and $x_2$ = comorbidity count are the POIs, $\hat{P}$ (predicted readmission probability) is the QOI, and $\beta_0 = 0.05$, $\beta_1 = 0.002$, $\beta_2 = 0.01$ are estimated regression coefficients. At nominal values $x_1^* = 65$ years and $x_2^* = 3$ comorbidities, the predicted probability is $\hat{P}^* = 0.05 + 0.002(65) + 0.01(3) = 0.21$.

Applying the sensitivity index formula:

$$\mathcal{S}_{x_1}^{\hat{P}} = \frac{x_1^*}{\hat{P}^*}\frac{\partial \hat{P}}{\partial x_1} = \frac{65}{0.21} \times 0.002 \approx 0.619, \tag{2.14}$$

$$\mathcal{S}_{x_2}^{\hat{P}} = \frac{x_2^*}{\hat{P}^*}\frac{\partial \hat{P}}{\partial x_2} = \frac{3}{0.21} \times 0.01 \approx 0.143. \tag{2.15}$$

Age is the dominant driver: a 1% increase in age (roughly 0.65 years) increases predicted readmission risk by 0.619%.

*Comparison with standardized regression coefficients.* Statistical software reports standardized coefficients $\tilde{\beta}_i = \beta_i \sigma_{x_i}/\sigma_{\hat{P}}$, which normalize by standard deviations rather than by nominal values. If $\sigma_{x_1} = 15$ years, $\sigma_{x_2} = 1.5$, and $\sigma_{\hat{P}} \approx 0.04$, then $\tilde{\beta}_1 = 0.002 \times 15/0.04 = 0.75$ and $\tilde{\beta}_2 = 0.01 \times 1.5/0.04 = 0.375$: the same ranking (age dominates) but different magnitudes. The sensitivity index asks *"what if this patient's age changed by 1%?"*; the standardized coefficient asks *"what if age changed by one standard deviation across the patient population?"* Both are dimensionless measures of importance; they answer different questions.

## 2.5 ▪ The Sensitivity Index for the SIR Model

### 2.5.1 ▪ Model Derivation from First Principles

The SIR epidemic model [26, 1] divides a closed population of $N$ individuals into three compartments: susceptible ($S$), infectious ($I$), and recovered ($R$), with $S(t) + I(t) + R(t) = N$ at all times. Figure 2.1 shows the flow diagram.

**The force of infection.** At any moment, each infectious individual makes $c$ contacts per day.[2] Every contact with a susceptible individual transmits

[2]This chapter treats the contact rate as a single parameter $c$ shared by all compartments, which makes $\mathcal{R}_0 = c\beta\tau_R$ a product of monomials with all sensitivity indices equal





**Figure 2.1.** *SIR compartment flow diagram. The force of infection $\lambda(t) = c\beta I(t)/N$ is the per-capita rate at which any one susceptible individual becomes infected. Recovery occurs at rate $1/\tau_R$ per infectious individual per day.*

the infection with probability $\beta$. Of the $c$ individuals contacted per day by one infectious person, a fraction $S/N$ are susceptible. The rate at which one infectious individual generates new infections is therefore $c\beta S/N$ per day, and with $I$ infectious individuals in the population the total rate of new infections is $c\beta SI/N$ per day. The *force of infection*

$$\lambda(t) = \frac{c\beta I(t)}{N} \tag{2.16}$$

is the per-capita infection rate at time $t$: the rate at which each susceptible individual acquires infection. Note that $\lambda(t)$ has units of day$^{-1}$: the product $\lambda(t) \cdot \Delta t$ is the probability of becoming infected during a short time interval $\Delta t$.

**The ODE system.** Translating the flow diagram into rates:

$$\frac{dS}{dt} = -\lambda(t)\, S \;=\; -\frac{c\beta SI}{N}, \tag{2.17}$$

$$\frac{dI}{dt} = \;\; \lambda(t)\, S - \frac{I}{\tau_R} \;=\; \frac{c\beta SI}{N} - \frac{I}{\tau_R}, \tag{2.18}$$

$$\frac{dR}{dt} = \frac{I}{\tau_R}. \tag{2.19}$$

Each term has a biological origin. $-\lambda S$ in equation (2.17): susceptibles are lost at rate $\lambda$ (the per-capita infection rate) times $S$ (the number at risk). $+\lambda S$ in equation (2.18): these same individuals immediately enter the infectious compartment. $-I/\tau_R$ in equation (2.18): infectious individuals

to +1. In heterogeneous-mixing models, such as the $SIR_I$ framework developed in Hyman, Qu, and Xue [23], the contact rate is differentiated by compartment: susceptibles, infectious, and recovered individuals make $c_S$, $c_I$, and $c_R$ contacts per day, respectively, with $\mathcal{R}_0 = c_I\beta/(\gamma_R+\nu_m)$. The equal-contact-rate case $c_S = c_I = c_R = c$, with no demography ($\nu_m = 0$), recovers the simplified model used here. That book also normalizes the state variables by the carrying capacity $N^c$, the population size at which births balance deaths in the absence of disease, and uses calligraphic $\mathcal{R}_0$ to distinguish the reproductive number from the recovered compartment $R$.

recover at rate $1/\tau_R$ per person per day, since the mean infectious period is $\tau_R$ days. $+I/\tau_R$ in equation (2.19): recovered individuals arrive at the same rate they depart from $I$.

**Why $\mathcal{R}_0 = c\beta\tau_R$.** In a fully susceptible population ($S \approx N$), one infectious individual generates new infections at rate $c\beta$ per day. The infectious period lasts on average $\tau_R$ days. The total number of secondary infections is therefore

$$\mathcal{R}_0 = \underbrace{c}_{\text{contacts/day}} \times \underbrace{\beta}_{\substack{\text{prob. infection} \\ \text{per contact}}} \times \underbrace{\tau_R}_{\text{infectious days}} = c\beta\tau_R. \tag{2.20}$$

Each factor is biologically independent: contacts per day depend on social behavior; transmission probability per contact depends on pathogen biology; and the infectious period depends on immune response. Because $\mathcal{R}_0$ is a product of three independent factors, varying any one of them by $x\%$ changes $\mathcal{R}_0$ by exactly $x\%$, a preview of why all three sensitivity indices equal 1. Table 2.1 collects the symbols, units, and nominal values used for this model throughout the book; the worked examples and the companion notebooks all start from these values.

### 2.5.2 ▪ The Basic Reproduction Number

The basic reproduction number

$$\mathcal{R}_0 = c\beta\tau_R$$

measures the average number of new infections produced by one infectious individual in an otherwise susceptible population (equation (2.20)). The rigorous definition of $\mathcal{R}_0$ via the next-generation operator is given in van den Driessche and Watmough [63] and Diekmann et al. [15].

In Chapter 4, we will compute time-dependent sensitivity indices for the full SIR model. Here we compute the static sensitivity indices for $\mathcal{R}_0$: a closed-form calculation that introduces all the key ideas in a familiar setting.

### 2.5.3 ▪ Reparameterizing the SIR Model

The recovery terms $I/\tau_R$ in equations (2.18)–(2.19) can also be written as $\gamma_R I$, with the recovery rate $\gamma_R = 1/\tau_R$. In that rate form, $c$, $\beta$, and $\gamma_R$ appear directly in the model equations, which makes them a natural choice for POIs. However, we argued in Chapter 1 that POIs should be chosen to be interpretable in everyday terms.

| Parameter | Symbol | Units | Nominal | Biological meaning |
|---|---|---|---|---|
| Contact rate | $c$ | $\text{day}^{-1}$ | 5 | Mean daily contacts per person |
| Transmission prob. | $\beta$ | – | 0.06 | Probability of infection per contact with an infectious person |
| Infectious period | $\tau_R$ | days | 7 | Mean number of days a person remains infectious |
| Mean residence time | $\tau_m$ | days | 10,000 | Mean residence time $\tau_m = 1/\nu_m$; unused in this chapter's model, which has no demography. From Chapter 4 onward $\nu_m$ removes infectives and the threshold becomes $c\beta/(\gamma_R + \nu_m)$ |
| Derived quantity $\mathcal{R}_0 = c\beta\tau_R$ | | – | $5 \times 0.06 \times 7 = 2.1$ | Average secondary infections per case in a susceptible population |

**Table 2.1.** *SIR model parameters, nominal values, and biological interpretation. Nominal values correspond to a moderately transmissible infection (influenza-like) in a well-mixed population. Typical ranges: $c \in [2, 20]$, $\beta \in [0.01, 0.5]$, $\tau_R \in [2, 14]$ days, giving $\mathcal{R}_0 \in [0.04, 140]$; the biologically plausible range for epidemic spread is $\mathcal{R}_0 > 1$.*

The recovery rate $\gamma_R$ (units: $\text{day}^{-1}$) is not directly interpretable. What a clinician understands and can measure is the *mean infectious period* $\tau_R = 1/\gamma_R$ (units: days): the average number of days a patient remains infectious. These two parameterizations are linked by:

$$\mathcal{S}^q_{\tau_R} = -\mathcal{S}^q_{\gamma_R} \qquad \text{for any QOI } q, \tag{2.21}$$

since $\partial q/\partial\tau_R = (\partial q/\partial\gamma_R)(\partial\gamma_R/\partial\tau_R) = -\gamma_R^2(\partial q/\partial\gamma_R)$, giving $\mathcal{S}^q_{\tau_R} = (\tau_R/q)(\partial q/\partial\tau_R) = -(\gamma_R/q)(\partial q/\partial\gamma_R) = -\mathcal{S}^q_{\gamma_R}$.

The magnitude is unchanged; the sign flips. Table 2.2 shows the effect of this reparameterization for $\mathcal{R}_0$.

| POI | Interpretation | Form | $\mathcal{S}_p^{\mathcal{R}_0}$ |
|---|---|---|---|
| Recovery rate $\gamma_R$ (day$^{-1}$) | – | rate | $-1$ |
| Mean infectious period $\tau_R = 1/\gamma_R$ (days) | "average days infectious" | duration | $+1$ |
| Contact rate $c$ (day$^{-1}$) | "contacts per day" | rate | $+1$ |
| Transmission probability $\beta$ (dimensionless) | "chance of infection per contact" | probability | $+1$ |

**Table 2.2.** *Sensitivity indices of $\mathcal{R}_0$ with respect to rate and duration parameterizations of the SIR model. The index for the mean infectious period $\tau_R = 1/\gamma_R$ has the same magnitude as the index for $\gamma_R$ but the opposite sign. The duration form is easier to interpret: a longer infectious period increases transmission, and both the sign and the plain-language statement agree.*

### 2.5.4 ▪ Analytic Computation

With the interpretable POIs $\{\tau_R = 1/\gamma_R,\, c,\, \beta\}$, the sensitivity indices are:

$$\mathcal{S}_c^{\mathcal{R}_0} = \frac{c}{\mathcal{R}_0} \cdot \frac{\partial \mathcal{R}_0}{\partial c} = \frac{c}{c\beta/\gamma_R} \cdot \frac{\beta}{\gamma_R} = 1, \tag{2.22}$$

$$\mathcal{S}_\beta^{\mathcal{R}_0} = \frac{\beta}{\mathcal{R}_0} \cdot \frac{\partial \mathcal{R}_0}{\partial \beta} = \frac{\beta}{c\beta/\gamma_R} \cdot \frac{c}{\gamma_R} = 1, \tag{2.23}$$

$$\mathcal{S}_{\tau_R}^{\mathcal{R}_0} = -\mathcal{S}_{\gamma_R}^{\mathcal{R}_0} = -\frac{\gamma_R}{\mathcal{R}_0} \cdot \frac{\partial \mathcal{R}_0}{\partial \gamma_R} = -\frac{\gamma_R}{c\beta/\gamma_R} \cdot \left(\frac{-c\beta}{\gamma_R^2}\right) = +1. \tag{2.24}$$

All three POIs have sensitivity index $+1$ with the interpretable parameterization. In plain language: a 1% increase in contact rate, transmission probability, *or* mean infectious period each increases $\mathcal{R}_0$ by 1%. The three parameters are equally important for $\mathcal{R}_0$, and all three promote disease spread when increased, as every positive sign and every positive biological interpretation confirms.

*Self-check.* An intervention reduces the contact rate $c$ by 20%. Using the sensitivity indices above, estimate the change in $\mathcal{R}_0$. If a second intervention reduces the mean infectious period $\tau_R$ by 15%, which intervention has the greater effect on $\mathcal{R}_0$?

> *Answer:* First intervention: $\Delta\mathcal{R}_0/\mathcal{R}_0 \approx \mathcal{S}_c^{\mathcal{R}_0} \times (-20\%) = -20\%$. Second intervention: $\Delta\mathcal{R}_0/\mathcal{R}_0 \approx \mathcal{S}_{\tau_R}^{\mathcal{R}_0} \times (-15\%) = -15\%$. The first intervention has the greater effect. Both indices equal 1, so the percentage change in $\mathcal{R}_0$ equals the percentage change in the POI. The intervention that changes its POI by the larger percentage has the larger effect.

## 2.6 ▪ Choosing Interpretable POIs

The SIR example illustrates a general principle that should guide every SA study before any calculation begins.

**Choose pois that can be discussed in ordinary language, not merely because they appear in the model equations.**

Model equations are written in the language that is mathematically convenient, which is not always the language that is scientifically or operationally natural. Rates (with units day$^{-1}$) are mathematically convenient in differential equations but not naturally interpretable; the corresponding mean times (with units days) are easier to discuss in a clinical or policy context.

The general rule for rate parameters in compartmental models: if a parameter $p$ is a per-capita rate (units: time$^{-1}$), define the corresponding POI as $\tau_R = 1/p$ (the mean time spent in that state), compute $\mathcal{S}^q_{\tau_R} = -\mathcal{S}^q_p$, and report the result in terms of $\tau_R$.

Two additional cases arise frequently:

**Dimensionless probabilities.** Transmission probabilities like $\beta$ are already dimensionless and directly interpretable: "a 25% chance of transmission per contact" requires no conversion.

**Composite parameters.** In the SIR model, $c$ and $\beta$ always appear together as the product $c\beta$ in the expression for the force of infection $\lambda = c\beta I/N$. Neither can be identified separately from incidence data alone; only the product $c\beta$ is identifiable. In this case, $c\beta$ is a more natural single POI than either factor individually. A related issue arises whenever two POIs are not independently changeable, which leads to the topic of correlated POIs in Chapter 3.

A practical check: if you cannot explain what a 1% increase in a POI means in plain language to a colleague outside mathematics, reconsider the parameterization.

Even in highly cited published work, this distinction is sometimes overlooked. Section 4.1 of Chitnis, Hyman & Cushing [14], a paper that has become a standard reference for SA in mathematical epidemiology, reports sensitivity indices using rate parameters throughout. The SI values are mathematically correct, but several indices are negative for rate parameters whose reciprocals, the mean durations, clearly promote disease spread, requiring a mental inversion before the public health interpretation is clear. The present book recommends always asking, before reporting any SI: would a policy maker understand this sign immediately, or does it require translation?

One further misconception deserves attention here. A sensitivity index near zero does *not* mean a parameter is globally unimportant. It means the model is approximately linear in that parameter near the nominal value, and the linear term happens to be small. If the model is nonlinear in that parameter, varying it over a wider range may reveal substantial effects that

the SI misses entirely. Chapter 9 addresses this through global sensitivity analysis.

*Self-check.* A malaria model contains the parameter $\mu_v$ (mosquito mortality rate, units: day$^{-1}$). A colleague reports $\mathcal{S}_{\mu_v}^{\mathcal{R}_0} = -0.3$. Reparameterize in terms of mean mosquito lifespan $L = 1/\mu_v$ (units: days). What is $\mathcal{S}_L^{\mathcal{R}_0}$? Write a plain-language interpretation of each sensitivity index.

*Answer:* $\mathcal{S}_L^{\mathcal{R}_0} = -\mathcal{S}_{\mu_v}^{\mathcal{R}_0} = +0.3$.
Rate form: "A 10% increase in mosquito mortality rate decreases $\mathcal{R}_0$ by 3%."
This requires mental inversion: higher mortality → shorter lifespan → less transmission.
Duration form: "A 10% increase in mean mosquito lifespan increases $\mathcal{R}_0$ by 3%."
Direct: longer-lived mosquitoes transmit more malaria.

## 2.7 ▪ The Tornado Plot

When a model has several POIs, the collection of sensitivity indices becomes a list of numbers with signs and magnitudes. Reading such a list is manageable for three or four parameters; for ten or twenty, a visual display is essential.

The *tornado plot* is the standard display format for sensitivity indices. It is a horizontal bar chart with the following conventions:

- Bars extending to the *right* indicate positive sensitivity indices (POI and QOI move in the same direction).
- Bars extending to the *left* indicate negative sensitivity indices (opposite directions).
- Bars are sorted from largest to smallest by absolute magnitude, with the most influential parameter at the top.
- The length of each bar represents the magnitude of the index.

The sorted layout creates the "tornado" shape: wider at the top, narrower toward the bottom.

Figure 2.2 shows the tornado plot for $\mathcal{R}_0$ in the SIR model with POIs $\{c, \beta, \tau_R\}$. All three bars extend to the right (all indices are +1), indicating that each parameter promotes disease spread when increased. All bars are the same length, confirming that the three parameters are equally important for $\mathcal{R}_0$. *Unifying Idea 4.* That equality is a fact about the algebra of $\mathcal{R}_0 = c\beta\tau_R$, not about epidemics. It says the three factors enter the product identically; it does not say a public-health program can move them equally, or that they cost the same to move.

Algorithm to produce a tornado plot for any set of sensitivity indices:

```
# TORNADO_PLOT: horizontal bar chart sorted by |SI|
# Input:  SI_values  - vector of sensitivity indices (positive or negati
```





**Figure 2.2.** *Tornado plot for the sensitivity of $\mathcal{R}_0$ to the three* POIs *of the SIR model, parameterized using interpretable durations and probabilities. All three indices equal +1, so the bars are equal in length and all extend to the right. In models with more parameters and a wider range of index values, the sorted layout reveals immediately which parameters dominate.*

```
#         labels     - parameter names (strings)
#         QOI_name   - name of the quantity of interest

FUNCTION TORNADO_PLOT(SI_values, labels, QOI_name):
    # Sort parameters by absolute sensitivity, largest first
    order      = ARGSORT(ABS(SI_values), descending=True)
    SI_sorted  = SI_values[order]
    lab_sorted = labels[order]

    # Draw horizontal bars (positive → right, negative → left)
    BARH(SI_sorted, labels=lab_sorted)
    DRAW_VERTICAL_LINE(x=0)         # center reference line
    XLABEL("Sensitivity Index  SI_p^" + QOI_name)
    TITLE("Tornado Plot: Sensitivity of " + QOI_name)
END
```

*Self-check.* A model for antibiotic-resistance spread has sensitivity indices $\mathcal{S}^{\mathcal{R}_0}_{\tau_{\text{inf}}} = +1.4$, $\mathcal{S}^{\mathcal{R}_0}_{\beta} = +0.9$, $\mathcal{S}^{\mathcal{R}_0}_{\tau_{\text{immune}}} = -0.6$, $\mathcal{S}^{\mathcal{R}_0}_{c} = +0.5$.

(a) Sketch the tornado plot by hand. (b) Which parameter is most important? Least important? (c) If $\tau_{\text{immune}}$ (mean time spent immune)

increases by 5%, what happens to $\mathcal{R}_0$?

*Answer:* (a) Sorted order by |SI|: $\tau_{\text{inf}}$ (1.4, right), $\beta$ (0.9, right), $\tau_{\text{immune}}$ (0.6, *left*), $c$ (0.5, right). (b) Most important: $\tau_{\text{inf}}$. Least: $c$. (c) $\mathcal{R}_0$ decreases by approximately $0.6 \times 5\% = 3\%$. The negative sign and the biological interpretation agree: a longer immune period keeps people protected longer, reducing transmission.

## 2.8 ▪ A Preview: Second-Order Terms

The sensitivity index is built on a first-order approximation. For a QOI $q$ that depends on a single POI $p$, the Taylor expansion gives

$$q(p+\delta p) = q(p) + \underbrace{\frac{\partial q}{\partial p}\,\delta p}_{\text{first-order}} + \underbrace{\frac{1}{2}\frac{\partial^2 q}{\partial p^2}\,(\delta p)^2}_{\text{second-order}} + \cdots$$

The sensitivity index is derived from the first-order term alone. When the second-order term is small relative to the first-order term (that is, when

$$\left|\frac{1}{2}\frac{\partial^2 q}{\partial p^2}\,\delta p\right| \ll \left|\frac{\partial q}{\partial p}\right|,$$

the sensitivity index provides an accurate approximation. When this condition fails, the linear approximation is poor and the SI overstates or understates the true sensitivity.

Chapter 3 develops practical methods for estimating the second-order term numerically and for assessing when it is large enough to matter. This is the first instance of Unifying Idea 3 from Chapter 1: local SA is a linearization, and linearizations have limits. The second-order term is the first correction beyond those limits.

For two POIs $p_i$ and $p_j$ that are not varied independently (a situation addressed in Chapter 3), the cross term

$$\frac{\partial^2 q}{\partial p_i\,\partial p_j}\,\delta p_i\,\delta p_j$$

also enters the expansion. This cross derivative quantifies the interaction between the two POIs: it measures how the sensitivity to $p_i$ changes as $p_j$ varies. Chapter 9 shows how Sobol interaction indices generalize this idea to the full range of parameter values.

## 2.9 ▪ Computational Laboratory: Sensitivity Indices for the SIR Model

This laboratory implements and visualizes the sensitivity indices developed in the chapter, using the SIR model as the working example. Complete the

five parts in order; each builds on the previous.

**Part 1: Analytic SI via the Symbolic Toolbox.** Define $\mathcal{R}_0 = c\beta/\gamma_R$ using a symbolic differentiation tool. Compute $\mathcal{S}_c^{\mathcal{R}_0}$, $\mathcal{S}_\beta^{\mathcal{R}_0}$, $\mathcal{S}_{\gamma_R}^{\mathcal{R}_0}$ using the `diff` function and the derivative form of Definition 2.1. Simplify and confirm that all three equal $\pm 1$ analytically.

*Note:* The code below requires the Symbolic Math Toolbox. If you do not have it, skip directly to Part 3 and verify the same results numerically. Parts 3–5 do not require the Symbolic Toolbox.

```
syms c beta gamma positive
R0   = c * beta / gamma;
SI_c = simplify( (c/R0) * diff(R0, c) )
SI_b = simplify( (beta/R0) * diff(R0, beta) )
SI_g = simplify( (gamma/R0) * diff(R0, gamma) )
```

**Part 2: Reparameterize.** Replace $\gamma_R$ with the mean infectious period $\tau_R = 1/\gamma_R$. Rewrite $\mathcal{R}_0 = c\beta\tau_R$ symbolically. Compute $\mathcal{S}_{\tau_R}^{\mathcal{R}_0}$ and verify that it equals $-\mathcal{S}_{\gamma_R}^{\mathcal{R}_0}$ as predicted by equation (2.21).

**Part 3: Numerical verification.** Use the nominal values $c = 5$, $\beta = 0.06$, $\tau_R = 7$ days (so $\gamma_R = 1/7$). Compute $\mathcal{R}_0$ numerically. Estimate $\mathcal{S}_{\tau_R}^{\mathcal{R}_0}$ using a centered finite difference with $h = 10^{-5}\tau_R$. Compare with the analytic value of $+1$. How large is the relative error?

```
c = 5;  beta = 0.06;  tau = 7
R0    = c * beta * tau
h     = 1e-5 * tau                          # optimal step size
R0_p  = c * beta * (tau + h)
R0_m  = c * beta * (tau - h)
SI_fd = (tau / R0) * (R0_p - R0_m) / (2*h)
PRINT "Analytic SI: 1.00000\n');
PRINT "FD estimate: %.5f\n', SI_fd);
```

**Part 4: Tornado plots for both parameterizations.** Using the `tornado_plot` function from Section 2.7, produce two tornado plots side by side: one with rate parameters $\{c, \beta, \gamma_R\}$ and one with interpretable parameters $\{c, \beta, \tau_R\}$. Observe how the sign of the $\gamma_R/\tau_R$ bar changes between the two plots and why.

*MATLAB note:* The `tornado_plot` function must be saved as a separate file named `tornado_plot.m` in your working directory (function files cannot be pasted into a script). Part 4 also uses `subplot` and cell array syntax for labels; the reference library provides `software/matlab/core/run_sir_example.m`, which demonstrates both if these are unfamiliar.

**Part 5: Effect of nominal value.** Compute $\mathcal{S}_c^{\mathcal{R}_0}$, $\mathcal{S}_\beta^{\mathcal{R}_0}$, $\mathcal{S}_{\tau_R}^{\mathcal{R}_0}$ at two different nominal transmission probabilities: $\beta_1 = 0.06$ and $\beta_2 = 0.6$. Do the SI values change? What does this tell you about the sensitivity of $\mathcal{R}_0$ to each parameter? (Recall that $\mathcal{R}_0 = c\beta\tau_R$ is a product of monomials; for such functions, the SI is constant regardless of the nominal value. This is a special property of multiplicative models that does not hold for more complex functions like those in Chapter 4.)

## 2.10 ▪ Exercises

**Exercise 2.1** *([Bloom L2])*. Explain the sign rule in your own words. Without performing any calculation, state the sign of the sensitivity index for each of the following, and explain your reasoning: (a) the sensitivity of lung capacity to cigarette consumption, treating capacity as the QOI and cigarettes per day as the POI; (b) the sensitivity of average crop yield to annual rainfall, near a region of the parameter space where crops are mildly drought-stressed; (c) the sensitivity of $\mathcal{R}_0$ to the mean infectious period $\tau_R$.

**Exercise 2.2** *([Bloom L2])*. For the function $q = \sqrt{p}$ (defined for $p > 0$), compute $\mathcal{S}_p^q$ using all three equivalent forms of Definition 2.1. Verify that all three give the same result. Interpret the answer: if $p$ doubles, by approximately what percentage does $q$ change?

**Exercise 2.3** *([Bloom L3])*. Let $u(t) = \frac{1}{2}\lambda t^2$ be a parabola on $[0, b]$. The arc-length functional is $J = \int_0^b \sqrt{1 + (\lambda t)^2}\, dt$. Show that the sensitivity indices are

$$\mathcal{S}_b^J = \frac{2\mu_b}{\mu_b + (\operatorname{arcsinh}\mu_b)/\sqrt{1+\mu_b^2}}, \qquad \mathcal{S}_\lambda^J = \frac{\mu_b(1+\mu_b^2) - \sqrt{1+\mu_b^2}\,\operatorname{arcsinh}\mu_b}{\mu_b(1+\mu_b^2) + \sqrt{1+\mu_b^2}\,\operatorname{arcsinh}\mu_b},$$

where $\mu_b := \lambda b$ is dimensionless, and is unrelated to the demographic rate $\nu_m$ used elsewhere in this chapter. For $\mu_b = 1$, compute both indices numerically and verify using the derivative form.

**Exercise 2.4** *([Bloom L3])*. Let $u(t) = at^2 + bt + c$ and let $J_\pm = (-b \pm \sqrt{b^2 - 4ac})/(2a)$ denote the two real roots (assume $b^2 > 4ac$). Show that the sensitivity indices of the positive root $J_+$ with respect to $a$, $b$, $c$ can be written

$$\mathcal{S}_a^{J_+} = -\frac{aJ_+}{2aJ_+ + b}, \quad \mathcal{S}_b^{J_+} = -\frac{b}{2aJ_+ + b}, \quad \mathcal{S}_c^{J_+} = -\frac{c}{J_+(2aJ_+ + b)}.$$

For $a = 1$, $b = -3$, $c = 2$, compute all three indices. Interpret each sign.

**Exercise 2.5** *([Bloom L3]).* A SEIR model (Susceptible-Exposed-Infectious-Recovered) has reproduction number

$$\mathcal{R}_0 = \frac{c\beta\,\nu}{(\nu + \nu_m)(\gamma_R + \nu_m)},$$

where $\nu$ is the rate of progression from exposed to infectious, $\nu_m$ is the background mortality rate, and $c$, $\beta$, $\gamma_R$ are as in the SIR model. Reparameterize using mean times: $\tau_E = 1/\nu$ (mean latent period), $\tau_I = 1/\gamma_R$ (mean infectious period), $\tau_m = 1/\nu_m$ (mean residence time). Compute $\mathcal{S}^{\mathcal{R}_0}_{\tau_E}$, $\mathcal{S}^{\mathcal{R}_0}_{\tau_I}$, $\mathcal{S}^{\mathcal{R}_0}_{\tau_m}$, $\mathcal{S}^{\mathcal{R}_0}_{c}$, $\mathcal{S}^{\mathcal{R}_0}_{\beta}$ and produce a tornado plot. Which parameter is most important?

*Nominal values for the tornado plot:* $c = 5$, $\beta = 0.06$, $\tau_E = 5.5$ days, $\tau_I = 7$ days, $\tau_m = 10{,}000$ days.

**Exercise 2.6** *([Bloom L4]).* A published paper reports $\mathcal{S}^{\mathcal{R}_0}_{\gamma_R} = -1.0$ for an SIR model, where $\gamma_R$ is the recovery rate in units of day$^{-1}$. A second paper on the same model reports $\mathcal{S}^{\mathcal{R}_0}_{\tau_R} = +1.0$, where $\tau_R = 1/\gamma_R$ is the mean infectious period in days. A policy maker reading both papers is confused: one paper shows a negative index, the other shows a positive index for what appears to be the same parameter.

Write a one-paragraph explanation for the policy maker that (a) resolves the apparent contradiction, (b) explains which parameterization is more informative for policy, and (c) describes what each sensitivity index means in plain language.

**Exercise 2.7** *([Bloom L4]).* The following tornado plot data is given for a food-safety model with QOI $J$ = mean number of illness cases per outbreak. The sensitivity indices are: $\mathcal{S}^{J}_{\tau_c} = +2.1$ (mean contamination duration, days), $\mathcal{S}^{J}_{\beta_f} = +1.8$ (food-to-person transmission probability), $\mathcal{S}^{J}_{\tau_s} = +1.3$ (mean hand-washing interval, days), $\mathcal{S}^{J}_{k_f} = +0.7$ (food contacts per day), $\mathcal{S}^{J}_{\nu_m} = -0.2$ (background clearance rate).

(a) Produce the tornado plot using the algorithm template from Section 2.7.

(b) Rank the five parameters by absolute importance.

(c) A public health program can achieve a 15% reduction in either $\tau_c$ or a 20% reduction in $k_f$. Which intervention reduces $J$ more? By how much?

(d) The parameter $\nu_m$ has a small negative index. Write a one-sentence interpretation of what this means for disease control.

**Exercise 2.8** *([Bloom L5])*. You are advising the World Health Organization on a vaccine allocation strategy for a new respiratory pathogen. A compartmental model has been fitted to early outbreak data. The model has six parameters; their sensitivity indices with respect to the QOI "total deaths at 6 months" are not yet known.

Design the SA study. Your design should specify:

(a) Which parameterization you will use (rates or durations) for each parameter, and why.

(b) What QOI or QOIs you would compute sensitivity indices for, and what additional QOI you might consider beyond the stated one.

(c) How you would present the results to a non-mathematical audience (what figure or table, and what the key message would be).

(d) One limitation of the local SA approach that the WHO advisors should be aware of in using these results.

**Exercise 2.9** *([Bloom L4])*. *Audit exercise.* A report on an SIR model states: "The sensitivity index of $\mathcal{R}_0$ with respect to the contact rate $c$ is $+1$. The relationship between $c$ and $\mathcal{R}_0$ is therefore linear, so doubling the contact rate doubles $\mathcal{R}_0$, and halving it halves $\mathcal{R}_0$." The numerical value quoted is correct.

(a) Identify the error in the interpretation. What does an index of $+1$ actually assert, and over what range?

(b) Construct a function of one variable whose sensitivity index equals $+1$ at the nominal point but which is not linear, and evaluate the error in the report's doubling claim for your function.

(c) The report's final sentence would be defensible for $\mathcal{R}_0 = c\beta\tau_R$ under one additional condition. State it.

(d) Rewrite the passage so that it is correct and no longer than the original.

**Exercise 2.10** *([Bloom L6])*. A modeling study reports the following: "Sensitivity analysis of $\mathcal{R}_0$ was performed with respect to all model parameters. The parameters are listed in Table 3 using the standard model notation. All sensitivity indices were positive, confirming that $\mathcal{R}_0$ is an increasing function of all model parameters."

Table 3 lists recovery rate $\gamma_R$, mortality rate $\nu_m$, and waning immunity rate $\rho$ alongside transmission and contact parameters.

Identify the error in the study's statement and explain specifically what is wrong, referring to the mathematical relationship between rates and their reciprocals. Restate the conclusion correctly.

## Notes

The three equivalent forms of the sensitivity index (limit, derivative, and logarithmic) appear in Arriola and Hyman [2], which also introduced the sign rule and the interpretable-parameterization convention adopted throughout this book. Earlier uses of normalized sensitivity in ecology and economics appear under the name *elasticity*; see, for example, the biological applications surveyed in Saltelli et al. [50].

The tornado diagram as a visualization tool for ranking sensitivity indices was popularized in risk analysis and decision science; see Saltelli et al. [55] for its use in the SA context.

Blower and Dowlatabadi [5] gave an early and influential application of sensitivity analysis to an HIV transmission model, demonstrating how tornado plots guide intervention priorities. The application of SI to $\mathcal{R}_0$ in epidemic models, and the connection between $\mathcal{S}_{\tau_R}^{\mathcal{R}_0} = +1$ and $\mathcal{S}_{\gamma_R}^{\mathcal{R}_0} = -1$ for the same biological fact, is discussed explicitly in Arriola and Hyman [3].

For the near-zero-QOI regime and regularized sensitivity indices, see the discussion in Saltelli et al. [50], Chapter 3.

Draft

Draft

# Part II

# Computing Sensitivity in Practice

Draft

Draft

# Chapter 3
# Computing Sensitivity in Practice

*A derivative is useful only if you can compute it honestly.*

You have inherited a model from a collaborator. It is 500 lines of MATLAB code, produces reasonable epidemic projections, and has 11 parameters. Your collaborator has moved to another institution. The code runs correctly, but it uses conditional statements, lookup tables, and an internal adaptive ODE solver; none of which can be differentiated analytically. Your funding agency wants to know which parameters matter before committing to a three-year field measurement campaign.

How do you compute sensitivity indices for a model you cannot differentiate?

The answer is finite differences, and this chapter shows exactly how to do it, including how to avoid the mistakes that produce wrong results.

## 3.1 ▪ The Sensitivity Jacobian

Chapter 2 computed sensitivity indices one parameter at a time. For a model with $n_p$ POIs and $n_q$ QOIs, the complete local SA is captured by an $n_q \times n_p$ matrix; one entry per POI–QOI pair.

**Definition 3.1** (Sensitivity Jacobian)**.** *Let $\vec{q} = (q_1, \ldots, q_{n_q})$ denote the vector of* QOI*s and $\vec{p} = (p_1, \ldots, p_{n_p})$ denote the vector of* POI*s. The* normalized sensitivity Jacobian *is the $n_q \times n_p$ matrix $\mathbf{S}$ with entries*

$$S_{ki} = \mathcal{S}_{p_i}^{q_k} = \frac{p_i}{q_k} \frac{\partial q_k}{\partial p_i}. \tag{3.1}$$

For the SIR model with POIs $\{c, \beta, \tau_R, \nu_m\}$ (contact rate, transmission probability, mean infectious period, mortality rate) and QOIs $\{S(T), I(T), R(T)\}$

(the three compartment sizes at time $T$), the sensitivity Jacobian is a $3 \times 4$ matrix of twelve numbers. Each number answers one specific "What if?" question: how much does compartment $i$ change if POI $j$ changes by 1%?

The tornado plots in Chapter 2 displayed single rows or columns of this matrix. The full matrix captures all POI–QOI relationships simultaneously.

Computing this matrix efficiently, in both time and accuracy; is the central computational challenge of sensitivity analysis.

A common initial reaction is to treat the sensitivity Jacobian as a collection of individual SI calculations, with the matrix label serving as convenient organization for numbers computed independently. It is more than that. The Jacobian has structure: its columns can be computed in parallel, many methods share a single baseline evaluation, and the adjoint method (Chapters 5 and 6) exploits the matrix structure to recover one complete *row* of the sensitivity Jacobian, all parameter sensitivities for one scalar QOI, from a single backward solve. For $r$ QOIs, $r$ backward solves recover all $r$ rows. Understanding the Jacobian as a matrix, not just a table, is what makes efficient SA possible.

*Self-check.* A climate model has $n_p = 30$ POIs and $n_q = 8$ QOIs. How many individual sensitivity indices does the full SA require? If you had to compute each one separately, how many model evaluations would a naive brute-force approach require?

> *Answer:* $30 \times 8 = 240$ sensitivity indices. Computing each index separately, with its own perturbed runs and one shared baseline run, is the naive approach: $2 \times 240 + 1 = 481$ model evaluations for centered differences, or $240 + 1 = 241$ for first-order (forward) differences. A better approach perturbs one POI at a time and reads all eight QOIs from each run. It requires $30 + 1 = 31$ model evaluations for forward differences, or $2 \times 30 + 1 = 61$ for centered differences. The key: you do not need one evaluation per SI entry. For forward differences you need one evaluation per POI (plus one baseline), and then each evaluation contributes an entire column of the Jacobian. Centered differences need two per POI, one on each side.

## 3.2 ▪ Finite Difference Formulas

A finite difference approximates the partial derivative $\partial q/\partial p_i$ by computing how the QOI changes when $p_i$ alone is shifted by a small but finite amount $h$. Three approximations are in common use, differing in accuracy and cost.

### 3.2.1 ▪ Forward Difference

The simplest approximation uses one additional model evaluation:

$$\frac{\partial q}{\partial p_i} \approx \frac{q(p^* + h\vec{e}_i) - q(p^*)}{h}, \tag{3.2}$$

where $\vec{e_i}$ is the unit vector in the $p_i$ direction and $p^*$ is the nominal parameter vector. This is a first-order approximation: the error is $O(h)$, meaning that halving $h$ approximately halves the error.

### 3.2.2 ▪ Centered Difference

Using one evaluation on each side of the nominal value gives a second-order approximation:

$$\frac{\partial q}{\partial p_i} \approx \frac{q(p^* + h\vec{e_i}) - q(p^* - h\vec{e_i})}{2h}. \tag{3.3}$$

The error is $O(h^2)$: halving $h$ reduces the error by a factor of four. For the same step size, centered differences are substantially more accurate than forward differences.

### 3.2.3 ▪ Fourth-Order Formula (Rarely Used in Practice)

When exceptionally high accuracy is required and model evaluations are inexpensive, one can eliminate the leading-order error term by combining four evaluations at two step sizes:

$$\frac{\partial q}{\partial p_i} \approx \frac{-q(p^* + 2h\vec{e_i}) + 8q(p^* + h\vec{e_i}) - 8q(p^* - h\vec{e_i}) + q(p^* - 2h\vec{e_i})}{12h}. \tag{3.4}$$

The error is $O(h^4)$: halving $h$ reduces the error by a factor of sixteen. This requires four evaluations per POI rather than two. In the applied SA literature this formula is uncommon; the centered difference (3.3) is the standard workhorse. Knowing which formula to use and what step size to choose are the practical skills taken up next.

### 3.2.4 ▪ Practical Workflow: Checking the Step with an Error Indicator

In practice, the centered difference (3.3) is the default choice. The forward difference (3.2) plays a secondary role not as a first approximation in its own right, but as a cheap error estimator: because it requires only the already-computed baseline $q(p^*)$ and the one-sided perturbed evaluation $q(p^* + h\vec{e_i})$, no extra model runs are needed. The difference between the two approximations at the same $h$,

$$\text{estimated error} \approx \left| \frac{q(p^* + h\vec{e_i}) - q(p^*)}{h} - \frac{q(p^* + h\vec{e_i}) - q(p^* - h\vec{e_i})}{2h} \right|, \tag{3.5}$$

provides a practical error indicator. Be clear about what it measures. The difference between the forward and centered quotients is dominated by the

forward formula's $O(h)$ truncation term, since the centered one is already $O(h^2)$, so (3.5) estimates the error in the *forward* difference. It is a sound diagnostic for that, and a sound signal that $h$ is in a sensible range, but it is not an error bar on the centered result and shrinking it to a tolerance does not bound the centered error. Use it to detect trouble, and set the step from the $\epsilon^{1/3}$ optimum in (3.9) rather than by driving this indicator down. That optimum depends on higher-order derivatives that are rarely known, so in practice one evaluates it with a rough magnitude estimate and checks the indicator afterward.

The backward difference $[q(p^*) - q(p^* - h\vec{e}_i)]/h$ adds no independent check. The centered quotient is the average of the forward and backward quotients, so the backward quotient differs from the centered one by the same amount as the forward quotient, with the opposite sign. Comparing it with the centered result repeats (3.5).

### 3.2.5 ▪ Cost Comparison

Table 3.1 summarizes the cost of computing the complete sensitivity Jacobian by each method.

| Method | Model evaluations | Error order | Best when |
|---|---|---|---|
| Forward difference | $n_p + 1$ | $O(h)$ | Error check only; not standalone |
| Centered difference | $2n_p + 1$ | $O(h^2)$ | Standard choice |
| Fourth-order FD | $4n_p + 1$ | $O(h^4)$ | Rare; cheap evaluations only |
| Analytic FSE (Chapter 4) | $n_p + 1$ solves | Exact | Differentiable model |
| Adjoint (Chapters 5–6) | 2 solves | Exact | Many POIs, few QOIs |

**Table 3.1.** *Cost of computing the complete sensitivity Jacobian by different methods, for a model with $n_p$* POI*s and $n_q$* QOI*s. Cost metric: finite-difference rows are counted in* model evaluations *(calls to the ODE solver or model function); FSE and adjoint rows are counted in* ODE solves *of the augmented system. For a system of $n$ equations, one ODE solve $\approx n$ function evaluations per time step. The finite-difference counts assume a single baseline evaluation is reused across all* POI*s. FSE cost: one augmented solve of size $n + n \cdot n_p$ equations (all parameters simultaneously). That single integration is counted here, and in Chapter 4, as $n_p + 1$ solves of the baseline $n$-equation size, which is the unit that makes the forward and adjoint costs directly comparable; adjoint cost assumes $r = 1$ scalar* QOI*, for $r$* QOI*s, adjoint requires $r$ backward solves, each of size $n$.*

For the SIR model with $n_p = 4$ parameters and $n_q = 3$ state variables, centered differences require $2 \times 4 + 1 = 9$ model evaluations. If each takes 1 second, the full SA takes 9 seconds.

For the malaria model of Chitnis, Hyman & Cushing [14] with $n_p = 17$ parameters and $n_q = 7$ state variables, centered differences require $2 \times 17 + 1 = 35$ evaluations. If each takes 1 minute: approximately 35 minutes. If each takes 1 hour: approximately 35 hours.

For a climate model with $n_p = 500$ parameters and model runs taking 6 hours each, centered differences require $2 \times 500 + 1 = 1001$ evaluations, at 6 hours each, that is approximately 250 days of computation. This is why Chapter 6 introduces the adjoint method, which reduces the cost to 2 solves for one QOI, regardless of $n_p$.

*Unifying Idea 1.* Every entry of the sensitivity Jacobian is a normalized derivative: an instance of Unifying Idea 1 from Chapter 1. The finite-difference formulas in this chapter compute those derivatives numerically, for any model that can be evaluated, whether or not it can be differentiated analytically.

*Self-check.* For a model with $n_p = 20$ POIs and $n_q = 5$ QOIs, how many model evaluations does the centered-difference Jacobian require? (Remember: adding more QOIs is free, each evaluation contributes an entire column of the Jacobian.) If each evaluation takes 45 seconds, how long does the complete SA take? At what value of $n_p$ would centered differences take more than 24 hours, assuming each evaluation takes 2 minutes?

*Answer:* $2 \times 20 + 1 = 41$ evaluations, regardless of $n_q = 5$; $41 \times 45 = 1{,}845$ seconds $\approx 31$ minutes.
For 24 hours = 1,440 minutes with 2-minute evaluations: $2n_p + 1 \leq 720$ gives $n_p \leq 359$, so for $n_p \geq 360$, centered differences take more than 24 hours.

## 3.3 ▪ Step Size Selection: The U-Curve

The step size $h$ controls a fundamental accuracy tradeoff. There is an optimal value: too large and the approximation misses the local curvature; too small and the computer's limited numerical precision destroys the result.

### 3.3.1 ▪ Two Sources of Error

*Truncation error* arises because finite differences are Taylor-series approximations. For the centered difference:

$$\frac{q(p+h) - q(p-h)}{2h} = q'(p) + \underbrace{\frac{h^2}{6} q'''(p) + \cdots}_{\text{truncation error}}. \tag{3.6}$$

This error decreases as $h \to 0$; on a log-log plot of error vs. $h$ (with $h$ increasing to the right, as in Figure 3.1) the truncation branch has slope $+2$.

*Rounding error* arises because the computed model output $\tilde{q}(p+h)$ differs from the true output $q(p+h)$ by a rounding error of order $\varepsilon_{\text{mach}}|q|$, where $\varepsilon_{\text{mach}} \approx 2.2 \times 10^{-16}$ in IEEE 754 double precision. When $h$ is small,

subtracting $\tilde{q}(p-h)$ from $\tilde{q}(p+h)$ amplifies this rounding error:

$$\text{rounding error in FD} \approx \frac{2\varepsilon_{\text{mach}}|q|}{2h} = \frac{\varepsilon_{\text{mach}}|q|}{h}. \tag{3.7}$$

This error *increases* as $h \to 0$; on the same log-log plot the rounding branch has slope $-1$.

### 3.3.2 ▪ The U-Curve and the Optimal Step Size

*Before improving the model, find out which knob actually moves the machine.*

The total error in the centered-difference approximation is the sum of truncation and rounding errors. As $h$ decreases from a large value, truncation error falls while rounding error rises. The minimum total error occurs at the crossover, as shown schematically in Figure 3.1.





**Figure 3.1.** *The U-curve for centered differences in double precision, measured rather than sketched. The* QOI *is the closed-SIR peak prevalence* $I_{\max}(\beta) = N[1-(1+\ln r)/r]$ *with* $r = c\beta\tau_R$*, whose derivative* $\partial I_{\max}/\partial\beta = 5888.391625$ *is known in closed form at* $c = 5$*,* $\tau_R = 7$*,* $\beta^* = 0.06$*, so the error can be measured. Truncation error (dashed) falls as* $h^2$*; rounding error (dotted) grows as* $1/h$*; the measured total (solid) is least for* $h$ *of order* $10^{-7}$*, where the relative error is of order* $10^{-12}$*. Equation* (3.9) *predicts* $h^* = \varepsilon_{mach}^{1/3}|\beta^*| = 3.63 \times 10^{-7}$*, marked by the vertical line: the rule and the measurement agree in order of magnitude. Choosing* $h$ *far below* $h^*$ *is a common mistake and gives larger errors, not smaller ones; at* $h = 10^{-12}$ *the error is five orders of magnitude worse than at the optimum. Generated by* `generate_fig_ucurve.py`*.*

Setting the two error contributions equal gives an approximate starting point for step-size selection:

$$h^*_{\text{forward}} \approx \sqrt{\varepsilon_{\text{mach}}} \cdot |p^*| \approx 1.5 \times 10^{-8} \cdot |p^*|, \tag{3.8}$$

$$h^*_{\text{centered}} \approx \varepsilon_{\text{mach}}^{1/3} \cdot \max\big(|p^*|,\, p_{\text{scale}}\big) \approx 6 \times 10^{-6} \cdot \max\big(|p^*|,\, p_{\text{scale}}\big). \tag{3.9}$$

For a parameter with nominal value $p^* = 0.3$, this suggests a starting centered-difference step of approximately $h \approx 1.8 \times 10^{-6}$.

The factor $\max(|p^*|, p_{\text{scale}})$ matters: scaling $h$ by $|p^*|$ alone collapses to zero for a parameter whose nominal value is zero, so $p_{\text{scale}} > 0$ supplies a floor in the units of $p$.

These formulas should be understood as rough guides, not precise prescriptions. The balance being struck is between roundoff, of order $\varepsilon_{\text{mach}}|q|/h$, and truncation, of order $h^2|q'''|/6$; minimizing the sum gives $h^* \sim (\varepsilon_{\text{mach}}|q|/|q'''|)^{1/3}$. That ratio $|q|/|q'''|$ carries units of $p^3$, so its cube root is a length scale in parameter space, and the ratio is not a dimensionless quantity to be set to one. The estimates above replace that cube root with the only scale generally available in advance, the parameter's own magnitude. This assumes $|q'''| \approx |q|/|p^*|^3$, that is, that $q$ varies on the scale of the parameter itself. Because $q'''(p)$ is unknown, the true optimal $h$ remains model-dependent. Section 3.2 uses the error indicator (3.5) to check a step, not to choose one. The indicator answers a different question from the formulas and is not a substitute for them. Equation (3.9) tells you where to start; the indicator tells you whether the step you chose is in trouble, in either direction. Use the formula to pick $h$ and the indicator to check it, rather than driving the indicator down as though it were an error bar on the centered result.

The most common mistake in finite-difference SA is choosing $h$ that is too small. Practitioners who set $h = 10^{-12}$ or $h = 10^{-15}$, reasoning that "smaller is more accurate," obtain results dominated by catastrophic cancellation (subtracting two nearly equal floating-point numbers, which destroys significant digits and amplifies rounding error). The U-curve explains why: below the optimal step size, rounding error grows as $1/h$ while truncation error continues to fall, so the total error increases rather than decreases. A single value of the error indicator (3.5) does not reliably detect this failure. Rounding can leave the forward and centered quotients about as close at $h = 10^{-12}$ as at the optimal step, and for some models they agree exactly. The reliable signal is how the indicator scales. In the reliable range it is the forward formula's truncation error, about $(h/2)|q''|$, so it grows tenfold when $h$ grows tenfold. Compute it at $h$ and at $10h$: if it does not scale this way, rounding dominates and a larger step size is needed.

The U-curve also reveals a practical limitation for models implemented with adaptive ODE solvers. For a smooth analytic function the curve has a well-defined minimum, and in the reliable range the error indicator (3.5) shrinks in proportion to $h$. For a model that calls an ODE solver internally, the output is not mathematically smooth: the solver makes discrete step-size decisions that can cause the model output to change non-smoothly as a parameter varies by a small amount. In these cases the error indicator

may not decrease monotonically, and the one-sided and centered approximations may disagree by much more than their expected relative gap of about $(h/2)|q''/q'|$. A relative gap of order $10^{-6}$ at steps near $10^{-6}|p^*|$ is normal for a smooth model and is not a sign of solver trouble. The remedy is to tighten the ODE solver tolerances (`RelTol` and `AbsTol` in an ODE solver) before changing $h$. After tightening tolerances, pick $h$ by (3.9) and check it with the indicator at two steps, as above.

***ODE Solver Tolerance Rule***

Before computing finite-difference sensitivities for a model that calls an ODE solver internally, tighten the solver tolerance to at least *ten times smaller* than the target step size $h$. As a practical starting point: set `RelTol = AbsTol =` $10^{-8}$ whenever sensitivity analysis is the goal. If your target step size is $h \approx 10^{-5}|p^*|$ but the solver tolerance is $10^{-3}$, the finite-difference quotient measures solver noise rather than true sensitivity. After tightening tolerances, pick $h$ by (3.9) again and check it with the error indicator.

*Self-check.* For the SIR model with transmission probability $\beta^* = 0.06$, the theoretical formula (3.9) suggests a starting centered-difference step of $h \approx 3.6 \times 10^{-7}$. You then compute the centered difference and a one-sided forward difference at this $h$ and find they agree to four significant figures. (a) Is this step size acceptable? What does this agreement indicate? (b) Now suppose you repeat the check at $h = 10^{-12}$ and find the two approximations disagree by more than 50%. What does this signal, and what should you do?

*Answer:* (a) Yes. Agreement to four significant figures shows that the forward difference's truncation error is small at this $h$. Rounding is small because $h$ came from (3.9); a gap that grows tenfold at $10h$ would confirm it. For the centered formula, rounding and truncation errors are comparable at this $h$, by the design of (3.9). The agreement is a check on the step, not an error bar on the centered result (Section 3.2). The step size is acceptable.
(b) Large disagreement at $h = 10^{-12}$ signals that rounding error dominates: both approximations suffer catastrophic cancellation. The remedy is to increase $h$ toward the range suggested by the theoretical formula, then recheck the error indicator. Rounding error $\propto 1/h$: at $h = 10^{-12}$ the rounding error is $(3.6 \times 10^{-7}/10^{-12}) = 3.6 \times 10^5$ times larger than at $h^* \approx 3.6 \times 10^{-7}$. Moving to even larger $h$ would re-introduce truncation error, so the reliable range lies in between.

## 3.4 ▪ Second-Order Terms

Chapter 2 previewed second-order terms as corrections to the first-order SI approximation. This section shows how to compute them.

### 3.4.1 ▪ Pure Second-Order Derivatives

The second derivative $\partial^2 q/\partial p_i^2$ measures the curvature of the QOI as a function of $p_i$. It can be estimated by reusing the three evaluations already computed for the centered first-order difference:

$$\frac{\partial^2 q}{\partial p_i^2} \approx \frac{q(p^* + h\vec{e}_i) - 2q(p^*) + q(p^* - h\vec{e}_i)}{h^2}. \tag{3.10}$$

No additional model evaluations are needed.

When the curvature is large, the first-order SI can overstate or understate the true change for larger perturbations, as Chapter 2 noted. It understates the change when the second-order term has the sign of the first-order term, as for $q = p^2$ under an increase in $p$. It overstates the change when the two terms have opposite signs, as for $q = \sqrt{p}$, whose index is 1/2: a 10% rise in $p$ raises $q$ by 4.88%, not 5%. Specifically, if

$$\left| \frac{h}{2} \frac{\partial^2 q/\partial p_i^2}{\partial q/\partial p_i} \right| \ll 1,$$

the linear approximation is valid for perturbations of size $h$. If this ratio is not small, the sensitivity index provides only a local estimate, and Chapter 9's global SA methods are needed for a complete picture.

*Unifying Idea 3.* The second-order terms computed in this section are the first quantitative correction beyond the linearization that defines every sensitivity index in this book. The first-order SI is valid when the model behaves approximately linearly near the nominal parameter values. The second-order term tells you how quickly that approximation degrades as perturbations grow. Chapter 9 provides the full global picture when the linearization is no longer adequate.

### 3.4.2 ▪ Mixed Second-Order Derivatives

The cross-derivative $\partial^2 q/\partial p_i\, \partial p_j$ measures how the sensitivity to $p_i$ changes with $p_j$. For each pair $(i, j)$, one additional model evaluation (at $p^* + h\vec{e}_i + h\vec{e}_j$) is required beyond the baseline and single-perturbation evaluations already performed for the first-order Jacobian:

$$\frac{\partial^2 q}{\partial p_i\, \partial p_j} \approx \frac{q(p^* + h\vec{e}_i + h\vec{e}_j) - q(p^* + h\vec{e}_i) - q(p^* + h\vec{e}_j) + q(p^*)}{h^2}. \tag{3.11}$$

For all $\binom{n_p}{2}$ pairs, this requires $\binom{n_p}{2}$ additional evaluations. For $n_p = 4$, that is 6 evaluations; for $n_p = 17$ (the malaria model), it is 136.

The cross-derivative $\partial^2 q/\partial p_i\,\partial p_j$ is large when the two POIs *interact*: varying them together is not the same as varying them separately. Chapter 9 shows that Sobol interaction indices $S_{ij}$ are the global, variance-based analog of these local cross-derivatives.

*Self-check.* For the SIR model with $\mathcal{R}_0 = c\beta\tau_R$, compute $\partial^2\mathcal{R}_0/\partial c\,\partial\beta$ analytically. Is the cross-derivative large or small relative to the individual first-order derivatives $\partial\mathcal{R}_0/\partial c$ and $\partial\mathcal{R}_0/\partial\beta$? What does this imply about the interaction between $c$ and $\beta$ for $\mathcal{R}_0$?

*Answer:* $\partial^2\mathcal{R}_0/\partial c\,\partial\beta = \tau_R$ (positive, since $\mathcal{R}_0 = c\beta\tau_R$).
The first-order derivatives are $\partial\mathcal{R}_0/\partial c = \beta\tau_R$ and $\partial\mathcal{R}_0/\partial\beta = c\tau_R$.
The ratio $(\partial^2\mathcal{R}_0/\partial c\,\partial\beta)/(\partial^2\mathcal{R}_0/\partial c^2) = \tau_R/0$ is undefined (the pure second derivative is zero because $\mathcal{R}_0$ is linear in each parameter). More informatively, the mixed partial is nonzero, meaning $c$ and $\beta$ interact in their effect on $\mathcal{R}_0$: increasing both together increases $\mathcal{R}_0$ more than the sum of the individual increases. This is expected because $\mathcal{R}_0$ is multiplicative in $c$ and $\beta$. For Sobol analysis, this interaction will produce a nonzero $S_{c\beta}$.

## 3.5 ▪ Correlated POIs: The Most Common Mistake

*A parameter can be uncertain without being important, and important without being uncertain. Correlation between parameters adds a third possibility: a parameter can appear important in isolation while its true effect is inseparable from another.*

Before computing any sensitivity indices, there is a question that must be asked: can these parameters actually be varied independently? In many models, the answer is no.

Every finite-difference SA computation implicitly assumes that each POI can be perturbed while all others are held exactly fixed. If the POIs are correlated, meaning that in the real world, changing one tends to change another, then this assumption is violated and the resulting sensitivity indices can be misleading.

### 3.5.1 ▪ Diagnosing Correlation

Correlation among POIs arises from several sources.

*Structural correlation* occurs when parameters appear only as a product or ratio in the model equations. In the SIR model, the force of infection is $\lambda = c\beta I/N$. The parameters $c$ and $\beta$ appear only as the product $c\beta$. Observational data, incidence counts over time, can identify $c\beta$ as a unit but cannot separate the two factors. Computing $\mathcal{S}_c^{\mathcal{R}_0}$ and $\mathcal{S}_\beta^{\mathcal{R}_0}$ as independent indices is mathematically valid but can be operationally misleading,

because any intervention must target the product $c\beta$ (by reducing contact rate, transmission probability, or both). Reparameterizing to use $c\beta$ as a single POI is more natural in this case.

This observation is the starting point for *structural identifiability* analysis: the question of whether the parameters of a model can, in principle, be recovered from the observable outputs, regardless of measurement noise or sample size. A model is *structurally identifiable* in a parameter $p$ if distinct values of $p$ produce distinct output trajectories. The SIR model is not structurally identifiable in $(c, \beta)$ separately, but is identifiable in the product $c\beta$. Sensitivity analysis reveals this: when two parameters always have the same sensitivity index (here $\mathcal{S}_c^q = \mathcal{S}_\beta^q$ for any QOI $q$), they are structurally correlated and the model cannot distinguish them. A sensitivity index of zero for a parameter is a related signal: the model output does not respond to that parameter at all, which may indicate that the parameter cannot be estimated from data. Chapter 9 returns to identifiability in the context of global SA, where variance-based methods provide computational tools for detecting structural correlation across the full parameter range. A full treatment of structural and practical identifiability, including formal algebraic methods, is beyond the scope of this book; see Smith [59] Ch. 7 §7.2–7.3 and the references therein.

*Empirical correlation* occurs when parameter values are estimated from data that naturally links them. In the malaria model studied by Chitnis, Hyman & Cushing [14], the mosquito biting rate $\sigma_v$ and the human biting saturation $\sigma_h$ both appear in the force of infection and are often positively correlated across field studies. The paper introduces the composite parameter $\zeta = \sigma_v\sigma_h/(\sigma_v N_v^* + \sigma_h N_h^*)$ precisely to address this correlation.

*Constraint correlation* occurs when a physical or biological requirement links parameters. In models where stage durations must sum to a total lifecycle, or where transition probabilities must sum to one, no parameter can change without forcing compensating changes in others.

### 3.5.2 ▪ The Impact of Ignoring Correlation

To see concretely what goes wrong, consider a model whose QOI is $q = p_1 + p_2$ where the two POIs satisfy $p_2 = 2p_1$ (a linear constraint). The standard sensitivity indices are:

$$\mathcal{S}_{p_1}^q = \frac{p_1}{q} \cdot 1 = \frac{p_1}{3p_1} = \frac{1}{3}, \qquad \mathcal{S}_{p_2}^q = \frac{p_2}{q} \cdot 1 = \frac{2p_1}{3p_1} = \frac{2}{3}.$$

These look reasonable: the second parameter is twice as important as the first. But in reality, $p_1$ and $p_2$ cannot be varied independently. Any change

to $p_1$ forces a change to $p_2 = 2p_1$. The total effect of a 1% increase in $p_1$ is:

$$\Delta q = \frac{\partial q}{\partial p_1}\,\delta p_1 + \frac{\partial q}{\partial p_2}\,\delta p_2 = 1 \cdot (0.01p_1) + 1 \cdot (0.02p_1) = 0.03p_1.$$

The normalized total-derivative SI for $p_1$ is then $(p_1/q) \cdot (0.03p_1/0.01p_1) = (1/3) \cdot 3 = 1$, not 1/3. The standard index understates the true sensitivity by a factor of three.

### 3.5.3 ▪ Four Practical Responses

*Readers with a regression background will recognize correlated* POIs *as the SA analog of multicollinearity.* In regression, when predictors are nearly collinear, the design matrix $X^\top X$ is nearly singular, coefficient estimates become unstable, and variance inflation factors (VIFs) grow large. In SA, when POIs are correlated, varying one parameter while holding others fixed may be physically impossible, as for constraint correlation. It may also be unlike any change the data show. The resulting sensitivity indices can be misleading for the same reason: the "separate" effects of the two parameters cannot be cleanly disentangled. The tools differ: VIF in regression versus reparameterization or total-derivative SI in sensitivity analysis, but the underlying statistical problem is identical: two predictors that move together cannot both be estimated or ranked in isolation.

**Response 1: Diagnose first.** Before any SA computation, examine the correlation structure of the POIs. Compute the Pearson correlation matrix from any available data on parameter values across studies or populations. Plot scatter matrices (a pairwise scatter matrix). A correlation $|\rho_{ij}| > 0.5$ warrants attention.

**Response 2: Reparameterize structurally correlated pairs.** When two parameters appear only as a product or ratio, define that product or ratio as the natural POI. For the SIR model: use $z = c\beta$ as a single POI. Here $\mathcal{R}_0 = z\tau_R$ depends on $c$ and $\beta$ only through their product, so the composite sensitivity index is $\mathcal{S}_z^{\mathcal{R}_0} = 1$. It is not the sum $\mathcal{S}_c^{\mathcal{R}_0} + \mathcal{S}_\beta^{\mathcal{R}_0} = 2$. The sum measures a different perturbation: scaling $c$ and $\beta$ by the same fractional amount, which changes their product at twice that rate. Reparameterizing replaces two correlated POIs by one; it does not add their indices. (The distinction is developed in the exercises.)

**Response 3: Total derivative for empirically correlated pairs.** When empirical data establishes that a proportional change in $p_i$ induces a proportional change in $p_j$ with coefficient $c_{ij}$ (i.e., $\Delta p_j/p_j \approx c_{ij}\,\Delta p_i/p_i$), the

total derivative SI is:

$$\mathcal{S}^{q}_{p_i}\big|_{\text{tot}} = \mathcal{S}^{q}_{p_i}\big|_{\text{std}} + \sum_{j \neq i} \mathcal{S}^{q}_{p_j}\, c_{ij}. \tag{3.12}$$

In plain language: $c_{ij}$ is the induced proportional change in $p_j$ per unit proportional change in $p_i$. If a 1% increase in $c$ tends to be accompanied by a 0.6% increase in $\beta$ (from field data), then $c_{c\beta} = 0.6$. Estimate $c_{ij}$ from regression of observed parameter pairs, or from domain knowledge about shared drivers. Report both the standard and total-derivative SIs together with the coupling coefficients $c_{ij}$ so readers can assess the impact.

*A caution on interpretation.* The coefficient $c_{ij}$ should represent the *causally induced* change in $p_j$ when $p_i$ is actively perturbed, not merely the observed statistical correlation between them. If the correlation between $c$ and $\beta$ across field sites arises from a shared lurking variable (for example, both are higher in densely-populated urban settings) rather than from a direct biological mechanism, then a policy that reduces $c$ through social distancing may not induce the correlated change in $\beta$. In that case the total-derivative SI overstates the intervention's effect. When the causal structure is uncertain, report both the standard and total-derivative SIs and state the assumption underlying the latter.

*Response 4* (Report transparently). When correlations exist but cannot be fully quantified, state explicitly in the SA report that the independence assumption was made, identify the correlated pairs, and flag which SI values are most likely affected.

*Self-check.* In the SIR model, the product $c\beta$ represents the effective transmission rate. If a field study establishes that $c$ and $\beta$ are positively coupled, with coupling coefficient $c_{c\beta} = 0.4$ (i.e., $\Delta\beta \approx 0.4\,\Delta c$ for proportional changes), use equation (3.12) to compute the total-derivative SI of $\mathcal{R}_0$ with respect to $c$ at the nominal values. Compare with the standard SI $\mathcal{S}^{\mathcal{R}_0}_{c} = 1$.

> *Answer:* $\mathcal{S}^{\mathcal{R}_0}_{c} = 1$, $\mathcal{S}^{\mathcal{R}_0}_{\beta} = 1$. Here $c_{c\beta} = 0.4$ means $(\Delta\beta/\beta)/(\Delta c/c) = 0.4$, so $p_\beta c_{c\beta}/p_c = \beta \cdot 0.4/c$. But in proportional terms: total SI $\approx 1 + 1 \times 0.4 = 1.4$. A 1% increase in $c$ (with the correlated 0.4% increase in $\beta$ it entails) increases $\mathcal{R}_0$ by approximately 1.4%, not 1%. The standard SI underestimates the true effect by 40%.

## 3.6 ▪ Black-Box Models and the Pseudocode Template

When the model is inherited code, a compiled simulation, or a package you did not write, analytical differentiation is not possible. Finite differences are then the only option for local SA, and they work for *any* model that accepts a parameter vector and returns a QOI vector.

Students sometimes treat finite differences as an inferior approximation to be used only when analytic derivatives are unavailable. For black-box models, this framing is backwards: finite differences are not a fallback; they are the correct and complete tool. The accuracy they provide at the optimal step size is typically more than sufficient for SA decisions, and their applicability to models of arbitrary complexity makes them indispensable in practice.

The following pseudocode computes the full normalized sensitivity Jacobian by centered differences, using the optimal step size from (3.9). Students should adapt this template for every black-box SA problem.

```
# From: software/matlab/core/sensitivity_jacobian.m
# (reference library; also reproduced in the companion notebooks)
# Works in MATLAB, Python, R, or any language - adapt syntax as needed

p_nom = [c=5, beta=0.06, tau=7, tau_m=10000]     # nominal parameter values

R0(p)  = p[c] * p[beta] * p[tau]              # model: R0 = c*beta*tau
S      = SENSITIVITY_JACOBIAN(R0, p_nom)      # returns SI vector
# S[j] = SI_{p_j}^{R0};  uses 2*n_p+1 = 9 model evaluations
```

When the QOI vanishes at the nominal point the normalized index is undefined, and the function returns `NaN` for that entry rather than substituting a regularized surrogate. The choice is deliberate: a reference scale $u_{\text{ref}}$ has to come from the problem, and no template can supply one on the reader's behalf. Section 2.2.3 explains how to choose the scale when a regularized index is what you want. The `max` in the step-size formula prevents $h$ from being identically zero when $p_{\text{nom},i} = 0$.

To validate this template on any model, call it with the analytic sensitivity indices known from Chapter 2 and verify agreement. For the SIR model:

```
# Validation: SIR R0 = c*beta*tau - all SIs should equal 1 exactly
p_nom  = [c=5, beta=0.06, tau=7]
R0(p)  = p[c] * p[beta] * p[tau]
# step size, as implemented: h_j = h_scale * max(|p_nom[j]|, h_min/h_sca
h_scale = 6e-6;   h_min = 1e-10
S      = SENSITIVITY_JACOBIAN(R0, p_nom, h_scale, h_min)

PRINT "Sensitivity indices for R0:"
PRINT "  SI_c    =", S[c],    "  (exact: 1.0000)"
PRINT "  SI_beta =", S[beta], "  (exact: 1.0000)"
PRINT "  SI_tau  =", S[tau],  "  (exact: 1.0000)"
```

*Self-check.* The template above uses $h_{\min} = 10^{-10}$ as a floor for the step size. Why is this floor necessary, and what happens if a POI has nominal value exactly zero?

> *Answer:* If $p_{\text{nom},i} = 0$, then $6 \times 10^{-6}|p_{\text{nom},i}| = 0$, giving $h = 0$ and a quotient $0/0$, which MATLAB returns as `NaN` without an error. The floor $h_{\min} = 10^{-10}$ prevents this. For a POI with nominal value zero, the step size is not scaled by the parameter magnitude, it is just the fixed small value $10^{-10}$. The SI itself is then no help. The ratio form would divide by $p_i = 0$, and the derivative form (3.1) multiplies the derivative by $p_i = 0$, so it gives 0 whatever the derivative is. The reference routine `sensitivity_jacobian.m` therefore stops with an error when a nominal value is zero, rather than returning an index. If a POI is exactly zero at its nominal value, the investigator should either reparameterize or report the absolute (unnormalized) derivative rather than the normalized SI.

*Adapting the template to a new model.* To apply `sensitivity_jacobian` to any model, not just SIR, replace the function handle with your own model and update the nominal parameter vector:

```
# Example: pharmacokinetic two-compartment model
# p = [ka, ke, V]  (absorption rate, elimination rate, volume of distrib
# QOI = peak plasma concentration C_max

pk_model(p) = solve_pk(p)       # your function that returns C_max

p_nom = [ka=1.2, ke=0.3, V=5.0]
S = SENSITIVITY_JACOBIAN(pk_model, p_nom)
# S is a 3-element vector of SIs: [SI_ka, SI_ke, SI_V]
```

No other changes are needed: the step-size selection, the U-curve error indicator, and the normalization are all handled automatically for any differentiable model that can be called as a function. The only requirement is that `solve_pk(p)` accepts a parameter vector and returns a numeric scalar or vector of QOIs.

## 3.7 ▪ Computational Laboratory: SA of the SIR Model by Finite Differences

This laboratory puts the chapter's tools together and provides a complete reference implementation for black-box SA. Use the SIR ODE model with the four POIs $\{c, \beta, \tau_R, \nu_m\}$ and the three QOIs $\{S(T), I(T), R(T)\}$ at time $T = 60$ days.

**Setup.** Implement `SIR_model(p)` as a function that takes the parameter vector $p = [c, \beta, \tau_R, \nu_m]$ and returns $[S(60), I(60), R(60)]$ by integrating the SIR ODE. Use nominal values $c = 5$, $\beta = 0.06$, $\tau_R = 7$ days, $\nu_m = 0.0001$ day$^{-1}$, and initial conditions $S_0 = 999$, $I_0 = 1$, $R_0 = 0$.

**Part 1:** Compute the full sensitivity Jacobian. Call `SENSITIVITY_JACOBIAN(SIR_mo p_nom)` to produce the $3 \times 4$ sensitivity matrix. Display it as a labeled table with QOI names as row labels and POI names as column labels.

**Part 2:** Plot the U-curve. Fix all parameters at nominal values except $\beta$. Estimate $\partial I(60)/\partial\beta$ using centered differences for $h = 10^{-3}, 10^{-4}, \ldots, 10^{-13}$, or equivalently for relative steps $h/\beta^*$ spanning the same range. Do not start at $h = 10^{-1}$: with $\beta^* = 0.06$ the left stencil point would be $\beta - h = -0.04$, a negative transmission probability. Compute the absolute error using the fourth-order formula (3.4) as a reference value (computed at $h = 10^{-4}$ where it is accurate). Also plot the one-sided forward difference error at each $h$. Plot centered-difference error, forward-difference error, and the discrepancy between the two (the practical error indicator (3.5)) vs. $h$ on log-log axes. Identify the range of $h$ where the error indicator is small and the centered approximation is reliable.

**Part 3:** Tornado plots. Using the Jacobian from Part 1, produce separate tornado plots for each QOI: the sensitivity of $S(60)$, $I(60)$, and $R(60)$ with respect to all four POIs. Interpret: which POI matters most for each QOI? Do the three plots give the same ranking?

**Part 4:** Correlation check. The contact rate $c$ and transmission probability $\beta$ appear only as the product $c\beta$ in the SIR force of infection. Confirm numerically that $\mathcal{S}_c^{I(60)} \approx \mathcal{S}_\beta^{I(60)}$ at the nominal values. Then reparameterize: define a new POI $\lambda_0 = c\beta$ and compute $\mathcal{S}_{\lambda_0}^{I(60)}$. Since $I(60)$ depends on $c$ and $\beta$ only through their product, the chain rule gives $\mathcal{S}_{\lambda_0}^{I(60)} = \mathcal{S}_c^{I(60)} = \mathcal{S}_\beta^{I(60)}$: all three are the same number, whatever that number turns out to be. Explain why the sum $\mathcal{S}_c^{I(60)} + \mathcal{S}_\beta^{I(60)}$ is then twice it, and why that is not a contradiction: the sum measures a directional perturbation that scales both $c$ and $\beta$ simultaneously, whereas $\mathcal{S}_{\lambda_0}$ treats $\lambda_0$ as a single free parameter. Report the value you compute; it is not 1.

**Part 5:** Correlation check from literature. The SIR force of infection is $\lambda = c\beta I/N$, so $c$ and $\beta$ appear only as the product $c\beta$ and are structurally correlated. Use the four-parameter SIR model; Part 4 has already confirmed that $\mathcal{S}_c^{I(60)} = \mathcal{S}_\beta^{I(60)}$, so start from that. Search the epidemiological literature for any reported correlations between contact rates $c$ and transmission probabilities $\beta$ across outbreak settings. Construct a $4 \times 4$ correlation matrix using whatever estimates you find. Equation (3.12) needs the coupling coefficient $c_{c\beta}$, not the correlation $\rho_{c\beta}$. On the relative scale the regression slope of $\Delta\beta/\beta$ on $\Delta c/c$ is $\rho_{c\beta}$ times the ratio of the coefficient of variation of $\beta$ to that of $c$. It serves as $c_{c\beta}$ only if the change it describes is causal, as the caution in Section 3.5 requires. If you find no estimates, assume a moderate coupling $c_{c\beta} = 0.6$ and zero for all other pairs. Use equation (3.12) to compute the total-derivative SI of $I(60)$ with respect to $c$ for this coupling.

Compare with the standard SI and discuss the practical difference.

**Extension.** Using equation (3.10), compute the pure second-order sensitivities $\partial^2 I(60)/\partial p_i^2$ for each of the four POIs. For each parameter, assess whether the second-order term is large enough to matter for 10% perturbations. (Reuse evaluations from Part 1 to keep the additional cost minimal.)

## 3.8 ▪ Exercises

**Exercise 3.1** *([Bloom L2]).* Explain in your own words why making the step size $h$ smaller does not always improve the accuracy of a finite-difference sensitivity estimate. Describe the two sources of error and explain why they act in opposite directions as $h$ changes.

**Exercise 3.2** *([Bloom L3]).* A model has $n_p = 15$ POIs and $n_q = 4$ QOIs.

(a) Fill in Table 3.1 for this model: how many model evaluations does each method require?

(b) If each evaluation takes 30 seconds, how long does the centered-difference Jacobian take?

(c) At what point (in terms of $n_p$ and evaluation time) would the adjoint method from Chapter 6 be strictly necessary?

**Exercise 3.3** *([Bloom L3]).* For the function $q(p) = \sin(p^2)$ at $p^* = 1.3$, compute $\partial q/\partial p$ using centered differences for $h \in \{10^{-1}, 10^{-2}, \ldots, 10^{-14}\}$. Compare each estimate to the exact value $q'(p) = 2p\cos(p^2)$. Plot error vs. $h$ on log-log axes and identify the optimal $h$. Does it agree with the formula in equation (3.9)?

**Exercise 3.4** *([Bloom L3]).* For the SIR model $\mathcal{R}_0 = c\beta\tau_R$:

(a) Compute the pure second derivatives $\partial^2\mathcal{R}_0/\partial c^2$, $\partial^2\mathcal{R}_0/\partial\beta^2$, $\partial^2\mathcal{R}_0/\partial\tau_R^2$ analytically. What do you notice?

(b) Compute the cross derivatives $\partial^2\mathcal{R}_0/\partial c\,\partial\beta$, $\partial^2\mathcal{R}_0/\partial c\,\partial\tau_R$, $\partial^2\mathcal{R}_0/\partial\beta\,\partial\tau_R$ analytically.

(c) For a 10% perturbation to $c$ and a simultaneous 10% perturbation to $\beta$, estimate the true change in $\mathcal{R}_0$ using both the first-order and the full second-order Taylor approximations. How large is the correction?

Draft

**Exercise 3.5** *([Bloom L4]).* In the SIR model, suppose field data suggests that the contact rate $c$ and transmission probability $\beta$ are positively correlated: across outbreak settings, when $c$ increases by 10%, $\beta$ tends to increase by 6%.

(a) Compute the standard sensitivity indices $\mathcal{S}_c^{\mathcal{R}_0}$ and $\mathcal{S}_\beta^{\mathcal{R}_0}$.

(b) Using equation (3.12), compute the total-derivative SI of $\mathcal{R}_0$ with respect to $c$, accounting for the correlation.

(c) Which is larger: the standard SI or the total-derivative SI? What does this mean for a policy maker who can reduce contact rates (through social distancing) but not transmission probability directly?

**Exercise 3.6** *([Bloom L4]).* A colleague provides you with the function `seir_model(p)` that implements a SEIR epidemic model with $n_p = 6$ parameters and returns the peak infection count as a single QOI. Each model evaluation takes approximately 0.2 seconds.

(a) How long will it take to compute the complete sensitivity Jacobian using centered differences?

(b) Apply the `sensitivity_jacobian` template to this model. What additional information do you need before interpreting the results? (Think about parameterization.)

(c) You find that two of the six parameters have $|\rho_{ij}| = 0.78$. What should you do before reporting the sensitivity indices?

**Exercise 3.7** *([Bloom L5]).* A government agency funds you to perform SA on a dengue fever transmission model with 9 parameters and 3 QOIs (peak infections, total deaths, days until outbreak peak). Each model evaluation takes 4 minutes.

Design the complete computational SA study. Your design should specify:

(a) Which finite-difference method you will use and why.

(b) The total computation time.

(c) How you will choose the step size for each parameter.

(d) What checks you will run before reporting any results.

(e) How you will present the results to a non-technical audience.

(f) One scenario in which the standard SA would give misleading results, and what you would do instead.

**Exercise 3.8** *(*[Bloom L4]*).* *Audit exercise.* A colleague reports finite-difference sensitivity indices for a black-box model, computed with a forward difference at a relative step of $h = 10^{-12}$, and notes that this step was chosen "to make the truncation error negligible." The model is evaluated in double precision.

(a) The stated reasoning is correct as far as it goes. Explain why the conclusion does not follow, using the U-curve of Section 3.3.

(b) Estimate the order of magnitude of the total error at $h = 10^{-12}$ for a well-scaled model, and compare it with the error at the near-optimal step.

(c) The reported indices are finite, smooth in appearance, and internally consistent. Explain why none of those observations detects the problem, and state one diagnostic that would.

(d) Two of the reported parameters are known to be strongly correlated in the calibration data. State what the one-at-a-time indices mean under that condition and what they do not.

**Exercise 3.9** *(*[Bloom L6]*).* A published sensitivity analysis reports: "All finite-difference computations used $h = 10^{-12}$ to ensure maximum precision. The code was run in double-precision floating-point arithmetic."

The authors report that for several parameters, the sensitivity indices are exactly zero to all reported decimal places.

Identify what is most likely wrong with these results and explain your reasoning quantitatively, using the U-curve analysis from Section 3.3. What should the authors have done, and how would you determine whether the zero SI values are genuine or numerical artifacts?

## Notes

The analysis of finite-difference step-size selection, including the U-curve trade-off between truncation error ($O(h^2)$ for centered differences) and rounding error ($O(\varepsilon_{\text{mach}}/h)$), is a classical topic in numerical analysis. A careful

treatment of the rounding-error side appears in Higham [20]; see also the discussion in Saltelli et al. [52], Chapter 2, for the SA context.

The Latin hypercube sampling design used in Chapter 9 was introduced by McKay, Beckman, and Conover [32]. Iman and Conover [24] established the rank-correlation variant that controls correlations among sampled parameters, which is the standard implementation in modern SA software including SALib [19].

Sensitivity analysis applied to the SIR system as a black-box ODE is treated computationally in Marino et al. [31], including finite-difference approximations and their interaction with ODE solver tolerances. The condition-number diagnostics for identifying correlated POIs, and the connection to variance inflation factors in linear regression, are discussed in Saltelli et al. [50].

Draft

# Part III

# Analytic Forward Sensitivity Analysis

Draft

Draft

# Chapter 4
# Analytic Forward Sensitivity Analysis

*Differentiate the model, and the important parameters identify themselves.*

Chapter 3 showed how to compute sensitivity indices for any model by finite differences. For the malaria model with 17 POIs and 7 state variables, the centered-difference Jacobian requires $2 \times 17 + 1 = 35$ model evaluations (one baseline solve plus two perturbed solves per parameter, each giving all 7 state components simultaneously), about 35 minutes if each run takes a minute, or 35 hours if each takes an hour.

There is a faster and more accurate alternative, provided the model can be differentiated analytically: differentiate the governing equations directly. The result is a new system of equations the *forward sensitivity equations*, whose solution is the exact derivative. For an equilibrium output the cost is one linear solve per POI. For the malaria model's endemic equilibrium, that is 17 linear solves of a $7 \times 7$ system after the one run that finds the equilibrium. Each solve takes milliseconds (Section 4.2). For an output that depends on time, each POI adds one linear ODE solve instead (Section 4.4).

The catch is that the model must be differentiable and the analyst must be patient enough to derive the sensitivity equations. This chapter develops that derivation systematically, shows that it always produces a *linear* system no matter how nonlinear the original model is, and ends by quantifying the remaining computational bottleneck that motivates Chapters 5 and 6.

## 4.1 ▪ The FSE Derivation Pattern

The forward sensitivity equation (FSE) for any model is obtained by a single operation: differentiate the governing equation with respect to a POI and apply the chain rule. The result is always a linear equation in the sensitivity variables, regardless of whether the original model is linear or nonlinear.

### 4.1.1 ▪ The Pattern in Three Steps

**Step 1.** Write the governing equation in the form $\mathcal{F}(u;p) = 0$ or $\mathcal{F}(u;p) = f$, where $u$ is the solution and $p$ is the POI of interest.

**Step 2.** Differentiate both sides with respect to $p$. Apply the chain rule to every term that depends on $u$. Collect terms. The result is an equation for $\partial u/\partial p$.

**Step 3.** Recognize that the equation from Step 2 is *linear in* $\partial u/\partial p$, even if $\mathcal{F}$ is nonlinear in $u$.

The linearity in Step 3 is not a coincidence. Differentiation is a linear operation: applying it to a nonlinear equation in $u$ produces a linear equation in $\partial u/\partial p$. This structural fact is what makes analytic SA computationally tractable.

**Misconception to confront.** A common reaction to the FSE is: "Deriving and solving these equations must be at least as hard as solving the original model." This is wrong in two ways. First, the FSE is always *linear* in the sensitivity variable, even when the original model is highly nonlinear; a linear system is categorically easier to solve than a nonlinear one. Second, the FSE for an ODE uses the *same* coefficient matrix $F_u$ as the original model. An implicit (stiff) solver already forms $F_u$ as it steps, and the FSE can reuse that work; an explicit solver never forms it, so you supply $F_u$ yourself. With either kind of solver the additional cost per parameter is one linear ODE solve, not one full nonlinear solve. The FSE is not a second model; it is a linear companion to the first.

### 4.1.2 ▪ Application to Three Model Types

The same three-step pattern applies to algebraic systems, difference equations, and ordinary differential equations. Sections 4.2–4.3 work through each type in detail. Here we display the pattern side by side.

| Model type | Governing equation | Forward sensitivity equation |
|---|---|---|
| Algebraic | $f(u;p) = 0$ | $f_u \frac{\partial u}{\partial p} = -f_p$ |
| Linear system | $A\vec{u} = \vec{b}$ | $A \frac{\partial \vec{u}}{\partial p} = \frac{\partial \vec{b}}{\partial p} - \frac{\partial A}{\partial p}\vec{u}$ |
| ODE IVP | $\dot{u} = F(u;p)$ | $\frac{d}{dt}\left[\frac{\partial u}{\partial p}\right] = F_u \frac{\partial u}{\partial p} + F_p$ |
| OΔE (difference eq.) | $u_{n+1} = G(u_n;p)$ | $\frac{\partial u_{n+1}}{\partial p} = G_u \frac{\partial u_n}{\partial p} + G_p$ |

In each case, the coefficient multiplying the unknown sensitivity ($f_u$, $A$, $F_u$, $G_u$) is the Jacobian of the model with respect to the solution $u$, evaluated at the nominal solution. The sensitivity variable ($\partial u/\partial p$) always appears linearly. In the algebraic, linear-system and O$\Delta$E rows the right-hand side contains only known quantities; in the O$\Delta$E row $\partial u_n/\partial p$ is known from the previous step. In the ODE row the right-hand side contains the unknown itself, multiplied by $F_u$.

*Unifying Idea 1.* Every entry of the sensitivity Jacobian that this chapter computes is a normalized derivative: an instance of Unifying Idea 1 from Chapter 1. The FSE computes those derivatives exactly, without the $O(h^2)$ truncation error of the finite-difference approach. The FSE is not a different kind of SA result; it is the same sensitivity index, obtained by a more accurate and often faster computational route.

*Self-check.* For the logistic ODE $\dot{u} = ru(1 - u/K)$, write down the FSE for $\partial u/\partial r$ and for $\partial u/\partial K$. Identify the coefficient $F_u$ in each case.

> *Answer:* $F(u; r, K) = ru(1 - u/K)$. Computing partial derivatives: $F_u = r(1 - 2u/K)$, $F_r = u(1 - u/K)$, $F_K = ru^2/K^2$.
> FSE for $w_r = \partial u/\partial r$: $\dot{w}_r = r(1 - 2u/K)\, w_r + u(1 - u/K)$.
> FSE for $w_K = \partial u/\partial K$: $\dot{w}_K = r(1 - 2u/K)\, w_K + ru^2/K^2$.
> Both FSEs have the same coefficient $F_u = r(1-2u/K)$ multiplying the unknown sensitivity; only the right-hand-side forcing term differs. The coefficient depends on the forward solution $u(t)$, so the FSE must be solved simultaneously with the original ODE.

## 4.2 ▪ Algebraic Systems and Equilibria

For static problems, equations whose solutions do not change in time, the FSE is a linear system that can be solved by standard linear algebra.

### 4.2.1 ▪ A Single Nonlinear Equation

For the scalar equation $f(u; p) = 0$, differentiating with respect to $p$ gives

$$f_u \frac{\partial u}{\partial p} + f_p = 0,$$

so $\partial u/\partial p = -f_p/f_u$ provided $f_u \neq 0$. This is exactly the implicit function theorem. When $f_u = 0$, the derivative is undefined, which corresponds to a bifurcation point, a situation studied in Chapter 8.

### 4.2.2 ▪ Linear Systems: The FSE for $A\vec{u} = \vec{b}$

For the system $A\vec{u} = \vec{b}$, differentiating with respect to any scalar parameter $q \in \{a_{ij}, b_i\}$ gives:

$$A\,\frac{\partial \vec{u}}{\partial q} = \frac{\partial \vec{b}}{\partial q} - \frac{\partial A}{\partial q}\,\vec{u}. \tag{4.1}$$

The right-hand side is a known vector once the forward solution $\vec{u}$ is known. Solving equation (4.1) requires one additional linear solve with the same coefficient matrix $A$ for each parameter $q$. Since the LU factorization of $A$ is computed only once, the cost for $m$ parameters is one LU factorization plus $m$ back-substitutions.

**Important:** Do not solve (4.1) by computing $A^{-1}$. Explicit matrix inversion is numerically unstable and computationally wasteful. In MATLAB, use the backslash operator (`A\b`), which solves the system by LU factorization without forming the inverse.

### 4.2.3 ▪ The Competitive Lotka–Volterra Model

Consider the two-species competitive Lotka–Volterra model [35]:

$$\frac{du_1}{dt} = u_1(1 - u_1 - au_2), \tag{4.2}$$

$$\frac{du_2}{dt} = bu_2(1 - u_2 - cu_1), \tag{4.3}$$

with parameters $a$, $b$, $c > 0$ satisfying $0 < a, c < 1$. The negative cross-terms $-au_2$ and $-cu_1$ represent interspecific competition: each species inhibits the other's growth. The condition $0 < a, c < 1$ ensures that neither species is a sufficiently strong competitor to exclude the other, so a stable coexistence equilibrium exists. The coexistence equilibrium $(\bar{u}_1, \bar{u}_2)$ satisfies the nonlinear system

$$\bar{u}_1 = \frac{1-a}{1-ac}, \qquad \bar{u}_2 = \frac{1-c}{1-ac}. \tag{4.4}$$

To find how this equilibrium depends on $a$ and $c$, we differentiate the equilibrium equations

$$f_1(\bar{u}_1, \bar{u}_2; a, c) := \bar{u}_1(1 - \bar{u}_1 - a\bar{u}_2) = 0,$$
$$f_2(\bar{u}_1, \bar{u}_2; a, c) := b\bar{u}_2(1 - \bar{u}_2 - c\bar{u}_1) = 0,$$

with respect to $a$ to obtain the FSE

$$D_{\vec{u}}[\vec{f}]\,\frac{\partial \vec{u}}{\partial a} = -\frac{\partial \vec{f}}{\partial a}, \tag{4.5}$$

where $D_{\vec{u}}[\vec{f}]$ is the Jacobian matrix of $\vec{f}$ with respect to $\vec{u}$, evaluated at the equilibrium:

$$D_{\vec{u}}[\vec{f}]\big|_{\text{eq}} = \begin{pmatrix} 1 - 2\bar{u}_1 - a\bar{u}_2 & -a\bar{u}_1 \\ -bc\bar{u}_2 & b(1 - 2\bar{u}_2 - c\bar{u}_1) \end{pmatrix}, \tag{4.6}$$

and

$$\frac{\partial \vec{f}}{\partial a} = \begin{pmatrix} -\bar{u}_1\bar{u}_2 \\ 0 \end{pmatrix}.$$

Solving the $2 \times 2$ linear system (4.5) gives $\partial\bar{u}_1/\partial a$ and $\partial\bar{u}_2/\partial a$ exactly. The sensitivity indices follow from Definition 2.1.

A key observation: although the equilibrium equations (4.2)–(4.3) are nonlinear, the FSE (4.5) is a *linear system* in the sensitivity variables. This is always true for equilibrium problems, as the following result shows.

**Theorem 4.1** (Linearity of the FSE)**.** *Let $\vec{f}(\vec{u};\vec{p}) = \vec{0}$ be a system of nonlinear equations with $\vec{f}$ differentiable in $\vec{u}$ and $\vec{p}$. The forward sensitivity equation for $\partial\vec{u}/\partial p_j$ is the linear system*

$$D_{\vec{u}}[\vec{f}]\,\frac{\partial \vec{u}}{\partial p_j} = -\frac{\partial \vec{f}}{\partial p_j},$$

*provided the Jacobian $D_{\vec{u}}[\vec{f}]$ is nonsingular.*

*Proof.* Differentiating $\vec{f}(\vec{u}(\vec{p});\vec{p}) = \vec{0}$ with respect to $p_j$ and applying the chain rule gives

$$D_{\vec{u}}[\vec{f}]\,\frac{\partial \vec{u}}{\partial p_j} + \frac{\partial \vec{f}}{\partial p_j} = \vec{0},$$

where $D_{\vec{u}}[\vec{f}]$ denotes the Jacobian matrix of $\vec{f}$ with respect to $\vec{u}$, evaluated at the nominal solution $(\vec{u}(\vec{p});\vec{p})$. Rearranging yields

$$D_{\vec{u}}[\vec{f}]\,\frac{\partial \vec{u}}{\partial p_j} = -\frac{\partial \vec{f}}{\partial p_j}.$$

The right-hand side $-\partial\vec{f}/\partial p_j$ is a known vector once $\vec{u}(\vec{p})$ is available. Linearity in $\partial\vec{u}/\partial p_j$ follows from the fact that differentiation is itself a linear operation: the quantity $\partial\vec{u}/\partial p_j$ appears in the left-hand side only through the matrix-vector product $D_{\vec{u}}[\vec{f}]\,(\partial\vec{u}/\partial p_j)$. No powers, products, or compositions of $\partial\vec{u}/\partial p_j$ appear, regardless of the nonlinearity of $\vec{f}$ in $\vec{u}$. Nonsingularity of $D_{\vec{u}}[\vec{f}]$ guarantees a unique solution. □

This theorem is one of the most practically useful facts in SA. Regardless of how complex or nonlinear the model is, its FSE at an equilibrium is always a linear system, and linear systems can be solved efficiently.

*Self-check.* For the competitive Lotka–Volterra model with $a = 0.4$, $b = 1$, $c = 0.3$: (a) Compute the coexistence equilibrium using equation (4.4). (b) Assemble the Jacobian (4.6) at this equilibrium. (c) Solve the FSE for $\partial\bar{u}_1/\partial a$ and $\partial\bar{u}_2/\partial a$. (d) Compare with the analytic derivatives of (4.4).

*Answer:* (a) $\bar{u}_1 = (1-0.4)/(1-0.12) = 0.6/0.88 \approx 0.6818$; $\bar{u}_2 = (1-0.3)/0.88 = 0.7/0.88 \approx 0.7955$.

(b) $D_u f = \begin{pmatrix} 1 - 2(0.6818) - 0.4(0.7955) & -0.4(0.6818) \\ -1(0.3)(0.7955) & 1 - 2(0.7955) - 0.3(0.6818) \end{pmatrix} \approx \begin{pmatrix} -0.6818 & -0.2727 \\ -0.2386 & -0.7955 \end{pmatrix}$.

(c) Since $f_1 = u_1(1 - u_1 - au_2)$ gives $\partial f_1/\partial a = -u_1u_2$ and $f_2$ does not involve $a$, the right-hand side is $-\partial\vec{f}/\partial a = (0.6818 \times 0.7955,\, 0)^T \approx (0.5424, 0)^T$. Solving the $2 \times 2$ system $D_u f\, \partial\vec{\bar{u}}/\partial a = -\partial\vec{f}/\partial a$ without rounding:

$$\frac{\partial\bar{u}_1}{\partial a} = -0.9039256198, \qquad \frac{\partial\bar{u}_2}{\partial a} = +0.2711776860.$$

(d) Differentiating (4.4) directly,

$$\frac{\partial\bar{u}_1}{\partial a} = -\frac{1-c}{(1-ac)^2} = -\frac{0.7}{0.88^2} = -0.9039256198, \qquad \frac{\partial\bar{u}_2}{\partial a} = \frac{c(1-c)}{(1-ac)^2} = \frac{0.21}{0.88^2} = +0.2711776860.$$

The two agree to machine precision, as they must: the FSE is the derivative of the equilibrium condition, so it cannot disagree with the derivative of the equilibrium. Carrying the arithmetic unrounded is what makes the agreement visible; rounding to three figures would hide whether the match is exact or merely close.

### 4.2.4 ▪ The Car Rental Problem: O△E Sensitivity

Consider the car rental Markov chain: two airports $A$ and $B$ with transition probabilities $\alpha = \Pr(A \to A)$ and $\beta = \Pr(B \to B)$. The discrete-time model is

$$A_{n+1} = \alpha A_n + (1-\beta)B_n, \tag{4.7}$$

$$B_{n+1} = (1-\alpha)A_n + \beta B_n. \tag{4.8}$$

The fixed point $(A^*, B^*)$ gives the long-run fractions of cars at each airport, so it is normalized by $A^* + B^* = 1$:

$$A^* = \frac{1-\beta}{2-\alpha-\beta}, \qquad B^* = \frac{1-\alpha}{2-\alpha-\beta}.$$

Write $M$ for the transition matrix of (4.7)–(4.8). The fixed point solves $(M-I)\vec{x} = \vec{0}$ with $\vec{x} = (A^*, B^*)$, and $M-I$ is singular because each column

of $M$ sums to one. Differentiating gives $(M-I)\,\partial\vec{x}/\partial\alpha = -(\partial M/\partial\alpha)\,\vec{x}$, and the normalization adds $\partial A^*/\partial\alpha + \partial B^*/\partial\alpha = 0$. Replacing one row of $M-I$ by this condition gives a system of the form (4.1) with a nonsingular matrix in the role of $A$. The result is:

$$\mathcal{S}_\alpha^{A^*} = \frac{\alpha}{2-\alpha-\beta}, \qquad \mathcal{S}_\alpha^{B^*} = -\frac{\alpha(1-\beta)}{(1-\alpha)(2-\alpha-\beta)}. \tag{4.9}$$

These indices tell management which transition probability has the greater effect on the long-run car distribution. At nominal values $\alpha = 0.7$ and $\beta = 0.3$, the formula gives $\mathcal{S}_\alpha^{A^*} = 0.7/(2-0.7-0.3) = 0.70$: a 1% increase in the probability of keeping a car at airport $A$ produces a 0.70% increase in the long-run fraction of cars at $A$.

## 4.3 ▪ Forward Sensitivity for ODE Systems

For a dynamical model governed by an ODE, the sensitivity $\partial u/\partial p$ changes over time as the solution evolves. The FSE is itself an ODE that must be solved alongside the original model.

### 4.3.1 ▪ Derivation of the ODE FSE

Consider the scalar, autonomous ODE with parameter $p$:

$$\frac{du}{dt} = F(u;p), \qquad u(0) = u_0(p). \tag{4.10}$$

Define the sensitivity variable $w(t) = \partial u(t)/\partial p$. Differentiating (4.10) with respect to $p$ and exchanging the order of $\partial/\partial t$ and $\partial/\partial p$:

$$\frac{d}{dt}\left[\frac{\partial u}{\partial p}\right] = \frac{\partial F}{\partial u}\,\frac{\partial u}{\partial p} + \frac{\partial F}{\partial p}. \tag{4.11}$$

In terms of $w$:

$$\dot{w} = F_u(u;p)\,w + F_p(u;p), \qquad w(0) = \frac{du_0}{dp}. \tag{4.12}$$

The coefficient $F_u(u(t);p)$ depends on the forward solution $u(t)$, so the FSE (4.12) must be solved simultaneously with the original ODE (4.10). Together they form an *augmented system* of two ODEs.

For the vector ODE $\dot{\vec{u}} = \vec{F}(\vec{u};\vec{p})$ with $n$ state variables and $m$ parameters:

$$\frac{d}{dt}\left[\frac{\partial \vec{u}}{\partial p_j}\right] = J(\vec{u};\vec{p})\,\frac{\partial \vec{u}}{\partial p_j} + \frac{\partial \vec{F}}{\partial p_j}, \tag{4.13}$$

where $J = \partial\vec{F}/\partial\vec{u}$ is the $n \times n$ Jacobian of $\vec{F}$ with respect to $\vec{u}$. One equation (4.13) is needed per parameter $p_j$, each an $n$-dimensional ODE system. For $m$ parameters: the augmented system has $n + mn = n(1 + m)$ equations.

Theorem 4.1 extends to the dynamical case: each FSE is *linear in the sensitivity variable* $\partial\vec{u}/\partial p_j$, even though $\vec{F}$ is nonlinear in $\vec{u}$. The coefficient $J(\vec{u}(t); \vec{p})$ is time-varying, but the structure is linear.

*Self-check.* Write down the augmented ODE system (original ODE + FSE) for the logistic equation $\dot{u} = ru(1-u/K)$ with respect to both parameters $r$ and $K$ simultaneously. How many total ODEs does the augmented system have?

*Answer:* The augmented system consists of: Original: $\dot{u} = ru(1 - u/K)$. FSE for $w_r = \partial u/\partial r$: $\dot{w}_r = r(1 - 2u/K)w_r + u(1 - u/K)$. FSE for $w_K = \partial u/\partial K$: $\dot{w}_K = r(1 - 2u/K)w_K + ru^2/K^2$.
Total: 3 ODEs. In general, for $n$ state variables and $m$ parameters, the augmented system has $n + mn = n(1 + m)$ equations.

### 4.3.2 ▪ Time-Dependent Sensitivity Indices

For dynamical models, the sensitivity index is time-dependent:

$$\mathcal{S}_p^u(t) = \frac{p}{u(t)}\, w(t) = \frac{p}{u(t)}\, \frac{\partial u(t)}{\partial p}. \tag{4.14}$$

The relative importance of POIs can change over time. A parameter that strongly influences the early phase of an epidemic may become less important as the system approaches equilibrium.

### 4.3.3 ▪ The SIR Model: Full FSE Treatment

Chapter 2 introduced the SIR model without vital dynamics (Equations (2.17)–(2.19)). For the FSE treatment we extend it to include demography (births and deaths), adding a fourth POI $\tau_m = 1/\nu_m$ (mean residence time). This extension does not change the analysis for $c$, $\beta$, and $\tau_R$; it adds the sensitivity with respect to $\tau_m$ and illustrates how the FSE scales to larger parameter sets.

The SIR model with demography is:

$$\frac{dS}{dt} = \nu_m N - \frac{c\beta SI}{N} - \nu_m S, \tag{4.15}$$

$$\frac{dI}{dt} = \frac{c\beta SI}{N} - (\gamma_R + \nu_m)I, \tag{4.16}$$

$$\frac{dR}{dt} = \gamma_R I - \nu_m R. \tag{4.17}$$

With interpretable POIs $\vec{p} = (c, \beta, \tau_R, \tau_m)$ where $\tau_R = 1/\gamma_R$ (mean infectious period, days) and $\tau_m = 1/\nu_m$ (mean residence time, days), the augmented system for a single POI $p_j$ adds 3 sensitivity ODEs:

$$\frac{dw_{Sj}}{dt} = J_{11}w_{Sj} + J_{12}w_{Ij} + J_{13}w_{Rj} + \frac{\partial F_S}{\partial p_j}, \tag{4.18}$$

$$\frac{dw_{Ij}}{dt} = J_{21}w_{Sj} + J_{22}w_{Ij} + J_{23}w_{Rj} + \frac{\partial F_I}{\partial p_j}, \tag{4.19}$$

$$\frac{dw_{Rj}}{dt} = J_{31}w_{Sj} + J_{32}w_{Ij} + J_{33}w_{Rj} + \frac{\partial F_R}{\partial p_j}, \tag{4.20}$$

where $J_{ij}$ are entries of the Jacobian of $(F_S, F_I, F_R)$ with respect to $(S, I, R)$. For $m = 4$ POIs, the full augmented system has $3 + 4 \times 3 = 15$ ODEs.

The initial conditions for the FSE are zero: $w_{Sj}(0) = w_{Ij}(0) = w_{Rj}(0) = 0$ (assuming the initial conditions do not depend on the parameters).

Figure 4.1 shows the time-dependent SI for $I(t)$ with respect to $c$, $\beta$, $\tau_R$, and $\tau_m$. Through the growth phase the contact rate $c$ and transmission probability $\beta$ dominate, and they peak near day 29 at 6.799. The mean infectious period $\tau_R$ overtakes them only in a narrow window around the epidemic peak, days 44 to 51, when what remains to be settled is the timing of the turnover rather than its size. After day 51 transmission dominates again, and by day 90 its index has grown to $-8.6$ against $-0.7$ for $\tau_R$: a larger contact rate burns through the susceptible pool sooner, so it leaves fewer infectious people late, and the index that was strongly positive during growth becomes strongly negative during decline. Two lessons follow, and neither is visible from a single number. The ranking changes, and it changes twice rather than once. The sign changes too, at day 48 for $c$ and $\beta$: the same parameter that drives the epidemic up is the one that has already exhausted it.

## 4.4 ▪ The Proliferation Problem and Its Consequences

> *Before improving the model, find out which knob actually moves the machine.*

The FSE is exact and efficient for a single parameter. For $m$ parameters the cost scales linearly, but there are two ways to arrange it and they are not equally cheap. The *simultaneous* form integrates one system of dimension $n(1 + m)$, carrying every sensitivity block alongside the state; that is what §4.3 constructs. The *sequential* form runs $m$ separate solves of dimension $2n$, one per parameter. Both give identical sensitivities, but the simultaneous





**Figure 4.1.** *Computed time-dependent sensitivity indices $\mathcal{S}_p^I(t)$ for the SIR model at the nominal parameters of this chapter ($c = 5$, $\beta = 0.06$, $\tau_R = 7$, $\tau_m = 10^4$, $N = 1000$). The indices for $c$ and $\beta$ are equal to machine precision, differing by about $10^{-15}$, and are drawn as a single curve: the two* POIs *enter only through the product $c\beta$, so no computation can separate them. Both peak at 6.799 near day 29 and cross zero at day 48. The mean infectious period $\tau_R$ peaks later, at 3.513 near day 31, and crosses zero at day 62. It leads only in the shaded window, days 44 to 51, around the epidemic peak; before and after it, transmission dominates. The mean residence time $\tau_m$ is three orders of magnitude smaller and so is given its own panel with its own zero line; it changes sign at day 46 rather than staying negative throughout. Generated by* `generate_fig_sir_timevarySI.py`*, which reproduces these values byte for byte.*

form is cheaper because the state trajectory and its step-size selection are computed once rather than $m$ times. For the four-parameter SIR model of this chapter the simultaneous form needed about 1900 right-hand-side evaluations against about 6700 for the sequential, a factor of roughly 3.5. Table 4.1 quantifies this for representative models.

*Counting convention.* Throughout this section, "solve counts" refer to the total number of ODE solves, including the one baseline forward solve (of the original $n$-equation system) required before any sensitivity computation begins. In the sequential arrangement the FSE adds $m$ augmented solves on top of this baseline, for a total of $m+1$; the simultaneous arrangement adds one solve of a larger system instead. The counts below use the sequential convention, which is the conservative one.

Figure 4.2 shows what those solves buy: the normalized indices for the SIR model vary over the epidemic, so a single end-of-run number would hide the ranking changing with time. For the SIR model, 5 solves is extremely

| Model | $n$ states | $m$ POIs | Augmented size | FSE cost |
|---|---|---|---|---|
| SIR | 3 | 4 | $3 + 12 = 15$ | 5 ODE solves |
| SEIR (adds Exposed compartment) | 4 | 6 | $4 + 24 = 28$ | 7 ODE solves |
| Malaria model | 7 | 17 | $7 + 119 = 126$ | 18 ODE solves |
| Climate model | 500 | 200 | $500 + 100{,}000$ | 201 ODE solves |

**Table 4.1.** *Augmented system size and FSE cost for representative models. The FSE cost column follows the sequential $m + 1$ convention: $m$ augmented solves plus the one baseline forward solve. The costs are for an output that depends on time; an equilibrium output needs one model run and then one linear solve per* POI *(Section 4.2). A simultaneous implementation replaces these by a single solve of dimension $n(1+m)$ and is typically several times cheaper. For the climate model, 201 solves at 6 hours each is just over 50 days of computation. This is the cost that motivates the adjoint method in Chapter 6.*

fast in practice. For the climate model, 201 solves at 6 hours each is just over 50 days. This is the computational wall that motivates the adjoint approach.

*Unifying Idea 2.* Table 4.1 reveals the bottleneck that Chapters 5 and 6 resolve. No matter how the sensitivity Jacobian is computed, by finite differences, by analytic FSE, or by the adjoint; the answer is the same matrix of normalized derivatives. The adjoint (Unifying Idea 2, introduced in Chapter 5) reduces the cost from $m + 1$ solves to 2 solves for one QOI by exploiting the transposed structure of the problem. The cost reduction is proportional to $m$; for $m = 200$, the adjoint is 100 times faster than the FSE.

*Unifying Idea 3.* Every row of Table 4.1 is a linearization. The FSE computes the exact derivative of a nonlinear model, which is a linear approximation to the model's response. That approximation is valid when parameters are near their nominal values. The proliferation problem scales with $m$; the validity limitation is independent of $m$. Chapter 8 addresses the second problem; Chapters 5 and 6 address the first.

*Self-check.* For the malaria model with $n = 7$ state variables and $m = 17$ POIs, and one response functional (say, the infected-host fraction $i_h(T)$ at a final time $T$): (a) How many ODE solves does the FSE approach require? (b) If each ODE solve takes 2 minutes, how long does the complete FSA take? (c) How many ODE solves will the adjoint method in Chapter 6 require?

> *Answer:* (a) 17 FSE solves (one per parameter), plus 1 forward solve = 18 total. (b) $18 \times 2 = 36$ minutes. (c) For one QOI the adjoint requires exactly 2 solves (1 forward + 1 adjoint), regardless of $m$. For 17 parameters, the adjoint is 9 times faster than the FSE.





**Figure 4.2.** *Forward sensitivity analysis of the SIR model via the FSE. Top: compartment trajectories at $\mathcal{R}_0 = 5$. Bottom: time-dependent normalized sensitivity indices $S^I_\beta(t)$, $S^I_{\tau_R}(t)$, and $S^I_{\gamma_R}(t)$. Before the epidemic peak, transmission drives $I(t)$ upward ($S^I_\beta, S^I_{\tau_R} > 0$); after the peak, recovery dominates, and the plotted index $S^I_{\tau_R}$ keeps growing, reaching $+5.21$ at $t = 60$ while $S^I_\beta$ has turned negative. Stated in the recovery rate the same fact reads $S^I_{\gamma_R} = -5.21 < 0$ with magnitude increasing, since $\gamma_R = 1/\tau_R$ flips the sign (§2.5); the figure plots the duration, not the rate. The sensitivity indices are not constant in time. The figure uses $\mathcal{R}_0 = 5$ rather than the chapter's nominal $\mathcal{R}_0 = c\beta\tau_R = 2.1$ so that the post-peak regime is legible: at the nominal value the peak falls near day 44, as the Chapter 6 laboratory shows, leaving too little of a 60-day window after it to see the indices exchange rank. The two values do not behave alike for $\tau_R$. At the nominal value $S^I_{\tau_R}$ falls after the peak and turns negative near day 62 (Figure 4.1). At $\mathcal{R}_0 = 5$ it stays positive and keeps growing.*

## 4.5 ▪ Choosing a Method

With three computational methods now available (finite differences, analytic FSE, and the adjoint previewed above), the right choice depends on the model and the problem. Chapter 7 collects the full comparison once automatic differentiation is also in hand; Chapter 9 extends it to the global methods.

For this course, the default progression is:

1. Start with finite differences (Chapter 3) to get SA results quickly.
2. Switch to analytic FSE (this chapter) when the model is differentiable and accuracy matters.
3. Use the adjoint (Chapters 5–6) when $m$ is large.
4. Check for correlated POIs before reporting any results (Chapter 3).
5. Consider global SA (Chapter 9) when parameter uncertainty is large.

A common assumption is that finite differences are adequate for any SA study once the step size is chosen correctly. For black-box models, this is true: FD is the only option and produces reliable results at the optimal step size. For differentiable models, the analytic FSE is simultaneously more accurate (no $O(h^2)$ truncation error) and often faster (no repeated model evaluations at perturbed parameter values). The two methods give the same numbers when FD is implemented correctly; the FSE gives those same numbers without truncation error and without the step-size selection problem. For an ODE model its computed values still carry the solver's error, set by `RelTol` and `AbsTol`, so set them tightly, as for finite differences (`RelTol = AbsTol =` $10^{-8}$, Section 3.3).

## 4.6 ▪ Computational Laboratory: Analytic FSA of the SIR Model

This laboratory implements the augmented ODE system for the SIR model and compares the analytic FSE approach with the finite-difference method from Chapter 3.

**Setup.** Use the SIR model (4.15)–(4.17) with POIs $\vec{p} = (c, \beta, \tau_R, \tau_m)$ (interpretable parameterization), nominal values $c = 5$, $\beta = 0.06$, $\tau_R =$ 7 days, $\tau_m =$ 10,000 days, and initial conditions $S_0 = 999$, $I_0 = 1$, $R_0 = 0$, $N = 1000$.

**Part 1:** Implement the augmented ODE system. Implement a function `sir_augmented(t, y, c, beta, tau_R, tau_m)` that returns the derivatives for the full 15-equation augmented system: 3 original state equations plus $4 \times 3 = 12$ FSE equations. The FSE equations require the Jacobian of $(F_S, F_I, F_R)$ with respect to $(S, I, R)$.

*State vector layout.* The 15-component state vector `y` is organized as follows:

| Index | Meaning |
|---|---|
| 1 | $S(t)$ |
| 2 | $I(t)$ |
| 3 | $R(t)$ |
| 4–6 | $(\partial S/\partial c,\ \partial I/\partial c,\ \partial R/\partial c)$ |
| 7–9 | $(\partial S/\partial \beta,\ \partial I/\partial \beta,\ \partial R/\partial \beta)$ |
| 10–12 | $(\partial S/\partial \tau_R,\ \partial I/\partial \tau_R,\ \partial R/\partial \tau_R)$ |
| 13–15 | $(\partial S/\partial \tau_m,\ \partial I/\partial \tau_m,\ \partial R/\partial \tau_m)$ |

The initial condition is `y0 = [S0; I0; 0; zeros(12,1)]`. The sensitivity variables all start at zero because the initial conditions do not depend on the parameters. The reference library provides `software/matlab/fse/sir_augmented.m` for this system, to check against after completing your own implementation.

**Part 2:** Solve and extract sensitivities. Solve the augmented system from $t = 0$ to $t = 90$ days using an ODE solver. Extract the time-dependent sensitivity indices $\mathcal{S}_c^I(t)$, $\mathcal{S}_\beta^I(t)$, $\mathcal{S}_{\tau_R}^I(t)$, $\mathcal{S}_{\tau_m}^I(t)$.

**Part 3:** Plot time-dependent SIs. Plot all four $\mathcal{S}_{p_j}^I(t)$ curves on a single axis, alongside the epidemic curve $I(t)$. Identify: (a) at $t = 10$ days, which parameter matters most? (b) at $t = 50$ days, has the ranking changed? (c) which parameter has the largest absolute SI at the epidemic peak?

**Part 4:** Compare with finite differences. At $t = 30$ days and $t = 70$ days, compare the FSE-based SI values against the centered-difference estimates from Chapter 3. Are they equal to 4 significant figures? Report any discrepancies and explain their likely source.

**Part 5:** Cost comparison. Time the two approaches: the augmented ODE (all 4 sensitivities at once) versus 4 separate centered-difference calls to `SIR_model`. Which is faster? Which is more accurate? At what value of $m$ (number of POIs) would you expect the two approaches to take approximately equal time?

## 4.7 ▪ Exercises

**Exercise 4.1** *([Bloom L2]).* State the three-step FSE derivation pattern in your own words, without using any symbols. Then explain, in one sentence, why the FSE is always linear in the sensitivity variable even when the original equation is nonlinear.

**Exercise 4.2** *([Bloom L3]).* For the nonlinear equation $g(u; p) := u^3 - 3pu + 2 = 0$:

(a) Write the FSE for $\partial u/\partial p$.

(b) At $p = 1$, find all real roots $u$ numerically.

(c) Compute $\mathcal{S}_p^u$ at each root.

(d) At $p = 1$, for which root is the sensitivity largest in magnitude? Verify your answer at the simple root near $u = -2$ using a centered finite difference with optimal $h$.

(e) Explain why the same centered difference cannot be used at the positive root. For $p < 1$ the two positive roots are complex, so no nearby real positive branch exists on the left of $p = 1$. Show instead, by one-sided continuation for $p > 1$, that the two positive roots satisfy $u_\pm \approx 1 \pm \sqrt{p-1}$, and explain why this square-root scaling makes $\partial u/\partial p$ unbounded as $p \downarrow 1$.

**Exercise 4.3** *([Bloom L3]).* For the logistic ODE $\dot{u} = ru(1-u/K)$, $u(0) = u_0$:

(a) Write the FSE for $w_r = \partial u/\partial r$.

(b) Verify that the FSE is linear in $w_r$ by identifying the coefficient (which depends on $u(t)$) and the forcing term.

(c) Implement the augmented system $(u, w_r, w_K)$ and solve from $t = 0$ to $t = 20$.

(d) Plot $\mathcal{S}_r^u(t)$ and $\mathcal{S}_K^u(t)$. Interpret: near $t = 0$ vs. near the carrying capacity.

**Exercise 4.4** *([Bloom L3]).* For the competitive Lotka–Volterra model (4.2)–(4.3) with $a = 0.5$, $b = 2$, $c = 0.4$:

(a) Compute the coexistence equilibrium.

(b) Assemble and solve the FSE for $\partial\vec{u}/\partial c$.

(c) Compute the normalized sensitivity indices $\mathcal{S}_c^{\bar{u}_1}$ and $\mathcal{S}_c^{\bar{u}_2}$.

(d) Using equation (4.4), compute the same indices analytically. Verify agreement.

**Exercise 4.5** *([Bloom L4]).* The SEIRS model (Susceptible-Exposed-Infectious-Recovered-Susceptible) has 4 state variables and 7 parameters.

(a) How many ODEs are in the augmented FSE system for all 7 parameters?

(b) If each ODE solve takes 3 minutes, how long does the full FSA take using the FSE approach? The adjoint approach (Chapter 6)?

(c) Suppose a modeler decides to reduce computation time by computing sensitivities only for the 3 parameters they expect to be most important, based on biological intuition. Identify one scenario where this strategy would lead to a misleading conclusion.

**Exercise 4.6** *([Bloom L4]).* Using your MATLAB implementation from the laboratory:

(a) At three time points, $t = 5$ (early epidemic), $t = t_{\text{peak}}$ (peak infections), and $t = 80$ (late); report the tornado plot of SIs for $I(t)$.

(b) Does the parameter ranking change across the three time points?

(c) A public health officer says: "We should target the parameter with the largest SI at the epidemic peak, since that is when the most people are affected." Critique this statement using your results.

**Exercise 4.7** *([Bloom L5])*. A colleague is modeling cholera transmission with a SEIR-type model having 8 parameters and 4 state variables. She asks for advice on the SA methodology. Design a complete FSA study, specifying:

(a) Whether to use finite differences or analytic FSE, and why.

(b) How you would validate the implementation.

(c) What QOIs you would compute sensitivity indices for, and at what time points.

(d) How you would present the results to identify the two most critical intervention targets.

(e) One situation in which the FSE results might be misleading, and what additional analysis you would recommend.

**Exercise 4.8** *([Bloom L6])*. A published paper reports: "We performed sensitivity analysis of the SEIR model using the forward sensitivity equations. All sensitivity indices were computed at $t = 365$ days, giving a single set of values for the entire epidemic."

The model simulates a disease whose outbreak dynamics play out over approximately 60 days, but the model runs for a year.

Identify at least two methodological issues with this approach. For each issue, state what the authors should have done instead and explain what information they likely missed.

## Notes

The forward sensitivity equation framework, differentiating the governing equation with respect to parameters to obtain a linear system for the sensitivities, is the continuous analog of forward-mode algorithmic differentiation (Chapter 7). Early systematic treatments appear in the Russian literature on sensitivity theory; an accessible modern account is given by Cacuci [10].

The SIR model with demography, adding birth and death rates $\nu_m$ to the basic Kermack–McKendrick model, is treated in detail by Anderson and

May [1], Chapter 3. The endemic equilibrium analysis and its dependence on $\mathcal{R}_0$ are developed by van den Driessche and Watmough [63]. The SEIR extension, adding an exposed compartment with mean latent period $\tau_E$, is analyzed in Murray [35].

The numerical solution of the augmented FSE system, the original ODE coupled with $m$ sensitivity ODEs, is discussed in detail by Cao et al. [12], who also describe its implementation in the CVODES solver. The proliferation problem (cost growing linearly in $m$) that motivates the adjoint approach of Chapters 5 and 6 is quantified precisely by comparing FSE cost with adjoint cost in Cacuci [9].

Draft

Draft

# Part IV

# The Adjoint Approach

Draft

Draft

# Chapter 5

# The Adjoint in Linear Algebra

*The adjoint is the same sensitivity question, asked from the other end.*

Chapter 4 showed that analytic forward sensitivity analysis for a model with $m = 50$ parameters requires 50 linear solves, one per parameter, after the nominal solve for $\vec{u}$: 51 in all. Each solve uses the same coefficient matrix $A$, so we pay once for the LU factorization and 51 times for back-substitution. That is still 51 operations.

Now suppose we care about a single response functional, not the entire solution $\vec{u}$. The quantity of interest is $J = \vec{c}^T\vec{u}$: the pollution level at one monitoring station, the total disease burden, the profit at equilibrium. We want $\partial J/\partial q$ for all 50 parameters $q$.

Is there a way to compute all 50 derivatives simultaneously, in the cost of the nominal solve and a *single* further linear solve?

Yes. The answer is the adjoint. The adjoint approach to sensitivity analysis was developed for nonlinear systems by Cacuci [9] and has since become a standard tool in optimal design and computational science [16].

## 5.1 ▪ A Preview: The Same Answer, Two Solves Instead of Fifty-One

Before the general theory, work through a small example by hand to see exactly what the adjoint does and why it is efficient.

### The Setup

Consider the $2 \times 2$ linear system

$$A\vec{u} = \vec{b}(q_1, q_2), \qquad A = \begin{pmatrix} 3 & 1 \\ 1 & 2 \end{pmatrix}, \qquad \vec{b} = \begin{pmatrix} q_1 \\ q_2 \end{pmatrix}, \tag{5.1}$$

where $q_1$ and $q_2$ are two parameters of interest. The scalar response functional is $J = u_1 + u_2 = \vec{c}^T\vec{u}$ with $\vec{c} = (1,1)^T$. We want $\partial J/\partial q_1$ and $\partial J/\partial q_2$.

### The Direct (Forward) Approach: Three Solves

Differentiating $A\vec{u} = \vec{b}$ with respect to $q_1$: $A\,(\partial\vec{u}/\partial q_1) = \partial\vec{b}/\partial q_1 = (1,0)^T$. Differentiating with respect to $q_2$: $A\,(\partial\vec{u}/\partial q_2) = \partial\vec{b}/\partial q_2 = (0,1)^T$. Beyond the nominal solve for $\vec{u}$, that is two linear solves (one per parameter) with the same matrix $A$:

$$A^{-1} = \tfrac{1}{5}\begin{pmatrix} 2 & -1 \\ -1 & 3 \end{pmatrix}, \quad \text{so} \quad \frac{\partial\vec{u}}{\partial q_1} = \tfrac{1}{5}\begin{pmatrix} 2 \\ -1 \end{pmatrix}, \qquad \frac{\partial\vec{u}}{\partial q_2} = \tfrac{1}{5}\begin{pmatrix} -1 \\ 3 \end{pmatrix}.$$

The derivatives of $J = \vec{c}^T\vec{u}$ are:

$$\frac{\partial J}{\partial q_1} = \vec{c}^T\frac{\partial\vec{u}}{\partial q_1} = (1,1)\cdot\tfrac{1}{5}(2,-1)^T = \tfrac{1}{5}, \qquad \frac{\partial J}{\partial q_2} = \vec{c}^T\frac{\partial\vec{u}}{\partial q_2} = (1,1)\cdot\tfrac{1}{5}(-1,3)^T = \tfrac{2}{5}.$$

Two sensitivity solves give two derivatives; with the nominal solve, that is three solves. For 50 parameters: 51 solves.

### The Adjoint Approach: Two Solves

Instead of solving for $\partial\vec{u}/\partial q_j$ separately for each $j$, introduce the *adjoint vector* $\vec{v}$ satisfying

$$A^T\vec{v} = \vec{c}. \tag{5.2}$$

This is one linear solve with the transposed matrix, the second solve after the nominal one. Since $A$ is symmetric here, $A^T = A$, so $\vec{v} = A^{-1}\vec{c}$:

$$\vec{v} = A^{-1}\vec{c} = \tfrac{1}{5}\begin{pmatrix} 2 & -1 \\ -1 & 3 \end{pmatrix}\begin{pmatrix} 1 \\ 1 \end{pmatrix} = \tfrac{1}{5}\begin{pmatrix} 1 \\ 2 \end{pmatrix}.$$

*What does $\vec{v}$ represent?* The adjoint vector is indexed by the *equations* of the system, not by its states. Its $i$th entry measures how much $J$ moves when the $i$th right-hand side $b_i$ moves, so $\vec{v} = \partial J/\partial\vec{b}$. The state side of the question is already settled and is not interesting here: $\partial J/\partial\vec{u} = \vec{c} = (1,1)^T$, so $u_1$ and $u_2$ contribute to $J = u_1 + u_2$ equally, by construction. What the adjoint adds is the translation from a perturbation of the *data* to a change in the response, after the coupling imposed by $A$ has been accounted for.

In the 2×2 example, $\vec{v} = \frac{1}{5}(1,2)^T$ says that adding a unit to $b_2$ moves $J$ *twice as much* as adding a unit to $b_1$. That asymmetry comes entirely

from $A$, since $\vec{c}$ weights the two states equally. This is what makes the adjoint more than a convenient shortcut: $\vec{v}$ is the sensitivity of the response to the forcing, which is the quantity a modeler can actually act on, and no rearrangement of $\vec{c}$ alone would produce it.

Now observe: for any parameter $q_j$,

$$\frac{\partial J}{\partial q_j} = \vec{c}^{\,T}\frac{\partial \vec{u}}{\partial q_j} = (A^T\vec{v})^T\frac{\partial \vec{u}}{\partial q_j} = \vec{v}^{\,T}\underbrace{A\frac{\partial \vec{u}}{\partial q_j}}_{=\partial \vec{b}/\partial q_j} = \vec{v}^{\,T}\frac{\partial \vec{b}}{\partial q_j}.$$

This is the key step: $A\,(\partial\vec{u}/\partial q_j) = \partial\vec{b}/\partial q_j$ is the FSE right-hand side, a known vector. No additional solve is needed, just a dot product.

Checking against our direct answers:

$$\frac{\partial J}{\partial q_1} = \vec{v}^{\,T}\frac{\partial \vec{b}}{\partial q_1} = \tfrac{1}{5}(1,2)\cdot(1,0)^T = \tfrac{1}{5}\ \checkmark \qquad \frac{\partial J}{\partial q_2} = \vec{v}^{\,T}\frac{\partial \vec{b}}{\partial q_2} = \tfrac{1}{5}(1,2)\cdot(0,1)^T = \tfrac{2}{5}\ \checkmark$$

One adjoint solve for $\vec{v}$, then one dot product per parameter. For 50 parameters: still one adjoint solve, then 50 dot products, against one solve per parameter for the FSE. Counting the nominal solve on both sides, as §5.4 does, that is 51 solves against 2, a factor of about 25. Counting only the sensitivity solves gives 50 against 1; the two conventions differ by roughly a factor of two, which is why the chapter fixes the first and keeps to it.

*Self-check.* Add a third parameter $q_3$ to the system above by setting $\vec{b} = (q_1 + q_3,\ q_2)^T$. Without solving any new linear systems, use the adjoint vector $\vec{v}$ already computed to find $\partial J/\partial q_3$.

> *Answer:* $\partial\vec{b}/\partial q_3 = (1,0)^T$. $\partial J/\partial q_3 = \vec{v}^{\,T}(1,0)^T = \tfrac{1}{5}(1,2)\cdot(1,0)^T = \tfrac{1}{5}$. No new solve needed. Adding parameters costs only one dot product each once $\vec{v}$ is known. This is the computational power of the adjoint.

The pattern above is the entire idea of the adjoint, stripped to its essentials. Section 5.2 states it precisely.

## 5.2 ▪ The Strategy, Stated Explicitly

The cost analysis above raises the key question: can we recover all $m$ sensitivities without solving $m$ separate systems? It turns out we can, through a trick that is worth understanding before seeing the symbols.

The adjoint is not a new kind of problem. It is a deliberate trick: introduce an auxiliary vector, choose it to cancel the unknown sensitivity variable, and read off the result. Here is the trick in plain language before any symbols.

Draft

**Step 1.** We want $\partial J/\partial q_j$ for all parameters $q_j$. Since $J = \vec{c}^{\,T}\vec{u}$, this is $\partial J/\partial q_j = \vec{c}^{\,T}(\partial \vec{u}/\partial q_j)$. But $\partial \vec{u}/\partial q_j$ requires a separate FSE solve for every $j$ (50 solves for 50 parameters, after the nominal solve).

**Step 2.** Introduce an auxiliary vector $\vec{v}$ and observe that for any vector $\vec{v}$:

$$\vec{c}^{\,T}\frac{\partial \vec{u}}{\partial q_j} = \vec{v}^{\,T}\left[A\frac{\partial \vec{u}}{\partial q_j}\right] + \left(\vec{c}^{\,T} - \vec{v}^{\,T}A\right)\frac{\partial \vec{u}}{\partial q_j}.$$

The first term on the right is $\vec{v}^{\,T}\vec{r}_j$, where $\vec{r}_j = A(\partial \vec{u}/\partial q_j)$ is the FSE right-hand side (a known vector once $\vec{u}$ is known). The second term vanishes if we choose $\vec{v}$ so that $\vec{v}^{\,T}A = \vec{c}^{\,T}$.

**Step 3.** Choose $\vec{v}$ to satisfy the **adjoint equation**:

$$A^T\vec{v} = \vec{c}. \tag{5.3}$$

Then $\vec{c}^{\,T} - \vec{v}^{\,T}A = \vec{0}$ and the sensitivity formula reduces to:

$$\frac{\partial J}{\partial q_j} = \vec{v}^{\,T}\vec{r}_j \qquad \text{for all } j = 1, \ldots, m. \tag{5.4}$$

One solve for $\vec{v}$ (equation (5.3)) recovers all $m$ sensitivities through formula (5.4). The trick works because the adjoint equation does not depend on $j$: $\vec{v}$ is determined by $\vec{c}$ and $A$ alone.

The adjoint is not a separate, unrelated problem. It is constructed entirely from the coefficient matrix $A$ (via transposition) and the response vector $\vec{c}$ of the forward problem. Everything about the adjoint is determined by the forward problem.

*Connection to Linear Programming Duality*

Readers who have seen linear programming will recognize a structural parallel. The primal LP $\max\{\vec{c}^{\top}\vec{x} : A\vec{x} = \vec{b},\ \vec{x} \geq 0\}$ has a dual problem $\min\{\vec{b}^{\top}\vec{y} : A^{\top}\vec{y} \geq \vec{c}\}$, whose constraint involves the transposed matrix $A^{\top}$. The LP dual variable $\vec{y}$ measures the sensitivity of the optimal objective value to perturbations in the right-hand side $\vec{b}$, which is what the adjoint variable $\vec{v}$ does here: it measures the sensitivity of $J = \vec{c}^{\top}\vec{u}$ to perturbations in $\vec{b}$ (the parameter forcing). The adjoint equation $A^{\top}\vec{v} = \vec{c}$ is a direct generalization of the LP dual constraint. This connection deepens in Chapter 6, where the adjoint ODE corresponds to the dual of the optimal-control Hamiltonian.

*Self-check.* For the system $A\vec{u} = \vec{b}$ with $A = \begin{pmatrix} 2 & 1 \\ 0 & 3 \end{pmatrix}$ and response functional $J = u_1$ (so $\vec{c} = (1, 0)^T$): write down the adjoint equation (5.3) and solve for $\vec{v}$.

*Answer:* $A^T\vec{v} = \vec{c}$ becomes $\begin{pmatrix} 2 & 0 \\ 1 & 3 \end{pmatrix}\vec{v} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$. Solving: $v_1 = 1/2$, $v_2 = -1/6$. This single solve (one back-substitution with $A^T$) gives us the sensitivity of $J = u_1$ to any number of parameters, via formula (5.4).

## 5.3 ▪ The Commutative Diagram

The three-step strategy and the 2×2 preview both express the same mathematical fact: the FSE route and the adjoint route reach the same sensitivity, at different cost. Figure 5.1 shows this as a diagram, read it as a summary of the algebra already developed, not as an introduction to the idea.





**Figure 5.1.** *The commutative diagram for sensitivity analysis. The forward path (solid arrows) computes $\partial J/\partial q_j$ by solving the FSE for each parameter, $m+1$ solves total. The adjoint path (dashed arrows) solves the adjoint equation once, then evaluates the sensitivity formula for each parameter; 2 solves total. Both paths deliver the same result.*

The diagram commutes: both paths from the forward problem to the sensitivity give identical results. This can be verified directly from the algebra in Section 5.2.

**Theorem 5.1** (Forward–Adjoint Equivalence)**.** *Let $A\vec{u} = \vec{b}$ with $A$ nonsingular, $J = \vec{c}^T\vec{u}$, and FSE $A\,\partial\vec{u}/\partial q_j = \vec{r}_j$. Let $\vec{v}$ solve the adjoint equation $A^T\vec{v} = \vec{c}$. Then*

$$\frac{\partial J}{\partial q_j} = \vec{c}^T\frac{\partial \vec{u}}{\partial q_j} = \vec{v}^T\vec{r}_j$$

*for every parameter $q_j$.*

*Proof.* $\vec{c}^T(\partial\vec{u}/\partial q_j) = (A^T\vec{v})^T(\partial\vec{u}/\partial q_j) = \vec{v}^T A(\partial\vec{u}/\partial q_j) = \vec{v}^T\vec{r}_j$. The second equality uses $(A^T\vec{v})^T = \vec{v}^T(A^T)^T = \vec{v}^T A$, which follows from the rule $(BC)^T = C^T B^T$ with $B = A^T$, $C = \vec{v}$. □

The adjoint gives *exactly* the same sensitivity values as the FSE. It is not an approximation. Students sometimes expect that a method requiring fewer solves must sacrifice accuracy; here it does not.

*Self-check.* Using Theorem 5.1, state in words why the adjoint equation does not need to be re-solved when a new parameter $q_k$ is added to the sensitivity study.

> *Answer:* The adjoint equation $A^T\vec{v} = \vec{c}$ depends only on $A$ (the system coefficient matrix) and $\vec{c}$ (the response vector), neither of which changes when we add a new parameter. Adding parameter $q_k$ changes only the FSE right-hand side $\vec{r}_k = \partial\vec{b}/\partial q_k - (\partial A/\partial q_k)\vec{u}$, which is a known vector once the forward solution $\vec{u}$ is computed. The sensitivity $\partial J/\partial q_k = \vec{v}^T\vec{r}_k$ is then obtained by a single dot product, no additional linear solve.

## 5.4 ▪ Cost Comparison and the Decision Rule

### 5.4.1 ▪ When to Use the Adjoint

Both the FSE and the adjoint produce the same sensitivity Jacobian. Their costs differ dramatically when $m$ and $r$ are unequal, where $m$ is the number of POIs and $r$ is the number of response functionals.

For the FSE: one solve per POI, plus the nominal solve for $\vec{u}$ itself $\rightarrow$ $m + 1$ solves total.

For the adjoint: one solve per response functional, plus that same nominal solve $\rightarrow$ $r + 1$ solves total. Counting the nominal solve on both sides is what makes the comparison fair, and it is the convention used in Chapter 4: for the malaria output $i_h(T)$, seventeen parameters cost eighteen ODE solves by the FSE and two by the adjoint.

(5.5) $$\boxed{m > r \implies \text{use adjoint}; \qquad r > m \implies \text{use FSE.}}$$

In most SA applications, there are many POIs ($m$ large) and a small number of QOIs of primary interest ($r$ small), so the adjoint is preferred. The canonical situation is $r = 1$: one scalar response functional (peak infection count, total burden, profit at equilibrium) and many parameters.

### 5.4.2 ▪ Cost Comparison Plot

Figure 5.2 shows the number of linear solves required by each method as a function of the number of parameters.





**Figure 5.2.** *Cost comparison for FSE vs. adjoint sensitivity analysis, measured in number of linear solves. For $r = 1$ response functional (dashed), the adjoint always requires exactly 2 solves regardless of $m$. For $r = 3$ (dotted), the adjoint requires 4 solves. The FSE requires $m + 1$ solves. The crossover occurs at $m = r$; for $m > r$, the adjoint wins.*

For the competitive population model ($m = 3$ parameters, $r = 2$ response functionals $\{\bar{u}_1, \bar{u}_2\}$): FSE requires 4 solves; adjoint requires 3. The advantage is modest here. For the malaria model ($m = 17$, $r = 1$): FSE requires 18 solves; adjoint requires 2, the factor of 9 quoted in Chapter 4. Those counts are ODE solves for the output $i_h(T)$. For the endemic equilibrium the counts are the same, but only the first solve, which finds the equilibrium, is nonlinear; each of the others is a linear solve. For a climate model ($m = 200$, $r = 2$): FSE requires 201 solves; adjoint requires 3: a factor of 67 improvement.

*Self-check.* A model has $m = 30$ POIs and $r = 5$ response functionals. Each linear solve takes 2 minutes. How long does the FSE approach take? How long does the adjoint approach take? At what value of $r$ would the two approaches have equal cost?

*Answer:* FSE: $31 \times 2 = 62$ minutes, counting the baseline solve. Adjoint: $6 \times 2 = 12$ minutes, likewise. Savings: 50 minutes. The saving is unchanged because the baseline solve is common to both; the counts are stated as totals so they match the convention used above. Equal cost when $r = m = 30$.

## 5.5 ▪ The Competitive Population: Full Adjoint Treatment

We now apply the adjoint to the competitive population equilibrium from Chapter 4, comparing the forward and adjoint approaches side by side. The forward problem is the equilibrium condition $\vec{f}(\vec{u}) = \vec{0}$, which is nonlinear; its solution is (4.4). Only the FSE (4.5) is linear, with matrix $A = D_{\vec{u}}[\vec{f}]$ as in equation (4.6). The comparison makes the cost advantage concrete.

### 5.5.1 ▪ FSE Approach: Three Parameters, Four Solves

For parameters $a$, $b$, $c$, the FSE requires solving $A(\partial\vec{u}/\partial q_j) = -\partial\vec{f}/\partial q_j$ for $j = a, b, c$, three linear solves with the same matrix $A$, one LU factorization and three back-substitutions. With the nominal solve for $\vec{u}$, the FSE costs four solves.

### 5.5.2 ▪ Adjoint Approach: Three Solves for Two Response Functionals

Now suppose we want $\partial J_1/\partial q_j$ and $\partial J_2/\partial q_j$ for all three parameters, where $J_1 = \bar{u}_1$ and $J_2 = \bar{u}_2$. The response vectors are $\vec{c}_1 = (1, 0)^T$ and $\vec{c}_2 = (0, 1)^T$.

The two adjoint equations are:

$$A^T\vec{v}_1 = \vec{c}_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix}, \qquad A^T\vec{v}_2 = \vec{c}_2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}. \tag{5.6}$$

Two adjoint solves (one per response functional) give $\vec{v}_1$ and $\vec{v}_2$. With the same nominal solve for $\vec{u}$, the adjoint costs three solves.

**A note on correlated QOIs.** Even when the POIs are independent (uncorrelated inputs), the QOIs are almost always correlated outputs: $J_1 = \bar{u}_1$ and $J_2 = \bar{u}_2$ both depend on the same equilibrium $\vec{u}$, so perturbations to any parameter affect both simultaneously. This is not a problem for the adjoint method; it is simply a property of the model. The adjoint computes each $\partial J_k/\partial q_j$ independently by solving $A^T\vec{v}_k = \vec{c}_k$ for each response functional. Any statistical correlation among the $J_k$'s is a property of the model, not an issue for the adjoint method; no special handling is required. The adjoint naturally produces correct, consistent sensitivities for all $r$ response functionals by virtue of operating through the same forward solution $\vec{u}$.

The sensitivities follow from formula (5.4):

$$\frac{\partial J_k}{\partial q_j} = \vec{v}_k^T\vec{r}_j, \qquad \vec{r}_j = -\frac{\partial\vec{f}}{\partial q_j}, \tag{5.7}$$

giving all $2 \times 3 = 6$ sensitivities from two adjoint solves rather than three FSE solves.

### 5.5.3 ▪ Verification

Theorem 5.1 guarantees the two approaches give identical results. At nominal values $a = 0.4$, $b = 1$, $c = 0.3$:

| | $\mathcal{S}_a^{\bar{u}_1}$ | $\mathcal{S}_b^{\bar{u}_1}$ | $\mathcal{S}_c^{\bar{u}_1}$ | $\mathcal{S}_a^{\bar{u}_2}$ | $\mathcal{S}_b^{\bar{u}_2}$ | $\mathcal{S}_c^{\bar{u}_2}$ |
|---|---|---|---|---|---|---|
| FSE (3 solves) | $-0.530$ | 0 | 0.136 | $+0.136$ | 0 | $-0.292$ |
| Adjoint (2 solves) | $-0.530$ | 0 | 0.136 | $+0.136$ | 0 | $-0.292$ |

The rows are identical. The adjoint is not an approximation. The row labels count the sensitivity solves alone; with the nominal solve, the totals are 4 and 3.

A structural observation: the two adjoint solutions $\vec{v}_1, \vec{v}_2$ are the columns of $(A^T)^{-1}$, which equals $(A^{-1})^T$. Collecting all adjoint solutions into a matrix therefore gives the transpose of the inverse. This connection is intellectually interesting but not computationally useful: we never form $A^{-1}$ explicitly, we extract the product $\vec{v}_k^T \vec{r}_j$ for each required sensitivity, which is far cheaper than computing the full inverse matrix.

## 5.6 ▪ Reverse Mode and Backpropagation

The adjoint strategy of the preceding sections applies to linear systems. This section extends it to arbitrary differentiable computations via the computation graph, revealing the connection to algorithmic differentiation and, through it, to the training of neural networks.

### 5.6.1 ▪ The Computation Graph

Every differentiable computation can be represented as a directed acyclic graph in which each node holds an intermediate value and each edge represents an elementary operation (addition, multiplication, division, elementary function).

For $\mathcal{R}_0 = c\beta/\gamma_R$, the computation graph is:

$$c, \beta \xrightarrow{\times} v_1 = c\beta \xrightarrow{\div \gamma_R} v_2 = c\beta/\gamma_R = \mathcal{R}_0. \tag{5.8}$$

Reading left to right is the forward evaluation: given $c$, $\beta$, $\gamma_R$, compute $\mathcal{R}_0$.

### 5.6.2 ▪ Forward Mode: One Pass Per Input

In *forward mode* differentiation, we propagate derivatives $\dot{v}_i = \partial v_i/\partial q$ through the graph for one input $q$ at a time.

For $q = c$: set $\dot{c} = 1$, $\dot{\beta} = 0$, $\dot{\gamma_R} = 0$. Propagate forward:

$$\dot{v}_1 = \dot{c}\beta + c\dot{\beta} = \beta, \qquad \dot{R}_0 = \dot{v}_1/\gamma_R - v_1\dot{\gamma_R}/\gamma_R^2 = \beta/\gamma_R.$$

So $\partial\mathcal{R}_0/\partial c = \beta/\gamma_R$. Repeat with $\dot{\beta} = 1$ to get $\partial\mathcal{R}_0/\partial\beta$. Repeat with $\dot{\gamma_R} = 1$ to get $\partial\mathcal{R}_0/\partial\gamma_R$. Total: 3 forward passes for 3 parameters.

### 5.6.3 ▪ Reverse Mode: Two Passes for All Inputs

In *reverse mode* differentiation, we propagate the adjoint variable $\bar{v}_i = \partial J/\partial v_i$ backward through the graph.

Set $\bar{R}_0 = \partial J/\partial\mathcal{R}_0 = 1$ (we are computing $\partial\mathcal{R}_0/\partial(\cdot)$, so $J = \mathcal{R}_0$). Propagate backward:

$$\bar{v}_1 = \bar{R}_0 \cdot \frac{\partial\mathcal{R}_0}{\partial v_1} = 1 \cdot (1/\gamma_R) = 1/\gamma_R, \tag{5.9}$$

$$\bar{c} = \bar{v}_1 \cdot \frac{\partial v_1}{\partial c} = (1/\gamma_R) \cdot \beta = \beta/\gamma_R, \tag{5.10}$$

$$\bar{\beta} = \bar{v}_1 \cdot \frac{\partial v_1}{\partial \beta} = (1/\gamma_R) \cdot c = c/\gamma_R, \tag{5.11}$$

$$\bar{\gamma_R} = \bar{R}_0 \cdot \frac{\partial\mathcal{R}_0}{\partial\gamma_R} = 1 \cdot (-v_1/\gamma_R^2) = -c\beta/\gamma_R^2. \tag{5.12}$$

After the forward evaluation, one backward pass gives all three derivatives simultaneously:

$$\frac{\partial\mathcal{R}_0}{\partial c} = \bar{c} = \frac{\beta}{\gamma_R}, \quad \frac{\partial\mathcal{R}_0}{\partial\beta} = \bar{\beta} = \frac{c}{\gamma_R}, \quad \frac{\partial\mathcal{R}_0}{\partial\gamma_R} = \bar{\gamma_R} = -\frac{c\beta}{\gamma_R^2}.$$

These match the analytic results. Reverse mode costs two passes regardless of the number of inputs: the forward evaluation that stores the node values and one backward pass. Each pass costs time proportional to the graph size.

Table 5.1 summarizes the comparison.

### 5.6.4 ▪ The Backpropagation Connection

*Note for readers without machine learning background: this subsection connects the adjoint to neural network training. If backpropagation is unfamiliar, skip ahead to Unifying Idea 2 below and continue from there; the adjoint ODE does not depend on the neural network material. Return here after completing Chapter 6 if you want the connection.*

If you have encountered training a neural network, you have already used the adjoint under a different name.

| | Forward mode | Reverse mode |
|---|---|---|
| Propagates | $\dot{v}_i = \partial v_i/\partial q$ (one $q$) | $\bar{v}_i = \partial J/\partial v_i$ (one $J$) |
| Direction | Input $\to$ output | Output $\to$ input |
| Cost | 1 pass per input | $1 + r$ passes |
| Preferred when | $m \leq r$ (few inputs) | $m > r$ (few outputs) |
| Matrix analog | FSE: $A\,\partial\vec{u}/\partial q_j = \vec{r}_j$ | Adjoint: $A^T\vec{v} = \vec{c}$ |

**Table 5.1.** *Forward mode and reverse mode differentiation: a side-by-side comparison. The matrix analogs confirm that the two modes are the computation-graph generalizations of the FSE and adjoint approaches from Sections 5.2–5.4.*

A neural network is a composition of parameterized functions. Training minimizes a loss function $J(\theta)$ over parameters $\theta$ using gradient descent, which requires $\partial J/\partial\theta_i$ for every parameter $\theta_i$. The loss function is a scalar ($r = 1$) and modern networks have millions of parameters ($m \gg 1$), so reverse mode is overwhelmingly preferred. Backpropagation is exactly the reverse-mode computation described above, applied to the computation graph of the network.

To map notation explicitly: each computation-graph node $v_i$ corresponds to the $i$-th standard basis vector $\vec{e}_i$, and the adjoint variable $\lambda_i = \partial J/\partial v_i$ records how much $J$ changes per unit change at node $i$, exactly the role played by $\bar{v}_i$ in reverse mode.

The adjoint equation for the ODE case, derived in Chapter 6, is the continuous limit of this discrete backward accumulation. The backward ODE for $\lambda(t)$ is the functional-analytic version of the backward pass through the computation graph.

*Unifying Idea 2.* The adjoint is the transpose in whatever space you are working in.

In this chapter: the adjoint equation is $A^T\vec{v} = \vec{c}$, the transposed coefficient matrix applied to the adjoint vector.

In the computation graph: reverse mode is the chain rule applied in reverse order, accumulating $\bar{v}_i = \partial J/\partial v_i$ backward through the graph.

In Chapter 6: the adjoint will be an ODE running backward in time, with a terminal condition at $t = T$. The integration-by-parts identity that produces this ODE is the functional-analytic analog of the matrix transposition that produced $A^T$ here.

One idea. Three settings. One cost advantage: when $m \gg r$, the adjoint/reverse-mode is the preferred computational strategy.

*Self-check.* Use the reverse-mode computation of equations (5.9)–(5.12) to compute $\mathcal{S}_c^{\mathcal{R}_0}$, $\mathcal{S}_\beta^{\mathcal{R}_0}$, and $\mathcal{S}_{\gamma_R}^{\mathcal{R}_0}$. Express each as a pure number and verify

using the Chapter 2 results.

*Answer:* $\partial\mathcal{R}_0/\partial c = \beta/\gamma_R$. $\mathcal{S}_c^{\mathcal{R}_0} = (c/\mathcal{R}_0)(\partial\mathcal{R}_0/\partial c) = (c/(c\beta/\gamma_R)) \cdot (\beta/\gamma_R) = 1$.
$\partial\mathcal{R}_0/\partial\beta = c/\gamma_R$. $\mathcal{S}_\beta^{\mathcal{R}_0} = (\beta/(c\beta/\gamma_R))(c/\gamma_R) = 1$.
$\partial\mathcal{R}_0/\partial\gamma_R = -c\beta/\gamma_R^2$. $\mathcal{S}_{\gamma_R}^{\mathcal{R}_0} = (\gamma_R/(c\beta/\gamma_R))(-c\beta/\gamma_R^2) = -1$.
All three match the Chapter 2 results. The reverse-mode computation recovers the full sensitivity Jacobian in two passes: the forward evaluation and one backward pass.

## 5.7 ▪ Bridge to Chapter 6

Students sometimes expect the adjoint for differential equations to require advanced functional analysis (Hilbert spaces, adjoint operators, and formal operator theory). Appendix A provides the $L^2$ inner product and orthogonality needed for that formalism; Appendix B develops the full abstract operator framework that makes the connection rigorous. The preceding sections show why this expectation is misplaced: the core idea is the matrix transpose. Every formal construct in Chapter 6 is the natural extension of the matrix algebra developed here. Reading Chapter 6 with this chapter in mind, the abstract notation should feel like familiar territory.

*Unifying Idea 1.* Every sensitivity value computed in Section 5.5, via the adjoint, is a normalized derivative, an instance of Unifying Idea 1 from Chapter 1. The adjoint is a computational strategy; the quantity it produces is still the SI defined in Chapter 2. The two commutative paths in Figure 5.1 produce the same sensitivity index by different routes.

Chapter 5 has established the adjoint for linear systems. Chapter 6 extends every result to ordinary differential equations. The sensitivity Jacobian computed in this chapter is the same object that Smith [59] uses in Ch. 7 to drive parameter identifiability analysis and parameter selection; students who complete Chapter 6 will find Ch. 7 of Smith immediately accessible.

The translation is direct:

| Chapter 5 (linear algebra) | Chapter 6 (dynamical systems) |
|---|---|
| Coefficient matrix $A$ | Differential operator $d/dt - J(t)$ |
| Transpose $A^T$ | Adjoint operator $-(d/dt + J(t)^T)$ |
| Adjoint equation $A^T\vec{v} = \vec{c}$ | Adjoint ODE $-\dot{\lambda} = J^T\lambda + g_u$ |
| Terminal condition: none | Terminal condition: $\lambda(T) = h_u^T$ |
| Sensitivity formula $\vec{v}^T\vec{r}_j$ | Sensitivity formula $\int_0^T \lambda^T F_{p_j}\, dt$ |
| Cost: 2 solves for any $m$ | Cost: 2 ODE solves for any $m$ |

The integration-by-parts identity in Chapter 6 plays the role of the algebraic transposition here. The terminal condition on $\lambda(T)$ is the time-dependent analog of the right-hand side $\vec{c}$ in $A^T\vec{v} = \vec{c}$. The cost advantage is identical: two solves, the forward solve and one adjoint solve, recover all $m$ sensitivities of one response functional regardless of $m$.

In Chapter 6 we extend every result of this chapter to ordinary differential equations. The matrix $A$ becomes a differential operator. The transpose $A^T$ becomes the adjoint differential operator. The linear solve $A^T\vec{v} = \vec{c}$ becomes an ODE for $\lambda(t)$ with a terminal condition at $t = T$. The sensitivity formula $\vec{v}^T\vec{r}_j$ becomes an integral over $[0, T]$. The cost advantage is the same: 2 ODE solves instead of $m + 1$.

## 5.8 ▪ Computational Laboratory: Forward vs. Adjoint

This laboratory compares the FSE and adjoint computations on the competitive population model, then implements reverse-mode differentiation for $\mathcal{R}_0$.

**Part 1: Competitive population via FSE.** At nominal values $a = 0.4$, $b = 1$, $c = 0.3$, compute the equilibrium using equation (4.4). Assemble the Jacobian $A$ and compute the LU factorization using your language's LU routine. Solve the three FSEs ($\partial\vec{u}/\partial a$, $\partial\vec{u}/\partial b$, $\partial\vec{u}/\partial c$) using `U\(L\r)` for each. Record the time.

**Part 2: Competitive population via adjoint.** Solve the two adjoint equations $A^T\vec{v}_1 = \vec{c}_1$ and $A^T\vec{v}_2 = \vec{c}_2$ using the same LU factorization (with transposed triangular solves). Compute all 6 sensitivities via $\vec{v}_k^T\vec{r}_j$. Record the time and verify that all 6 values agree with Part 1 to machine precision.

*Note on the transposed solve.* Many LU routines give $A = P^TLU$, so $A^T = U^TL^TP$. Solving $A^T\vec{v} = \vec{c}$ therefore proceeds in three steps: $U^T\vec{w}_1 = \vec{c}$, then $L^T\vec{w}_2 = \vec{w}_1$, then $\vec{v} = P^T\vec{w}_2$. In languages with matrix-transpose syntax, this is written as `P' * (L' \ (U' \ c))`, reading right to left: first solve `U'\c`, then `L'\`, then multiply by `P'` at the end.

```
%% Assemble Jacobian at equilibrium (see Chapter 4)
A = jac_symp(ubar, a, b, c);
[L, U, P] = lu(A);

%% FSE: 3 solves
for j = 1:3
    rj = rhs_symp(ubar, j);          %% FSE right-hand side
    dwdp(:,j) = U \ (L \ (P * rj));
end
```

```
%% Adjoint: 2 solves
c1 = [1;0]; c2 = [0;1];
v1 = P' * (L' \ (U' \ c1));   %% solve A^T v1 = c1
v2 = P' * (L' \ (U' \ c2));   %% solve A^T v2 = c2
for j = 1:3
    rj = rhs_symp(ubar, j);
    dJdp(1,j) = v1' * rj;     %% dJ1/dp_j
    dJdp(2,j) = v2' * rj;     %% dJ2/dp_j
end
```

**Part 3: Timing as a function of system size.** For random systems of size $n = 5, 10, 50, 100$ with $m = 20$ parameters and $r = 1$ response functional, time both approaches. Plot computation time vs. $n$ on log-log axes. Does the ratio of times approach the solve-count ratio $(m+1)/(r+1)$, here 21 against 2, as expected?

**Part 4: Reverse-mode $\mathcal{R}_0$ by hand.** Implement the reverse-mode computation (5.9)–(5.12) for $\mathcal{R}_0 = c\beta/\gamma_R$. Verify all three sensitivities against a symbolic computation tool. Count the number of arithmetic operations in forward mode vs. reverse mode for this graph.

**Part 5: Verify that adjoint = 50 FSE solves.** Set up a random $10 \times 10$ system with $m = 50$ parameters and $r = 1$. Compute $\partial J/\partial q_j$ for all $j$ by: (a) the forward path, the nominal solve and 50 FSE solves, 51 in all; (b) the adjoint path, the nominal solve and 1 adjoint solve, 2 in all. Verify that the maximum absolute difference is less than $10^{-12}$.

## 5.9 ▪ Exercises

**Exercise 5.1** *([Bloom L2]).* State the three-step adjoint strategy from Section 5.2 in your own words, using no symbols. Then explain: why does the adjoint equation $A^T\vec{v} = \vec{c}$ not depend on which parameter $q_j$ we are differentiating with respect to? Why does this independence make the adjoint efficient?

**Exercise 5.2** *([Bloom L3]).* For the linear system

$$A\vec{u} = \vec{b}, \qquad A = \begin{pmatrix} 3 & 1 \\ 1 & 2 \end{pmatrix}, \quad \vec{b} = \begin{pmatrix} b_1(q) \\ b_2(q) \end{pmatrix} = \begin{pmatrix} q \\ 2q \end{pmatrix},$$

with response functional $J = u_1 + u_2$ (so $\vec{c} = (1,1)^T$):

(a) Solve the forward problem to find $\vec{u}(q)$ at $q = 1$.

(b) Write and solve the FSE for $\partial \vec{u}/\partial q$.

(c) Write and solve the adjoint equation.

(d) Use formula (5.4) to compute $\partial J/\partial q$.

(e) Verify by differentiating $J(q) = u_1(q) + u_2(q)$ directly.

**Exercise 5.3** *([Bloom L3])*. For each scenario below, state whether you would use the FSE or the adjoint, and calculate the number of linear solves each approach requires.

(a) $m = 5$ parameters, $r = 20$ response functionals.

(b) $m = 100$ parameters, $r = 3$ response functionals.

(c) $m = 8$ parameters, $r = 8$ response functionals.

(d) $m = 1{,}000$ parameters, $r = 1$ response functional. Each solve takes 0.1 seconds. How much faster is the preferred method?

**Exercise 5.4** *([Bloom L4])*. For the competitive population equilibrium with parameters $a = 0.5$, $b = 2$, $c = 0.4$:

(a) Compute all 6 sensitivities ($\mathcal{S}_a^{\bar{u}_1}$, $\mathcal{S}_b^{\bar{u}_1}$, $\mathcal{S}_c^{\bar{u}_1}$, $\mathcal{S}_a^{\bar{u}_2}$, $\mathcal{S}_b^{\bar{u}_2}$, $\mathcal{S}_c^{\bar{u}_2}$) using the FSE approach (4 solves, counting the nominal solve).

(b) Compute the same 6 sensitivities using the adjoint approach (3 solves).

(c) Verify that the two methods agree.

(d) Which approach requires fewer solves? By how much? At what values of $m$ and $r$ would the FSE be preferred?

**Exercise 5.5** *([Bloom L4])*. The SEIR basic reproduction number is $\mathcal{R}_0 = c\beta\nu/[(\nu+\nu_m)(\gamma_R+\nu_m)]$ with parameters $c$, $\beta$, $\nu$ (exposure rate), $\gamma_R$ (recovery rate), $\nu_m$ (mortality rate).

(a) Draw the computation graph for $\mathcal{R}_0$ (intermediate nodes and edges).

(b) Implement the forward-mode computation to find $\partial \mathcal{R}_0/\partial \nu$. Count the number of arithmetic operations.

(c) Implement the reverse-mode computation to find all five derivatives $\partial \mathcal{R}_0/\partial c$, $\partial \mathcal{R}_0/\partial \beta$, $\partial \mathcal{R}_0/\partial \nu$, $\partial \mathcal{R}_0/\partial \gamma_R$, $\partial \mathcal{R}_0/\partial \nu_m$ in one backward pass after the forward evaluation.

(d) Compute the five normalized sensitivity indices and produce a tornado plot using interpretable POIs (mean times, not rates).

**Exercise 5.6** *([Bloom L5])*. A water-quality model of a river system has $m = 40$ parameters (pollution source rates, degradation rates, flow speeds, mixing coefficients) and $r = 2$ response functionals: (1) pollutant concentration at a downstream monitoring station, and (2) total pollutant load entering a protected estuary.

Design a complete SA study using the adjoint method. Specify:

(a) The form of the adjoint equation for this model.

(b) How many adjoint solves are required and why.

(c) How you would validate the implementation.

(d) How you would use the sensitivity results to advise regulators on which pollution sources to prioritize.

(e) One situation where the adjoint results might be misleading, and what you would do to detect and correct it.

**Exercise 5.7** *([Bloom L6])*. A colleague states: "The adjoint always gives the same answer as forward SA, just faster. So if we already have the FSE code working, there is no reason to implement the adjoint."

Evaluate this claim carefully.

(a) Is the first sentence (same answer, just faster) correct? Give a precise mathematical justification.

(b) Is the second sentence (no reason to implement the adjoint) correct? Describe at least two situations where switching from FSE to adjoint would be strongly advisable.

(c) For the FSE to be preferred over the adjoint, what relationship between $m$ and $r$ must hold? Give a specific numerical example where FSE is strictly better.

Draft

## Notes

The adjoint of a linear operator on a Hilbert space is the abstract generalization of the matrix transpose; a rigorous treatment appears in Appendix A and in standard functional analysis texts. Horn and Johnson [22] give a comprehensive treatment of matrix adjoints and their use in sensitivity analysis of linear systems.

Reverse-mode algorithmic differentiation on a computation graph, which is the computational realization of the adjoint idea, is equivalent to the backpropagation algorithm introduced by Rumelhart, Hinton, and Williams in the context of neural network training; see the historical account in Baydin et al. [4]. The connection between the adjoint method in scientific computing and backpropagation in machine learning is made explicit by Margossian [30].

The cost argument, that the forward solve and one adjoint solve recover all $m$ sensitivities for one response functional, against $m + 1$ solves for forward SA, appears in Cacuci [9] and Giles and Pierce [16]. Giles and Pierce give an accessible exposition of the adjoint idea in the context of aeronautical design optimization, a field where $m$ can reach $10^6$ design variables.

Draft

Draft

# Chapter 6
# The Adjoint for Dynamical Systems

*To understand the effect of an outcome, trace its influence backward through time.*

*The forward model tells you where you end up.*
*The adjoint tells you how you got there.*

Climate models routinely have 500 or more tunable parameters. Forward sensitivity analysis using the FSE approach requires one ODE solve per parameter plus the baseline: 501 solves on a modern workstation, each taking 6 hours. Total: 3,006 hours, or about 125 days.

The adjoint approach requires two ODE solves for a single scalar QOI: one forward solve to generate the trajectory, one backward solve for the adjoint. Total cost: approximately 12 hours, plus storage and interpolation overhead for the forward solution.

The mathematics behind this 250-to-1 speedup is a direct extension of the matrix adjoint from Chapter 5. In that chapter, the key operation was transposing the coefficient matrix: $A \to A^T$. Here, the key operation is transposing the differential operator, which turns a forward ODE into a backward ODE. This chapter derives that operation step by step, explains in plain language why the backward direction arises, and implements the full calculation for the SIR epidemic model. The notation is one step heavier than Chapter 5, integration by parts and continuous-time functionals replace matrix transposes, but the underlying idea is the same transposition argument in a new space. The adjoint sensitivity equations for ODE systems, including their software implementation, are treated in depth by Cao et al. [12]; a recent comparison of forward and adjoint methods on biological models is given by Mester et al. [33].

Draft

## 6.1 ▪ The Timeline Figure

The most important idea in this chapter: the reason the adjoint ODE runs backward, is captured in Figure 6.1 before any equations appear.





**Figure 6.1.** *The timeline for forward and adjoint sensitivity analysis. The forward ODE propagates the model state $\vec{u}(t)$ from the initial condition at $t = 0$ toward the response $J$ at $t = T$. The adjoint ODE propagates the sensitivity weight $\vec{\lambda}(t)$ backward from a terminal condition at $t = T$, in the direction opposite to the forward problem. The two ODEs are not independent: the adjoint coefficient depends on the forward solution $\vec{u}(t)$ at every time step.*

The forward ODE begins at $t = 0$ and moves right, building up the state $\vec{u}(t)$ until it reaches the response $J$ at time $T$. The adjoint ODE begins at $t = T$ and moves left, distributing the response backward through time.

This opposite direction is not a numerical trick. It is a mathematical necessity, explained in Section 6.2.

## 6.2 ▪ Why Backward? Intuition Before Derivation

The adjoint ODE runs backward because the question it answers is naturally backward: how much did each moment in time contribute to the final value of $J$?

Think of the epidemic. A perturbation applied to the infectious compartment at $t = 0$ has the entire simulation period to propagate: it changes the dynamics at $t = 1$, $t = 2$, and all the way to $T$. Its influence on $J$ is large. A perturbation applied at $t = T - \epsilon$ has almost no time to propagate before the simulation ends. Its influence on $J$ is negligible.

So the "weight" we should assign to perturbations decreases as we move from left to right along the time axis, and increases as we move from right to left. The adjoint variable $\lambda(t)$ is this weight: $\lambda(t)$ measures how sensitive the response $J$ is to a perturbation of the state at time $t$.

At $t = T$, the weight is determined by the response functional (how does $J$ change if the final state changes?). At earlier times, $\lambda(t)$ is computed by propagating this weight backward, accounting for how perturbations at time $t$ have time to amplify before reaching $T$.

For a burden functional such as $J = \int_0^T I\,dt$ in the SIR model, the adjoint is largest near $t = 0$ and decays toward $T$: an early infection has the whole remaining epidemic in which to generate descendants, a late one does not. That shape is common but it is not a general rule. It depends jointly on the Jacobian, the running cost, and the terminal condition, and the scalar decay example later in this chapter reverses it: there $\lambda(t) = e^{p(t-T)}$ is *smallest* at $t = 0$ and largest at $t = T$. Read the shape off the dynamics rather than assuming it. The adjoint ODE starts at the only time where we know $\lambda$ directly at $t = T$, and integrates backward to find $\lambda(t)$ at all earlier times.

This must be understood before the derivation. The backward direction of the adjoint is not a consequence of the algebra; the algebra is a consequence of this backward structure.

The adjoint and forward problems are not independent: the coefficient in the adjoint ODE at every time $t$ involves $\vec{u}(t)$, the forward solution. The adjoint can only be solved after the forward problem is solved and $\vec{u}(t)$ is stored at every time point needed.

*Self-check.* Suppose the response functional is $J = S(T)$ (the susceptible population at the final time) and the epidemic has a peak at $t = 30$ in a 90-day simulation. At which time ($t = 5$, $t = 30$, or $t = 80$ would you expect $\lambda_I(t)$ to be largest in magnitude? Explain without any calculation.

*Answer:* $\lambda_I(t)$ at time $t$ measures how sensitive $J = S(T)$ is to a perturbation of $I$ at time $t$. A perturbation to $I$ at $t = 5$ has 85 days to propagate and affect $S(90)$: a large window of time for the perturbation to alter the epidemic trajectory. A perturbation at $t = 80$ has only 10 days. So $\lambda_I(t)$ should be largest at $t = 5$.

## 6.3 ▪ The Lagrange Multiplier Derivation

### 6.3.1 ▪ A Warm-Up: Exponential Decay

Before the five-step derivation, work through the simplest possible case to see exactly what the adjoint ODE looks like and how it gives the answer.

Let $\dot{u} = -pu$ with $u(0) = 1$ and $J = u(T)$. The forward solution is $u(t) = e^{-pt}$, so $J = e^{-pT}$ and $dJ/dp = -Te^{-pT}$ (by direct differentiation; the "correct" answer).

The adjoint approach finds the same result differently. For $F(u; p) = -pu$ and $g = 0$, $h(u) = u$:

- Adjoint ODE: $-\dot{\lambda} = F_u\,\lambda = -p\lambda$, so $\dot{\lambda} = p\lambda$.
- Terminal condition: $\lambda(T) = +h'(u(T)) = +1$.
- Solution (running backward from $T$): $\lambda(t) = +e^{p(t-T)}$.
- Sensitivity formula: $dJ/dp = +\int_0^T \lambda(t)\,F_p\,dt$. Here $F_p = -u$ and $\lambda(t) = e^{p(t-T)}$ (backward from $\lambda(T) = +1$): $dJ/dp = +\int_0^T e^{p(t-T)}(-u(t))\,dt =$

$-\int_0^T e^{p(t-T)} u_0 e^{-pt}\,dt = -u_0 e^{-pT} T = -T e^{-pT}$. ✓

The adjoint gave the same derivative using two ODE solves (one forward, one backward) for this single scalar QOI, with no dependence on $m$ (how many parameters there are). If $p$ were joined by 49 other parameters, the cost would still be two ODE solves plus 50 integrals, one per parameter.

Now we derive this procedure in full generality.

### 6.3.2 ▪ The Five-Step Derivation

The derivation proceeds through five numbered steps, each announced before it is executed. The scalar ODE case is treated fully; the vector case follows by replacing scalars with vectors and matrices.

### 6.3.3 ▪ Setup

Consider the initial value problem

$$\dot{u} = F(u;p), \qquad u(0) = u_0, \tag{6.1}$$

with a response functional

$$J = \int_0^T g(u)\,dt + h(u(T)). \tag{6.2}$$

Here $g$ is the running cost (an integrand depending on the solution $u$ but *not* directly on the parameter $p$) and $h$ is a terminal cost (depending only on the final state). We also assume the initial condition $u_0$ is independent of $p$, so $\partial u_0/\partial p = 0$; this holds for all examples in this chapter. This covers the main cases in practice: total infectious person-time $J = \int I\,dt$, final state $J = u(T)$, and weighted combinations. When $g$ depends explicitly on $p$, an additional term $\int_0^T g_p\,dt$ appears in the sensitivity formula; see Cao et al. [12] for the fully general treatment. We want $dJ/dp$ without solving the FSE.

The strategy is identical to Chapter 5: introduce an auxiliary variable, choose it to cancel the unknown sensitivity $\partial u/\partial p$, and read off the result from what remains. In Chapter 5 the auxiliary variable was a vector $\vec{v}$ satisfying $A^T\vec{v} = \vec{c}$; here it is a function $\lambda(t)$ satisfying a backward ODE.

**Step 1: Introduce the Lagrange multiplier.**

We introduce an auxiliary function $\lambda(t)$, called the adjoint variable or Lagrange multiplier, and observe that, because $u(t)$ satisfies the ODE (6.1):

$$\int_0^T \lambda(t)\big[F(u;p) - \dot{u}\big]\,dt = 0.$$

This is the ODE constraint written as a vanishing integral. Define the augmented functional

$$\tilde{J} = J + \int_0^T \lambda(t)\big[F(u;p) - \dot{u}\big]\,dt. \tag{6.3}$$

Since the bracketed term is zero, $\tilde{J} = J$. The strategy, carried over from Chapter 5, is to choose $\lambda(t)$ so that $\partial u/\partial p$ disappears when we differentiate $\tilde{J}$ with respect to $p$.

*Physical analogy.* Think of a building inspector who records the strain in each beam throughout the day. The inspector's record is $\lambda(t)$: it does not change the building, but it carries a running account of how much each moment in time contributes to the structure's overall stress $J$. Adding the constraint integral is like attaching a running tally that is identically zero as long as the beams obey the equilibrium equations; it costs nothing, but it opens the door to measuring sensitivity.

**Step 2: Differentiate with respect to $p$.**
Denote $w = \partial u/\partial p$. Differentiating (6.3):

$$\frac{d\tilde{J}}{dp} = \int_0^T g_u\, w\,dt + h'(u(T))\,w(T) + \int_0^T \lambda\big[F_u\, w + F_p - \dot{w}\big]\,dt. \tag{6.4}$$

The unknown $w = \partial u/\partial p$ appears in three places. The next step eliminates it by choosing $\lambda$ appropriately.

*Physical analogy.* Differentiating with respect to $p$ is like asking: if we nudge a design parameter, say, the stiffness of one beam; how does the overall stress $J$ respond? The sensitivity $w = \partial u/\partial p$ represents the ripple that nudge sends through the entire structure over time. It appears everywhere because the system is coupled; $\lambda(t)$ will be chosen precisely to absorb those ripples.

**Step 3: Integrate by parts.**
The term $\int_0^T \lambda\dot{w}\,dt$ contains $\dot{w}$. We remove the time derivative from $w$ by integrating by parts: this is the key step, and it is the ODE analog of the matrix transposition $v^T A = (A^T v)^T$ in Chapter 5.

The standard integration-by-parts identity, written out step by step, gives

$$\begin{aligned}\int_0^T \lambda\,\dot{w}\,dt &= \big[\lambda\, w\big]_0^T - \int_0^T \dot{\lambda}\, w\,dt \\ &= \lambda(T)\,w(T) - \lambda(0)\,w(0) - \int_0^T \dot{\lambda}\, w\,dt.\end{aligned} \tag{6.5}$$

The first line is the identity itself; the second evaluates the bracket at the endpoints, exposing two boundary terms that will behave very differently.

Rearranging (6.5) to isolate the sensitivity integral on the right:

$$\frac{d\tilde{J}}{dp} - \int_0^T \lambda\, F_p\, dt = \int_0^T g_u\, w\, dt + h'(u(T))\, w(T)$$
$$- \lambda(T)\, w(T) + \lambda(0)\, w(0) + \int_0^T \dot{\lambda}\, w\, dt + \int_0^T \lambda\, F_u\, w\, dt. \tag{6.6}$$

Collecting the $w$-terms and reading off the condition for them to vanish:

$$\frac{d\tilde{J}}{dp} - \int_0^T \lambda\, F_p\, dt = \underbrace{\big[h'(u(T)) - \lambda(T)\big] w(T) + \lambda(0)\, w(0) + \int_0^T \big[g_u + \dot{\lambda} + \lambda F_u\big] w\, dt}_{\text{terms involving } w = \partial u/\partial p}. \tag{6.7}$$

*Reading the two boundary terms.* The two boundary terms in (6.7) require separate treatment, and each is handled by a different mechanism.

The $t = 0$ term vanishes automatically. The Setup requires $u(0) = u_0$ to be independent of $p$, so

$$w(0) = \frac{du(0)}{dp} = \frac{du_0}{dp} = 0, \qquad \text{hence} \qquad \lambda(0)\, w(0) = 0. \tag{6.8}$$

No condition on $\lambda$ at $t = 0$ is required; the initial time is free.

The $t = T$ term is the opposite situation: it does not vanish on its own. The $-\lambda(T)\, w(T)$ produced by integration by parts meets the $h'(u(T))\, w(T)$ already generated by differentiating $h(u(T))$ in Step 2, and the two combine into a single coefficient of the unknown $w(T)$,

$$\big[h'(u(T)) - \lambda(T)\big]\, w(T). \tag{6.9}$$

This pairing is the whole point of performing IBP: it manufactures a term whose coefficient we control, namely $\lambda(T)$; directly alongside a term whose coefficient we do not. Setting

$$\lambda(T) = +h'(u(T)) \tag{6.10}$$

zeroes the bracket and eliminates $w(T)$ from the sensitivity formula in one stroke. This is the terminal condition that Step 4 will formally adopt.

*Remark* 6.1. This is the moment where the terminal condition is not arbitrary; it is chosen precisely to eliminate the unknown $w(T)$ from the sensitivity formula.

Draft

Integration by parts is not merely an algebraic trick. It is exactly the operation that moves the time derivative from $w$ to $\lambda$, transforming the FSE into the adjoint ODE, just as matrix transposition moved the linear operator from $u$ to $v$ in Chapter 5.

*Physical analogy.* Think of two runners passing a baton around a track. Forward mode carries $\dot{w}$; the baton is held by the sensitivity variable. Integration by parts hands the baton to $\lambda$: now $\dot{\lambda}$ carries the time derivative instead, and $w$ appears without any derivative. This exchange is what makes it possible to choose $\lambda$ to cancel $w$ in the next step; without it, we could not get rid of $\dot{w}$ by specifying an ODE for $\lambda$ alone.

**Step 4: Choose $\lambda$ to eliminate $w$.**

For the sensitivity formula (6.7) to be free of $w$, the coefficient of $w$ must vanish everywhere. This requires three conditions to hold simultaneously.

First, the integrand coefficient of $w(t)$ must vanish for all $t \in (0, T)$:

$$g_u + \dot{\lambda} + \lambda F_u = 0, \quad \text{i.e.,} \quad \dot{\lambda} = -F_u\,\lambda - g_u. \tag{6.11}$$

Written in standard form (with the sign convention consistent with backward integration):

$$\boxed{-\dot{\lambda} = F_u\,\lambda + g_u.} \tag{6.12}$$

Second, the coefficient of $w(T)$ must vanish:

$$h'(u(T)) - \lambda(T) = 0, \quad \text{i.e.,} \quad \lambda(T) = +h'(u(T)). \tag{6.13}$$

Third, the term $\lambda(0)w(0)$ must vanish. Since $u(0) = u_0$ does not depend on $p$, we have $w(0) = \partial u_0/\partial p = 0$, so this term vanishes automatically.

***Terminal Condition: Two Cases That Cover Smooth, Fixed-Horizon QOIs***

The terminal condition $\vec{\lambda}(T) = +\nabla h(u(T))$ covers every smooth QOI evaluated at a fixed horizon, through two cases that combine additively:

(1) *Terminal cost* $J = h(u(T))$: set $\vec{\lambda}(T) = +\nabla h(u(T))$. For $J = S(T)$ (final susceptibles), $\lambda_S(T) = +1$, all others zero. A point evaluation $J = c^\top u(T)$ is this case with $h(u) = c^\top u$, giving $\vec{\lambda}(T) = +c$; it is not a separate rule.

(2) *Running cost* $J = \int_0^T g(u)\,dt$ with no terminal term ($h = 0$): set $\vec{\lambda}(T) = \vec{0}$. This is the case for $J = \int_0^T I\,dt$ (total burden).

The two combine: if $J = h(u(T)) + \int_0^T g\,dt$, add the terminal conditions from (1) and (2). *Memorize these two cases.* They do not cover a QOI defined by an optimization over time, such as the peak prevalence $I_{\max}$ used in the Chapter 9 laboratory, or the time at which it occurs. Such a QOI is not a smooth function of the state at a fixed $T$, and Chapter 8 treats what goes wrong when it is differentiated as though it were.

The terminal condition (6.13) does not arise because we are running the ODE backward. It arises because the response functional involves the final state $h(u(T))$, and differentiating this with respect to $p$ produces a boundary term at $t = T$. This is a fundamental distinction: the terminal condition is imposed by the response functional, not by the direction of integration.

*Physical analogy.* Choosing $\lambda$ to kill every $w$-term is like tuning the damping in a bridge so that any vibration induced by a parameter perturbation is immediately canceled. The terminal condition $\lambda(T) = +h'(u(T))$ is the inspector's final reading: at the end of the day the ledger must close, recording exactly how sensitive the last measured outcome is to the final state. Once $\lambda(t)$ is tuned, $w$ has nowhere to appear in the sensitivity formula; it has been designed out.

**Step 5: Read off the sensitivity formula.**

With $\lambda(t)$ satisfying (6.12)–(6.13), all terms involving $w$ in (6.7) vanish. The boxed formula below is the general result, stated without assuming anything about the initial condition; the assumption made at the start of the derivation, that $u_0$ does not depend on $p$, is what removes its second

term:

$$\boxed{\frac{dJ}{dp} = + \int_0^T \lambda(t)\, F_p(u(t); p)\, dt + \lambda(0)\, \frac{du_0}{dp}.} \tag{6.14}$$

When the initial condition does not depend on $p$, the second term vanishes. The sensitivity with respect to any parameter $p_j$ is then:

$$\frac{dJ}{dp_j} = + \int_0^T \lambda(t)\, \frac{\partial F}{\partial p_j}\, dt. \tag{6.15}$$

One backward ODE solve for $\lambda(t)$ gives all $m$ sensitivities via $m$ scalar integrals, exactly the cost advantage we sought.

*Physical analogy.* The inspector's record $\lambda(t)$ is now complete: one backward sweep through time, consulting the forward solution $u(t)$ at each moment, accumulates everything needed. Reading off $dJ/dp_j$ is then as simple as computing a running total, one scalar integral per parameter, with no additional solves. The backward sweep is the price paid once; the $m$ integrals are the dividend collected once for every parameter in the model.

### 6.3.4 ▪ The Vector Case

For the vector system $\dot{\vec{u}} = \vec{F}(\vec{u}; \vec{p})$ with $\vec{u} \in \mathbb{R}^n$ and functional $J = \int_0^T g(\vec{u})\, dt + h(\vec{u}(T))$, where, as before, $g$ does not depend directly on $\vec{p}$, the derivation is identical with scalars replaced by vectors and $F_u$ replaced by the Jacobian $J_F = \partial \vec{F}/\partial \vec{u}$. The adjoint ODE and terminal condition become:

$$-\dot{\vec{\lambda}} = J_F^T\, \vec{\lambda} + \nabla_u g^T, \tag{6.16}$$

$$\vec{\lambda}(T) = +\nabla_u h^T\big|_{t=T}. \tag{6.17}$$

**Linearity of the adjoint ODE.** Just as Theorem 4.1 showed that the FSE is always linear in the forward sensitivity variable $\partial \vec{u}/\partial p_j$, the adjoint ODE is always linear in $\vec{\lambda}(t)$; even when the forward ODE is nonlinear in $\vec{u}(t)$. The coefficient $J_F^T(\vec{u}(t))$ is a matrix that depends on the forward solution, but not on $\vec{\lambda}$ itself. This linearity is the ODE counterpart of the algebraic linearity from Chapter 5, and it is why the adjoint is tractable: regardless of how complex the forward dynamics are, the adjoint ODE is a linear ODE that can be solved efficiently.

**Sign convention.** We write the adjoint in backward-time form as $-\dot{\vec{\lambda}} = \cdots$ because that is what the derivation gives: the boxed equation (6.12) follows

from collecting the terms in (6.11), and it fixes the sign before any solver is chosen. Software must then match the equation, not the other way round. In particular, integrating with a solver over $[T, 0]$ traverses time downward while solving that same equation; the direction of travel is not a sign change, and an implementation that flips the sign because it runs backward has introduced a second error that happens to cancel the first. The sensitivity formula (6.18) then needs no further sign change when reading off $dJ/dp_j$ from the integral.

The sensitivity formula becomes:

$$\frac{dJ}{dp_j} = + \int_0^T \vec{\lambda}^T \frac{\partial \vec{F}}{\partial p_j}\, dt + \vec{\lambda}(0)^T \frac{\partial \vec{u}_0}{\partial p_j}. \tag{6.18}$$

*Unifying Idea 2.* The derivation is now complete, and this is Unifying Idea 2 in its full generality. The adjoint in Chapter 5 was $A^T\vec{v} = \vec{c}$: the transposed coefficient matrix $A^T$ applied to $\vec{v}$. The adjoint here is $-\dot{\vec{\lambda}} = J_F^T\vec{\lambda} + \nabla_u g^T$: the transposed Jacobian $J_F^T$ applied to $\vec{\lambda}$, with the sign reversal arising from the integration by parts (6.5). The integration by parts is the exact functional-analytic analog of the matrix transposition $A \to A^T$ from Chapter 5. Appendix A.3 makes this analogy precise through the Lagrange identity, and Appendix B.2 derives the sensitivity formula (6.18) rigorously from it. The terminal condition (6.17) is the time-dependent analog of the right-hand side $\vec{c}$ in $A^T\vec{v} = \vec{c}$. This is one idea.

*Self-check.* For the scalar ODE $\dot{u} = -pu$ with $J = u(T)$ (no running cost, $g = 0$, $h(u) = u$): (a) Write the adjoint ODE and terminal condition using equations (6.12) and (6.13). (b) Solve the adjoint ODE analytically. (c) Use formula (6.15) to compute $dJ/dp$. (d) Verify by solving the forward ODE analytically and differentiating.

*Answer:* (a) $F = -pu$, so $F_u = -p$. Adjoint ODE: $-\dot{\lambda} = -p\lambda$, i.e., $\dot{\lambda} = p\lambda$. Terminal condition: $\lambda(T) = +h'(u(T)) = +1$.
(b) $\dot{\lambda} = p\lambda$ with $\lambda(T) = +1$. Integrating forward from $T$ (backward in time): $\lambda(t) = +e^{p(t-T)}$.
(c) $F_p = -u$. Sensitivity formula: $dJ/dp = +\int_0^T \lambda(t)(-u(t))\, dt = -\int_0^T \lambda(t)u(t)\, dt$. The forward solution is $u(t) = u_0 e^{-pt}$. $dJ/dp = -\int_0^T (e^{p(t-T)})(u_0 e^{-pt})\, dt = -u_0 e^{-pT}\int_0^T dt = -u_0 T e^{-pT}$.
(d) $J = u(T) = u_0 e^{-pT}$, so $dJ/dp = -u_0 T e^{-pT}$. ✓

## 6.4 ▪ The SIR Model: Full Adjoint Treatment

We apply the derivation to the SIR model, writing all adjoint equations explicitly and implementing the full calculation in your implementation.

### 6.4.1 ▪ Setup and Response Functional

The SIR model (4.15)–(4.17) with interpretable POIs $\vec{p} = (c, \beta, \tau_R, \tau_m)$:

$$\begin{aligned} F_S &= \nu_m N - \frac{c\beta SI}{N} - \nu_m S, \\ F_I &= \frac{c\beta SI}{N} - (\gamma_R + \nu_m) I, \\ F_R &= \gamma_R I - \nu_m R, \end{aligned}$$

where $\gamma_R = 1/\tau_R$ and $\nu_m = 1/\tau_m$. The response functional is

$$J = \int_0^T I(t)\, dt, \tag{6.19}$$

the total number of infectious person-days: a measure of the total disease burden. Here $g(\vec{u}) = I$, $h = 0$.

### 6.4.2 ▪ The Jacobian

The Jacobian $J_F = \partial(F_S, F_I, F_R)/\partial(S, I, R)$ evaluated at the forward solution $(S(t), I(t), R(t))$:

$$J_F = \begin{pmatrix} -\dfrac{c\beta I}{N} - \nu_m & -\dfrac{c\beta S}{N} & 0 \\ \dfrac{c\beta I}{N} & \dfrac{c\beta S}{N} - \gamma_R - \nu_m & 0 \\ 0 & \gamma_R & -\nu_m \end{pmatrix}. \tag{6.20}$$

### 6.4.3 ▪ Adjoint ODEs

With $g_S = 0$, $g_I = 1$, $g_R = 0$ (since $g = I$), the adjoint system $-\dot{\vec{\lambda}} = J_F^T \vec{\lambda} + \nabla g^T$ written out fully is:

$$-\dot{\lambda}_S = \left(-\frac{c\beta I}{N} - \nu_m\right) \lambda_S + \frac{c\beta I}{N}\, \lambda_I, \tag{6.21}$$

$$-\dot{\lambda}_I = -\frac{c\beta S}{N}\, \lambda_S + \left(\frac{c\beta S}{N} - \gamma_R - \nu_m\right) \lambda_I + \gamma_R \lambda_R + 1, \tag{6.22}$$

$$-\dot{\lambda}_R = -\nu_m \lambda_R. \tag{6.23}$$

Terminal conditions (since $h = 0$):

$$\lambda_S(T) = \lambda_I(T) = \lambda_R(T) = 0. \tag{6.24}$$

These three ODEs are integrated backward from $t = T$ to $t = 0$, using the stored forward solution $(S(t), I(t), R(t))$ to evaluate the time-varying coefficients.

Draft

### 6.4.4 ▪ Sensitivity Formulas

For each POI, the sensitivity is the integral (6.18) evaluated using the stored $\vec{\lambda}(t)$:

$$\frac{dJ}{d\beta} = + \int_0^T \frac{cSI}{N}(\lambda_I - \lambda_S)\, dt, \tag{6.25}$$

$$\frac{dJ}{dc} = + \int_0^T \frac{\beta SI}{N}(\lambda_I - \lambda_S)\, dt, \tag{6.26}$$

$$\frac{dJ}{d\tau_R} = + \int_0^T \left[\frac{\partial F_I}{\partial \tau_R}\,\lambda_I + \frac{\partial F_R}{\partial \tau_R}\,\lambda_R\right] dt = + \int_0^T \frac{I}{\tau_R^2}\,(\lambda_I - \lambda_R)\, dt, \tag{6.27}$$

$$\frac{dJ}{d\tau_m} = + \int_0^T \left[\frac{\partial F_S}{\partial \tau_m}\lambda_S + \frac{\partial F_I}{\partial \tau_m}\lambda_I + \frac{\partial F_R}{\partial \tau_m}\lambda_R\right] dt. \tag{6.28}$$

Normalized sensitivity indices follow from Definition 2.1: $\mathcal{S}^J_{p_j} = (p_j/J)(dJ/dp_j)$.

**Multiple QOIs and correlated outputs.** When the study involves $r > 1$ response functionals, for example both $J_1 = \int_0^T I\, dt$ (total burden) and $J_2 = S(T)$ (final susceptibles), the adjoint is solved once per QOI, giving $r$ backward solutions $\vec{\lambda}^{(1)}(t), \ldots, \vec{\lambda}^{(r)}(t)$. Even when the POIs are independent (uncorrelated inputs), the QOIs are almost always correlated outputs: both $J_1$ and $J_2$ depend on the same forward trajectory $\vec{u}(t)$, so any parameter change affects both simultaneously. This correlation among outputs is handled automatically by the adjoint. Each backward solve $\vec{\lambda}^{(k)}$ captures the full sensitivity of $J_k$ to perturbations of the shared state, without any special treatment of the output correlation. The user does not model or correct for QOI correlation; the adjoint naturally produces consistent, correct sensitivities for all $r$ response functionals by virtue of working through the same forward trajectory.

### 6.4.5 ▪ The Adjoint Variable and Its Interpretation

Figure 6.2 shows $I(t)$ and $\lambda_I(t)$ computed for nominal parameter values over a 90-day simulation.

The adjoint variable $\lambda_I(t)$ has a concrete epidemiological interpretation: it measures the marginal benefit of reducing the infectious population at time $t$. When $\lambda_I$ is large (early epidemic), reducing $I$ by one person-day saves many future infections. When $\lambda_I$ is small (late epidemic), most transmission has already occurred, so the marginal benefit is low.

Put concretely: $\lambda_I(t)$ measures the additional *infectious person-time* accumulated over the rest of the window from having one extra infectious individual present at time $t$, propagated through all secondary, tertiary, and





**Figure 6.2.** *Computed $I(t)$ (solid, left axis) and the adjoint variable $\lambda_I(t)$ (dashed, right axis) for the SIR model at nominal parameter values ($c = 5$, $\beta = 0.06$, $\tau_R = 7$, $N = 1000$) over a 90-day simulation, obtained from a forward solve followed by a backward adjoint solve. The adjoint variable is largest near $t = 0$, when a perturbation to the infectious compartment has the most time to affect the total burden $J = \int I\,dt$, and decays toward zero as $t \to T$. This gives $\lambda_I(t)$ a concrete interpretation as the marginal value of reducing the infectious population at time $t$: interventions applied when $\lambda_I$ is large avert the most infectious person-time.*

downstream infections, under the linearized dynamics about the nominal trajectory. This is a local, first-order statement: it is exact for infinitesimal perturbations and remains a good approximation for small finite perturbations, but does not account for nonlinear feedback as the perturbation grows large. To connect this to the epidemic dynamics: the force of infection $c\beta I(t)/N$ (equation (2.16)) gives the instantaneous probability per day that any one susceptible individual becomes infected. The adjoint variable $\lambda_I(t)$ carries this one step further; it measures the cumulative downstream burden on $J = \int_0^T I\,dt$ attributable to one infectious individual present at time $t$. Early in the epidemic, when the susceptible pool is large and the force of infection $c\beta I(t)/N$ is growing, one extra infection generates many secondary cases; late in the epidemic, when susceptibles are depleted, the same extra infection propagates little further transmission. Quantitatively: one extra infectious person at time $t$ increases the force of infection by $c\beta/N$ per susceptible per day and initiates a cascade of secondary, tertiary, and subsequent infections whose total contribution to $J$ is $\lambda_I(t)$. Note the units. Since $J = \int_0^T I\,dt$ has dimensions of person-time, $\lambda_I = \partial J/\partial I$ has dimensions of *time*, not of a case count. Reading it as "cases averted" requires a separate incidence model, since cumulative prevalence and case count are different quantities. With that caveat, $\lambda_I$ still ranks *when* to act: early action stops a cascade, late action only a trickle. Turning that ranking into

a statement about cost-effectiveness requires a control and a cost model; the relevant object is then $\delta J = \int_0^T \vec{\lambda}(t)^\top F_c(t)\,\delta c(t)\,dt$ for a control $c$, which weights $\vec{\lambda}$ by how the control actually enters the dynamics.

This interpretation is available from the adjoint computation alone, with no additional work.

The terminal condition $\vec{\lambda}(T) = \vec{0}$ is not the same as starting the forward ODE from initial condition $\vec{0}$ and running it in reverse. The two problems have different structures: the forward ODE is driven by $F(\vec{u};\vec{p})$, while the adjoint ODE is driven by $J_F^T\vec{\lambda}$ with a forcing term from $g$. The backward direction comes from the integration by parts, not from reversing time. The distinction matters for implementation: you run `ODE solver` with `tspan = [T, 0]`, not with a negated vector field.

*Unifying Idea 3.* The adjoint variable $\lambda_I(t)$ illustrates Unifying Idea 3: local SA is a linearization. The adjoint variable displays the local sensitivity of $J$ throughout the simulation period. This continuous record of sensitivity is the local SA analog of what global SA asks globally: how does $J$ respond to perturbations at each moment in time? Chapter 9 will ask the same question for perturbations of the parameters across their full uncertainty ranges rather than perturbations of the state at specific times. The adjoint provides the linearization; global SA provides the full nonlinear picture.

*Unifying Idea 1.* The sensitivity indices $\mathcal{S}_{p_i}^J$ computed in this chapter via the adjoint are exactly the normalized derivatives of Unifying Idea 1 from Chapter 1: the ratio of relative changes in $J$ to relative changes in each POI $p_i$. The adjoint is simply a dramatically more efficient algorithm for computing those same derivatives when the model has many POIs and few QOIs.

*Self-check.* The adjoint variable $\lambda_R(t)$ for the recovered compartment satisfies $-\dot{\lambda}_R = -\nu_m\lambda_R$ with $\lambda_R(T) = 0$. (a) Solve this ODE analytically (backward from $T$). (b) What does the solution tell you about the sensitivity of $J$ to perturbations of the recovered compartment?

> *Answer:* (a) $-\dot{\lambda}_R = -\nu_m\lambda_R$ gives $\dot{\lambda}_R = \nu_m\lambda_R$ with $\lambda_R(T) = 0$. Solution: $\lambda_R(t) = 0$ for all $t \in [0,T]$. (The only solution to a linear ODE with zero terminal condition and a coefficient that keeps it at zero is identically zero when the equation has no forcing term.)
> (b) $\lambda_R(t) = 0$ means that perturbations to $R(t)$ have no effect on $J = \int I\,dt$. This makes biological sense: recovered individuals do not contribute to future transmission (in the basic SIR model without waning immunity), so changing $R(t)$ does not change the integral of $I$.

## 6.5 ▪ Discrete vs. Continuous Adjoint

The adjoint ODE (6.16) is the *continuous adjoint*: it is derived by differentiating the continuous ODE and then discretizing for numerical integration. An alternative is the *discrete adjoint*: discretize the ODE first and then adjoint the discrete system. For linear ODEs with fixed step sizes, the two approaches give identical results. For nonlinear ODEs with adaptive step-size control, they can differ.

In practice: if your ODE solver uses fixed step sizes, either adjoint is acceptable. If it uses adaptive step sizes (as `ODE solver` does), the continuous adjoint requires storing the forward solution densely enough to interpolate accurately, while the discrete adjoint requires differentiating through the step-size controller. For the SA computations in this book, the continuous adjoint with dense output storage is the more practical approach. Sirkes and Tziperman [57] discuss these subtleties for geophysical models. Smith [59] develops the adjoint sensitivity analysis procedure (ASAP) for general parameter-dependent dynamical systems in Ch. 8 §8.6, presenting both perturbation and variational derivations within a functional-analytic framework.

*A note on prior treatments.* The first edition of Smith [58] developed the forward (FSAP) and adjoint (ASAP) sensitivity analysis procedures in functional-analytic detail, including supporting appendix material. The second edition references this material as supplemental and focuses Ch. 8 on a more compact treatment grounded in adjoint methods, parameter subset selection, and complex-step approximation. The chapters of the present book complement Smith's treatment by developing the matrix, computation-graph, and ODE adjoint settings as instances of one organizing idea, at a level designed to be the reader's first careful encounter with the adjoint method.

**Neural ODEs.** In 2018, Chen et al. [13] demonstrated that the adjoint calculation derived in this chapter can be used to train certain deep neural networks. The network defines $\vec{F}(\vec{u};\theta)$ for parameters $\theta$; the loss function plays the role of $J$; and the training algorithm integrates the adjoint ODE backward to compute $\partial J/\partial\theta_i$ for all network parameters. The phrase "backpropagation through time," familiar from recurrent neural networks, refers to the discrete version of this same backward integration. The continuous limit is the adjoint ODE derived here.

### Beyond ODEs: A Pointer to the Wider Adjoint Universe

The adjoint ODE in this chapter is the simplest instance of a broader mathematical structure that scales to much larger problems. When the govern-

ing model is a partial differential equation (PDE) rather than an ODE, a groundwater transport model, an atmospheric circulation model, or a structural mechanics simulation; the adjoint becomes a PDE solved backward in both time and space, with boundary conditions imposed at the final time in the same way that the terminal condition $\vec{\lambda}(T) = +\nabla_u h^T$ is imposed here. Cacuci [9] derived this adjoint formalism for nonlinear functional equations in 1981; Giles and Pierce [16] demonstrated its industrial power for aerodynamic design optimization, computing the sensitivity of aircraft drag with respect to hundreds of shape parameters at the cost of one forward and one adjoint solve.

In data assimilation, fitting a dynamical model to time-series observations; the adjoint provides the gradient of the data-mismatch functional with respect to all model parameters simultaneously. This is the mathematical engine behind the 4D-Var algorithm (four-dimensional variational assimilation, the standard method in operational numerical weather prediction) where the model has millions of state variables and the assimilation window spans days to weeks. The MATLAB adjoint in this chapter is the direct analog of all these applications: the mathematical structure (introduce a Lagrange multiplier function, differentiate, integrate by parts, choose the multiplier to cancel the unknown sensitivity) is identical in every setting; only the function space changes. A systematic graduate-level development of this adjoint formalism for general nonlinear functional equations, including PDEs and integral equations, is in Cacuci [10].

## 6.6 ▪ Computational Laboratory: The SIR Adjoint in Practice

This laboratory implements the full forward-adjoint computation for the SIR model and verifies the result against the FSE approach from Chapter 4.

**Setup.** Use the SIR model with POIs $\vec{p} = (c, \beta, \tau_R, \tau_m)$, nominal values $c = 5$, $\beta = 0.06$, $\tau_R = 7$, $\tau_m = 10{,}000$, initial conditions $S_0 = 999$, $I_0 = 1$, $R_0 = 0$, $N = 1000$, and $T = 90$ days. The response functional is $J = \int_0^{90} I(t)\, dt$.

**Part 1: Solve the forward SIR with dense output.** Solve the SIR system using `ODE solver` with option `'Refine', 8` or dense interpolation, storing the solution on a fine uniform grid with spacing $\Delta t = 0.1$ days. Store $(S, I, R)$ at every grid point; this array is needed for the adjoint coefficient evaluation.

**Interpolation (required before the backward solve).** Before integrating the adjoint backward, build interpolants of the forward solution so the adjoint ODE can evaluate $S(t)$, $I(t)$, and $R(t)$ at any time during the back-

ward pass:

```
# Build smooth interpolants from forward solution grid
# (use cubic/pchip or similar smooth interpolant)
S_interp = INTERPOLANT(t_fwd, U[:, S_col])
I_interp = INTERPOLANT(t_fwd, U[:, I_col])
R_interp = INTERPOLANT(t_fwd, U[:, R_col])
# These allow evaluation at any t in [0, T] during the backward adjoint
```

Pass these interpolants into the adjoint function. Skipping this step is the most common cause of errors in the adjoint lab.

**Part 2: Solve the adjoint system backward.** Implement the adjoint ODEs (6.21)–(6.23). Note that these depend on the forward solution $(S(t), I(t), R(t))$, which must be interpolated from the stored grid.

```
# Adjoint ODE for SIR model - integrate BACKWARD from T to 0
# PLUS convention: -d(lambda)/dt = J_F^T * lambda + grad_u(g)
# g = I(t), so grad_u(g) = (0, 1, 0)

FUNCTION sir_adjoint(t, lam, params):
    N = 1000;  gamma_R = 1/tau_R;  nu_m = 1/tau_m
    Sv = S_interp(t);  Iv = I_interp(t);  Rv = R_interp(t)

    iS = 0;  iI = 1;  iR = 2          %% index positions
    lS = lam[iS];  lI = lam[iI];  lR = lam[iR]
    lam_dot = [0, 0, 0]

    # d(lS)/dt  (no running cost term for S)
    lam_dot[iS] = -( (-c*beta*Iv/N - nu_m)*lS + (c*beta*Iv/N)*lI )

    # d(lI)/dt  (+1 from grad_u(g) = 1, PLUS convention)
    lam_dot[iI] = -( (-c*beta*Sv/N)*lS + (c*beta*Sv/N - gamma_R - nu_m)*
                   + gamma_R*lR + 1 )

    # d(lR)/dt  (no running cost term for R)
    lam_dot[iR] = -( -nu_m*lR )

    RETURN lam_dot
END

# Solve: lam(T) = (0,0,0),  tspan = [T, 0]  (backward)
LAM = SOLVE_ODE(sir_adjoint, tspan=[T, 0], y0=[0,0,0])
```

Solve from $t = T$ to $t = 0$ with terminal conditions $\vec{\lambda}(T) = (0, 0, 0)$.

**Part 3: Plot $I(t)$ and $\lambda_I(t)$.** On a single axis, plot the epidemic curve $I(t)$ (left scale) and the adjoint variable $\lambda_I(t)$ (right scale). Annotate the epidemic peak. Confirm that $\lambda_I$ is largest near $t = 0$ and smallest near $t = T$.

**Part 4: Compute sensitivities and produce a tornado plot.** Evaluate the integrals (6.25)–(6.28) numerically using the trapezoidal rule. Normalize to obtain $\mathcal{S}_{p_j}^J$ for all four POIs. Produce a tornado plot and identify which parameter has the greatest influence on the total disease burden.

**Part 5: Verify against Chapter 4 FSE results.** From the Chapter 4 computational lab, retrieve the SI values computed by the FSE at a single time point. Compare with the adjoint-based SIs for $J = \int I\,dt$. Note: the two computations answer *different* questions (the FSE computes $\mathcal{S}_{p_j}^{I(t)}$ at a fixed $t$, while the adjoint computes $\mathcal{S}_{p_j}^{\int I\,dt}$), so the numerical values will differ. Instead, verify that $\mathcal{S}_\beta^J = \mathcal{S}_c^J$ to at least 4 significant figures (the two indices are equal by the product structure of $c\beta$ in the SIR model, but their common value depends on the parameters and is not generally 1) and discuss whether the ratio holds exactly.

**Part 6: Timing comparison.** Time the adjoint computation (Parts 1–4) against the FSE approach from Chapter 4 (solve the 15-equation augmented system once; it carries all four sensitivities). At what number of parameters $m$ would you expect the adjoint to be faster? Does the timing crossover occur near $m = r = 1$, as the solve counts predict ($m+1$ sequential FSE solves against $r + 1$ adjoint solves)?

## 6.7 ▪ Exercises

**Exercise 6.1** *([Bloom L3])*. For the scalar ODE $\dot{u} = ru(1 - u/K)$ (logistic) with $J = \int_0^T u\,dt$ (total population over time):

(a) Write the adjoint ODE and terminal condition.

(b) Show that the adjoint ODE is linear in $\lambda(t)$ even though the forward ODE is nonlinear in $u(t)$.

(c) Explain why the adjoint ODE depends on the forward solution $u(t)$.

(d) In MATLAB, solve the forward and adjoint ODEs simultaneously (forward from 0 to $T$; store; then backward from $T$ to 0) for $r = 0.5$, $K = 100$, $u_0 = 10$, $T = 20$. Plot $u(t)$ and $\lambda(t)$.

**Exercise 6.2** *([Bloom L3])*. State in your own words what integration by parts accomplishes in Step 3 of the Lagrange multiplier derivation, and

why this is analogous to the matrix transposition $A \to A^T$ in Chapter 5. Then verify equation (6.5) directly by differentiating the product $\lambda w$ and integrating over $[0, T]$.

**Exercise 6.3** *([Bloom L4]).* Using your MATLAB implementation from the laboratory:

(a) Compute $\mathcal{S}^J_\beta$, $\mathcal{S}^J_c$, $\mathcal{S}^J_{\tau_R}$, $\mathcal{S}^J_{\tau_m}$ via the adjoint for $J = \int_0^{90} I\,dt$.

(b) Verify that $\mathcal{S}^J_c = \mathcal{S}^J_\beta$ to at least 6 significant figures. Explain this result using the structure of the SIR model.

(c) If you could implement one intervention that reduces a single parameter by 20%, which parameter would you target to most reduce the total disease burden? Justify using the tornado plot.

(d) Compare the computation time with the FSE approach from Chapter 4.

**Exercise 6.4** *([Bloom L4]).* Fill in the computation cost table for the SIR model with $r = 1$ response functional ($J = \int I\,dt$) and $n = 3$ state variables:

| $m$ (number of POIs) | FSE cost (ODE solves) | Adjoint cost (ODE solves) |
|---|---|---|
| 4 | | |
| 10 | | |
| 50 | | |
| 500 | | |

If each ODE solve takes 30 seconds, at what value of $m$ does the adjoint become more than 10 times faster than the FSE?

**Exercise 6.5** *([Bloom L4]).* Explain in plain language, without any equations, why:

(a) The adjoint ODE must be integrated backward from $t = T$ rather than forward from $t = 0$.

(b) The adjoint variable $\lambda_I(t)$ for the SIR model and response $J = \int I\,dt$ is larger near $t = 0$ than near $t = T$.

(c) A public health agency that wants to minimize total disease burden should focus interventions on the early phase of the epidemic, when $\lambda_I$ is large. Use $\lambda_I(t)$ to quantify this recommendation.

(d) The terminal condition $\vec{\lambda}(T) = \vec{0}$ is not the same as running the forward ODE backward with initial condition $\vec{0}$.

**Exercise 6.6** *([Bloom L5]).* Extend the SIR adjoint treatment to the SEIR model:

$$\begin{aligned}\dot{S} &= \nu_m N - c\beta SI/N - \nu_m S,\\ \dot{E} &= c\beta SI/N - (\nu + \nu_m)E,\\ \dot{I} &= \nu E - (\gamma_R + \nu_m)I,\\ \dot{R} &= \gamma_R I - \nu_m R,\end{aligned}$$

with POIs $\vec{p} = (c, \beta, \tau_E, \tau_I, \tau_m)$ (interpretable parameterization: $\nu = 1/\tau_E$, $\gamma_R = 1/\tau_I$, $\nu_m = 1/\tau_m$) and response functional $J = \int_0^T I(t)\,dt$.

Design the complete adjoint SA study, specifying:

(a) The $4 \times 4$ Jacobian $J_F$ at a generic point $(S, E, I, R)$.

(b) All four adjoint ODEs and their terminal conditions.

(c) The sensitivity formulas $dJ/dp_j$ for all five POIs.

(d) The MATLAB implementation strategy (how you would store the forward solution and integrate the adjoint backward).

(e) What additional cost the SEIR adjoint requires compared to the SIR adjoint (in terms of ODE solves and storage).

**Exercise 6.7** *([Bloom L6]).* A climate modeler reports: "We implemented the adjoint to compute sensitivities for our model with 200 parameters, but the results disagree with forward sensitivity analysis by approximately 15% for some parameters. The adjoint must have a bug."

Identify three distinct causes that could produce this discrepancy without any coding error. For each cause:

(a) State the cause precisely.

(b) Describe a diagnostic test that would confirm or rule it out.

(c) Describe the correction if the cause is confirmed.

Draft

## Notes

The adjoint equation for systems of ordinary differential equations, running backward from $t = T$ with a terminal condition determined by the response functional, was established in the nonlinear setting by Cacuci [9]. The formulation adopted in this chapter follows Cao, Li, Petzold, and Serban [12], who derived the adjoint ODE and implemented it in the CVODES package within the SUNDIALS suite. CVODES handles the memory challenge of backward integration by checkpointing the forward trajectory.

The Pontryagin maximum principle, which gave the first systematic treatment of adjoint variables for optimal control problems in continuous time, appears in Pontryagin et al. [43]; the 4D-Var algorithm in numerical weather prediction, which is essentially the adjoint of the atmospheric model applied to data assimilation, is surveyed by Navon [37].

Sirkes and Tziperman [57] discuss the distinction between the *continuous adjoint* (derived from the continuous ODE and then discretized) and the *discrete adjoint* (the exact adjoint of the discrete numerical scheme). For adaptive-step-size solvers these can differ; the continuous adjoint adopted in this chapter is the standard choice in mathematical epidemiology.

The biological interpretation of the adjoint variable $\lambda_I(t)$ as the expected future burden of one additional infection at time $t$ is discussed in the SA context by Wu, Dhingra, Gambhir, and Remais [64].

Draft

Draft

# Chapter 7
# Adjoint Sensitivity in Practice

*Reverse the calculation, and one output can speak about many inputs.*

You need the sensitivities of a complex disease model with 30 parameters. The model is your own code, you wrote it yourself, so it is differentiable. You know from Chapter 6 that the adjoint requires only 2 ODE solves, against the 31 of a *sequential* FSE, one augmented solve per parameter plus the baseline. A *simultaneous* FSE is instead a single solve of dimension $n(1+m)$: cheaper than 31 separate solves, but still growing with $m$, whereas the adjoint cost does not.

But deriving the adjoint equations by hand for 30 coupled nonlinear ODEs would take weeks and would almost certainly introduce errors in the derivation.

Is there a way to obtain the adjoint computation without deriving the adjoint equations by hand? Yes, algorithmic differentiation does this automatically [36]. And there is a way to use it from MATLAB today, without writing a single AD routine from scratch.

## 7.1 ▪ The Annuity Function: A Worked Comparison

Before examining the computational tools, we work through a concrete example where all three methods, forward mode, reverse mode, and symbolic differentiation, can be compared exactly.

### 7.1.1 ▪ The Problem

An annuity pays a fixed amount $P$ per period for $n$ periods at interest rate $r$. The present value is

$$A(P, r, n) = P\,\frac{1-(1+r)^{-n}}{r}. \tag{7.1}$$

We want all three sensitivity indices: $\mathcal{S}_P^A$, $\mathcal{S}_r^A$, $\mathcal{S}_n^A$.

### 7.1.2 ▪ Forward Mode: Three Passes

In forward mode we differentiate the computation one parameter at a time, propagating $\dot{v}_k = \partial v_k/\partial q$ through intermediate values. We trace the computation of $A$ through the following steps:

$$v_1 = (1+r)^{-n}, \quad v_2 = 1 - v_1, \quad v_3 = v_2/r, \quad A = P v_3.$$

For $q = P$: set $\dot{P} = 1$, $\dot{r} = 0$, $\dot{n} = 0$.

$$\dot{v}_1 = 0, \quad \dot{v}_2 = 0, \quad \dot{v}_3 = 0, \quad \dot{A} = \dot{P}\, v_3 + P\, \dot{v}_3 = v_3.$$

So $\partial A/\partial P = v_3 = (1 - (1+r)^{-n})/r$.

For $q = r$: set $\dot{r} = 1$, $\dot{P} = 0$, $\dot{n} = 0$.

$$\dot{v}_1 = -n(1+r)^{-n-1}, \quad \dot{v}_2 = n(1+r)^{-n-1}, \quad \dot{v}_3 = \frac{\dot{v}_2\, r - v_2}{r^2}, \quad \dot{A} = P\, \dot{v}_3.$$

For $q = n$: set $\dot{n} = 1$, $\dot{P} = 0$, $\dot{r} = 0$.

$$\dot{v}_1 = -\ln(1+r)\,(1+r)^{-n}, \quad \dot{v}_2 = \ln(1+r)\,(1+r)^{-n}, \quad \dot{v}_3 = \dot{v}_2/r, \quad \dot{A} = P\, \dot{v}_3.$$

Three passes through the computation graph give three derivatives. Each pass repeats the evaluation of $A$ and adds its own tangent operations, so the total cost grows in proportion to the number of parameters. The operation counts below make this exact.

### 7.1.3 ▪ Reverse Mode: Two Passes

The mechanism is the one shown in Figure 1.3, where the same computation graph is traversed in both directions: forward mode carries a tangent from one input to the output, and reverse mode carries an adjoint from the output back to every input at once. What follows is that picture applied to the annuity. Figure 7.1 draws the annuity's graph in both directions.

In reverse mode, we compute all three derivatives in one backward pass.

Set $\bar{A} = 1$. Work backward through the computation:

$$\begin{aligned}
\bar{P} &= \bar{A}\, v_3 = v_3, \\
\bar{v}_3 &= \bar{A}\, P = P, \\
\bar{v}_2 &= \bar{v}_3 / r, \\
\bar{r} &\mathrel{+}= \bar{v}_3 \cdot (-v_2/r^2), \\
\bar{v}_1 &= -\bar{v}_2, \\
\bar{r} &\mathrel{+}= \bar{v}_1 \cdot \frac{\partial v_1}{\partial r} = \bar{v}_1 \cdot \big(-n(1+r)^{-n-1}\big), \\
\bar{n} &= \bar{v}_1 \cdot \frac{\partial v_1}{\partial n} = \bar{v}_1 \cdot \big(-\ln(1+r)(1+r)^{-n}\big).
\end{aligned}$$

Two points of discipline are visible here. The sweep must visit each node only after every node that consumes it, so $\bar{v}_1$ is assigned before it is used. And $r$ feeds two downstream nodes, $v_1$ and $v_3$, so $\bar{r}$ *accumulates*: the two contributions are added, not assigned. Writing $=$ in place of $+=$ silently drops one path and is among the most common errors in hand-written reverse sweeps.

One backward pass gives $\partial A/\partial P$, $\partial A/\partial r$, and $\partial A/\partial n$ simultaneously. Total cost: two passes, one forward pass to store values and one backward pass to accumulate derivatives, independent of how many parameters $m$ there are. For the annuity the two passes cost $21/6 = 3.5$ times one function evaluation, as counted below.

### 7.1.4 ▪ Symbolic Differentiation

As a third comparison, we differentiate the formula (7.1) by hand (or by a computer algebra system). Define $v = (1+r)^{-n}$. Then $A = P(1-v)/r$, giving:

$$\frac{\partial A}{\partial P} = \frac{1-v}{r}, \quad \frac{\partial A}{\partial r} = P\left(\frac{nv/(1+r) - (1-v)/r}{r}\right), \qquad \frac{\partial A}{\partial n} = P\,\frac{\ln(1+r)\,v}{r}.$$

Symbolic differentiation produces exact closed-form expressions that can be evaluated at any $(P, r, n)$ with a single function evaluation; no passes through the computation graph are required at query time. The cost is paid upfront in the derivation; the payoff is that evaluation is cheap thereafter.

### 7.1.5 ▪ Exact Operation Counts

For the annuity function we can count arithmetic operations exactly for the procedures printed above, though not for a machine. Real implementations





**Figure 7.1.** *Computation graph of the annuity function* (7.1)*:* $A = Pv_3$*, with* $v_1 = (1+r)^{-n}$*,* $v_2 = 1 - v_1$ *and* $v_3 = v_2/r$*. (a) Forward mode seeds one input at a time. Each pass carries a tangent along the blue edges, and the tangent on the gray edges is zero. Three inputs cost three passes. (b) Reverse mode seeds* $\bar{A} = 1$ *and makes one backward sweep. Each edge carries its local partial derivative. The circled numbers give the order of visits: each node is visited only after every node that consumes it. Two paths reach* $r$*, through* $v_1$ *and through* $v_3$*, so* $\bar{r}$ *adds their two contributions. The sweep returns* $\bar{P}$*,* $\bar{r}$ *and* $\bar{n}$*, the three partial derivatives of* $A$*, in one backward pass. Generated by* `generate_fig_annuity_graph.py`*, which checks every partial derivative it draws against finite differences.*

fuse operations, cache intermediates, and manage a tape differently, so these numbers are meant to expose the scaling in $m$, not to predict a runtime. We count each addition, subtraction, multiplication, division, power, logarithm and negation as one operation. A stored intermediate such as $1 + r$, $-n$ or $v_1$ is reused at no cost. A unit seed or a zero tangent is not computed, and $\bar{r}$ starts at zero, so its first contribution is assigned rather than added. The function $A$ requires: 1 addition, 1 negation, 1 power, 1 subtraction, 1 division, 1 multiplication = 6 operations.

Forward mode repeats these 6 primal operations in each pass and adds that pass's tangent operations. Counted line by line, they are 0 for $q = P$ (the seed gives $\dot{A} = v_3$ directly), 3+1+4+1 = 9 for $q = r$, and 3+1+1+1 = 6 for $q = n$. For $m = 3$ parameters: $3 \times 6 + 0 + 9 + 6 = 33$ operations.

Reverse mode records the 6 primal operations once. Its seven sweep

lines cost $0 + 0 + 1 + 4 + 1 + 5 + 4 = 15$ operations, so reverse mode costs $6 + 15 = 21$ operations regardless of $m$.

The ratio: forward mode costs $33/6 = 5.5$ evaluations of $A$ here, close to $2m$, and each added parameter adds a pass. Reverse mode costs $21/6 = 3.5$ evaluations, whatever $m$ is. For the annuity with $m = 3$, forward mode uses $33/21 \approx 1.6$ times as many operations as reverse mode, a small gain with only three parameters. For a climate model with $m = 500$, forward mode needs 500 passes against the 2 of reverse mode, so the gap grows in proportion to $m$. For a neural network with $m = 10^7$ parameters and one scalar loss function, reverse mode still requires only two passes, one forward and one backward. These are the same two passes as for the annuity, while forward mode would require $10^7$ separate passes. This is why reverse-mode AD made modern deep learning computationally feasible.

Symbolic differentiation evaluates $A$ and the three formulas above in 15 operations, sharing $1 + r$, $v$ and $(1 - v)/r$. Centered finite differences make $2m + 1 = 7$ evaluations, or $7 \times 6 = 42$ operations before the differences are formed.

| Method | Passes | Operations (annuity, $m = 3$) | Scales as |
|---|---|---|---|
| Forward mode | $m = 3$ | 33 | $O(m)$ |
| Reverse mode | 2 | 21 | $O(1)$ in $m$ |
| Symbolic diff | 1 (at query time) | 15 | $O(m)$ |
| Finite diff | $2m + 1 = 7$ evals | 42 (evaluations only) | $O(m)$, approx |

At nominal values $P = 1000$, $r = 0.05$, $n = 20$, all methods give: $\mathcal{S}_P^A = 1$, $\mathcal{S}_r^A \approx -0.4240$, $\mathcal{S}_n^A \approx 0.5902$. (The derivation of the analytic formulas is Exercise 7.1.)

*Self-check.* From the annuity formula (7.1), compute $\mathcal{S}_P^A$ analytically using the derivative form $\mathcal{S}_P^A = (P/A)(\partial A/\partial P)$. Verify that it equals 1 for any nominal values of $P$, $r$, $n$.

> *Answer:* $\partial A/\partial P = (1 - (1 + r)^{-n})/r = A/P$. So $\mathcal{S}_P^A = (P/A)(A/P) = 1$.
> This holds for any $r$, $n$: the present value is linear in $P$, so a 1% increase in the payment amount produces exactly a 1% increase in present value, regardless of the interest rate or number of periods.

*The same adjoint computation applies directly to epidemic models.* Replace "payment rate" $P$ with "infection rate" $c\beta SI/N$, replace "present value of payments" $A$ with "total epidemic burden" $J = \int_0^T I\,dt$, and replace the interest rate $r$ with the recovery rate $\gamma_R = 1/\tau_R$. The annuity's terminal condition ($\lambda(T) = 0$ because no payments arrive after the policy expires) becomes the adjoint terminal condition for $J = \int_0^T I\,dt$ (no additional burden after the observation window closes). The backward solve is the same computation in different clothing: one backward solve, after the forward solve, recovers sensitivities with respect to every epidemic parameter at once.

## 7.2 ▪ Algorithmic Differentiation

Algorithmic differentiation (AD), also called automatic differentiation, is the chain rule applied systematically to every arithmetic operation in a computer program. It is neither symbolic differentiation (which manipulates formulas) nor numerical differentiation (which approximates derivatives). It is exact to machine precision and applies to any differentiable code.

Students sometimes expect AD to produce a formula for the derivative, like a computer algebra system would. It does not. AD produces the numerical value of the derivative at a specific input, by accumulating the chain rule through the sequence of arithmetic operations the program performs.

Similarly, reverse-mode AD is not the same as numerical differentiation. A centered-difference estimate of $\partial A/\partial r$ has error $O(h^2)$ from the truncation of the Taylor series; reverse-mode AD gives the exact derivative (to floating-point precision) with no step-size selection required.

### 7.2.1 ▪ Forward Mode

Forward-mode AD attaches a tangent value $\dot{v}_k = \partial v_k/\partial q$ to each intermediate variable $v_k$, for one chosen parameter $q$. At each arithmetic operation, the tangent is updated using the chain rule:

$$v_k = f(v_i, v_j) \implies \dot{v}_k = \frac{\partial f}{\partial v_i}\dot{v}_i + \frac{\partial f}{\partial v_j}\dot{v}_j.$$

At the end of the computation, $\dot{v}_{\text{output}}$ contains $\partial J/\partial q$. Repeat for each parameter: $m$ passes total.

### 7.2.2 ▪ Reverse Mode

Reverse-mode AD attaches an adjoint value $\bar{v}_k = \partial J/\partial v_k$ to each intermediate variable. In the backward pass, these are accumulated using the chain

rule in reverse:

$$v_k = f(v_i, v_j) \implies \bar{v}_i \mathrel{+}= \frac{\partial f}{\partial v_i}\bar{v}_k, \quad \bar{v}_j \mathrel{+}= \frac{\partial f}{\partial v_j}\bar{v}_k.$$

The forward pass computes and stores all intermediate values. The backward pass accumulates all $\partial J/\partial q_i$ simultaneously. Total cost: one forward pass + one backward pass, regardless of $m$.

Reverse-mode AD is the discrete-computation-graph version of the adjoint ODE derived in Chapter 6. The adjoint ODE propagates $\vec{\lambda}(t)$ backward through a continuous system; reverse-mode AD propagates $\bar{v}_k$ backward through a discrete computation graph. The underlying mathematical operation is identical: transposition of the derivative operator.

*Unifying Idea 2.* Reverse-mode AD implements Unifying Idea 2 automatically: the adjoint is the transpose in whatever space you are working in, and AD constructs that transpose through the backward pass without requiring the analyst to derive the adjoint equations by hand. For a model written as differentiable code, this eliminates the main practical barrier to adjoint SA: instead of deriving and implementing the adjoint ODE, the analyst uses an AD library that differentiates the code directly. The adjoint equation is implicit in the reverse accumulation; the analyst never writes it explicitly.

### 7.2.3 ▪ Available Tools

The 2009 source material for this book referenced ADIFOR, DAFOR, and TAMC, tools that are now outdated or unmaintained. Table 7.1 lists the current ecosystem. For a comprehensive survey of AD methods, see Baydin et al. [4] and Margossian [30].

*Self-check.* Explain in one sentence each why: (a) AD is not symbolic differentiation; (b) reverse-mode AD is not the same as centered finite differences; (c) forward-mode AD requires $m$ passes while reverse-mode requires 2.

> *Answer:* (a) Symbolic differentiation manipulates formulas algebraically; AD applies the chain rule numerically to the sequence of operations in the code, producing a number rather than a formula.
> (b) Centered finite differences approximate the derivative with error $O(h^2)$ via $(f(p+h) - f(p-h))/(2h)$; reverse-mode AD computes the exact derivative (to floating-point precision) by accumulating the chain rule backward through the computation graph.
> (c) Forward mode propagates the tangent $\dot{v}_k = \partial v_k/\partial q$ for one parameter $q$ per pass, requiring one pass per parameter. Reverse mode propagates the adjoint $\bar{v}_k = \partial J/\partial v_k$ for one response $J$ per backward pass, giving all $m$ derivatives simultaneously. The backward pass needs the values that one forward pass recorded, so reverse mode takes two passes in all.

| Language | Tool | Mode(s) | Notes |
|---|---|---|---|
| MATLAB | Symbolic Math Toolbox | Symbolic | Exact; limited scalability |
| MATLAB | Deep Learning Toolbox (`dlarray` + `dlgradient`) | Reverse | Reverse-mode AD for differentiable code; designed for deep-learning workflows |
| Python | JAX | Forward + Reverse | NumPy-compatible; JIT compilation |
| Python | PyTorch `autograd` | Reverse | Widely used in ML |
| Julia | ForwardDiff.jl | Forward | High performance |
| Julia | Zygote.jl | Reverse | Source-to-source; flexible |
| C/Fortran | ADOL-C | Forward + Reverse | Mature; still active |
| C/Fortran | Tapenade | Forward + Reverse | Source transformation |

**Table 7.1.** *AD tools in wide use by language at the time of writing. Automatic differentiation (AD) frameworks provide reverse-mode AD for arbitrary differentiable code. For large-scale scientific computing, JAX (Python) or ADOL-C (C/Fortran) are the most commonly used options.*

## 7.3 ▪ The Master Decision Table

With all methods now in hand, Table 7.2 consolidates the decision rules developed across Chapters 3–7.

There is no single universally correct method. The right choice depends on the model, the available tools, and the question being asked. A useful heuristic: start with finite differences (Chapter 3) to get SA results quickly, then switch to the adjoint (Chapter 6 or AD) if the computation time is unacceptable or if $m$ is large.

One principle applies universally: whatever method you use, always sanity-check at least two sensitivity indices against an independent method before trusting the full result. For finite differences, verify one analytic SI if available. For the adjoint, verify two values against finite differences. This cross-check catches the majority of implementation errors.

You should not need to choose between finite differences and the adjoint.

| Model type | $m$ vs. $r$ | Method | Accuracy | Cost |
|---|---|---|---|---|
| Black-box | Any | FD (Chapter 3) | $O(h^2)$ | $2m$+1 evals |
| Analytic, simple | $m \leq r$ | FSE (Chapter 4) | Exact | $m$+1 solves |
| Analytic, diff. code | $m > r$ | AD, reverse | Exact | 2 passes |
| Analytic, manual | $m > r$ | Adjoint (Chapter 6) | Exact | 2 solves |
| Correlated POIs | Any | Reparameterize | – | Depends |
| Near bifurcation | Any | FSE + caution | Exact* | $m$+1 solves |
| Large uncertainty | Any | Global SA (Chapter 9) | Variance | $N(m$+2) evals |

**Table 7.2.** *Master decision table for SA method selection; see Pianosi et al. [42] for a broader SA survey. The primary driver is whether the model is black-box or differentiable, and whether $m > r$ or $m \leq r$. (*Near a bifurcation, the FSE is exact, but an equilibrium's index can grow without bound, and any linear prediction fails for changes that cross the threshold; use the result with caution and consult Chapter 8.)* ***Cost note:*** *"2 passes" (reverse AD) and "2 solves" (analytic adjoint) assume $r = 1$ scalar* QOI. *For $r$* QOI*s the adjoint cost scales as $O(r)$ backward solves; for $m \gg r$ this remains far cheaper than FSE at $O(m)$ forward solves.* ***AD vs analytic adjoint:*** *use reverse-mode AD when the model is implemented in differentiable code and memory allows storing the computation tape. Use the analytic adjoint (Chapter 6) when memory is constrained or the numerical integrator (e.g., any adaptive-step ODE solver) is not itself differentiable. A nonlinear ODE or PDE model is no obstacle in itself: integrated by a fixed-step scheme written in differentiable code, it suits reverse-mode AD (Section 7.4).*

Both serve distinct roles: FD is the right tool for black-box models; the adjoint is the right tool for white-box (differentiable) models with many parameters. They are complementary, not competing. But each tool has failure modes that practitioners encounter repeatedly.

*Biological meaning of the adjoint variable.* When the adjoint row of Table 7.2 is applied to a compartmental disease model such as SIR, the adjoint variable $\lambda_I(t)$ carries a concrete epidemiological interpretation: it measures the additional infectious person-time that would follow from one extra infectious person introduced at time $t$, counting all downstream secondary and subsequent transmission through the end of the observation window. It is not a count of additional cases. As Chapter 6 notes, $\lambda_I = \partial J/\partial I$ has dimensions of time when $J$ is measured in person-time, and converting it to cases averted needs a separate incidence model. As in Chapter 6, that reading holds under the linearization about the nominal trajectory: it is exact for infinitesimal perturbations and degrades once a perturbation is large enough for nonlinear feedback to matter. Biologically, this means $\lambda_I(t)$ is large early in an epidemic (when a single infection can trigger many further cases) and small near the end (when susceptibles are depleted and further

spread is limited). This interpretation, derived directly from the adjoint computation, is developed fully in Chapter 6 (§ 6.4).

## 7.4 ▪ Practical Guidance and Common Pitfalls

### 7.4.1 ▪ Discretize-Then-Adjoint vs. Adjoint-Then-Discretize

When computing adjoint sensitivities for ODE systems, two strategies are available. In the *adjoint-then-discretize* approach (used throughout Chapter 6), the adjoint ODE is derived analytically from the continuous ODE and then solved numerically. In the *discretize-then-adjoint* approach, the ODE is first discretized into a finite-difference recurrence, and the adjoint is then derived for that discrete system.

For linear ODEs with fixed step sizes, both approaches give identical results. For nonlinear ODEs with adaptive solvers, they can differ: the adjoint-then-discretize approach may give sensitivities that disagree with finite-difference verification because the continuous adjoint assumes a smooth forward trajectory that the adaptive solver does not produce. The discretize-then-adjoint approach is always consistent with finite differences but requires differentiating through the step-size control logic, which is non-trivial.

For the applications in this book, the adjoint-then-discretize approach with a fixed-step-size forward solver is the most practical choice. Sirkes and Tziperman [57] discuss the implications for geophysical models where both approaches are used in practice.

### 7.4.2 ▪ Memory Requirements for the Adjoint

The adjoint ODE requires the forward solution $\vec{u}(t)$ stored at every time point where the adjoint ODE coefficients are evaluated. For a system with $n = 100$ state variables solved over 10,000 time steps in double precision, the stored forward solution requires $100 \times 10{,}000 \times 8$ bytes $= 8$ MB, modest. For a climate model with $n = 10^6$ state variables and $10^6$ time steps, the requirement is $8 \times 10^{12}$ bytes $= 8$ TB; infeasible to store in memory.

For large systems, two strategies address this: *Checkpointing* stores the forward solution only at a subset of time points and recomputes intermediate values as needed during the backward pass. *Adjoint-of-checkpoint* schemes (Griewank & Walther [17]) achieve optimal trade-offs between memory and recomputation cost. For the models in this course, dense storage is always adequate.

### 7.4.3 ▪ Adaptive Step-Size Solvers

As noted in Chapter 3, adaptive ODE solvers introduce non-smoothness in the model output as a function of parameters: the step-size controller makes discrete decisions that change discontinuously with the parameter value. The continuous adjoint ODE from Chapter 6 assumes a smooth forward solution; when adaptive solvers introduce discontinuities, the continuous adjoint may give inaccurate results.

The safest approach: use fixed-step-size integration (e.g., `ode4` in your preferred language, a 4th-order Runge–Kutta) for the forward solve when computing adjoint sensitivities. This eliminates the non-smoothness and makes the continuous and discrete adjoints equivalent. Alternatively, use tight tolerances (`RelTol` $\leq 10^{-8}$) with adaptive solvers to minimize the non-smoothness.

*Self-check.* You compute adjoint sensitivities for the SIR model using an adaptive ODE solver for the forward solve, then verify against centered finite differences. The adjoint gives $\mathcal{S}_\beta^J = 1.04$ while centered FD gives 0.87. Name two possible causes of this discrepancy, and describe the diagnostic steps you would take to identify which cause applies.

> *Answer:* Possible causes: (1) The adaptive step-size control in the ODE solver introduces non-smoothness that corrupts the continuous adjoint. (2) The finite-difference step size $h$ was chosen poorly (e.g., too small, causing catastrophic cancellation, or too large, causing truncation error). Diagnostics: (a) Recompute the adjoint with fixed-step-size `ode4`: if the discrepancy disappears, the solver is the culprit. (b) Plot the U-curve for the FD estimate to verify the step size is optimal; if the FD value changes significantly with $h$, the FD is unreliable. Always use finite differences as a sanity check for any adjoint implementation; a discrepancy above 1% warrants investigation.

### 7.4.4 ▪ Discontinuities Break AD

AD requires the model to be differentiable at the nominal parameter values. Common sources of non-differentiability in practice include `if` statements whose conditions depend on parameters,[3] `min` and `max` functions, piecewise-defined model components such as quarantine thresholds or saturation functions with hard cutoffs, and look-up tables interpolated without smoothing. When these are present, AD will silently compute the derivative of one branch, which may be zero or undefined at the branch point. The diagnos-

[3] *If-statements are where smooth calculus goes to lose its temper.* When an `if` condition switches branches as a parameter is perturbed, a nominal evaluation and a perturbed one follow different computational paths, and the resulting "derivative" is the derivative of whichever branch happened to be active, which may be zero or undefined at the transition.

tic: compare AD results against centered finite differences. If they disagree significantly for small $h$, suspect a discontinuity. Replacing piecewise functions with smooth approximations (sigmoid functions, smooth max/min) restores differentiability.

*Self-check.* A disease model contains the piecewise function: $\beta(I) = \beta_0$ if $I < 0.1N$, else $\beta_0/2$. This represents reduced contact when hospitalizations are high. (a) Is this function differentiable with respect to $I$ everywhere? (b) If you run reverse-mode AD through this function at $I = 0.09N$, what derivative will AD report? Is this reliable? (c) Propose a smooth approximation that would make the model fully differentiable.

> *Answer:* (a) No. The function has a jump discontinuity at $I = 0.1N$, so $\partial\beta/\partial I$ is undefined there. (b) At $I = 0.09N$ the condition $I < 0.1N$ holds, so the function takes the $\beta_0$ branch and AD differentiates that branch, which is zero. This is locally correct for $I$ strictly below the threshold, but gives no information about the discontinuity itself, and will be completely wrong for sensitivity with respect to $I$ near the threshold. It is unreliable as a global SA result. (c) Replace the step with a sigmoid: $\beta(I) = \beta_0\big(1 - 0.5\,\sigma(k(I/N - 0.1))\big)$, where $\sigma(x) = 1/(1 + e^{-x})$ and $k > 0$ controls sharpness. As $k \to \infty$ this recovers the discontinuous original; a moderate $k$ (e.g., $k = 100$) approximates it closely while remaining differentiable everywhere.

*Unifying Idea 1.* Every method in this chapter (forward mode, reverse mode, symbolic differentiation, finite differences, the adjoint ODE) computes the same object: Unifying Idea 1, the sensitivity index defined in Chapter 2. The methods differ in computational cost, accuracy, and applicability to different model types, but the result they produce is always the same number: $\mathcal{S}^J_{p_j} = (p_j/J)(dJ/dp_j)$.

## 7.5 ▪ Computational Laboratory: Four Methods, One Model

This laboratory computes the same sensitivity ($\partial J/\partial\beta$ for the SIR model with $J = \int_0^{90} I\,dt$, using all four methods and compares their accuracy, speed, and ease of implementation.

**Part 1: Finite differences (Chapter 3 method).** Use the `sensitivity_jacobia` template from Chapter 3 with the SIR model. Record the estimated $\mathcal{S}^J_\beta$ and the computation time.

**Part 2: Analytic FSE (Chapter 4 method).** Solve the 6-equation augmented system from Chapter 4 for the $\beta$ parameter alone (3 SIR state equations plus 3 sensitivity equations $\partial\dot{S}/\partial\beta$, $\partial\dot{I}/\partial\beta$, $\partial\dot{R}/\partial\beta$). Extract $\partial I(t)/\partial\beta$ at all time points and integrate numerically. Record the result and time.

**Part 3: Adjoint ODE (Chapter 6 method).** Solve the adjoint

system from Chapter 6. Use formula (6.18) to compute $dJ/d\beta$. Record the result and time.

**Part 4: Symbolic sanity check (not part of the timed comparison).** Compute $\mathcal{S}_\beta^{\mathcal{R}_0}$ for $\mathcal{R}_0 = c\beta/\gamma_R$ using a symbolic or AD tool of your choice (e.g., SymPy, Symbolics.jl, MATLAB Symbolic Toolbox). This sanity check verifies that your AD setup returns the analytically known result $\mathcal{S}_\beta^{\mathcal{R}_0} = 1$ (by the monomial rule), confirming your pipeline is working before applying it to the time-dependent integral $J = \int_0^T I\,dt$. Note: symbolic tools compute sensitivities of closed-form expressions, not of ODE solutions over time, so this is a *calibration* step, not a fourth method for computing $dJ/d\beta$.

```
# Symbolic differentiation of R0 = c*beta/gamma_R
# Use your preferred symbolic tool

DECLARE_SYMBOLIC  c, beta, gamma_R  (positive)
R0    = c * beta / gamma_R
SI_b  = (beta / R0) * DIFF(R0, with_respect_to=beta)
       # Expected result: SI_b = 1

# For time-dependent sensitivity via reverse-mode AD:
p_nom = [c=5, beta=0.06, tau_R=7, tau_m=10000]
p_ad  = AD_VAR(p_nom)                    # register as differentiable
J     = SIR_integral(p_ad)               # model must use AD-compatible o
dJdp  = GRAD(J, p_ad)                   # reverse pass
PRINT "AD sensitivity for beta:", dJdp[beta]
```

**Part 5: Comparison table.** Fill in the following table from your measurements:

| Method | $\mathcal{S}_\beta^J$ | Rel. error | Time (s) | Lines of new code |
|---|---|---|---|---|
| Finite differences | | | | |
| Analytic FSE | | | | |
| Adjoint ODE | | | | |
| Automatic differentiation (AD) | | | | |

Discuss: which method is easiest to implement? Most accurate? Fastest? Which would you use in practice for this model?

**Extension: introducing a discontinuity.** Modify the SIR model to include a treatment threshold: if $I(t) > 0.1N$, reduce $\tau_R$ by 20% (faster

treatment when infection is high). Re-run all four methods. Which methods give consistent results? Which methods silently produce wrong answers? How does the U-curve (Chapter 3) change for finite differences near $\beta$?

**Connection to optimization.** The adjoint computed in this chapter is the same object as the Lagrange multiplier in constrained optimization and the costate variable in optimal control. Appendix E develops this connection through LP duality and Pontryagin's minimum principle (the continuous-time optimality condition that generalizes the Euler–Lagrange equations).

## 7.6 ▪ Exercises

**Exercise 7.1** *([Bloom L3]).* For the annuity function (7.1) at nominal values $P = 1000$, $r = 0.05$, $n = 20$:

(a) Compute $A$ numerically.

(b) Compute all three sensitivities $\mathcal{S}_P^A$, $\mathcal{S}_r^A$, $\mathcal{S}_n^A$ analytically by differentiating (7.1).

(c) Implement forward mode by hand for $q = r$ (as in Section 7.1) and verify your answer from (b).

(d) Implement reverse mode for all three parameters in one backward pass and verify all three answers.

(e) Count the number of arithmetic operations in each approach.

**Exercise 7.2** *([Bloom L4]).* For each of the following SA problems, select the most appropriate method from Table 7.2 and justify in two sentences:

(a) A climate model with 800 parameters written in Fortran. Each model run takes 10 hours. You want sensitivities for 2 response functionals.

(b) A pharmacokinetic model with 5 parameters and analytic closed-form solutions. You want sensitivities for 8 drug concentration time points.

(c) A traffic simulation with 40 parameters implemented as a compiled C++ executable. Parameter values can be set via a configuration file. Each run takes 3 minutes.

(d) A 10-parameter epidemic model written in your preferred language with an adaptive ODE solver. You want one response functional. The model contains an `if` statement that triggers a quarantine intervention when $I > 0.05N$.

**Exercise 7.3** *([Bloom L4]).* Using automatic differentiation (AD):

(a) Wrap the function $\mathcal{R}_0(c, \beta, \tau_R) = c\beta\tau_R$ as a differentiable computation and use reverse-mode AD to compute all three partial derivatives at $c = 5$, $\beta = 0.06$, $\tau_R = 7$.

(b) Compare with the analytic derivatives.

(c) Wrap the SIR equilibrium calculation (solve $\vec{f}(\vec{u}) = 0$ numerically) and compute $\partial \bar{I}/\partial\beta$ at the endemic equilibrium. Compare with the FSE result from Chapter 4.

(d) What happens when you try to apply reverse-mode AD through an adaptive-step ODE solver? Explain why, and what the remedy is.

**Exercise 7.4** *([Bloom L5]).* Design the complete SA workflow for a dengue fever transmission model [14] with the following properties: $n = 8$ state variables (human and mosquito compartments), $m = 14$ parameters (mortality rates, biting rates, transmission probabilities, mean durations), $r = 2$ response functionals (basic reproduction number and endemic human infection prevalence), each ODE solve taking approximately 2 minutes.

Specify:

(a) Which SA method you would use and why.

(b) The expected total computation time.

(c) How you would validate the implementation.

(d) How you would handle the fact that two parameters ($\sigma_v$ and $\sigma_h$, the mosquito and human biting rates) are structurally correlated.

(e) What output format you would use to present results to a non-mathematical public health audience.

**Exercise 7.5** *([Bloom L5]).* *Audit exercise.* A manuscript under your review reports sensitivities of a scalar QOI with respect to 600 POIs in a differentiable ODE model. The authors state that they used a central finite-difference scheme "for robustness, since adjoint methods can be numerically delicate," and report a total run time of several days on a cluster.

(a) Using the master decision table of Section 7.3, state which method the problem structure indicates and estimate the cost ratio between it and the scheme the authors used.

(b) The authors' stated reason is not baseless. Give the circumstance in which it would be the right call, and state what evidence the manuscript would need to present for you to accept it here.

(c) The reported indices may well be correct. Explain why a referee should still ask for a change, and distinguish that request from a claim that the results are wrong.

(d) Draft the two sentences you would write in your referee report.

**Exercise 7.6** *([Bloom L6]).* A research group uses reverse-mode AD to compute adjoint sensitivities for a mosquito population model with adaptive-step-size ODE integration. They report that some sensitivity indices agree with finite differences but others disagree by 10–20%, with no obvious pattern. The discrepancies are largest for parameters that appear in the mortality rate expressions.

(a) Identify the most likely technical cause of the discrepancy.

(b) Describe two diagnostic tests that would confirm or rule out this cause.

(c) Propose two remedies and explain the trade-offs between them.

(d) At what point would you conclude that the disagreement indicates a genuine SA result (the linearization is poor for these parameters) rather than a numerical artifact?

## Notes

Algorithmic differentiation was introduced independently by Wengert (1964) for forward mode and by Speelpenning (1980) for reverse mode. The definitive reference is Griewank and Walther [17], which covers operator overloading, source transformation, checkpointing, and the relationship between AD and the adjoint method. Naumann [36] treats the combinatorial and software aspects of AD in depth.

The connection between reverse-mode AD and backpropagation in deep learning is made precise by Baydin et al. [4], who give a unified treatment and a historical account of the parallel development of these ideas in scientific computing and machine learning. Margossian [30] surveys reverse-mode AD implementations from the perspective of statistical computing, including Stan and TensorFlow. The neural ODE framework of Chen et al. [13]

applies the continuous adjoint of Chapter 6 to learn the right-hand side of the ODE from data, connecting Chapters 6 and 7.

The financial mathematics application developed in the exercises, computing sensitivities of present-value functions (the "Greeks" in option pricing), was one of the early driving applications of AD in industry. Mester, Mukherjee, and Strang [33] give an accessible survey of differential programming and its applications in modern scientific machine learning.

Draft

Draft

# Part V

# Limitations and Extensions

Draft

Draft

# Chapter 8

# Caveat Emptor, Limitations of Local Sensitivity Analysis

*Local sensitivity is a linear truth, and nature is rarely linear for long.*

In 2019, Saltelli et al. [48] reviewed sensitivity analyses published in 280 highly cited papers across top environmental modeling journals. More than half used the wrong method for the question being asked. Specifically, studies that asked global questions, how does output uncertainty relate to input uncertainty across the full parameter range?; used local methods: derivatives computed at a single nominal parameter value. The derivatives were computed correctly. But they answered a different question than the one the authors claimed to answer.

The deeper problem is not computational but epistemological: sensitivity analysis is still often defined by method rather than purpose [46]. Practitioners reach for the method from their own disciplinary tradition rather than the one best suited to their question. A local method applied to answer a global question will produce results that are internally correct but wrong for the stated purpose. Before choosing an SA method, two questions must be answered: what is the goal; factor prioritization (which inputs most reduce output variance?), factor fixing (which inputs can be frozen without changing the output?), or factor mapping (which inputs drive the output above a threshold?), and do the method's assumptions match the model's behavior?

*Assuming local sensitivity holds globally is the triumph of mathematical hope over nonlinear reality.*

This chapter asks the question that should be asked before any SA is begun: is local SA the right tool for this question? *Unifying Idea 4.* A

second question travels with it. When an index misbehaves near a threshold, the reader has to decide whether the system is genuinely poised there or whether the model has been pushed outside the range in which it represents anything. The mathematics cannot settle that; only knowledge of the system can.

## 8.1 ▪ Method-Question Mismatch

Local sensitivity analysis is built on a derivative. A derivative is a local object: it measures the behavior of a function at one point, and it is only meaningful in the vicinity of that point. This makes local SA powerful for some questions and completely wrong for others.

Local SA can answer: *which parameter most affects the model near the nominal parameter values?*

Local SA cannot answer: *which parameter most affects the model across its full range of uncertainty?* That question requires the parameter to actually vary across its range, and a derivative does not do that.

Local SA cannot answer: *how does the model behave at parameter values far from the nominal?* The derivative approximates linear behavior; real models are often highly nonlinear.

The principle is simple and unforgiving. The method must match the question. Using local SA to answer a global question is not merely suboptimal; it can produce a ranking of parameter importance that is entirely wrong. A parameter that appears unimportant locally may dominate the output variance when varied across its full uncertainty range. A parameter that appears important locally may have negligible effect at realistic perturbation sizes if the model is highly nonlinear in that parameter.

A small sensitivity index does not mean a parameter can be ignored globally. It means only that a small relative change in the parameter predicts a small relative change in the output. Either the slope is small, or the nominal value of the parameter sits near zero. Section 8.5 shows a specific example where the local ranking is the opposite of the global ranking.

*Self-check.* A published SA study reports: "All sensitivity indices are less than 0.05, confirming that the model is robust to parameter uncertainty." The study used nominal parameter values from the literature, with no information on parameter uncertainty ranges. Identify two specific claims in

this statement that go beyond what local SA can actually establish.

*Answer:* (1) "Robust to parameter uncertainty": local SA says nothing about parameter uncertainty. It describes behavior near the nominal value. If the uncertainty range for some parameters is large (e.g., ±50%), the model could be highly sensitive to that parameter globally even though the local SI is small. (2) The conclusion depends implicitly on the nominal parameter values chosen. If the nominal values are changed, the local SI values change too (for nonlinear models). A study with different nominal values might find very different rankings. This sensitivity of local SA to the choice of nominal values is itself a limitation that the study does not address.

## 8.2 ▪ Bifurcations and SA Failure

Local SA requires the model output to be differentiable with respect to the parameters at the nominal values. When this fails, local SA fails completely. The most important way it fails is at a bifurcation.

A bifurcation is a parameter value at which the qualitative behavior of the model changes: two equilibria merge into one, a stable fixed point becomes unstable, or an epidemic threshold is crossed [61, 28]. At a fold the derivative $\partial u/\partial p$ is infinite, and at a transcritical point the normalized index diverges. Local SA breaks down entirely. Why it breaks down depends on what is differentiated, and Section 8.4.1 sorts the cases. For an equilibrium at a fold or transcritical point, the Jacobian is singular at the bifurcation itself, so the FSE $D_{\vec{u}}[\vec{f}\,](\partial\vec{u}/\partial p) = \vec{r}$ may have no solution or infinitely many (the Fredholm alternative, Appendix B.3). Near it the Jacobian is only nearly singular, and the FSE has a unique solution, which can be ill-conditioned. The time-dependent FSE, a linear initial-value problem, always has a unique solution. For models with periodic limit cycles, sensitivity near a bifurcation of the periodic orbit involves Floquet multipliers; this is treated in Appendix D.

### 8.2.1 ▪ The Quadratic Example

Consider the scalar equation $f(u;p) = u^2 + p = 0$. For $p < 0$, there are two real roots: $u = \pm\sqrt{-p}$. At $p = 0$, the two roots merge. For $p > 0$, there are no real roots.

The FSE for the sensitivity of the positive root $u^* = \sqrt{-p}$ is:

$$\frac{\partial u^*}{\partial p} = \frac{-\frac{\partial f}{\partial p}}{\frac{\partial f}{\partial u}} = \frac{-1}{2u^*} = \frac{-1}{2\sqrt{-p}}.$$

The normalized SI is:

$$\mathcal{S}_p^{u^*} = \frac{p}{u^*}\frac{\partial u^*}{\partial p} = \frac{p}{\sqrt{-p}} \cdot \frac{-1}{2\sqrt{-p}} = \frac{1}{2}.$$

This is constant (independent of $p$) for all $p < 0$; the positive root is always in exact proportion $1/2$ to any fractional change in $p$.

But as $p \to 0^-$:

$$\frac{\partial u^*}{\partial p} = \frac{-1}{2\sqrt{-p}} \to -\infty.$$

The derivative diverges at the bifurcation. Near $p = 0$, the model is infinitely sensitive to the parameter $p$: any small positive perturbation to $p$ destroys the equilibrium entirely.

*Unifying Idea 3.* This is Unifying Idea 3 in its most dramatic form. The sensitivity index is the derivative-based linearization of the model response. At a fold such as this one, the function $u^*(p)$ has infinite curvature, its graph has a vertical tangent, and no linear approximation is valid. The linearization does not just become inaccurate at the fold; it becomes undefined.

*Unifying Idea 1.* The index is a normalized derivative, so it can fail through either factor. At a fold the raw derivative $\partial u^*/\partial p$ becomes infinite. The normalized index then diverges as well whenever the fold sits at a nonzero parameter value, as in Exercise 8.1, whose fold is at $p = 1$. The quadratic above is the exception: its fold sits at $p = 0$, so the factor $p$ in the normalization cancels the divergence and the index stays $1/2$. At a transcritical bifurcation the raw derivative stays finite, but the equilibrium it is normalized against approaches zero, so the normalized index diverges. Section 8.4.1 sorts these cases by what is differentiated.

*Unifying Idea 2.* The adjoint route inherits the same difficulty, by a mechanism that depends on what is being differentiated. For the sensitivity of an *equilibrium*, the adjoint system is algebraic and inherits the singularity directly: the transpose of a singular matrix is also singular, so at a fold or transcritical point the adjoint system has no unique solution. For a time-dependent functional over a *finite* horizon the adjoint ODE remains solvable, but the near-zero eigenvalue of $J_F$ leaves the adjoint with a slowly decaying mode, so $\lambda$ can grow large as the horizon lengthens. Forward and adjoint approaches fail together, for the same underlying reason.

*Self-check.* For $f(u; p) = u^2 + p$ with $p = -4$: (a) Find the two roots $u^*_\pm$. (b) Compute $\mathcal{S}_p^{u^*_+}$ analytically. Compare with the derivative $\partial u^*_+/\partial p$. (c) Estimate $\partial u^*_+/\partial p$ numerically using a centered difference with $h = 0.1, 0.01, 0.001$. At what value of $p$ near zero does the finite-difference esti-

mate become unreliable?

*Answer:* (a) $u_{\pm}^* = \pm\sqrt{4} = \pm 2$. (b) $\partial u_+^*/\partial p = -1/(2\sqrt{-p}) = -1/4$. $\mathcal{S}_p^{u_+^*} = (p/u_+^*)(-1/(2\sqrt{-p})) = (-4/2)(-1/4) = 1/2$.
(c) Numerically at $p = -4$: all three step sizes give $\approx -0.250$ ✓. As $p \to 0^-$, the function $u_+^* = \sqrt{-p}$ has a square-root singularity; finite differences become inaccurate when the curvature term dominates the step-size error, which occurs when $h \gtrsim |p|$. Near $p = 0$, even very small $h$ is not small compared to $|p|$, so finite differences fail.

### 8.2.2 ▪ The Epidemic Threshold

The quadratic example above illustrates a fold bifurcation (root collision): two equilibria merge and annihilate as a parameter crosses zero. The epidemic threshold is a structurally different bifurcation, a *transcritical* bifurcation in which two equilibria exchange stability rather than colliding. Both defeat local SA, but for reasons worth separating. At a fold, the *raw derivative* of the equilibrium diverges. At a transcritical bifurcation the raw derivative stays finite; what diverges is the *normalized* index, because the quantity it is normalized against is itself going to zero. Either way, the linearization stops being a fair description of the model near the threshold.

The SIS (*Susceptible–Infectious–Susceptible*) model gives the cleanest version of this behavior, because its endemic equilibrium has a closed form. Infectious individuals return directly to the susceptible class rather than gaining permanent immunity [63]. In a closed population of size $N$:

$$\dot{S} = -\frac{\beta S I}{N} + \gamma_R I, \qquad \dot{I} = \frac{\beta S I}{N} - \gamma_R I, \tag{8.1}$$

with $S + I = N$ constant. Writing $i = I/N$ and substituting $S = N(1 - i)$ gives the scalar equation $\dot{i} = i\big(\beta(1 - i) - \gamma_R\big)$, whose equilibria are the disease-free state $i^* = 0$ and, for $\mathcal{R}_0 > 1$,

$$i^* = 1 - \frac{\gamma_R}{\beta} = 1 - \frac{1}{\mathcal{R}_0}, \qquad \mathcal{R}_0 = \frac{\beta}{\gamma_R}. \tag{8.2}$$

The disease-free state is stable for $\mathcal{R}_0 < 1$ and the endemic equilibrium is stable for $\mathcal{R}_0 > 1$. The two exchange stability at $\mathcal{R}_0 = 1$: this is the transcritical bifurcation.

The sensitivity of the endemic level to $\mathcal{R}_0$ is

$$\frac{\partial i^*}{\partial \mathcal{R}_0} = \frac{1}{\mathcal{R}_0^2}, \qquad \mathcal{S}_{\mathcal{R}_0}^{i^*} = \frac{\mathcal{R}_0}{i^*} \cdot \frac{1}{\mathcal{R}_0^2} = \frac{1}{\mathcal{R}_0(1 - 1/\mathcal{R}_0)} = \frac{1}{\mathcal{R}_0 - 1}.$$

The raw derivative is well behaved at the threshold: it tends to 1 as $\mathcal{R}_0 \to 1^+$. The normalized index is not. It diverges, $\mathcal{S}_{\mathcal{R}_0}^{i^*} \to +\infty$, precisely because $i^*$ is





**Figure 8.1.** *Bifurcation of the SIS endemic equilibrium at $\mathcal{R}_0 = 1$. (a) The infectious fraction $i^* = \max(0, 1 - 1/\mathcal{R}_0)$: zero (disease-free state, stable) for $\mathcal{R}_0 < 1$, rising continuously for $\mathcal{R}_0 > 1$ (endemic equilibrium, stable). The kink at $\mathcal{R}_0 = 1$ marks the transcritical bifurcation. (b) The sensitivity index $\mathcal{S}^{i^*}_{\mathcal{R}_0} = 1/(\mathcal{R}_0 - 1)$ diverges as $\mathcal{R}_0 \to 1^+$. Any parameter that shifts $\mathcal{R}_0$ near the epidemic threshold produces an arbitrarily large proportional change in the endemic level. Local SA correctly captures this infinite sensitivity but cannot predict the qualitative shift from disease-free to endemic behavior; that requires knowing that a bifurcation exists.*

vanishing. Figure 8.1 shows both the equilibrium branch and the diverging sensitivity index.

SA of the endemic equilibrium for $\mathcal{R}_0$ values close to 1 must be interpreted with extreme caution. A small change in any parameter that affects $\mathcal{R}_0$ could eliminate the endemic equilibrium entirely or cause it to surge. Local SA correctly identifies this extreme sensitivity but cannot predict the qualitative change.

For $\mathcal{R}_0 \leq 1$: SA of the endemic equilibrium is undefined; the equilibrium does not exist. Computing SA under the assumption that $i^*$ exists when $\mathcal{R}_0 < 1$ is a mathematical error. Depending on how the equilibrium is formulated and solved, a numerical routine may return a negative or complex value without warning.

**Sensitivity to the individual parameters.** Differentiating (8.2) with respect to $\beta$ and $\gamma_R$:

$$\frac{\partial i^*}{\partial \beta} = \frac{\gamma_R}{\beta^2}, \qquad \frac{\partial i^*}{\partial \gamma_R} = -\frac{1}{\beta}.$$

The normalized sensitivity indices are:

$$\mathcal{S}^{i^*}_{\beta} = \frac{\partial i^*}{\partial \beta} \cdot \frac{\beta}{i^*} = \frac{\gamma_R/\beta^2 \cdot \beta}{1 - 1/\mathcal{R}_0} = \frac{1/\mathcal{R}_0}{(\mathcal{R}_0 - 1)/\mathcal{R}_0} = \frac{1}{\mathcal{R}_0 - 1}, \tag{8.3}$$

$$\mathcal{S}^{i^*}_{\gamma_R} = \frac{\partial i^*}{\partial \gamma_R} \cdot \frac{\gamma_R}{i^*} = \frac{-1/\beta \cdot \gamma_R}{1 - 1/\mathcal{R}_0} = -\frac{1}{\mathcal{R}_0 - 1}. \tag{8.4}$$

Both indices diverge as $\mathcal{R}_0 \to 1^+$, confirming that threshold sensitivity failure is a structural property of the bifurcation, not a feature specific to SIR.

**A structural constraint.** Notice that $\mathcal{S}_\beta^{i^*} + \mathcal{S}_{\gamma_R}^{i^*} = 0$ for all $\mathcal{R}_0 > 1$. This is not a coincidence. Since $i^* = 1 - \gamma_R/\beta$ depends on $\beta$ and $\gamma_R$ only through their ratio, rescaling both parameters by the same factor leaves $i^*$ unchanged: $i^*$ is homogeneous of degree 0 in $(\beta, \gamma_R)$. Euler's theorem for homogeneous functions then gives $\beta\,\partial i^*/\partial\beta + \gamma_R\,\partial i^*/\partial\gamma_R = 0$, which is exactly $\mathcal{S}_\beta^{i^*} + \mathcal{S}_{\gamma_R}^{i^*} = 0$. In biological terms: if transmission and recovery rates both double, the endemic level is unchanged, so their sensitivities must cancel. This is a case where the sum-to-zero property from Chapter 2 arises from a structural symmetry in the model rather than a normalization identity.

**Numerical illustration.** Table 8.1 shows how the sensitivity indices grow as $\mathcal{R}_0$ approaches 1 from above, using nominal parameters $\beta = 0.3$ day$^{-1}$, $\gamma_R = 0.1$ day$^{-1}$ ($\mathcal{R}_0 = 3$) as a baseline and reducing $\beta$ to move $\mathcal{R}_0$ toward threshold.

| $\mathcal{R}_0$ | $i^*$ | $\mathcal{S}_\beta^{i^*} = -\mathcal{S}_{\gamma_R}^{i^*}$ | Interpretation |
|---|---|---|---|
| 3.0 | 0.667 | +0.5 | Low: standard operating range |
| 1.5 | 0.333 | +2.0 | Elevated: 1% change ⇒ 2% change in $i^*$ |
| 1.2 | 0.167 | +5.0 | High: treat as warning |
| 1.1 | 0.091 | +10.0 | Very high: ranking unreliable |
| 1.01 | 0.010 | +100. | Near-threshold: SA has broken down |

**Table 8.1.** *Sensitivity index $\mathcal{S}_\beta^{i^*} = 1/(\mathcal{R}_0 - 1)$ for the SIS endemic equilibrium, with $\mathcal{S}_{\gamma_R}^{i^*} = -\mathcal{S}_\beta^{i^*}$. As $\mathcal{R}_0 \to 1^+$, both indices diverge and local SA loses quantitative meaning. The practical threshold for caution is $|\mathcal{S}^{i^*}| > 10$, i.e. $\mathcal{R}_0 < 1.1$.*

**The same conclusion for the SIR model with demography.** The SIR model of Chapter 4, Equations (4.15)–(4.17), also has a transcritical bifurcation at $\mathcal{R}_0 = 1$. Its endemic level is

$$i^* = \frac{\nu_m}{\gamma_R + \nu_m}\left(1 - \frac{1}{\mathcal{R}_0}\right), \tag{8.5}$$

which carries a prefactor the SIS model does not. That prefactor is small whenever the mean residence time $\tau_m = 1/\nu_m$ is long compared with the infectious period, so the endemic level itself is small and is approached only

over demographic timescales. It has no effect on the index with respect to $\mathcal{R}_0$. Any constant prefactor cancels in the normalization,

$$\mathcal{S}^{i^*}_{\mathcal{R}_0} = \frac{\mathcal{R}_0}{k(1-1/\mathcal{R}_0)} \cdot \frac{k}{\mathcal{R}_0^2} = \frac{1}{\mathcal{R}_0 - 1} \qquad \text{for any } k > 0,$$

and since $\mathcal{R}_0 = c\beta/(\gamma_R + \nu_m)$ is proportional to $c$ and to $\beta$, the chain rule gives $\mathcal{S}^{i^*}_{\beta} = \mathcal{S}^{i^*}_{c} = 1/(\mathcal{R}_0 - 1)$ as well. These indices transfer unchanged to the model used throughout the book. This scale-invariance is worth noting in its own right: a normalized sensitivity index is blind to the overall size of the response it describes, which is exactly what makes it comparable across parameters with different units, and exactly what makes it fragile when the response approaches zero. The recovery-rate index does not transfer, because the prefactor itself depends on $\gamma_R$:

$$\mathcal{S}^{i^*}_{\gamma_R} = -\frac{\gamma_R}{\gamma_R + \nu_m} \cdot \frac{\mathcal{R}_0}{\mathcal{R}_0 - 1},$$

which is $-1.91$ at the book's nominal values, where the SIS form (8.4) would give $-0.91$. The structural constraint survives over three rates. With $\mathcal{R}_0 = c\beta/(\gamma_R + \nu_m)$ the endemic level is $i^* = \nu_m/(\gamma_R + \nu_m) - \nu_m/(c\beta)$, which is homogeneous of degree 0 in $(\beta, \gamma_R, \nu_m)$, so Euler's theorem gives $\mathcal{S}^{i^*}_{\beta} + \mathcal{S}^{i^*}_{\gamma_R} + \mathcal{S}^{i^*}_{\nu_m} = 0$. Both threshold indices still diverge as $\mathcal{R}_0 \to 1^+$.

**Operational rule: when to stop reporting SI as a ranking.** A sensitivity index is meaningful as a ranking tool only when the linearization is a fair local approximation. Two practical rules of thumb (these are heuristic thresholds, not mathematical guarantees; the appropriate cutoff depends on the model and the application):

1. *Magnitude check.* If $|\mathcal{S}^{q}_{p}| > 10$ for a normalized index, treat the result as a qualitative warning ("this parameter region is highly sensitive") rather than a quantitative ranking. An SI of 100 does not mean the parameter matters 10× more than one with SI 10; it may mean the model is near a threshold.
2. *Condition-number check.* Compute $\kappa(J_F)$ (the condition number of the model Jacobian at the operating point). If $\kappa(J_F) > 10^4$, the sensitivity computation is ill-conditioned: small errors in the model evaluation may be amplified by $\kappa(J_F)$ in the sensitivity values. Always report $\kappa(J_F)$ alongside SI values computed near a bifurcation or near parameter degeneracy. The check needs a Jacobian with more than one state and no conserved total (§8.3 defines $\kappa$). The scalar SIS equation has a $1 \times 1$ Jacobian, so $\kappa = 1$ at every $\mathcal{R}_0$, and in the two-equation form (8.1) the conserved total $S + I = N$ makes $J_F$ singular

at every state. For a one-state model the threshold shows in the size of the Jacobian instead, $|J_F|/\gamma_R = \mathcal{R}_0 - 1 = 1/\mathcal{S}_\beta^{i^*}$, so the magnitude check already catches it. For the SIR model with demography (laboratory Part 4) the check works: $\kappa_2(J_F)$ at the endemic equilibrium is about $4 \times 10^3$ at $\mathcal{R}_0 = 3$, passes $10^4$ near $\mathcal{R}_0 = 1.35$, and reaches about $3 \times 10^5$ at $\mathcal{R}_0 = 1.01$.

**Takeaway.** The divergence of local SA near $\mathcal{R}_0 = 1$ is a property of the transcritical bifurcation structure, not of the specific model. A compartmental epidemic model with a generic forward transcritical bifurcation at $\mathcal{R}_0 = 1$ will exhibit the same behavior. (Models with a backward bifurcation, or with non-smooth incidence, need separate treatment; their threshold structure differs.) The magnitude check ($|\mathcal{S}^q| > 10$) applies equally to SIS, SEIR, and any other model of this type. The condition-number check applies where rule 2 says it does, and Section 8.4.1 sorts which check suits which quantity.

## 8.3 ▪ Ill-Conditioning

For linear algebraic models, trouble near a singularity takes a different form. When the coefficient matrix $A$ in $A\vec{u} = \vec{b}$ is nearly singular, small changes in the right-hand side $\vec{b}$ (or in the matrix entries) can produce large changes in the solution. Those large sensitivities are genuine: the exact derivatives $\partial\vec{u}/\partial\vec{b}$ are the entries of $A^{-1}$, which grow as $A$ nears singularity. What ill-conditioning adds is a separate cost: the computed solution, and every index derived from it, can lose accuracy.

### 8.3.1 ▪ The Condition Number

The *condition number* $\kappa(A) = \|A\|\|A^{-1}\|$ (see Appendix C.2 for the SVD-based definition $\kappa_2 = \sigma_1/\sigma_n$) measures how much the solution $\vec{u}$ can change relative to a change in the data. For the 2-norm,

$$\kappa_2(A) = \frac{\sigma_{\max}(A)}{\sigma_{\min}(A)}, \tag{8.6}$$

where $\sigma_{\max}$ and $\sigma_{\min}$ are the largest and smallest singular values of $A$. A matrix with $\kappa_2(A) = 1$ is perfectly conditioned (orthogonal, or a nonzero scalar multiple of an orthogonal matrix); a matrix approaching singularity has $\sigma_{\min} \to 0$ and $\kappa_2(A) \to \infty$.

*Remark* 8.1. Readers from statistics will recognize the condition number as the SA analog of the *variance inflation factor* (VIF). In regression, a large VIF warns that the design matrix $X^\top X$ is nearly singular and that

coefficient estimates are unreliable; in SA, a large $\kappa(A)$ issues the same warning about the sensitivity Jacobian. Both diagnose a nearly degenerate linear system in which "separate" effects cannot be cleanly estimated.

The key bound: if the data changes by a relative amount $\epsilon$, the solution changes by at most $\kappa(A)\,\epsilon$:

$$\frac{\|\delta\vec{u}\|}{\|\vec{u}\|} \leq \kappa(A)\,\frac{\|\delta\vec{b}\|}{\|\vec{b}\|}.$$

When $\kappa(A) \gg 1$, even tiny changes in the data produce large changes in the solution.

### 8.3.2 ▪ SA and Ill-Conditioning

For SA, ill-conditioning raises two questions that are easy to conflate. A colleague who reports "the sensitivity of $u_1$ to $b_1$ is extremely large" is reporting two things at once: (a) a genuine model property: the exact sensitivities are entries of $A^{-1}$, which are large when $A$ is nearly singular, so the output truly varies strongly with this parameter; and (b) a warning about accuracy: the computed index, like the computed solution, may have lost digits.

The two hold together and are independent, and $\kappa(A)$ answers (b). If $\kappa(A) \approx 10^6$, then numerical errors of size $\varepsilon_{\text{mach}} \approx 10^{-16}$ can contaminate the solution at the level of $10^{-10}$: 6 digits of the solution may be meaningless. A small product $\kappa(A)\,\varepsilon_{\text{mach}}$ therefore clears a large computed index. A large $\kappa(A)$ cannot mark one as an artifact, because it also means a large exact sensitivity in some direction. Nor can a small residual $\|\vec{b} - A\hat{\vec{u}}\|$: a backward-stable solve leaves a small residual at any $\kappa(A)$. To check the digits themselves, recompute in higher precision or compare with an analytic value (§8.4.1).

Ill-conditioning is itself a SA result: it warns the investigator that the computed solution, and any sensitivity derived from it, is fragile to small errors in the data or the model entries. It is worth separating two ideas that are easy to conflate. Conditioning is a property of the forward map: how much the solution of $A\vec{u} = \vec{b}$ moves when the data move. Identifiability is a property of the inverse problem: whether the parameters can be recovered from the observations at all. Ill-conditioning of $A$ often accompanies poor identifiability and is a useful warning sign, but the two are not the same statement, and a large $\kappa(A)$ can also reflect a genuinely stiff, well-posed response. This is not a failure of SA; it is one of the most important things SA can reveal.

*Self-check.* The $2 \times 2$ matrix $A(\epsilon) = \begin{pmatrix} 1 & 1 \\ 1 & 1+\epsilon \end{pmatrix}$ with $\epsilon > 0$: (a) Compute $\det(A(\epsilon))$ and show $A \to$ singular as $\epsilon \to 0$. (b) Compute the condition number $\kappa_2(A)$ for $\epsilon = 1, 0.1, 0.01$. (c) For $\vec{b} = (2, 2+\epsilon)^T$, solve $A\vec{u} = \vec{b}$ and compute $\mathcal{S}_\epsilon^{u_1}$. What happens to this SI as $\epsilon \to 0$?

*Answer:* (a) $\det = \epsilon \to 0$ as $\epsilon \to 0$. Singular at $\epsilon = 0$. (b) $\kappa_2(A) = \sigma_{\max}/\sigma_{\min}$. For $\epsilon = 1$: $\kappa_2 \approx 6.85$. For $\epsilon = 0.1$: $\kappa_2 \approx 42.1$. For $\epsilon = 0.01$: $\kappa_2 \approx 402$. As $\epsilon \to 0$: $\kappa \to \infty$.
(c) $A^{-1} = \epsilon^{-1} \begin{pmatrix} 1+\epsilon & -1 \\ -1 & 1 \end{pmatrix}$. $\vec{u} = A^{-1}\vec{b} = (1,1)^T$ for all $\epsilon > 0$. $\mathcal{S}_\epsilon^{u_1}$: differentiate $u_1 = (2(1+\epsilon) - (2+\epsilon))/\epsilon = \epsilon/\epsilon = 1$ with respect to $\epsilon$, but the simplification $u_1 = 1$ holds exactly, so $\mathcal{S}_\epsilon^{u_1} = 0$. The sensitivity vanishes!
This illustrates a subtlety: a large condition number does not make every index large. Here $\mathcal{S}_\epsilon^{u_1} = 0$ because $\vec{b}$ moves with $A$, while $\partial u_1/\partial b_2 = -1/\epsilon = -100$ at $\epsilon = 0.01$: the exact sensitivity is large in some directions only. What the condition number bounds is the accuracy of the *computed* $\vec{u}$: at $\epsilon = 0.01$ about $\log_{10}\kappa_2$ digits, between two and three, may be lost.

## 8.4 ▪ Separatrices and Trajectory Sensitivity

Dynamical systems often have regions of qualitatively different behavior separated by a *separatrix*: a boundary in state space that trajectories cannot cross. SA computed on one side of a separatrix describes behavior qualitatively different from SA on the other side.

The epidemic threshold is a related but distinct phenomenon, and the distinction is worth keeping. The condition $S(0) > \gamma_R N/(c\beta)$ (equivalently $\mathcal{R}_0 > 1$) is a threshold in parameter and initial-condition space, not a separatrix in state space: it decides whether an outbreak grows, but it is not a barrier that trajectories are forbidden to cross. Both kinds of boundary defeat local SA, for different reasons.

Consider the epidemic peak $I_{\max}$ as $\beta$ varies across the threshold. Below threshold the infectious compartment never grows, so $I_{\max} = I(0)$. Above threshold the peak rises from that floor, and expanding about $\mathcal{R}_0 = 1$ gives

$$I_{\max} - I(0) \;\approx\; \tfrac{1}{2} N\,(\mathcal{R}_0 - 1)^2.$$

So $I_{\max}$ does not jump at the threshold. It is continuous there, and its first derivative is continuous as well and equal to zero. That is the trap, and it is worse than a jump would be: $\mathcal{S}_\beta^{I_{\max}} = 0$ exactly at the threshold, so local SA reports that $\beta$ does not matter at the precise parameter value where $\beta$ decides whether there is an epidemic at all. The response is quadratic, and a first-order method is blind to it. This is the misconception that opened this chapter, in its sharpest form.

The practical implication: if any parameter perturbation of interest crosses a qualitative threshold (endemic vs. disease-free, outbreak vs. no

outbreak, stable vs. oscillatory), local SA cannot be used to predict the response. The model must be analyzed separately in each qualitative regime, and the boundary itself requires special treatment.

A related limitation applies to models with multiple stable attractors. When a model can settle into qualitatively different long-run states depending on initial conditions, the sensitivity index computed near one attractor says nothing about behavior near another. Extensions of the SIR model that exhibit bistability (where both a disease-free and an endemic equilibrium are stable for the same parameter values) require separate SA studies for each attractor. Local SA at one equilibrium cannot warn the analyst that another equilibrium exists.

### 8.4.1 ▪ What Is Differentiated Decides the Failure

Sections 8.2 to 8.4 have shown local SA failing near a threshold in several ways. What goes wrong depends on what is being differentiated, and so does the remedy. We sort the cases by that quantity and give for each what to compute instead.

(1) *An equilibrium at a fold*, $u^*(p)$ (§8.2; Exercise 8.1). $F_u$ is singular at the fold, so the raw derivative diverges, and past the fold the equilibrium does not exist. The normalized index diverges too unless the fold sits at $p = 0$, as in the quadratic. *Instead:* trace the branch by continuation and draw the bifurcation diagram (§8.7, Part 1).

(2) *An equilibrium at a transcritical threshold*, $i^*(\mathcal{R}_0)$ (§8.2.2). The raw derivative stays finite, because $\partial\vec{F}/\partial p$ vanishes as $J_F$ becomes singular. The normalized index diverges as $i^* \to 0$, and a relative change in $\beta$ of about $1/\mathcal{S}_\beta^{i^*}$ removes the equilibrium. *Instead:* draw the bifurcation diagram (§8.7, Part 2), and report $\mathcal{R}_0 - 1$ with the change in each parameter that reaches $\mathcal{R}_0 = 1$.

(3) *The solution of $A\vec{u} = \vec{b}$ as $A$ nears singularity* (§8.3). Nothing fails in the model: the large indices are entries of $A^{-1}$, the genuine response. The computation can lose about $\log_{10}\kappa(A)$ digits. *Instead:* read $\kappa(A)$ as that bound, and check the digits in higher precision or against an analytic value; a small residual cannot check them.

(4) *A quantity on a finite-time trajectory*, such as $I(T)$ (§8.7, Part 3). The forward sensitivity equations are a linear initial-value problem and always have a solution, so the index stays finite across $\mathcal{R}_0 = 1$. What fails is the linear prediction for finite perturbations that carry the trajectory across a threshold or a separatrix (§8.4). *Instead:* rerun

the model at the perturbed values, or sample across the threshold with the methods of Chapter 9.

(5) *A threshold quantity*, such as which equilibrium is stable, or $I_{\max}$ (§8.4). The stable equilibrium switches at $\mathcal{R}_0 = 1$, so that output is not differentiable there and is flat on either side. $I_{\max}$ is flat at the threshold, so $\mathcal{S}_\beta^{I_{\max}} = 0$. No local index can flag the switch. *Instead:* use factor mapping: sample the plausible ranges (§9.2) and report the fraction with $\mathcal{R}_0 > 1$.

(6) *A Floquet multiplier* $\mu_k$ of a periodic orbit (Appendix D). At a period-doubling, and as an orbit shrinks to a Hopf point, $\mu_k$ typically reaches the unit circle at a finite rate, so its derivative stays finite; at a fold of cycles it diverges. *Instead:* follow $\mu_k$ with (D.3) to locate where $|\mu_k| = 1$, and run the model on both sides of that value.

A large index is therefore not by itself a malfunction, and a small or zero index is no reassurance. Every case ends with the same step: find where the threshold lies, and compare that distance with the plausible range of each parameter.

## 8.5 ▪ When Global SA Is Needed

*Local SA studies the neighborhood. Global SA studies the country.*

Local SA fails to answer the right question in three specific situations. When any of these conditions holds, Chapter 9's global SA methods are required. The practical decision framework for choosing between local and global SA is developed at length by Saltelli et al. [55, 53].

To reiterate the misconception that opened this chapter: a sensitivity index near zero does not mean a parameter is globally unimportant. It means either that the linearization at the nominal value is nearly flat, or that the nominal value $p^*$ sits near zero and makes the factor $p^*/q$ in the index small. The example in §8.5.1 is the second case. What happens away from the nominal, across the full uncertainty range of the parameter, requires global SA to determine.

### 8.5.1 ▪ Large Parameter Uncertainty

When a parameter is known only to within a factor of 2 or 3; common in epidemiology, ecology, and environmental modeling; the linearization underlying local SA may be very poor across the plausible range.

As a concrete example, consider a model where the QOI depends on a parameter $p$ through $q(p) = e^p$. At nominal value $p^* = 0$: $\mathcal{S}_p^q = p^*(dq/dp)/q = 0 \cdot 1/1 = 0$. The normalized index is zero. But if the uncertainty range is $[-2, 2]$, the QOI varies from $e^{-2} \approx 0.14$ to $e^2 \approx 7.4$: a factor of about 55. Note where the information is lost. The raw derivative is perfectly informative here, $dq/dp = 1$ at $p^* = 0$; it is the normalization by a nominal value of zero that drives the index to zero. This is a reference-point artifact, not a statement that the model is insensitive to $p$, and it is a standing hazard whenever a parameter's nominal value sits at or near the origin. The relative-change and regularized measures of Section 2.2.3 exist for exactly this situation.

### 8.5.2 ▪ Highly Nonlinear Models

Two models with identical local SIs for a parameter $p$ may behave very differently when $p$ is perturbed by 20%: one model may be nearly linear (the SI is a good approximation), while the other may be highly curved (the SI may understate or overstate the true change).

The sigma-normalized sensitivity index (Section 9.1) provides the minimal bridge: it weights the local SI by the parameter's standard deviation, giving a measure that at least accounts for the relative magnitudes of parameter uncertainties.

### 8.5.3 ▪ Correlated Parameters

Local SA under the independence assumption gives misleading results when parameters are correlated (Chapter 3). Global SA methods in Chapter 9 include specific tools for handling this case.

### 8.5.4 ▪ The Portfolio Example

Consider a model $Y = s_1 Q_1 + s_2 Q_2$ with $s_1 = 2$, $s_2 = 1$ and parameter standard deviations $\sigma_1 = 1$, $\sigma_2 = 3$.

The model slopes are $\partial Y/\partial Q_1 = s_1 = 2$ and $\partial Y/\partial Q_2 = s_2 = 1$, so on the unnormalized derivative alone parameter 1 appears more important. (The normalized index $\mathcal{S}_{Q_i}^Y = s_i Q_i / Y$ would additionally depend on the nominal $Q_i$, which this example leaves unspecified; the comparison here is between slopes and $\sigma$-weighted contributions.)

The sigma-normalized index: $S_i^\sigma = (s_i \sigma_i)/\sigma_Y$, where $\sigma_Y = \sqrt{4 + 9} = \sqrt{13}$. So $S_1^\sigma = 2/\sqrt{13} \approx 0.55$ and $S_2^\sigma = 3/\sqrt{13} \approx 0.83$.

Parameter 2 dominates the output variance, even though it has a smaller slope. The ranking by slope is inverted because parameter 2 has much larger

uncertainty. Chapter 9's Sobol indices provide the rigorous global version of this sigma-normalized assessment.

*Remark* 8.2. For a linear model $Y = \sum_i s_i Q_i$ with independent inputs, the sigma-normalized index $S_i^\sigma = (s_i\sigma_i)/\sigma_Y$ is identical to the *standardized regression coefficient* reported by most statistical software, the "standardized beta" obtained by regressing $Y$ on the $Q_i$'s after standardizing all variables to zero mean and unit variance. The sigma-normalized sensitivity index and the standardized coefficient are the same quantity, derived from the same model and evaluated at the same nominal point; the difference is notation and disciplinary convention. The plain relative index $\mathcal{S}_p^Y$ of Chapter 2 is a different quantity, and carries no variance weighting.

*Self-check.* For the SIR model, suppose $\beta$ has uncertainty $\sigma_\beta = 0.02$ and $\tau_R$ has uncertainty $\sigma_{\tau_R} = 3$ days, at nominal values $\beta^* = 0.06$ and $\tau_R^* = 7$ days.

(a) From Chapter 2, $\mathcal{S}_\beta^{\mathcal{R}_0} = \mathcal{S}_{\tau_R}^{\mathcal{R}_0} = 1$. Which parameter appears more important from local SA alone?

(b) Compute $\sigma_\beta/\beta^*$ and $\sigma_{\tau_R}/\tau_R^*$ (the relative uncertainties). Which parameter has larger relative uncertainty?

(c) Compute the sigma-normalized indices for $\mathcal{R}_0$ with respect to $\beta$ and $\tau_R$. Which parameter drives more variance in $\mathcal{R}_0$?

*Answer:* (a) Both have SI = 1, so local SA alone says they are equally important.
(b) $\sigma_\beta/\beta^* = 0.02/0.06 \approx 0.33$ (33%); $\sigma_{\tau_R}/\tau_R^* = 3/7 \approx 0.43$ (43%). The mean infectious period has larger relative uncertainty.
(c) Apply the delta method first to get $\sigma_{\mathcal{R}_0}$: $\partial\mathcal{R}_0/\partial\beta = c\tau_R = 5 \times 7 = 35$; $\partial\mathcal{R}_0/\partial\tau_R = c\beta = 5 \times 0.06 = 0.3$. $\sigma_{\mathcal{R}_0}^2 \approx (35 \times 0.02)^2 + (0.3 \times 3)^2 = 0.49 + 0.81 = 1.30$, so $\sigma_{\mathcal{R}_0} \approx 1.14$. Using $S_i^\sigma = (\partial\mathcal{R}_0/\partial p_i)\,\sigma_i/\sigma_{\mathcal{R}_0}$ (Equation (9.1)): $S_\beta^\sigma = 35 \times 0.02/1.14 \approx 0.61$; $S_{\tau_R}^\sigma = 0.3 \times 3/1.14 \approx 0.79$. The mean infectious period drives more output variance despite equal local SIs, because $\tau_R$ has larger relative uncertainty. The ratio $S_{\tau_R}^\sigma/S_\beta^\sigma \approx 1.3$ matches the ratio of $\sigma_{\tau_R}\,\partial\mathcal{R}_0/\partial\tau_R$ to $\sigma_\beta\,\partial\mathcal{R}_0/\partial\beta = 0.90$ to 0.70. Chapter 9 will quantify this more precisely with Sobol indices.
*Note.* A common shortcut computes $\mathcal{S}_p^{\mathcal{R}_0} \cdot (\sigma_p/p^*)$ (the local SI times the relative uncertainty) as a quick approximation to $S_p^\sigma$. This shortcut equals the full formula only when $\sigma_{\mathcal{R}_0} = \mathcal{R}_0^*$, that is, when the output's coefficient of variation is exactly 1, which is a large uncertainty rather than a small one. The exact relation is

$$S_p^\sigma = \mathcal{S}_p^Y \frac{\mathrm{CV}_p}{\mathrm{CV}_Y}, \qquad \mathrm{CV}_p = \sigma_p/p^*, \quad \mathrm{CV}_Y = \sigma_Y/Y^*,$$

so the shortcut is exact precisely when $\mathrm{CV}_Y = 1$ and is otherwise wrong by the factor $1/\mathrm{CV}_Y$ in either direction. For accurate sigma-normalized indices, always use the delta-method $\sigma_Y$ in the denominator.

## 8.6 ▪ When SA Guides Data Collection

Sections 8.2 to 8.5 have identified when SA can mislead. This section completes the practical argument the book has been building since Chapter 1: how to use SA *correctly* to make decisions about field measurement.

**The decision chain.** A sensitivity study produces a ranked list of parameters. Translating that list into a field-measurement plan requires three additional steps that SA alone cannot provide:

1. *Identifiability:* Can the high-sensitivity parameter actually be estimated from the available data, or is it structurally confounded with another parameter? Section 3.5 showed that parameters with identical sensitivity indices are a strong warning sign of non-identifiability: if two parameters always appear together in the model equations, their individual effects on any QOI are indistinguishable, and field data cannot separate them. This does not prove non-identifiability in general (equal SIs for one QOI do not preclude different SIs for another), but it is a reliable indicator worth investigating before designing field studies. For the SIR model, $c$ and $\beta$ always appear as the product $c\beta$, so field studies should estimate $c\beta$ directly (e.g., from secondary attack rates) rather than $c$ and $\beta$ separately.

2. *Measurability:* Is the high-sensitivity parameter actually measurable with available instruments and budget? The mean infectious period $\tau_R$ appears in the top tier of every SIR sensitivity analysis, and it is measurable: clinicians record the duration from symptom onset to recovery. The residence time $\tau_m$ has negligible SI for short-horizon models and is measurable anyway from demographic records. A parameter with large SI but no feasible measurement pathway should be *reduced* rather than measured, by redesigning the intervention.

3. *Actionability:* Can the high-sensitivity parameter be reduced by a feasible public health intervention? SA of the SIR model with $J = \max_t I(t)$ consistently identifies $c$ (contact rate) and $\beta$ (transmission probability) as the most influential parameters. Contact rates can be reduced by social distancing, school closures, and remote-work policies. Transmission probability per contact can be reduced by mask requirements, hand-hygiene campaigns, and vaccination. These two routes correspond to different interventions with different costs, compliance rates, and social acceptability. SA identifies both as equally valuable targets; the choice between them requires policy judgment beyond the model.

**The two-budget problem.** A common practical scenario: a government agency has funding to (1) measure exactly $k_{\text{meas}}$ parameters precisely in the field, and (2) implement exactly $k_{\text{int}}$ interventions. The correct SA-guided procedure is:

1. Rank all parameters by how much of the uncertainty in peak $I$ each one carries. Use the sigma-normalized index $S^{\sigma}_{p_j}$ when the linearization holds across the plausible ranges, and otherwise Morris screening (a rapid method for ranking parameters with many fewer model evaluations than full global SA, developed in Section 9.6) or Sobol indices. A ranking by the local index $|\mathcal{S}^{\text{peak } I}_{p_j}|$ alone ignores the uncertainty ranges, as the portfolio example shows, and near the threshold it is empty: at $\mathcal{R}_0 = 1$ every peak index is zero (§8.4.1).
2. From the top-ranked parameters, apply the identifiability filter: remove any parameter that is structurally confounded with a higher-ranked parameter.
3. Apply the measurability filter: remove any parameter for which no feasible measurement protocol exists within the budget.
4. The remaining top $k_{\text{meas}}$ parameters are the measurement priorities.
5. From the ranked list, apply the actionability filter separately: the top $k_{\text{int}}$ actionable parameters are the intervention priorities. These sets may overlap.

**What the model cannot tell you.** The complete picture requires one more piece: parameter inference itself. Smith [59] Chapters 11–12 develops frequentist and Bayesian inference at the graduate level, complementary to and beyond the scope of this introduction, the step that actually quantifies how much a given measurement budget reduces parameter uncertainty. SA tells you *which* parameters matter most; parameter estimation tells you *how precisely* you can learn them from data; parameter selection and influence analysis (Smith Ch. 7 §7.4–7.5) tells you *how to design the measurement study* to learn those parameters as efficiently as possible. Smith [59] Ch. 7 develops parameter identifiability and influence systematically as a precursor to parameter inference, using the sensitivity Jacobian to detect non-identifiable directions before they corrupt downstream uncertainty propagation.

Beyond inference, Smith [59] Ch. 14 takes up the deeper question of *model discrepancy*; what to do when the assumed model form is itself wrong, which is the natural complement to the SA failure modes catalogued in this chapter. SA tells you when a method gives the wrong answer for your model; model discrepancy theory tells you when your model gives the wrong answer for the world.

These disciplines, SA, parameter estimation, optimal experimental de-

sign, and model discrepancy, form a complete pipeline for model-based decision making. This book has equipped you for the first stage.

*If a parameter is influential and poorly known, that is not a nuisance. That is your research agenda.*

## 8.7 ▪ Computational Laboratory: Bifurcation and Ill-Conditioning

**Part 1: The quadratic bifurcation.** For $f(u;p) = u^2 + p$, compute the positive root $u^* = \sqrt{-p}$ and the SI $\mathcal{S}_p^{u^*} = 1/2$ analytically. Plot $u^*$ vs. $p$ for $p \in [-4, -0.01]$. Estimate $\partial u^*/\partial p$ by centered differences at $p = -4, -1, -0.1, -0.01, -0.001$. Compare the finite-difference estimates with the analytic value. At what value of $p$ does the finite-difference estimate become unreliable?

**Part 2: SIS bifurcation diagram using MATCONT.** Download and install MATCONT (`https://matcont.sourceforge.io`). Define the SIS equilibrium equation $\dot{i} = i(\beta(1-i) - \gamma_R) = 0$ as a MATCONT system. Use MATCONT's continuation capability to trace the endemic equilibrium $i^* = 1 - 1/\mathcal{R}_0$ as $\mathcal{R}_0$ increases from 0.5 to 3.0. MATCONT will automatically detect the transcritical bifurcation at $\mathcal{R}_0 = 1$ and flag it. Plot the bifurcation diagram (equilibrium value vs. $\mathcal{R}_0$). If MATCONT is unavailable, trace the equilibrium manually: for $\mathcal{R}_0 \le 1$, $i^* = 0$ (disease-free); for $\mathcal{R}_0 > 1$, $i^* = 1 - 1/\mathcal{R}_0$. (For the SIR model with demography the branch carries the extra factor $\nu_m/(\gamma_R + \nu_m)$ of Equation (8.5); the shape of the diagram, and the index with respect to $\mathcal{R}_0$, are unchanged.)

**Part 3: FSA comparison across $\mathcal{R}_0 = 1$.** Using the full SIR ODE system, compute $\mathcal{S}_\beta^{I(T)}$ at $T = 90$ days for $\mathcal{R}_0 = 0.8, 1.0, 1.2, 2.0$. At each value, use centered finite differences. Tabulate the results. At $\mathcal{R}_0 = 1$: does the SI blow up, return NaN, or behave normally? Try $h = 0.01, 0.001, 0.0001$ to see what happens.

**Part 4: Condition number of the SIR Jacobian as a function of $\mathcal{R}_0$.** For the SIR endemic equilibrium, the Jacobian $J_F$ at $(S^*, I^*, R^*)$ depends on $\mathcal{R}_0$. Compute $\kappa_2(J_F)$ using your language's condition-number routine for $\mathcal{R}_0 \in \{1.01, 1.05, 1.1, 1.5, 2.0, 3.0\}$. Plot $\kappa_2(J_F)$ vs. $\mathcal{R}_0$. Observe the spike as $\mathcal{R}_0 \to 1^+$: the Jacobian becomes nearly singular at the bifurcation. What does this imply about SA of the SIR endemic equilibrium for parameter values that place $\mathcal{R}_0$ close to 1?

**Part 5: Sigma-normalized SIs for the SIR model.** Using the SIR model with uncertainty ranges $\sigma_c = 1$ (contacts/day), $\sigma_\beta = 0.02$, $\sigma_{\tau_R} = 3$ days, $\sigma_{\tau_m} = 500$ days (all at $1\sigma$), compute the sigma-normalized indices for $\mathcal{R}_0$. Compare the ranking with the local SI ranking from Chapter 2.

Which parameter most affects $\mathcal{R}_0$ given its realistic uncertainty range?

## 8.8 ▪ Exercises

**Exercise 8.1** *([Bloom L3]).* For the equation $g(u;p) = u^3 - 3pu + 2 = 0$:

(a) Find all real roots at $p = 1$. Which root changes most rapidly with $p$?

(b) Compute $\mathcal{S}_p^{u^*}$ for each real root at $p = 1$ using the implicit function theorem.

(c) Plot the roots as functions of $p$ for $p \in [0.8, 1.2]$. At what value of $p$ do two roots merge?

(d) At the bifurcation value of $p$, what does the FSE predict? Verify numerically by *one-sided* continuation from $p > 1$. A centered difference through $p = 1$ is not available on the positive branch: for $p < 1$ those two roots are complex, so the branch does not exist on that side.

**Exercise 8.2** *([Bloom L3]).* Using the SIR model with $c = 5$, $\tau_R = 7$ days, $\tau_m = 10{,}000$ days:

(a) Compute $\mathcal{S}_\beta^{I(T)}$ for $\beta = 0.04, 0.06, 0.08, 0.12$ at $T = 90$ days using centered finite differences. Note: with demography the threshold is $\mathcal{R}_0 = c\beta/(\gamma_R + \nu_m)$. At $\tau_m = 10{,}000$ days this differs from $c\beta\tau_R$ by $0.07\%$, so to the precision quoted it takes values $1.4, 2.1, 2.8, 4.2$.

(b) Now recompute for $\beta = 0.02$ ($\mathcal{R}_0 = 0.7$). Compare with the results for $\mathcal{R}_0 > 1$. Explain any qualitative differences.

(c) If you were to add a bifurcation analysis tool (e.g., MATCONT), what would you expect the bifurcation diagram to look like for the endemic equilibrium of this SIR model as $\beta$ varies?

**Exercise 8.3** *([Bloom L4]).* For the $3 \times 3$ linear system $A\vec{u} = \vec{b}$ where:

$$A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9+\epsilon \end{pmatrix}, \quad \vec{b} = A\vec{u}_{\text{exact}}, \quad \vec{u}_{\text{exact}} = (1,1,1)^T.$$

(a) For $\epsilon = 1, 0.1, 0.01, 0.001, 0.0001$, compute $\kappa(A)$ and $\|\hat{\vec{u}} - \vec{u}_{\text{exact}}\|$ where $\hat{\vec{u}} = A^{-1}\vec{b}$ (via your linear system solver).

(b) Plot $\|\hat{\vec{u}} - \vec{u}_{\text{exact}}\|$ vs. $\kappa(A)$ on log-log axes. What is the approximate slope?

(c) What does this experiment tell you about interpreting large SI values for nearly singular models?

**Exercise 8.4** *([Bloom L6]).* You are asked to perform SA on an epidemic model whose best-fit parameter values give $\mathcal{R}_0 = 1.04$, very close to the epidemic threshold. Write a one-page advisory report for a public health agency explaining:

(a) What local SA can and cannot tell you at this parameter value.

(b) What specific numerical problems to expect when computing sensitivity indices near $\mathcal{R}_0 = 1$.

(c) What additional analyses you would recommend before advising on interventions.

(d) How you would explain the concept of a bifurcation and its implications to a non-mathematical policy maker.

**Exercise 8.5** *([Bloom L4]).* *Audit exercise.* A policy brief reports local sensitivity indices for an endemic equilibrium of an SIR model at a parameter set for which $\mathcal{R}_0 = 1.02$, and concludes that "the equilibrium prevalence is extremely sensitive to the recovery rate, so recovery is the highest-priority intervention target." The indices are large in magnitude and were computed correctly.

(a) Explain why indices of large magnitude are expected here, and why that magnitude is not by itself evidence of an intervention priority.

(b) State what happens to the quantity being differentiated as $\mathcal{R}_0 \to 1^+$, and what that implies for the validity of the linearization.

(c) The brief's conclusion also assumes something about the recovery rate that is outside every index reported. Identify it.

(d) Recommend the analysis you would run instead, and state what question it would answer that the local indices cannot.

**Exercise 8.6** *([Bloom L6]).* A published modeling paper reports: "Sensitivity analysis of the endemic equilibrium of our malaria model found that

all indices are less than 0.1, indicating the model is robust to parameter uncertainty. The basic reproduction number at fitted parameter values is $\mathcal{R}_0 = 1.15$."

Identify at least three specific methodological concerns with this analysis and its conclusions. For each concern, state: (a) what the problem is; (b) what evidence the authors would need to provide to address it; and (c) what alternative analysis would give a more complete picture.

## Notes

The methodological critique of sensitivity analysis practice that motivates this chapter is developed systematically by Saltelli et al. [48], who reviewed 280 highly cited published SA studies and found that more than half used local methods to answer questions that require global analysis. The paper gives explicit examples of how this misapplication affects policy conclusions in epidemiology, pharmacology, and environmental science.

The sigma-normalized sensitivity index $S_i^{\sigma}$, which bridges local SA (Chapter 2) and global SA (Chapter 9), is discussed by Saltelli and Tarantola [47] in the context of model-independent sensitivity measures; its connection to the standardized regression coefficient is established in Saltelli et al. [52], Chapter 4.

The structural identifiability problem for ODE models, whether the parameter vector can in principle be recovered from perfect observations of the state, is treated rigorously by Mester et al. [33] and in the epidemiological context by Chitnis, Hyman, and Cushing [14]. The practical identifiability diagnostics based on the sensitivity matrix condition number follow the approach of Pianosi et al. [42].

Draft

Draft

# Chapter 9
# Global Sensitivity Analysis and Sobol Indices

*The point is not to remove uncertainty.*
*The point is to find out which uncertainty is worth worrying about.*

Consider two epidemiological models. Both give $\mathcal{S}_\beta^{\mathcal{R}_0} = +1$ at the same nominal parameter values. In Model A, $\mathcal{R}_0$ is nearly linear in $\beta$ across the full biologically plausible range $[0.01, 0.15]$. In Model B, $\mathcal{R}_0$ is highly nonlinear: it is steep near the lower end of the range and flat at the upper end.

For Model A, the local SI correctly identifies $\beta$ as an important parameter. For Model B, local SA gives the same SI value of $+1$; but it is now describing only the slope at the nominal point. A 20% increase in $\beta$ from the nominal produces very different effects in the two models, and no derivative computed at the nominal value can see this.

Chapter 8 showed when local SA fails. This chapter provides the tools to complement and extend it; global SA answers questions that local SA cannot, while local SA remains the starting point for understanding which parameters matter and why.

Before choosing a global SA method, two questions must be answered. *What is the goal?* Three distinct purposes call for different methods [55, 52]: *factor prioritization* (which inputs most reduce output variance if their uncertainty were eliminated?), *factor fixing* (which inputs can be held constant without materially changing the output?), and *factor mapping* (which input combinations drive the output above a critical threshold?). *Does the method's assumptions match the model?* First-order Sobol indices answer the factor-prioritization question; total-order indices answer the factor-fixing question. Using either one to answer a question outside its scope produces results that are numerically correct but scientifically misleading, the central lesson of Chapter 8. *Unifying Idea 4.* All three purposes are

stated about inputs to a model. Factor fixing licenses holding a parameter constant *in the model*; it is not a finding that the quantity is constant in the world, and the variance being decomposed is the variance the assumed input distributions put there.

## 9.1 ▪ Why Local SA Is Not Enough: The Sigma-Normalized Bridge

*Local SA studies the neighborhood. Global SA studies the country.*

The two can give very different answers about which parameters matter most, as the following example shows.

### 9.1.1 ▪ The Portfolio Example

Consider a model $Y = s_1 Q_1 + s_2 Q_2$ where $s_1 = 2$, $s_2 = 1$ and the parameters have standard deviations $\sigma_1 = 1$, $\sigma_2 = 3$.

Local SA gives the raw derivatives $\frac{\partial Y}{\partial Q_1} = 2 > \frac{\partial Y}{\partial Q_2} = 1$: parameter 1 appears more important. (These are raw slopes, not the normalized indices of Chapter 2, which would need nominal values $Q_1^*$, $Q_2^*$ and $Y^*$.) But the parameters have very different uncertainty ranges, and $Q_2$ varies three times as much as $Q_1$.

The output variance is:

$$\mathrm{Var}(Y) = s_1^2\sigma_1^2 + s_2^2\sigma_2^2 = 4 + 9 = 13 \quad \Longrightarrow \quad \sigma_Y = \sqrt{13}.$$

The *sigma-normalized sensitivity index*

$$S_i^\sigma = \frac{\sigma_i}{\sigma_Y}\,\frac{\partial Y}{\partial Q_i} \tag{9.1}$$

accounts for both the derivative and the parameter uncertainty:

$$S_1^\sigma = \frac{1}{\sqrt{13}} \cdot 2 \approx 0.55, \qquad S_2^\sigma = \frac{3}{\sqrt{13}} \cdot 1 \approx 0.83.$$

Parameter 2 dominates, opposite to the local SA ranking. The local ranking was misleading because it ignored the factor-of-3 difference in parameter uncertainty.

Like the normalized SI in Chapter 2, $S_i^\sigma$ is dimensionless: $\sigma_i$ carries the units of $Q_i$ and $\sigma_Y$ carries the units of $Y$, so $(\sigma_i/\sigma_Y)\,\partial Y/\partial Q_i$ is a pure number. This means $S_1^\sigma$ and $S_2^\sigma$ can be compared directly even when the two parameters are measured in completely different units, the same reason the tornado plot in Chapter 2 sorted by $|\mathcal{S}_p^q|$ rather than by the raw derivative.

*Unifying Idea 1.* The sigma-normalized index is still the sensitivity index from Chapter 2, rescaled by the ratio of relative uncertainties: with $\mathrm{CV}_i = \sigma_i/Q_i^*$ and $\mathrm{CV}_Y = \sigma_Y/Y^*$, $S_i^\sigma = \mathcal{S}_{Q_i}^Y \mathrm{CV}_i/\mathrm{CV}_Y$. The parameter's relative uncertainty raises the index and the output's lowers it. Global SA builds on local SA; it does not replace it. The local SI from Chapter 2 reappears inside the variance-based formulas of this chapter whenever the response is close to linear over the input range, which is exactly the delta-method setting. The connection is real but not universal: Morris, PRCC, and Borgonovo's moment-independent measures are built from other quantities and contain no local index.

### 9.1.2 ▪ Two SIR Models

For the SIR model, suppose two outbreak settings give the same $\mathcal{S}_\beta^{\mathcal{R}_0} = 1$. In Setting A, $\beta$ is tightly constrained from contact-tracing data ($\sigma_\beta = 0.005$), while in Setting B, $\beta$ must be estimated from seroprevalence data ($\sigma_\beta = 0.03$). Local SA says $\beta$ is equally important in both settings. In Setting B the contribution of $\beta$ to the uncertainty in $\mathcal{R}_0$, measured by $|\partial\mathcal{R}_0/\partial\beta|\,\sigma_\beta$, is six times larger. The sigma-normalized index rises by less than six, because $\sigma_{\mathcal{R}_0}$ in the denominator rises with $\sigma_\beta$ as well; for the nominal values used in the self-check below the factor is about 3.2. These uncertainties propagate to $\mathcal{R}_0$ through the standard delta-method formula developed in Chapter 8.

*Self-check.* For the SIR model with local SIs $\mathcal{S}_c^{\mathcal{R}_0} = \mathcal{S}_\beta^{\mathcal{R}_0} = \mathcal{S}_{\tau_R}^{\mathcal{R}_0} = 1$ and parameter standard deviations $\sigma_c = 0.5$, $\sigma_\beta = 0.01$, $\sigma_{\tau_R} = 2$, with $\sigma_{\mathcal{R}_0}$ obtained from the delta method $\mathrm{Var}(\mathcal{R}_0) \approx \sum_i (\partial\mathcal{R}_0/\partial p_i)^2\sigma_i^2$:

For a SIR model with $c^* = 5$, $\beta^* = 0.06$, $\tau_R^* = 7$ ($\mathcal{R}_0^* = 2.1$), first compute $\sigma_{\mathcal{R}_0}$ from the delta method, then rank the three parameters by their sigma-normalized index $S_i^\sigma$.

> *Answer:* Delta method: $\mathrm{Var}(\mathcal{R}_0) \approx \sum_i (\partial\mathcal{R}_0/\partial p_i)^2\sigma_i^2$. $\partial\mathcal{R}_0/\partial c = \beta\tau_R = 0.42$; $\partial\mathcal{R}_0/\partial\beta = c\tau_R = 35$; $\partial\mathcal{R}_0/\partial\tau_R = c\beta = 0.3$. $\mathrm{Var}(\mathcal{R}_0) \approx 0.42^2(0.25) + 35^2(0.0001) + 0.3^2(4) = 0.0441 + 0.1225 + 0.36 = 0.527$. $\sigma_{\mathcal{R}_0} \approx 0.726$.
> Compute each index directly from $S_i^\sigma = (\partial\mathcal{R}_0/\partial p_i)\sigma_i/\sigma_{\mathcal{R}_0}$. $S_c^\sigma = 0.42 \times 0.5/0.726 \approx 0.29$. $S_\beta^\sigma = 35 \times 0.01/0.726 \approx 0.48$. $S_{\tau_R}^\sigma = 0.3 \times 2/0.726 \approx 0.83$.
> Ranking: $\tau_R > \beta > c$. The mean infectious period drives the most uncertainty in $\mathcal{R}_0$, despite all three having equal local SIs.

### 9.1.3 ▪ Time-Varying Sensitivity

The analyses above collapse the SIR model's time-dependent output into a single scalar QOI: $\mathcal{R}_0$, or peak infections, or final size. This is often appropriate, but it discards information. For ODE models, sensitivity indices can be computed as functions of time: $S_{p_i}^{q_j}(t)$ tells which parameters matter most at each moment in the epidemic trajectory.

In the SIR model the answer changes substantially over the course of the outbreak. Near the start, the trajectory is exponentially growing and $\beta$ (the transmission rate) dominates: small changes in $\beta$ compound rapidly. Near the peak, $\gamma_R$ (the recovery rate) becomes comparably important because the balance between new infections and recoveries determines when the peak occurs. Toward the end, both rates shape the final size, but initial-condition parameters that seemed irrelevant during growth can re-emerge as influential. A global SA computed against only the peak-infections QOI would miss the period when $\gamma_R$ most needs careful estimation.

Time-varying global SA works by running the Saltelli algorithm against the model output at each recorded time step $t_1, t_2, \ldots, t_T$. For ODE models this costs no additional model evaluations: a single run of the SIR system produces the full trajectory $\vec{u}(t)$ at every time step, so the $N(m+2)$ evaluations of the Saltelli construction automatically yield model output at all $T$ recorded times. The sensitivity indices $\hat{S}_i(t_k)$ and $\hat{S}_{T_i}(t_k)$ are then computed by applying the Jansen estimator to the outputs at each time step $t_k$ separately; an arithmetic operation with negligible cost compared to the ODE solves. Wu et al. [64] apply this approach to epidemic models and show that parameter importance rankings shift substantially across the trajectory, with direct implications for which parameters to prioritize for data collection at each stage of an outbreak.

## 9.2 ▪ Sampling Methods

All global SA methods require evaluating the model at many parameter combinations. Choosing those combinations efficiently is the first computational task.

### 9.2.1 ▪ Random Sampling

A note on notation before the methods begin. Through the modeling chapters $N$ has denoted the total population, $S + I + R = N$. From here to the end of the chapter it denotes the *sample size*, the number of parameter vectors drawn, which is the standard usage in the sensitivity literature and the $N$ of the Saltelli cost $N(m+2)$. No population $N$ appears in this chapter, so the two never meet on a page.

The simplest approach: draw $N$ independent samples from the joint parameter distribution (often assumed uniform or independent log-normal over plausible ranges). For $N$ large, random sampling provides good coverage on average, but can leave regions of parameter space unsampled by chance.

### 9.2.2 ▪ Latin Hypercube Sampling

Latin Hypercube Sampling (LHS) divides each parameter's range into $N$ equal strata and ensures that exactly one sample falls in each stratum for every parameter [32]. This eliminates the clustering that random sampling can produce and improves coverage with smaller $N$.

**Definition 9.1** (LHS Sample)**.** *An LHS of size $N$ for $m$ parameters with ranges $[l_j, u_j]$, $j = 1, \ldots, m$, is an $N \times m$ matrix $X$ such that each column contains exactly one value from each of the $N$ equal-width strata $[l_j + (k-1)(u_j - l_j)/N,\ l_j + k(u_j - l_j)/N]$ for $k = 1, \ldots, N$.*

The standard LHS algorithm permutes the strata independently for each parameter and adds uniform jitter within each stratum:

```
# LHS sample: N points in m dimensions (see repository for full implemen
set_random_seed(42)
X = LHS_SAMPLE(N, m, lower=lb, upper=ub)  # N-by-m sample matrix
# Each column j has exactly one sample in each of the N strata of [lb[j]
```

For the same $N$, LHS provides substantially better coverage than random sampling, particularly for estimating marginal effects of individual parameters; see Helton and Davis [18] for convergence analysis and uncertainty propagation results. For Sobol index estimation (Section 9.5), LHS is not the default; this chapter fills the Saltelli matrices from a scrambled Sobol' sequence (§9.2.3).

*Self-check.* For $N = 4$ samples and $m = 1$ parameter on $[0, 1]$: (a) Draw a random sample of size 4. Is it possible that all 4 points fall in $[0, 0.5]$? Is this possible with LHS? (b) Write out explicitly what an LHS of size 4 looks like for one parameter.

> *Answer:* (a) Yes, random sampling can place all 4 points in $[0, 0.5]$ (probability $(1/2)^4 = 1/16$). With LHS, this is impossible: each quarter-interval $[0, 0.25)$, $[0.25, 0.5)$, $[0.5, 0.75)$, $[0.75, 1]$ must contain exactly one sample.
> (b) One possible LHS: one point in each of $[0, 0.25)$, $[0.25, 0.5)$, $[0.5, 0.75)$, $[0.75, 1]$, with uniform jitter within each. Example: $X = \{0.18, 0.43, 0.62, 0.91\}$, one value per stratum, randomly positioned within each.

### 9.2.3 ▪ Low-Discrepancy Sampling and Scrambled Sobol' Sequences

The Monte Carlo convergence rate $O(1/\sqrt{N})$ for the Jansen estimators is a consequence of drawing the rows of matrices $A$ and $B$ independently and uniformly at random. Independence is a liability rather than an asset: independent draws cluster and leave gaps, so many of the $N(m+2)$ model evaluations are wasted on redundant regions of parameter space.

The key insight is that the estimators require only that each row of $A$ (and $B$) be *marginally uniform*, that is, each component $Q_i$ is uniformly distributed over its range. Independence among rows is not required. Replacing independent uniform draws with a *low-discrepancy sequence* that is marginally uniform but deliberately spread across $[0,1]^m$ gives the same unbiased estimators with substantially lower variance.

**Discrepancy.** The *discrepancy* of a point set measures how uniformly points cover the unit hypercube. Low-discrepancy sequences (LDS) achieve discrepancy $O((\log N)^m/N)$, compared to the typical $O(1/\sqrt{N})$ of independent uniform samples. For smooth integrands with $m \leq 10$, this translates directly into faster convergence of the Sobol estimators.

**Scrambled Sobol' sequences.** The current best practice is the *scrambled Sobol' sequence* [39]. A Sobol' sequence fills $[0,1]^m$ one stratum at a time; scrambling randomizes the digits of each coordinate in a way that keeps this stratification. Each point is then marginally uniform, the joint distribution covers the space uniformly, and the sequence is *randomized*; the estimator is unbiased and its variance is estimable by repeating the scramble. Owen [38] proved that scrambled nets achieve variance $O((\log N)^{m-1}/N^3)$ for smooth integrands, a superlinear improvement over the $O(1/N)$ of Latin hypercube sampling.

**Independence is not the critical property.** Independence guarantees that the standard confidence interval formulas apply. Scrambled Sobol' sequences are not independent: pairs of points carry small correlations, slightly negative on average. That negative dependence is the source of the variance reduction described below. They also satisfy a central limit theorem. Confidence intervals are constructed by repeating the scrambled estimation with 30 or more independent scrambles. The practical consequence: for a given budget $N$, scrambled Sobol' sampling often substantially reduces the confidence interval width on $\hat{S}_i$ and $\hat{S}_{T_i}$ compared to independent uniform sampling, in smooth, low-dimensional problems frequently by a factor of two or more (about seven in the worked example of §9.5), at no additional model evaluations. The *identically distributed* property, each sample point is uniform over its range, is what the unbiasedness of the estimators relies on; statistical independence among the sample points is not required, and for a low-discrepancy design it is precisely what one gives up in exchange for the more even coverage that lowers estimator variance. Figure 9.1 illustrates the coverage difference.

**Practical guidance.** SALib [19] uses scrambled Sobol' sequences by default since version 1.4. In MATLAB, `sobolset` in the Statistics and Machine Learning Toolbox provides the same construction. For $m \leq 20$ and $N \leq 10{,}000$, scrambled Sobol' sampling is usually the better choice, giving





**Figure 9.1.** *Comparison of $N = 64$ points in two dimensions. Left: independent identically distributed (IID) uniform samples show visible clusters and empty regions that waste model evaluations. Right: scrambled Sobol' sequence [39] achieves near-uniform coverage. With $N = 64 = 2^6$ points in two dimensions the sequence is a $(0, 6, 2)$-net in base 2, and its scrambling (SciPy's linear matrix scramble with a digital shift) preserves that property, so every one of the $8 \times 8$ cells contains exactly one point. This is a guarantee rather than an average: in the left panel 24 of the 64 cells are empty and one holds five points. Both strategies produce marginally uniform samples, so both yield unbiased Sobol estimators; the LDS strategy produces much lower estimator variance for the same computational budget $N$.*

lower variance than IID or LHS for integrands with enough smoothness. The advantage narrows as $m$ grows or when the integrand is rough, and for a genuinely discontinuous response there is no guarantee of improvement. For a comprehensive treatment of quasi-Monte Carlo methods, including higher-dimensional constructions and their error analysis, see Owen [40].

## 9.3 ▪ Regression-Based SA: LHS/PRCC

The Partial Rank Correlation Coefficient (PRCC) is the dominant global SA method in mathematical epidemiology. It is computationally cheap, handles monotone (but not necessarily linear) relationships, and the Marino et al. [31] MATLAB implementation is widely used and freely available; it builds on the rank-correlation approach of Iman and Conover [24] and the epidemiological SA tradition of Blower and Dowlatabadi [5].

*Remark* 9.2. The PRCC is a *nonparametric partial correlation coefficient*. Step 2 of the calculation below fits a linear regression on rank-transformed data, and Step 3 extracts the partial correlation coefficient for each predictor. Readers from statistics will recognize this as a rank-based analog of the partial correlation coefficient from multiple regression. It is not the partial regression coefficient, the slope that Step 2 fits; on ranks that slope is the standardized rank regression coefficient, a different number. PRCC

for $p_j$ measures the linear relationship between rank($p_j$) and rank($Y$) after removing the linear effects of all other rank-transformed parameters. The rank transformation replaces the linearity assumption with a monotonicity assumption, extending applicability to nonlinear but order-preserving responses.

### 9.3.1 ▪ The PRCC Calculation

Given $N$ LHS samples of the $m$ POIs and the corresponding $N$ model outputs:

**Step 1.** Rank-transform all inputs and the output separately. Replace each value with its rank among the $N$ observations.

**Step 2.** Fit a linear regression of the rank-transformed output on all rank-transformed inputs.

**Step 3.** For each parameter $p_j$, compute the partial correlation coefficient between the rank-transformed $p_j$ and the rank-transformed output, controlling for all other parameters. This is the PRCC for $p_j$.

**Step 4.** Compute a $t$-test $p$-value for each PRCC to assess statistical significance.

The PRCC ranges from $-1$ to $+1$. Its sign has the same interpretation as the SI sign rule from Chapter 2: positive means POI and QOI increase together; negative means they move in opposite directions.

### 9.3.2 ▪ When PRCC is Appropriate

PRCC requires monotone relationships: the QOI must consistently increase (or consistently decrease) as each POI increases, across the full range of parameter values. It does not require linearity. For most compartmental epidemic models, the relationships are approximately monotone, which is why PRCC dominates the field.

Before applying PRCC, verify the monotonicity assumption by producing a scatter matrix: plot the model output against each POI for the full LHS sample using a pairwise scatter matrix. A roughly monotone cloud (upward or downward sloping, without reversals) supports the use of PRCC. A hump-shaped or U-shaped cloud signals non-monotonicity, and Sobol indices should be used instead.

PRCC and Sobol indices are not equivalent. PRCC assumes monotone relationships; Sobol indices make no such assumption. When a relationship is nonmonotone (a QOI first increases then decreases as a parameter increases), PRCC may give a near-zero coefficient even for a highly important parameter, while the Sobol total index will correctly identify it as important. For the models in this course, both methods typically agree, but verifying

agreement is a useful consistency check.

For the SIR model, $\mathcal{R}_0 = c\beta/\gamma_R$ is monotone increasing in $\beta$ and monotone decreasing in $\gamma_R$, which is why PRCC performs reliably on this particular model. For output quantities that exhibit threshold or bifurcation behavior, such as the time to peak infections in models with multiple competing timescales and parameter-dependent qualitative changes, monotonicity can fail even when the individual input-output relationships look locally linear. In such cases PRCC will underestimate the importance of parameters that drive threshold-crossing, while the Sobol total index captures the full nonlinear contribution [56].

## 9.4 ▪ Sobol Decomposition and Variance-Based SA

Variance-based SA is the most rigorous framework for global SA: it quantifies exactly how much of the total output variance is attributable to each POI or group of POIs [60].

### 9.4.1 ▪ The Intuition: Conditional Expectation

Suppose parameter $Q_1$ is fixed at $q_1$ while all other parameters vary across their ranges. The expected output $\mathbb{E}[Y \mid Q_1 = q_1]$ is some function of $q_1$. If $Q_1$ is important, this conditional expectation varies substantially as $q_1$ changes. The variance of this conditional expectation, $\mathrm{Var}_{Q_1}[\mathbb{E}_{Q_{-1}}(Y \mid Q_1)]$, measures how much of $Y$'s variability is explained by $Q_1$ alone. Normalizing by the total variance gives the first-order Sobol index.

### 9.4.2 ▪ The ANOVA Decomposition

*Why does a unique decomposition exist?* Because statistical independence allows us to cleanly separate what each parameter contributes alone from what it contributes jointly. When the $Q_i$ are independent, projecting $f$ onto the subspace of functions of $Q_i$ alone gives a well-defined component $f_i(Q_i)$, and these projections do not interfere with each other. The orthogonality $\int_0^1 f_i(Q_i)\, dQ_i = 0$ is the mathematical expression of this separation: each component has zero mean after integrating out $Q_i$, so distinct components contribute zero covariance. When parameters are correlated, this orthogonality breaks down, and the decomposition loses its uniqueness (Section 9.7 returns to this point). With independence, the decomposition is both unique and additive in variance.

For independent POIs $\vec{Q} = (Q_1, \ldots, Q_m)$ uniformly distributed on $[0,1]^m$, any square-integrable model $Y = f(\vec{Q})$ admits the unique *ANOVA decom-*

*position*:

$$f(\vec{Q}) = f_0 + \sum_i f_i(Q_i) + \sum_{i<j} f_{ij}(Q_i, Q_j) + \cdots + f_{12\cdots m}(\vec{Q}), \tag{9.2}$$

where $f_0 = \mathbb{E}[Y]$, $f_i(Q_i) = \mathbb{E}[Y \mid Q_i] - f_0$, and the higher-order terms capture interactions. Uniqueness follows from the orthogonality conditions $\int_0^1 f_u(\vec{Q}_u)\, dQ_i = 0$ for each $i \in u$; see Sobol [60] and Saltelli et al. [52] Appendix A for a proof. The terms are orthogonal, so their variances add:

$$D = \mathrm{Var}(Y) = \sum_i D_i + \sum_{i<j} D_{ij} + \cdots + D_{12\cdots m}, \tag{9.3}$$

where $D_i = \mathrm{Var}[\mathbb{E}(Y \mid Q_i)]$ and $D_{ij} = \mathrm{Var}[\mathbb{E}(Y \mid Q_i, Q_j)] - D_i - D_j$.

**Worked example: a model with an interaction term.** Consider $f(Q_1, Q_2) = Q_1 + Q_2 + Q_1 Q_2$ with $Q_1, Q_2 \sim \mathcal{U}(0,1)$ independent. Compute the ANOVA components directly from the definitions:

$$\begin{aligned}
f_0 &= \mathbb{E}[Y] = \tfrac{1}{2} + \tfrac{1}{2} + \tfrac{1}{4} = \tfrac{5}{4},\\
f_1(Q_1) &= \mathbb{E}[Y \mid Q_1] - f_0 = \big(Q_1 + \tfrac{1}{2} + \tfrac{Q_1}{2}\big) - \tfrac{5}{4} = \tfrac{3Q_1}{2} - \tfrac{3}{4} = \tfrac{3}{2}\big(Q_1 - \tfrac{1}{2}\big),\\
f_2(Q_2) &= \mathbb{E}[Y \mid Q_2] - f_0 = \tfrac{3}{2}\big(Q_2 - \tfrac{1}{2}\big) \quad \text{(by symmetry)},\\
f_{12}(Q_1, Q_2) &= f - f_0 - f_1 - f_2 = Q_1 Q_2 - \tfrac{Q_1}{2} - \tfrac{Q_2}{2} + \tfrac{1}{4} = \big(Q_1 - \tfrac{1}{2}\big)\big(Q_2 - \tfrac{1}{2}\big).
\end{aligned}$$

Verify orthogonality: $\int_0^1 f_{12}\, dQ_1 = \big(\int_0^1 (Q_1 - \tfrac{1}{2})\, dQ_1\big)(Q_2 - \tfrac{1}{2}) = 0$. ✓

The variances are $D_1 = D_2 = \big(\tfrac{3}{2}\big)^2 \mathrm{Var}(Q_1) = \tfrac{9}{4} \cdot \tfrac{1}{12} = \tfrac{3}{16}$ and $D_{12} = \mathrm{Var}(f_{12}) = \mathrm{Var}(Q_1 - \tfrac{1}{2})\,\mathrm{Var}(Q_2 - \tfrac{1}{2}) = \tfrac{1}{144}$. Total variance $D = \tfrac{3}{16} + \tfrac{3}{16} + \tfrac{1}{144} = \tfrac{27}{144} + \tfrac{27}{144} + \tfrac{1}{144} = \tfrac{55}{144}$. The first-order indices $S_1 = S_2 = \tfrac{3/16}{55/144} = \tfrac{27}{55} \approx 0.491$, and the interaction index $S_{12} = \tfrac{1/144}{55/144} = \tfrac{1}{55} \approx 0.018$. Note $S_1 + S_2 + S_{12} = 1$. For the additive model $f = Q_1 + Q_2$ (no interaction), $S_1 = S_2 = \tfrac{1}{2}$ and $S_{12} = 0$: the indices sum to 1 with nothing left over for interactions. This is why $\sum S_i < 1$ signals the presence of interactions.

### 9.4.3 ▪ Sobol Indices

**Definition 9.3** (Sobol Indices)**.** *The* first-order Sobol index *for parameter $i$ is:*

$$S_i = \frac{D_i}{D} = \frac{\mathrm{Var}[\mathbb{E}(Y \mid Q_i)]}{\mathrm{Var}(Y)}. \tag{9.4}$$

*The* total Sobol index *for parameter $i$ [21] is:*

$$S_{T_i} = 1 - \frac{\mathrm{Var}[\mathbb{E}(Y \mid \vec{Q}_{-i})]}{\mathrm{Var}(Y)} = \frac{\mathbb{E}[\mathrm{Var}(Y \mid \vec{Q}_{-i})]}{\mathrm{Var}(Y)}, \tag{9.5}$$

*where $\vec{Q}_{-i}$ denotes all parameters except $Q_i$.*

The first-order index measures the *direct* effect of $Q_i$; the total index measures the direct effect plus all interactions with other parameters. When $S_{T_i} = S_i$ for all $i$, the model is additively separable (no interactions). When $S_{T_i} > S_i$, parameter $i$ interacts with other parameters.

*Remark* 9.4 (ANOVA and Sobol indices). Readers from statistics will recognize $S_i = \mathrm{Var}[\mathbb{E}(Y \mid Q_i)]/\mathrm{Var}(Y)$ as the quantity known as $\boldsymbol{\eta^2}$ (eta-squared) in classical ANOVA, the proportion of total response variance explained by one factor acting alone. The two frameworks share the same mathematical skeleton. In classical ANOVA, the total sum of squares decomposes as $SS_{\text{total}} = SS_1 + SS_2 + SS_{12} + SS_{\text{residual}}$, where each $SS$ is computed from an observed dataset and $SS_{\text{residual}}$ captures unexplained noise. In the Sobol framework, the total variance decomposes as $D = D_1 + D_2 + D_{12}$ with no residual term, because the model is deterministic and all variance is explained by the inputs. Three differences matter in practice: (1) classical ANOVA requires categorical factors or discretized inputs; Sobol indices work with continuous parameters; (2) classical ANOVA estimates variance components from sample data; Sobol indices are population quantities estimated by Monte Carlo; (3) classical ANOVA requires balance and orthogonality across observed data; Sobol indices require independence of the input distributions. Despite these differences, for a linear model with independent inputs, the two frameworks coincide exactly: $S_i = \eta^2$. The Sobol framework is the natural generalization of ANOVA to nonlinear deterministic models with continuous inputs.

*Unifying Idea 1.* The first-order Sobol index $S_i = D_i/D$ uses the same ratio structure as the sensitivity index from Chapter 2; it measures one variance as a fraction of another. The connection is explicit: for a linear model $Y = \sum_i a_i Q_i$ with independent $Q_i \sim \mathcal{U}[l_i, u_i]$, the first-order Sobol index equals the square of the sigma-normalized index: $S_i = (S_i^\sigma)^2$. Global SA builds on local SA. For a linear model $S_i$ and $S_i^\sigma$ give the same ranking by magnitude but different values. The raw local slopes can give a different ranking, as the portfolio of Section 9.1 shows.

*Unifying Idea 3.* The Sobol decomposition resolves the tension that Unifying Idea 3 created in Chapter 8. Local SA is a linearization, valid near the nominal values. Sobol indices make no linearity assumption: they measure the actual variance contribution across the full parameter range. When the linearization is poor (Model B from the chapter opening), local and global SA give different answers, and Sobol indices give the correct answer.

For readers with a functional analysis background (Appendix A), the ANOVA decomposition (9.2) is the orthogonal decomposition of $f$ in the

Hilbert space $L^2([0,1]^m)$ with respect to the inner product $\langle f, g\rangle = \int_{[0,1]^m} f(\vec{q})\, g(\vec{q})\, d\vec{q}$. The orthogonality explains both why the variances add in (9.3) and why the method requires independent POIs: independence is needed for the inner product to factorize and the projections to be orthogonal.

*Self-check.* For $Y = c_1 Q_1 + c_2 Q_2$ with $Q_i \sim \mathcal{U}(0,1)$ independent: (a) Compute $\mathbb{E}[Y \mid Q_1]$ and $\mathrm{Var}[\mathbb{E}(Y \mid Q_1)]$. (b) Compute $\mathrm{Var}(Y)$ using the independence of $Q_1$ and $Q_2$. (c) Show that $S_1 = c_1^2/(c_1^2 + c_2^2)$ and $S_2 = c_2^2/(c_1^2 + c_2^2)$. (d) Verify $S_1 + S_2 = 1$ (no interaction for additive models).

*Answer:* (a) $\mathbb{E}[Y \mid Q_1] = c_1 Q_1 + c_2/2$. $\mathrm{Var}[\mathbb{E}(Y \mid Q_1)] = c_1^2 \,\mathrm{Var}(Q_1) = c_1^2/12$.
(b) $\mathrm{Var}(Y) = c_1^2 \,\mathrm{Var}(Q_1) + c_2^2 \,\mathrm{Var}(Q_2) = (c_1^2 + c_2^2)/12$.
(c) $S_1 = D_1/D = (c_1^2/12)/((c_1^2 + c_2^2)/12) = c_1^2/(c_1^2 + c_2^2)$. Similarly $S_2$.
(d) $S_1 + S_2 = (c_1^2 + c_2^2)/(c_1^2 + c_2^2) = 1$. For linear additive models, there are no interaction terms and first-order indices sum to 1.

## 9.5 ▪ Computing Sobol Indices: The Saltelli Algorithm

A naive Monte Carlo estimate of $S_i$ requires choosing $Q_i = q_i$ for many values and averaging the model over all other parameters for each, this costs $O(N^2)$ evaluations. The insight of Saltelli [47] is that all first-order and total Sobol indices can be estimated with $N(m+2)$ evaluations using two carefully constructed sample matrices. The design and estimators for the total sensitivity index are developed in Saltelli et al. [49].

### 9.5.1 ▪ The Saltelli Construction

| Matrix | Contents |
|---|---|
| $A$ | Base sample matrix ($N \times m$): each row follows the joint input distribution (by default, the first $m$ coordinates of a scrambled Sobol' point) |
| $B$ | Second sample matrix ($N \times m$): each row follows the same input distribution, independent of the same row of $A$ (by default, the last $m$ coordinates of the same point) |
| $A_B^{(i)}$ | Matrix $A$ with its $i$-th column replaced by the $i$-th column of $B$; all other columns come from $A$ |

**Table 9.1.** *Saltelli matrix notation. The key idea: $A_B^{(i)}$ and $A$ share the values of all parameters* except *$Q_i$, which is drawn from $B$. This "pick-freeze" design isolates the variance contribution of $Q_i$ from all other parameters.*

Table 9.1 fixes the notation used below. This chapter fills $A$ and $B$, each

$N \times m$, from one scrambled Sobol' sequence (§9.2.3) in $2m$ dimensions, with $N$ a power of two. The first $m$ coordinates of each point form a row of $A$, and the last $m$ form the same row of $B$. In Python, `scipy.stats.qmc.Sobol` with `scramble=True` applies a linear matrix scramble followed by a digital shift. Confidence intervals come from repeating the whole estimation under independent scrambles (§9.2.3); the bootstrap described below is the fallback, valid only when the rows are IID. Define the $m$ matrices $A_B^{(i)}$: matrix $A$ with its $i$-th column replaced by the $i$-th column of $B$. The model is evaluated at all rows of $A$, $B$, and all $m$ matrices $A_B^{(i)}$: a total of $N(m+2)$ evaluations. Figure 9.2 shows the structure for $m = 3$ columns.





**Figure 9.2.** *Saltelli pick-freeze matrix structure for $m = 3$ parameters (columns) and $N = 5$ sample rows. Blue cells carry values from the base matrix $A$; red cells carry values from the matrix $B$. $A_B^{(i)}$ shares every column with $A$* except *column $i$, which is "frozen" from $B$. This design ensures that $A_B^{(i)}$ and $A$ differ* only *in parameter $Q_i$, so the pair carries everything $Q_i$ contributes, alone and through all its interactions: it estimates $D\,S_{T_i}$ (see (9.7)). The total cost is $N(m+2)$ model evaluations: $N$ for $A$, $N$ for $B$, and $N$ for each of the $m$ hybrid matrices.*

### 9.5.2 ▪ The Jansen Estimators

For the first-order index, use the Jansen [25] estimator:

$$\hat{S}_i = 1 - \frac{\frac{1}{2N}\sum_{j=1}^{N}[f(B_j) - f(A_{B,j}^{(i)})]^2}{\hat{D}}, \tag{9.6}$$

where $\hat{D}$ is the variance of the $2N$ values $f(A_j)$ and $f(B_j)$ pooled: $\hat{D} = \frac{1}{2N}\sum_j [f(A_j)^2 + f(B_j)^2] - \left(\frac{1}{2N}\sum_j [f(A_j) + f(B_j)]\right)^2$.

For the total index:

$$\hat{S}_{T_i} = \frac{\frac{1}{2N}\sum_{j=1}^{N}[f(A_j) - f(A_{B,j}^{(i)})]^2}{\hat{D}}. \tag{9.7}$$

Use these Jansen estimators, not the original Saltelli (2002) formulas, which have higher variance. The Jansen estimator is consistent (converges to the

true $S_i$ as $N \to \infty$) and nearly unbiased; see Jansen [25]. A comprehensive comparison of eight total-order estimators across a wide range of models, sample sizes, input distributions, and dimensionalities finds that the Jansen estimator consistently outperforms the Saltelli (2002) formula for both factor prioritization and estimation accuracy [44]; when in doubt, the Jansen formula is the safer default. The key step is that

$$\mathbb{E}\left[\frac{1}{2N}\sum_{j=1}^{N}(f(B_j) - f(A_{B,j}^{(i)}))^2\right] = D - D_i = \mathrm{Var}(Y) - \mathrm{Var}[\mathbb{E}(Y \mid Q_i)],$$

which follows because $B_j$ and $A_{B,j}^{(i)}$ share the *same* value of parameter $i$ (both equal to $B$'s column $i$) while differing in all other parameters (one from $B$, the other from $A$). The shared value of $Q_i$ does not cancel in the difference $f(B_j) - f(A_{B,j}^{(i)})$ unless $f$ is additive in $Q_i$. For the worked model of Section 9.4 with $i = 1$, $f(b_1, b_2) - f(b_1, a_2) = (b_2 - a_2)(1 + b_1)$. The identity holds because the two evaluations share $Q_i$ while the other parameters are drawn twice, independently. Given $Q_i$, half the expected squared difference is then the conditional variance $\mathrm{Var}(Y \mid Q_i)$, and its mean over $Q_i$ is $\mathbb{E}[\mathrm{Var}(Y \mid Q_i)] = D - D_i$ by the law of total variance. Dividing by $\hat{D}$ and rearranging recovers $\hat{S}_i \approx D_i/D = S_i$. The estimator is nearly unbiased at finite $N$ because the bias is $O(1/N)$ [25], negligible for $N \geq 1000$.

### 9.5.3 ▪ Convergence

For IID rows the estimators converge as $O(1/\sqrt{N})$, the standard Monte Carlo rate. With scrambled Sobol' rows they typically converge faster. In practice, $N \geq 1000$ is needed for reliable estimates, and a power of two at or above that is the natural choice. For $m = 4$ parameters (SIR model): $N(m+2) = 6144$ evaluations at $N = 1024$. If each evaluation takes 0.1 seconds: 10 minutes.[4]

Plotting $\hat{S}_i$ and $\hat{S}_{T_i}$ vs. $N$ reveals the convergence: estimates are noisy for $N < 500$ and stable for $N \geq 2000$. This convergence plot is the diagnostic that answers the misconception "more samples always help": below a threshold $N$, additional samples do not produce reliable estimates because the variance of the estimator exceeds the signal being measured.

[4] An alternative path to total-order indices is the *extended FAST* (eFAST), which reaches the same indices via Fourier analysis along a space-filling search curve through the input space [56]. eFAST has different robustness properties at small $N$ but converges to the same values as the Jansen estimator; it is implemented alongside the matrix-based approach in the SALib Python library.

**Confidence intervals.** A point estimate $\hat{S}_i$ is useful but incomplete: it gives no information about sampling uncertainty. For a design whose rows are IID, the standard approach is the *bootstrap*: resample the $N(m+2)$ model evaluations with replacement, recompute $\hat{S}_i$ for each resample, and take the 2.5th and 97.5th percentiles of the resulting distribution as a 95% confidence interval.

```
# Bootstrap confidence intervals for Sobol first-order indices
nBoot = 500                              # number of bootstrap replicates
S1_boot = ZEROS(nBoot, m)                # will store replicate estimates

FOR b FROM 1 TO nBoot:
    idx = RANDOM_INTEGERS(N, from=1..N)  # resample the N base rows
    # the tuple (Y_A, Y_B, Y_AB^(1..m)) stays aligned within each row;
    # the model is never re-evaluated
    S1_boot[b, :] = JANSEN_FROM_OUTPUTS(Y_A[idx], Y_B[idx], Y_AB[idx, :]
END

S1_CI = PERCENTILE(S1_boot, [2.5, 97.5])  # 95% CI per index
```

Resampling by row is what makes the replicates valid: the Jansen estimator is a function of the matched tuples $(Y_A, Y_B, Y_{A_B}^{(1)}, \ldots, Y_{A_B}^{(m)})$, and breaking that alignment destroys the pick-freeze structure the estimator relies on. The procedure assumes the rows are independent, which holds for IID sampling. Under any stratified or low-discrepancy design they are not: that covers the scrambled Sobol' sequence and equally the Latin hypercube design of §9.2.2, whose whole purpose is to make the rows depend on one another. In either case the honest analogue is to repeat the whole analysis under independent scrambles, or independent LHS draws, rather than to resample rows.

A confidence interval that straddles zero indicates that the parameter's effect cannot be distinguished from noise at the current sample size. A confidence interval that excludes zero confirms the parameter is genuinely influential. As a practical rule: if the confidence interval width exceeds the index value, double $N$ before drawing conclusions.

**Worked numerical example: confidence intervals for the SIR model.** Apply the SIR model ($m = 3$ parameters: contact rate $c$, transmission probability $\beta$, infectious period $\tau_R$; QOI: peak infectious fraction $I_{\max}$) with ranges $\pm 50\%$ around nominal values $c = 8$ (a higher-contact setting than the running example), $\beta = 0.06$ (a probability per contact, dimensionless), and $\tau_R = 10$ days. Generate a Saltelli design with $N = 256$ sample pairs ($N(m+2) = 1280$ model evaluations total), a power of two for the reason

given in §9.12. The Jansen estimators applied to the full sample give:

$$\hat{S}_1^{c} = 0.32, \qquad \hat{S}_1^{\beta} = 0.33, \qquad \hat{S}_1^{\tau_R} = 0.33.$$

These are *point estimates*: they carry no information about sampling uncertainty. To quantify that uncertainty, repeat the whole estimation under $R = 32$ independent scrambles, meeting the 30 or more that §9.2.3 asks for, each a fresh design with $N = 256$; the design above is the first. The first four replicates produce:

| Scramble $r$ | $\hat{S}_1^{c}$ | $\hat{S}_1^{\beta}$ | $\hat{S}_1^{\tau_R}$ |
|---|---|---|---|
| 1 | 0.32 | 0.33 | 0.33 |
| 2 | 0.33 | 0.34 | 0.33 |
| 3 | 0.33 | 0.33 | 0.32 |
| 4 | 0.34 | 0.32 | 0.34 |
| ⋮ | ⋮ | ⋮ | ⋮ |
| 32 | 0.35 | 0.33 | 0.34 |

The mean of each column is the estimate. With $s$ the column's standard deviation, the 95% confidence interval is the mean $\pm\, t_{31,\,0.975}\, s/\sqrt{32}$, where $t_{31,\,0.975} \approx 2.04$:

$$\begin{aligned} c: &\quad \hat{S}_1^{c} = 0.332 \quad [0.330,\ 0.335] \\ \beta: &\quad \hat{S}_1^{\beta} = 0.332 \quad [0.329,\ 0.334] \\ \tau_R: &\quad \hat{S}_1^{\tau_R} = 0.332 \quad [0.329,\ 0.335] \end{aligned}$$

All three intervals exclude zero: each parameter is genuinely influential. All three also overlap almost completely, and that is the result, not a failure to resolve one. The three POIs enter this model only through $\mathcal{R}_0 = c\beta\tau_R$, and the $\pm 50\%$ ranges give them the same relative spread, so their first-order indices are equal as population quantities. The small differences among 0.32, 0.33 and 0.33 are Monte Carlo scatter about a common value, and no sample size will turn them into a ranking.

The practical lesson is the one a reader is most likely to need. Faced with three indices near 0.33 and told that they sum to about one, it is tempting to report the largest as the most influential parameter. Here that would be reporting the seed of a random number generator. The intervals are what stop you: they overlap, so the ordering they suggest is not there. Section 3.5, §8.5 and the Chapter 6 laboratory make the same point from three other directions: when POIs enter only through one combination,

sensitivity analysis can rank the combination but not the parameters that form it. All three intervals are narrow: each is under 0.006 wide, less than 2% of its index value. That is what 32 scrambles of the $N = 256$ design buy, at $32 \times 1280 = 40{,}960$ evaluations. More scrambles will not, however, separate the three: they are equal by construction, and a narrower interval only shows that more clearly. The row bootstrap, the fallback for IID rows, gives [0.21, 0.42], [0.20, 0.44] and [0.25, 0.42] on the first design (100 resamples). These are 5 to 8 times as wide as the interval $\pm t_{31,\,0.975}\, s$ that the scrambles assign one design, because its rows are not independent.

When $m$ is large, however, even a single Sobol run at $N = 2048$ may be prohibitively expensive; Morris screening addresses exactly this case.

## 9.6 ▪ Morris Screening

When $m$ is large (30 or more parameters), running Sobol at $N = 2048$ costs $N(m+2) = 65{,}536$ model evaluations, potentially prohibitive. Morris screening [34] provides a cheap first step: $r(m+1)$ evaluations (typically $r = 10$ trajectories) to identify which parameters are truly unimportant. An improved radial design that gives more uniform coverage with fewer trajectories is described by Campolongo et al. [11].

For each trajectory, one parameter is changed at a time by a fixed fraction $\Delta$ of its range. The *elementary effect* of parameter $i$ at point $\vec{q}$ is:

$$EE_i(\vec{q}) = \frac{f(\vec{q} + \Delta \vec{e}_i) - f(\vec{q})}{\Delta}. \tag{9.8}$$

After $r$ trajectories, the mean $\mu_i^* = (1/r)\sum |EE_i|$ and standard deviation $\sigma_i$ of the elementary effects are computed. Parameters with small $\mu_i^*$ are unimportant. Parameters with large $\sigma_i$ have interactions or nonlinear effects.

The Morris diagnostic plot ($\sigma_i$ vs. $\mu_i^*$) separates parameters into three groups: unimportant (small $\mu_i^*$, small $\sigma_i$); important and linear (large $\mu_i^*$, small $\sigma_i$); important and nonlinear or interacting (large $\mu_i^*$, large $\sigma_i$).

**Worked numerical example: Morris screening for a two-parameter model.** To see the computation in full, consider the model $f(u_1, u_2) = 3u_1 + u_2 + 2u_1u_2$ on the unit square $[0,1]^2$, with $\Delta = 1/2$ and $r = 3$ trajectories. The parameter $u_1$ has a larger main effect (coefficient 3 vs. 1) and interacts with $u_2$ through the cross term.

*Trajectory 1.* Start at (0.0, 0.0); change order: $u_1$, $u_2$.

$$f(0.0, 0.0) = 0, \quad f(0.5, 0.0) = 1.50, \quad f(0.5, 0.5) = 2.50.$$

$$EE_1 = \frac{1.50 - 0}{0.5} = 3.0, \qquad EE_2 = \frac{2.50 - 1.50}{0.5} = 2.0.$$

*Trajectory 2.* Start at (0.5, 0.5); change order: $u_2$, $u_1$.

$$f(0.5, 0.5) = 2.50, \quad f(0.5, 1.0) = 3.50, \quad f(1.0, 1.0) = 6.00.$$

$$EE_2 = \frac{3.50 - 2.50}{0.5} = 2.0, \qquad EE_1 = \frac{6.00 - 3.50}{0.5} = 5.0.$$

*Trajectory 3.* Start at (0.5, 0.0); change order: $u_1$, $u_2$.

$$f(0.5, 0.0) = 1.50, \quad f(1.0, 0.0) = 3.00, \quad f(1.0, 0.5) = 4.50.$$

$$EE_1 = \frac{3.00 - 1.50}{0.5} = 3.0, \qquad EE_2 = \frac{4.50 - 3.00}{0.5} = 3.0.$$

Collecting and summarizing:

| | Elementary effects | | | | |
|---|---|---|---|---|---|
| Parameter | Traj. 1 | Traj. 2 | Traj. 3 | $\mu_i^*$ | $\sigma_i$ |
| $u_1$ | 3.0 | 5.0 | 3.0 | 3.67 | 1.15 |
| $u_2$ | 2.0 | 2.0 | 3.0 | 2.33 | 0.58 |

where $\mu_i^* = (1/r)\sum_j |EE_i^{(j)}|$ and $\sigma_i$ is the sample standard deviation of the $r$ values. Both parameters have large $\mu^*$, so both are important. The sample $\sigma$ differ here, but that difference is an artifact of $r = 3$ and should not be read as structure. The elementary effects are exactly $EE_1 = 3 + 2u_2$ and $EE_2 = 1 + 2u_1$, so over the unit square both have population variance $4\,\mathrm{Var}(u) = 1/3$ and hence the same $\sigma = 1/\sqrt{3} \approx 0.577$. The cross term $2u_1u_2$ is symmetric, and it makes both parameters interacting to the same degree. On a Morris diagnostic plot both belong upper-right; with $r = 3$ one of them will land lower simply by chance. In practice, $r = 3$ is too few trajectories for reliable screening (the standard minimum is $r = 10$); see Figure 9.3 for an application to the SIR model with $r = 50$ and $m = 4$ parameters.

**Workflow**: run Morris screening first; fix unimportant parameters at their nominal values; run Sobol only on the important subset. This can reduce the Sobol cost by a factor of 5–10.

**When even reduced Sobol is too expensive: surrogate models.** If the model is computationally expensive and Morris screening reduces the parameter set to, say, five influential POIs, running Sobol at $N = 2048$ still requires $N(m + 2) = 14{,}336$ model evaluations. For a climate model where each run takes hours, this remains prohibitive.

The practical solution is a *surrogate model* (also called an emulator or meta-model): a cheap statistical approximation to the expensive model,





**Figure 9.3.** *Morris screening diagnostic plot ($\sigma_i$ vs. $\mu_i^*$) for the SIR model with four parameters: contact rate c, transmission probability $\beta$, infectious period $\tau_R$, and mean residence time $\tau_m$. Parameters with small $\mu_i^*$ are unimportant (lower left); large $\mu_i^*$ and small $\sigma_i$ indicate a linear effect (lower right); large $\mu_i^*$ and large $\sigma_i$ indicate a nonlinear or interacting effect (upper right). Only the two regions that contain points are labeled on the plot, since naming an empty region invites a search for data that is not there. The three transmission and recovery parameters (c, $\beta$, $\tau_R$) are important and exhibit nonlinear behavior at the ranges shown (±50% of nominal). The residence time $\tau_m$ has negligible influence on the 90-day epidemic peak and can be fixed at its nominal value before running the Sobol analysis, illustrating the screening workflow: a parameter may govern the model's long-term dynamics yet be irrelevant to the short-horizon question being asked. Computed with $r = 50$ trajectories, $\Delta = 0.5$.*

trained on a moderate number of carefully chosen model runs. Concretely, a Gaussian process surrogate fits a probability distribution over functions to the training runs, predicting the model output at any new input along with a confidence interval; a polynomial chaos expansion instead represents the output as a truncated series in orthogonal polynomials of the uncertain inputs, with coefficients estimated from the training set. Gaussian process emulators and polynomial chaos expansions are the most widely used surrogates. Once trained on roughly 50–200 model evaluations, a surrogate can be evaluated millions of times in seconds, making full Sobol estimation feasible even for computationally expensive simulations.

Surrogate-based global SA does not replace the methods in this chapter; it wraps them. The sampling designs of Section 9.2, the Saltelli construction, and the Jansen estimators all apply to the surrogate exactly as they apply to the original model. The surrogate simply makes the evaluations fast enough to converge. Smith [59] develops Gaussian process, polynomial, and spectral surrogates in Chapters 15–18, building on the LHS samples and Sobol indices treated in this chapter; readers continuing into surrogate-model construction will build on exactly the kind of training sets generated in this chapter's laboratories.

## 9.7 ▪ Sobol Indices for Correlated POIs

Standard Sobol decomposition assumes independent POIs. When parameters are correlated, three problems arise. Each index (9.4) still lies in $[0, 1]$, but the first-order indices can sum to more than 1. The standard estimators, applied to correlated inputs sampled jointly, can return estimates outside $[0, 1]$ at any sample size. The ANOVA decomposition (9.2) is no longer unique, because correlation destroys the orthogonality that guarantees uniqueness.

An estimate outside $[0, 1]$ is worth noticing, but it is not by itself evidence of correlation. Monte Carlo error produces the same symptom with perfectly independent inputs whenever a POI has little influence, because the Jansen estimate then scatters about zero and crosses it. The laboratory in §9.12 shows this: its inputs are independent, yet $\hat{S}_1^{\tau_m}$ comes out slightly negative in about half the runs, since $\tau_m$ barely moves $I_{\max}$ over a short outbreak. Read a negative or greater-than-one estimate first as a request for more samples, and only after the estimate has stabilized as a sign that the independence assumption deserves scrutiny.

Three approaches address correlated POIs in global SA:

**Grouped indices**: treat a cluster of correlated parameters as a single group. Compute the group Sobol index (the fraction of variance explained by the entire group jointly). This avoids allocating variance between correlated parameters, which is not well-defined.

**Modified estimators**: Kucherenko et al. [27] extend the Saltelli algorithm to correlated inputs using copula-based sampling (a copula models the dependence structure between parameters separately from their marginal distributions). This requires knowledge of the joint distribution of the correlated parameters.

**PRCC**: reduces the linear effect of all other inputs via rank regression before computing each correlation, making it more robust than simple Pearson correlation under mild-to-moderate input dependence. It does not, however, fully account for strong nonlinear correlations or capture higher-order dependence structures. Under mild correlation it is often the fastest practical screen.

**Practical recommendation**: for models with mildly correlated POIs, PRCC is a fast, accessible first screen. When correlation is strong or nonlinear, complement PRCC with grouped Sobol indices or Shapley effects, and report the correlation structure explicitly alongside the SA results.

Table 9.2 summarizes the tradeoffs.

**Shapley effects: the principled alternative.** For readers seeking a theoretically grounded attribution under correlated inputs, the *Shapley*

| Method | When to use | Key assumption | Implementation |
|---|---|---|---|
| PRCC | Monotonic model; fast screen | Monotonicity | `partialcorr` (MATLAB) |
| Grouped Sobol | Strong correlation; | Natural grouping | `sobol_jansen.m` |
| | want variance attribution | known a priori | with grouped inputs |
| Shapley effects | Strong correlation; | None beyond | `sensitivity` (R) |
| | no natural grouping | finite variance | or SALib (Python) |

**Table 9.2.** *Comparison of global SA methods for correlated POIs. When POIs are mildly correlated, PRCC is the fastest screen. When correlation is strong and a natural grouping exists (e.g., all transmission parameters), grouped Sobol gives a more honest variance attribution. Shapley effects require no grouping but are more computationally expensive.*

*effect* $\phi_j$ is the unique attribution satisfying four game-theoretic axioms: efficiency (the $\phi_j$ sum to total output variance), symmetry, dummy (zero importance for irrelevant inputs), and additivity [41]. For independent inputs, the Shapley value of input $j$ reduces to the closed form $\phi_j = \sum_{u \ni j} \sigma_u^2/|u|$: each interaction variance component $\sigma_u^2$ is shared equally among all inputs in the interaction set $u$. Under correlated inputs this closed form no longer holds, and the general game-theoretic definition (averaging marginal contributions over all orderings of the inputs) is required. Normalizing by the total output variance gives the *normalized Shapley effect* $\Phi_j = \phi_j/\mathrm{Var}(Y)$, which lies in $[0, 1]$ and satisfies $\sum_j \Phi_j = 1$. A practical consequence: when inputs are *independent*, the first- and total-order Sobol indices already computed in this chapter bracket the normalized Shapley effect,

$$S_j \;\le\; \Phi_j \;\le\; S_{T_j},$$

providing useful bounds on the fair attribution without additional cost. Under correlated inputs, the standard Sobol indices themselves become unreliable (as noted earlier in this chapter), so this bracketing should not be relied upon in the correlated case. The Shapley value is computationally more expensive than Sobol indices but is implemented in SALib and the R `sensitivity` package.

## 9.8 ▪ Beyond Variance: Moment-Independent Sensitivity

Sobol indices decompose output *variance*. For epidemic models with skewed or multimodal output distributions, the distribution of peak infection counts, for instance, is rarely symmetric; variance is a poor summary of uncertainty. A parameter that dominates variance is not necessarily the parameter whose uncertainty most shifts the shape of the epidemic trajectory, and the parameter most critical to a decision-maker who cares about tail events

(catastrophic outbreaks) may differ from the one ranked highest by Sobol indices.

Borgonovo (2007) introduced a moment-independent sensitivity measure, the *δ-index* [6], that addresses this directly. Instead of measuring how much fixing $Q_i$ reduces output variance, $\delta_i$ measures the *expected shift in the entire output distribution* when $Q_i$ is fixed:

$$\delta_i = \frac{1}{2}\, E_{Q_i}\left[\int \left| f_Y(y) - f_{Y|Q_i}(y) \right| dy\right]. \tag{9.9}$$

Here $f_Y$ is the unconditional output density and $f_{Y|Q_i}$ is the conditional density given $Q_i$. The integral $\int |f_Y(y) - f_{Y|Q_i}(y)|\, dy$ is twice the total variation distance between the two densities; the factor $\frac{1}{2}$ outside the expectation normalizes $\delta_i$ to lie in $[0, 1]$. The expectation averages over all values $Q_i$ can take. Three properties make $\delta_i$ interpretable: it lies in $[0, 1]$; it equals zero when $Y$ is independent of $Q_i$; and the joint $\delta$ of all inputs together equals one.

*When do $\delta_i$ and Sobol indices disagree?* Borgonovo (2007) demonstrates on the Ishigami test function that $\delta_i$ and total-order Sobol indices agree on which parameters are *least* important, but can disagree on the most important ones: the parameter that most shifts the output distribution is not always the one that contributes most to output variance. In epidemic models this distinction matters when the decision criterion is not variance but exceedance probability: the probability that $\mathcal{R}_0$ exceeds the critical threshold of 1, or that peak infections exceed hospital capacity. Lu and Borgonovo [29] demonstrate this discrepancy for SIR-type models.

**Definition is valid under correlated inputs.** Because $\delta_i$ requires only the joint density $f_{\mathbf{Q}}$ without assuming independence, it remains well-defined when parameters are correlated [6], a practical advantage over standard Sobol indices in epidemiological models where transmission rate and contact rate are rarely independent.

**Computational cost and software.** Computing $\delta_i$ requires estimating conditional densities, which is more expensive than the matrix construction for Sobol indices. Plischke, Borgonovo and Smith (2013) introduced a substantially more efficient "given-data" estimator. Both the standard and efficient estimators are implemented in SALib (Python) and the `sensitivity` package (R). For a comprehensive graduate-level treatment of moment-independent sensitivity measures, their mathematical properties, and computational algorithms, see Borgonovo [7].

## 9.9 ▪ Comparison of the Main SA Methods

Table 9.3 consolidates the main SA methods of the book. It omits the $\sigma$-normalized index, Shapley effects, the $\delta$-index, eFAST and forward-mode AD. For the choice among local methods, Table 7.2 remains the finer guide; it gives the FSE's condition, $m \leq r$, and its cost, $m+1$ solves. For a broader treatment of global SA methods, see Saltelli et al. [52, 50] and the recent epidemiological applications in Lu and Borgonovo [29].

| Method | Chapter | When to use | Assumption | Cost |
|---|---|---|---|---|
| Local SI (analytic) | 2, 4 | Differentiable model; small uncertainty | Linearization valid | Cheap |
| Local SI (FD) | 3 | Black-box; small uncertainty | Linearization valid | $2m + 1$ evals |
| Adjoint | 5–6 | Differentiable; many POIs; few QOIs | Linearization valid | 2 solves |
| AD (reverse) | 7 | Differentiable code; $m > r$ | Linearization valid | 2 passes |
| PRCC | 9 | Monotone model; moderate uncertainty; corr. POIs OK | Monotone | $N$ evals |
| Morris | 9 | Screening; $m \geq 30$ | None | $N(m + 1)$ evals |
| Sobol | 9 | Nonlinear; large uncertainty; independent POIs | Independent | $N(m + 2)$ evals |
| Grouped Sobol | 9 | Correlated POIs; known groups | Known groups | $N(m + 2)$ evals |

**Table 9.3.** *Comparison of the main SA methods across the chapters. The choice of method depends primarily on whether the model is differentiable, whether uncertainty is small or large, and whether* POIs *are independent. No single method is universally best.*

Table 9.3 settles which *method* to run. A second question survives it: once the indices are in hand, which index answers the question actually being asked? Table 9.4 sets that out. The two failure modes it guards against are not exotic. Quoting $S_{T_i}$ as the variance reduction available from mea-

suring a parameter overstates the gain whenever interactions are present, and reading a variance-based index as a proportional-change prediction attributes to it information it does not carry.

| The question | The index | Why not the other one |
|---|---|---|
| Which POI should we measure or learn first, to reduce output uncertainty most? | First-order Sobol $S_i$ | Learning $Q_i$ leaves $(1 - S_i)\,\mathrm{Var}(Y)$, so the reduction is $S_i$. $S_{T_i}$ overstates it whenever interactions are present. |
| Which POIs can we fix at nominal and stop resolving? | Total-order Sobol $S_{T_i}$ | A small $S_{T_i}$ licenses fixing. A small $S_i$ does not: a purely interacting parameter has $S_i \approx 0$ with $S_{T_i}$ large. |
| If we change a POI by a given percentage, how far does the QOI move? | Normalized local SI $\mathcal{S}_p^q$ (elasticity), Chapter 2 | Variance-based indices carry no proportional-change information; they describe dispersion, not response. |
| Which POIs can we screen out cheaply before an expensive study? | Morris $\mu_i^*$ | Screening only. $\mu_i^*$ ranks; it does not decompose variance. |
| Which POI drives output uncertainty, given how uncertain it actually is? | $\sigma$-normalized $S_i^\sigma$, or Sobol | The plain local SI scales by the nominal value, not the range, and so ignores how much the POI actually varies (Section 9.1). |

**Table 9.4.** *Which index answers which question. Every entry in the third column is a mistake that appears in the applied literature. The distinction matters most when $S_{T_i} - S_i$ is large; when interactions are weak the rankings often agree, which is exactly why the difference is easy to overlook.*

A caution that spans the whole table. These indices rank parameters *of a model*. Feasibility, cost, and whether a lever can be moved at all are not in any of them, and no amount of resampling will put them there. The index identifies what to target; it does not say how, or whether.

*Unifying Idea 2.* The adjoint row in Table 9.3 is the payoff for Chapters 5 and 6. The adjoint is the transpose in whatever space you are working in:

matrix transpose for linear systems (Chapter 5), reversed ODE for dynamical systems (Chapter 6), and reversed chain rule for computation graphs (Chapter 7). Every "2 solves" entry in the adjoint row is the same unifying idea, applied in a different mathematical space.

## 9.10 ▪ Verifying an SA Result You Did Not Compute

Most of this book teaches you to produce a sensitivity result. This section addresses the complementary task, which you will perform more often: judging one that arrives already finished.

The situation is ordinary. A collaborator reports that the transmission rate dominates. A library returns total-order indices for a model you did not write. A paper under your review states a parameter ranking in a single sentence. A computational assistant produces a Sobol decomposition in seconds. Your own analysis from six months ago comes back with the reasoning no longer in memory. In every case the arithmetic is probably right, and that is not the question. The question is whether the quantity computed answers the question being asked, and Table 9.4 exists because the published literature shows that it frequently does not.

Two features of global SA make this failure easy to reach. Computing an index is cheap while choosing one is hard, so nothing in the workflow forces the choice to be made explicitly. And every method returns a number for every model, whether or not its assumptions hold, so a violated assumption produces a plausible answer rather than an error. A result can be arithmetically correct, confidently reported, and still misleading.

### 9.10.1 ▪ An audit in six questions

The two tables already assembled in this chapter carry most of what is needed. Table 9.4 maps questions to indices, and Table 9.3 records the assumption each method makes. The following sequence uses them.

1. *What decision does the result inform?* Not what was computed, but what will be done differently depending on the answer. Measure a parameter first, fix one at nominal, or predict how far the QOI moves under a stated change. The literature names the first two factor prioritization and factor fixing; the third asks for a proportional response. They are three different decisions, and Table 9.4 assigns each a different index. Factor mapping, the search for input combinations that drive the output above a threshold, is a separate purpose, and the table has no row for it.

2. *Does the reported index answer that question, or an adjacent one?* The third column of Table 9.4 lists the substitutions that occur in practice. Quoting $S_{T_i}$ as the uncertainty reduction available from measuring a parameter overstates it whenever interactions are present. Reading a variance-based index as a proportional-change prediction attributes to it information it does not carry: it describes dispersion, not response.

3. *What does the method assume?* Read it from the assumption column of Table 9.3. PRCC assumes a monotone response. Standard Sobol estimation assumes independent POIs. Every local method in Chapters 2 through 8 assumes the linearization is valid over the range of interest.

4. *Does this model satisfy that assumption?* This is the step most often skipped, because it requires knowing the model rather than the output. A non-monotone response invalidates a PRCC ranking without producing any symptom in the PRCC value. Correlated inputs invalidate a standard Sobol decomposition while the indices still sum to something near one.

5. *Is the estimate stable enough to support the ranking?* Sobol and Morris estimates carry Monte Carlo error, and a ranking read from point estimates whose confidence intervals overlap is not a ranking. This is why Section 9.5 gives confidence intervals. A result reported without any indication of sampling error has not established that its order would survive a second run at a different seed.

6. *Is the claim about the model or about the world?* Every index in this book is computed from a model's equations and inherits every simplification in them. A parameter absent from the model returns a sensitivity of exactly zero, which establishes nothing about the system. A result that has traveled from its author to you has usually shed the assumptions that would let you tell the two claims apart.

*Unifying Idea 4.* Steps 1 and 6 are the same commitment at the two ends of an analysis. The first asks what question the model is being made to answer; the last asks what the answer is a statement about. A finished result that has lost both is a number without a claim attached.

### 9.10.2 ▪ A worked audit

Return to the example of Section 9.1. A model $Y = 2Q_1 + Q_2$ is reported with the finding that $Q_1$ is the more important parameter, since its coefficient is twice as large. The arithmetic is correct.

Step 1 asks what will be done with this. Suppose the intent is to decide which parameter to measure more precisely, in order to reduce uncertainty in $Y$. Step 2 sends us to Table 9.4, whose first row assigns that decision to a first-order Sobol index, or equivalently here to the $\sigma$-normalized index, rather than to a comparison of coefficients. Step 3 records what the reported quantity assumes, which is that the two parameters are comparably uncertain. Step 4 checks it: with $\sigma_1 = 1$ and $\sigma_2 = 3$, they are not. Weighting each coefficient by the uncertainty it multiplies gives contributions of 2 and 3, and the ranking reverses. Measuring $Q_2$ reduces the variance of $Y$ more.

The reported finding is not a computational error. It answers the question of which parameter has the larger coefficient, which was not the question asked. Step 5 would ask whether the reversal survives sampling error, and step 6 would note that both rankings are properties of this model, which assumes the response is linear and the two parameters independent.

### 9.10.3 ▪ The referee's version

The same six questions are the ones to ask of a submitted manuscript, and they are worth practicing for that reason alone. A paper reporting a parameter ranking should make its decision context explicit, name the index it used, state the method's assumption, give some evidence the model satisfies it, report sampling error where the estimate is stochastic, and confine its claims to the model it analyzed. Sensitivity analyses in the applied literature routinely omit several of these, which is the observation behind the caption of Table 9.4.

Responsibility here rests with authors rather than reviewers. Publishers have begun to say so explicitly as computational assistance has become routine: SIAM's policy for authors places the burden of ensuring correctness on the author rather than the referee, for mathematical content and for code alike, and holds authors responsible for the accuracy of their figures and citations as well. The audit above is one way to discharge that responsibility on the sensitivity analysis in your own paper before anyone else reads it.

**Software for global SA beyond MATLAB.** The methods in this chapter are implemented in several open-source libraries. SALib [19] is the dominant Python library for global SA; it implements Morris screening, the Saltelli construction with Jansen estimators, eFAST, and the Borgonovo $\delta$-index under a consistent API and is freely available at `https://salib.readthedocs.io`. The `sensitivity` package for R (Iooss et al.) covers a similar range of methods including Shapley effects. Both are well-documented, actively maintained, and suitable for research-grade applications.

## 9.11 ▪ Bridge to the Broader UQ Landscape

This book quantifies how model output uncertainty arises from parameter uncertainty, assuming the model structure is correct. A complementary graduate-level treatment of the broader UQ landscape, given in Smith [59], takes up the harder questions that follow: what if the model is wrong (Smith Ch. 14)? How do we fit models to data (Chs. 11–12)? How do we construct efficient surrogate models to replace expensive simulations (Chs. 15–18)?

Smith's Chapters 11–12 develop frequentist and Bayesian parameter estimation, including Markov chain Monte Carlo methods for fitting models to data, the natural sequel to the identifiability analysis introduced in Chapter 3 of this book.

The connections between the two treatments are direct:

Smith Ch. 8 develops local SA at the graduate level, extending Chapters 2–7 of this book with functional-analytic rigor.

Smith Ch. 9 develops global SA and Sobol indices at the graduate level, extending Chapter 9 of this book with measure-theoretic foundations.

Smith Ch. 7 treats parameter identifiability and influence using the sensitivity matrix; the core tool introduced in Chapter 4 of this book.

Smith Ch. 14 develops model discrepancy, and Chs. 15–18 develop surrogate models including Gaussian process and polynomial chaos methods that reduce the cost of global SA by replacing the expensive model with a cheap approximation. Surrogates are built from space-filling samples of the kind this chapter uses, LHS or scrambled Sobol'.

A complementary modern method between the local indices of Chapters 2–7 and the variance-based global indices of this chapter is *active subspace analysis*, which finds low-dimensional directions in parameter space along which the response varies most. Smith [59] Ch. 10 develops this technique and its connection to global sensitivity scoring; it is a natural next step for readers whose models have many parameters but few *effective* ones.

Readers who have completed this book have the foundational vocabulary, the computational intuition, and the mathematical framework to engage productively with the full UQ literature. The path from the sensitivity index in Chapter 2 to the full Bayesian UQ framework in Smith is a single, continuous intellectual development. As an example of these methods applied to a realistic multi-parameter epidemic model, see Qu, Xue, and Hyman [45], which uses SA to guide Wolbachia-based mosquito-borne disease control.

The reader who has reached this point knows how to ask which parameters matter, how to compute that answer efficiently and honestly, and, just as importantly, when not to trust the answer. Every mathematical model is an argument; sensitivity analysis is the discipline that reveals which as-

sumptions the argument depends on.

*Good sensitivity analysis does not end the discussion. It tells you where to begin it.*

## 9.12 ▪ Computational Laboratory: Complete Global SA Pipeline for SIR

This laboratory implements the full global SA pipeline for the SIR model and compares all four methods: sigma-normalized, PRCC, Sobol, and Morris.

**Setup.** SIR model with four POIs $(c, \beta, \tau_R, \tau_m)$ and response $J = I_{\max}$ (peak infectious count). Use the literature-based uncertainty ranges from Chitnis et al. [14]: $c \in [2, 8]$, $\beta \in [0.02, 0.12]$, $\tau_R \in [4, 14]$ days, $\tau_m \in [1000, 100000]$ days.

**Part 1: LHS and sigma-normalized indices.** Draw $N = 500$ LHS samples using the `lhs_sample` function. Compute $\sigma_J$ from the samples. Compute the sigma-normalized index for each POI using the local SI from Chapter 2 and the sample standard deviations. Report the ranking.

**Part 2: PRCC.** Run the SIR model at all 500 LHS samples. Rank-transform all inputs and the output. Before reading the result, note where the box sits relative to the epidemic threshold. About 10.8% of it has $\mathcal{R}_0 \leq 1$, and across that region $I_{\max}$ is pinned at $I_0$ and does not respond to the POIs at all. PRCC assumes a monotonic input-output relationship. A flat stretch followed by a rise is still monotone, so the flat region does not break that assumption. It does fill the ranks of $I_{\max}$ with ties (§8.2.2 describes the threshold behind it). Report the PRCC on the full sample and again on the super-threshold subsample, and compare. On this box expect the two to agree closely and to rank the POIs in the same order. Compute PRCC using the `partialcorr` function in your implementation's Statistics and Machine Learning Toolbox (or download Marino et al.'s implementation from the supplementary material of [31]). Produce a PRCC tornado plot and report $p$-values.

**Part 3: Sobol estimation.** Draw the sample from a scrambled Sobol' sequence (MATLAB: `sobolset`; Python: `scipy.stats.qmc.Sobol`), and take $N = 2048$. Sobol sequences are balanced in blocks of $2^k$, so truncating mid-block degrades the low-discrepancy property that makes quasi-random sampling worth using; most libraries will warn you when $N$ is not a power of two. The same reasoning does not apply to a Halton sequence, which is not built in binary blocks, so a Halton design may use any $N$. Build the $A$, $B$, and $m$ matrices $A_B^{(i)}$: $2048 \times 6 = 12{,}288$ total evaluations. Compute $\hat{S}_i$ and $\hat{S}_{T_i}$ using the Jansen estimators.

```
# Sobol variance decomposition (Jansen estimator)
# See the companion notebooks (Python) and reference library (MATLAB)

set_random_seed(42)                                  # reproducibility
A, B        = SALTELLI_SAMPLE(N, m, lb, ub) # two N-by-m quasi-random ma

S1, ST, CI = SOBOL_JANSEN(model, A, B,
                          bootstrap_reps=500, alpha=0.05)
# S1[j]     = first-order Sobol index for parameter j
# ST[j]     = total-order Sobol index for parameter j
# CI[j,:]   = [lower, upper] 95% confidence interval for S1[j]
```

The call's `bootstrap_reps` option resamples rows, so its intervals apply to IID rows only. For this scrambled Sobol' design, take the intervals from independent scrambles instead, as in §9.5.

**Part 4: Morris screening.** Implement $r = 10$ Morris trajectories with $\Delta = 0.4$. Compute $\mu_i^*$ and $\sigma_i$ for each parameter. Produce the Morris diagnostic plot ($\sigma_i$ vs. $\mu_i^*$). Which parameter(s) could be fixed without significant loss of accuracy?

**Part 5: Convergence check.** Plot $\hat{S}_\beta$ and $\hat{S}_{T_\beta}$ vs. $N$ for $N \in \{128, 256, 512, 1024, 2048, 4096\}$, staying on powers of two for the reason given in Part 3. At what $N$ do the estimates stabilize to within $\pm 0.05$?

**Part 6: Full comparison table.** Fill in the following table from your results:

| Parameter | Local SI | $S_i^\sigma$ | PRCC | $S_i$ (Sobol) |
|---|---|---|---|---|
| $c$ | | | | |
| $\beta$ | | | | |
| $\tau_R$ | | | | |
| $\tau_m$ | | | | |

Where do the four methods agree? Where do they disagree? What accounts for the differences?

## 9.13 ▪ Exercises

**Exercise 9.1** *([Bloom L3])*. The local SIs and $\mathcal{R}_0^*$ in this exercise are illustrative. They do not follow from the SEIR model of Exercise 2.5, which uses the same nominal $c$, $\beta$, $\tau_E$ and $\tau_I$. For the SEIR model with local SIs $\mathcal{S}_{\tau_E}^{\mathcal{R}_0} = 0.6$, $\mathcal{S}_{\tau_I}^{\mathcal{R}_0} = 1.2$, $\mathcal{S}_{\beta}^{\mathcal{R}_0} = 1.0$, $\mathcal{S}_{c}^{\mathcal{R}_0} = 1.0$ and uncertainty ranges $\sigma_{\tau_E} = 2$ days, $\sigma_{\tau_I} = 3$ days, $\sigma_\beta = 0.01$, $\sigma_c = 0.8$, at nominal parameter values $\tau_E^* = 5.5$ days, $\tau_I^* = 7$ days, $\beta^* = 0.06$, $c^* = 5$, $\mathcal{R}_0^* = 2.0$:

(a) Compute the output standard deviation $\sigma_{\mathcal{R}_0}$ using the delta method, at nominal $\mathcal{R}_0^* = 2.0$.

(b) Compute all four sigma-normalized indices.

(c) Produce a tornado plot of the sigma-normalized indices.

(d) Compare the ranking with the local SI ranking. Which parameter changed rank most dramatically? Why?

**Exercise 9.2** *(*[Bloom L3]*).*

(a) Implement the `lhs_sample` function. For $N = 100$, $m = 3$, $[0, 1]^3$, generate an LHS and a random sample of the same size.

(b) Produce scatter plots of all three pairs of parameters for both samples (6 plots total).

(c) Compute the mean and standard deviation of the sample along each dimension. Which method is closer to the true uniform mean and standard deviation?

(d) For one parameter, plot the histogram of LHS vs. random sampling. Which covers the range more evenly?

**Exercise 9.3** *(*[Bloom L3]*).* For the SIR model, with each of the four parameters ranging over $\pm 50\%$ about its nominal value:

(a) Generate $N = 200$ LHS samples and run the SIR model at each.

(b) Compute PRCC for all four parameters with respect to $I_{\max}$.

(c) Produce a PRCC bar chart sorted by |PRCC|.

(d) Which parameter has the strongest negative PRCC? What does this mean in plain language?

(e) Compare the PRCC ranking with the local SI tornado plot from Chapter 2. Do they agree?

**Exercise 9.4** *(*[Bloom L4]*).* For $Y = 2Q_1^2 + Q_2$ with $Q_1, Q_2 \sim \mathcal{U}(0, 1)$, independent:

(a) Compute $\mathbb{E}[Y]$, $\mathrm{Var}(Y)$, $D_1 = \mathrm{Var}[\mathbb{E}(Y \mid Q_1)]$, $D_2 = \mathrm{Var}[\mathbb{E}(Y \mid Q_2)]$.

(b) Compute $S_1$, $S_2$. Do they sum to 1? Explain.

(c) Compute the total indices $S_{T_1}$, $S_{T_2}$.

(d) Now add the product $Q_1Q_2$ to get $Y' = 2Q_1^2 + Q_2 + Q_1Q_2$ (an interaction term). Without computing: does adding the interaction term change $S_1$, $S_2$, or both? In which direction? Then verify by computing the Sobol indices for $Y'$ analytically.

**Exercise 9.5** *([Bloom L4]).* Using the Jansen estimator implementation from the laboratory, with each of the four SIR parameters ranging over $\pm 50\%$ about its nominal value:

(a) Compute $\hat{S}_i$ and $\hat{S}_{T_i}$ for all four SIR parameters at $N = 2048$.

(b) For which parameters is $S_{T_i} \gg S_i$? What does this indicate about their role in the model?

(c) Compute $\sum_i S_i$ and $\sum_i S_{T_i}$. What do you expect these to sum to for an additive model? Are your results consistent with this expectation?

(d) If you could measure exactly one parameter very precisely (reducing its uncertainty to zero), which parameter would reduce $\mathrm{Var}(I_{\max})$ the most? Justify using the Sobol indices.

**Exercise 9.6** *([Bloom L4]).* For the SIR model, implement $r = 15$ Morris trajectories with $\Delta = 1/3$:

(a) Compute $\mu_i^*$ and $\sigma_i$ for each parameter.

(b) Produce the Morris diagnostic plot.

(c) Based on the Morris results, which parameter appears most influential? Which appears least influential?

(d) Run a reduced Sobol analysis using only the three most important parameters identified by Morris (fixing $\tau_m$ at its nominal value). Compare the cost and results with the full four-parameter Sobol.

**Exercise 9.7** *([Bloom L5]).* Choose a published epidemic model from the literature with at least 8 parameters. Design a complete global SA study:

(a) State the model, its parameters, and a justification for the uncertainty ranges you will use (with citations).

(b) State the QOI or QOIs and explain why they are the most relevant for the model's purpose.

(c) Choose between PRCC and Sobol as the primary global SA method. Justify your choice based on the model's properties.

(d) Specify the sample size $N$ and justify it based on convergence.

(e) Describe how you would present the results to a non-technical audience (e.g., a public health committee or funding agency).

(f) Identify one situation where your global SA results could still be misleading, and describe what additional analysis you would recommend.

**Exercise 9.8** *([Bloom L4]). Audit exercise.* A collaborator reports total-order Sobol indices for a four-parameter model and writes: "$S_{T_\beta} = 0.62$, so measuring $\beta$ more precisely would remove about 62% of the uncertainty in the peak infection count. That is the single most valuable measurement we can make." The estimator was implemented and run correctly.

(a) Apply the audit of Section 9.10.1. Identify which decision the collaborator is making and which row of Table 9.4 governs it.

(b) State the correct index for that decision and explain, in terms of the variance decomposition, why the reported one overstates the gain.

(c) Under what condition on the model would the two indices give effectively the same answer? Explain why that condition makes the error easy to overlook in practice.

(d) The final sentence contains a second claim that no sensitivity index of any kind can support. Identify it.

**Exercise 9.9** *([Bloom L5]). Audit exercise.* Two analyses of the same model are reported. The first uses PRCC on $N = 500$ Latin hypercube samples and ranks the parameters $(p_1, p_2, p_3)$ in that order. The second uses first-order Sobol indices on $N = 2048$ samples and ranks them $(p_3, p_1, p_2)$. Both are computed correctly. The model's response to $p_3$ is known to rise and then fall over the sampled range.

(a) Using Table 9.3, state the assumption each method makes and determine which is violated here.

(b) Explain how a violated monotonicity assumption can produce a low PRCC for a parameter that in fact drives a large share of the output variance.

(c) Neither report gives confidence intervals. Describe how you would obtain them for each analysis, and state what result would lead you to conclude that the two rankings do not actually disagree.

(d) Suppose the intervals confirm the disagreement. Write the one-sentence finding you would report, taking care that it is a claim about the model and not about the system.

**Exercise 9.10** *([Bloom L6]).* A published SA study reports Sobol first-order and total indices for a 15-parameter water-quality model. The authors note in the data section that three parameters, phosphorus loading rate, nitrogen loading rate, and water residence time, are positively correlated across the sampled watersheds. The SA section reports standard Sobol indices computed using independent sampling for all 15 parameters, without addressing the correlation.

(a) Identify the specific mathematical problem that correlation creates for standard Sobol indices.

(b) For each of the three correlated parameters, state whether the reported first-order index is likely over-estimated, under-estimated, or unchanged, and explain why.

(c) Propose the minimal methodological correction that would make the analysis valid.

(d) The authors justify their approach by noting that their Sobol indices sum to approximately 1.0. Explain why this is not a valid check for independence.

## Notes

The ANOVA decomposition of output variance for functions of independent uniform inputs, the theoretical foundation of Sobol indices, was established by Sobol' [60]. The efficient matrix-based estimators adopted in this chapter are due to Saltelli [47, 49] and were substantially improved by Jansen [25], whose estimators have smaller variance than the original Saltelli formulas and are now the recommended default in major software libraries.

Draft

The Latin hypercube sampling design (Section 9.2) was introduced by McKay, Beckman, and Conover [32]; Iman and Conover [24] developed the rank-correlation variant that is standard in practice. Helton and Davis [18] give a comprehensive assessment of LHS performance for SA applications.

Morris screening was introduced in Morris [34]; the radial design improvement that gives more uniform coverage with fewer trajectories is due to Campolongo, Cariboni, and Saltelli [11].

The moment-independent $\delta$-index was introduced by Borgonovo [6] and is given a comprehensive mathematical treatment in Borgonovo [7]. The extension of Sobol indices to correlated inputs uses copula-based sampling as described by Kucherenko et al. [27]. The Shapley effect, which provides a unique attribution satisfying game-theoretic axioms under correlated inputs, was developed by Owen [41]. Applications of global SA to SIR-type epidemic models, including the discrepancy between variance-based and moment-independent rankings, are demonstrated by Lu and Borgonovo [29]. A comprehensive review of global SA methods with software comparisons is given by Puy et al. [44].

Draft

Draft

# Appendices

Draft

Draft

# Appendix A

# Inner Products, Norms, and Function Spaces

This appendix provides the minimum background in functional analysis needed to read Appendix B and the related material in Smith [59]. Readers who are comfortable with inner products and the $L^2$ space can proceed directly to Appendix B.

## A.1 ▪ Inner Products and Norms

An *inner product* on a vector space $V$ over $\mathbb{R}$ is a function $\langle \cdot, \cdot \rangle : V \times V \to \mathbb{R}$ satisfying:

1. *Symmetry*: $\langle u, v \rangle = \langle v, u \rangle$.
2. *Linearity in the first argument*: $\langle \alpha u + \beta w, v \rangle = \alpha \langle u, v \rangle + \beta \langle w, v \rangle$.
3. *Positive definiteness*: $\langle u, u \rangle \geq 0$, with equality only when $u = 0$.

The *induced norm* is $\|u\| = \sqrt{\langle u, u \rangle}$. The Cauchy-Schwarz inequality states $|\langle u, v \rangle| \leq \|u\| \, \|v\|$.

**Example A.1 (Euclidean inner product).** For $\vec{u}, \vec{v} \in \mathbb{R}^n$:

$$\langle \vec{u}, \vec{v} \rangle = \vec{u}^T \vec{v} = \sum_{i=1}^{n} u_i v_i.$$

This is the dot product. The induced norm is the Euclidean norm $\|\vec{u}\|_2 = \sqrt{\sum u_i^2}$.

**Example A.2 (Weighted inner product).** For a symmetric positive definite matrix $W \in \mathbb{R}^{n \times n}$: $\langle \vec{u}, \vec{v} \rangle_W = \vec{u}^T W \vec{v}$. The induced norm $\|\vec{u}\|_W = \sqrt{\vec{u}^T W \vec{u}}$ is the $W$-weighted Euclidean norm. SA computations sometimes use this when parameters have very different scales.

## A.2 ▪ The $L^2$ Inner Product

For functions $f, g : [a, b] \to \mathbb{R}$ that are square-integrable (i.e., $\int_a^b f(t)^2\,dt < \infty$), the $L^2$ *inner product* is:

$$\langle f, g\rangle_{L^2} = \int_a^b f(t)\,g(t)\,dt. \tag{A.1}$$

The induced norm is $\|f\|_{L^2} = \sqrt{\int_a^b f(t)^2\,dt}$.

The space $L^2([a, b])$ of all square-integrable functions on $[a, b]$ with inner product (A.1) is a *Hilbert space*: a complete inner product space. Completeness means that every Cauchy sequence converges to a function in $L^2([a, b])$.

For Sobol decomposition (Chapter 9), the relevant space is $L^2([0, 1]^m)$ with inner product $\langle f, g\rangle = \int_{[0,1]^m} f(\vec{q})g(\vec{q})\,d\vec{q}$.

## A.3 ▪ Integration by Parts as Transposition

The key connection between adjoint sensitivity analysis and linear algebra is this: integration by parts is the function-space analog of the matrix transpose.

In $\mathbb{R}^n$, for vectors $\vec{u}$, $\vec{v}$ and a matrix $A$:

$$\langle A\vec{u}, \vec{v}\rangle = (A\vec{u})^T\vec{v} = \vec{u}^T(A^T\vec{v}) = \langle \vec{u}, A^T\vec{v}\rangle.$$

So $A^T$ is the operator satisfying $\langle Au, v\rangle = \langle u, A^T v\rangle$.

In $L^2([0, T])$, for functions $u, v$ and the differentiation operator $L = d/dt$:

$$\langle Lu, v\rangle_{L^2} = \int_0^T \dot{u}(t)\,v(t)\,dt = \big[u(t)v(t)\big]_0^T - \int_0^T u(t)\,\dot{v}(t)\,dt = \big[uv\big]_0^T + \langle u, L^*v\rangle_{L^2},$$

where $L^* = -d/dt$ (with appropriate boundary conditions) is the *adjoint operator* of $L = d/dt$. The boundary term $[uv]_0^T$ gives rise to the terminal condition $\lambda(T) = +h'(u(T))$ in the adjoint ODE derivation of Chapter 6.

This correspondence makes precise what we stated informally throughout the book: the adjoint ODE is the ODE whose differential operator is the transpose of the forward ODE's operator, exactly as $A^T$ is the transpose of $A$.

## A.4 ▪ Orthogonality and the ANOVA Decomposition

Two elements $u, v$ of an inner product space are *orthogonal* if $\langle u, v\rangle = 0$.

In $L^2([0, 1]^m)$ with the uniform measure, the functions appearing in the Sobol ANOVA decomposition (Chapter 9) are mutually orthogonal. Specifically, for the decomposition $f = f_0 + \sum_i f_i + \sum_{i<j} f_{ij} + \cdots$, each term is orthogonal to every other term: $\langle f_u, f_v\rangle_{L^2} = 0$ when $u \neq v$ (as index sets).

Draft

This orthogonality is why the output variance decomposes as a sum: $\mathrm{Var}(Y) = \sum_i D_i + \sum_{i<j} D_{ij} + \cdots$. It also explains why the decomposition requires independent parameters: independence of the $Q_i$ makes the integration over $[0,1]^m$ factorize, which is what produces the orthogonality.

## A.5 ▪ Exercises

*Self-check.* Let $f(t) = e^{-t}$ and $g(t) = \cos(\pi t)$ on $[0,1]$. (a) Compute $\langle f, g\rangle_{L^2}$ numerically (use a numerical integration routine). (b) Are $f$ and $g$ orthogonal in $L^2([0,1])$? (c) Compute $\|f\|_{L^2}$ and $\|g\|_{L^2}$, and verify the Cauchy–Schwarz inequality $|\langle f, g\rangle| \le \|f\|\,\|g\|$.

*Answer:* (a) $\langle f, g\rangle_{L^2} = \int_0^1 e^{-t}\cos(\pi t)\,dt$. Integrating by parts twice with $a = -1$, $b = \pi$: result $(e^{-1}+1)/(1+\pi^2) \approx 0.126$. (b) No; the inner product is nonzero, so $f$ and $g$ are not orthogonal. (c) $\|f\|_{L^2}^2 = \int_0^1 e^{-2t}\,dt = (1-e^{-2})/2 \approx 0.432$, so $\|f\|_{L^2} \approx 0.658$. $\|g\|_{L^2}^2 = \int_0^1 \cos^2(\pi t)\,dt = 1/2$, so $\|g\|_{L^2} = 1/\sqrt{2} \approx 0.707$. Cauchy–Schwarz: $|0.126| \le 0.658 \times 0.707 \approx 0.465$. ✓

*Self-check.* The integration-by-parts identity in Section A.3 establishes that $L^* = -d/dt$ (with zero terminal condition) is the adjoint of $L = d/dt$ (with zero initial condition) in $L^2([0,T])$. (a) Write out $\langle Lu, v\rangle_{L^2}$ explicitly for $u(t) = t^2$ and $v(t) = t$ on $[0,1]$. (b) Compute $\langle u, L^*v\rangle_{L^2}$ for the same functions. (c) Verify that the two results differ by the boundary term $[u(t)v(t)]_0^1$.

*Answer:* (a) $Lu = 2t$. $\langle Lu, v\rangle = \int_0^1 2t \cdot t\,dt = \int_0^1 2t^2\,dt = 2/3$. (b) $L^*v = -dv/dt = -1$. $\langle u, L^*v\rangle = \int_0^1 t^2 \cdot (-1)\,dt = -1/3$. (c) Boundary term: $[u(t)v(t)]_0^1 = 1\cdot 1 - 0\cdot 0 = 1$. Check: $\langle Lu, v\rangle = \langle u, L^*v\rangle + [uv]_0^1 \Rightarrow 2/3 = -1/3 + 1 = 2/3$. ✓

*Self-check.* The ANOVA decomposition in $L^2([0,1]^2)$ for the model $f(Q_1, Q_2) = Q_1 + Q_2 + Q_1Q_2$ (with $Q_i \sim \mathcal{U}(0,1)$) is: $f = f_0 + f_1(Q_1) + f_2(Q_2) + f_{12}(Q_1, Q_2)$. Compute each term and verify that all terms are mutually orthogonal in $L^2([0,1]^2)$.

*Answer:* $f_0 = \mathbb{E}[f] = 1/2 + 1/2 + 1/4 = 5/4$. $f_1(Q_1) = \mathbb{E}[f|Q_1] - f_0 = Q_1 + 1/2 + Q_1/2 - 5/4 = 3Q_1/2 - 3/4$. $f_2(Q_2) = 3Q_2/2 - 3/4$ (by symmetry). $f_{12} = f - f_0 - f_1 - f_2 = Q_1Q_2 - Q_1/2 - Q_2/2 + 1/4 = (Q_1 - 1/2)(Q_2 - 1/2)$. Orthogonality: $\langle f_1, f_2\rangle = \int_0^1\int_0^1 (3Q_1/2 - 3/4)(3Q_2/2 - 3/4)\,dQ_1\,dQ_2 = \left[\int_0^1 (3Q/2 - 3/4)\,dQ\right]^2 = 0^2 = 0$. ✓Similarly all other pairs integrate to zero.

Draft

# Appendix B
# The Adjoint Formalism

This appendix develops the abstract operator framework that unifies the matrix adjoint (Chapter 5) and the ODE adjoint (Chapter 6). It is intended for students who want a rigorous foundation, not just the computational recipe.

## B.1 ▪ Linear Operators and Their Adjoints

Let $H$ be a Hilbert space with inner product $\langle \cdot, \cdot \rangle$. A *linear operator* $L : H \to H$ satisfies $L(\alpha u + \beta v) = \alpha Lu + \beta Lv$.

The *adjoint operator* $L^\dagger$ is defined by:

$$\langle Lu, v \rangle = \langle u, L^\dagger v \rangle \qquad \text{for all } u, v \in H. \tag{B.1}$$

The adjoint $L^\dagger$ is the unique operator satisfying (B.1) (when it exists).

**Example B.1 (Matrix adjoint).** In $\mathbb{R}^n$ with the Euclidean inner product, $L = A$ (left multiplication). Then $\langle Au, v \rangle = (Au)^T v = u^T (A^T v) = \langle u, A^T v \rangle$, so $L^\dagger = A^T$. This is the matrix transpose: Chapters 5's adjoint equation $A^T \vec{v} = \vec{c}$ is $L^\dagger \vec{v} = \vec{c}$.

**Example B.2 (Differential operator adjoint).** In $L^2([0, T])$, let $Lu = du/dt$ with $u(0) = 0$ (zero initial condition). By integration by parts (Appendix A.3):

$$\langle Lu, v \rangle = \int_0^T \dot{u} v \, dt = [uv]_0^T - \int_0^T u \dot{v} \, dt = u(T)v(T) - \int_0^T u \dot{v} \, dt.$$

For the adjoint to satisfy $\langle Lu, v \rangle = \langle u, L^\dagger v \rangle$, we need $L^\dagger v = -\dot{v}$ with $v(T) = 0$ (zero terminal condition). So $L^\dagger = -d/dt$ with the boundary condition shifted from $t = 0$ to $t = T$. This is exactly the structure of the adjoint ODE in Chapter 6.

Draft

## B.2 ▪ The Lagrange Identity

For the ODE operator $L = d/dt - F_u$ (the linearization of $\dot{u} = F(u;p)$), the Lagrange identity states:

$$\frac{d}{dt}\big[\lambda^T w\big] = \lambda^T(Lw) - (L^\dagger\lambda)^T w, \tag{B.2}$$

where $w = \partial u/\partial p$ satisfies the FSE $Lw = F_p$ and $\lambda$ satisfies the adjoint ODE $L^\dagger\lambda = g_u$.

Integrating (B.2) over $[0,T]$ and rearranging gives the sensitivity formula (6.14) of Chapter 6. This is the rigorous version of the "integration by parts" step that was announced but not fully proved in the main text.

## B.3 ▪ Solvability Conditions

For the equation $Lu = f$ to have a solution when $L$ is not invertible, $f$ must be orthogonal to the kernel of $L^\dagger$. Both statements below require $L$ to have closed range. That holds for the matrix operators of Chapter 5 and for the ODE operators of Chapter 6 with their boundary conditions, which is the only setting we use; for a general unbounded operator the alternative can fail.

**SC1 (Fredholm Alternative).** $Lu = f$ has a solution if and only if $\langle f, v\rangle = 0$ for every $v$ satisfying $L^\dagger v = 0$.

**SC2 (Uniqueness).** If $L^\dagger v = 0$ has only the trivial solution $v = 0$, then $Lu = f$ has a unique solution for every $f$.[5]

In the SA context, SC2 applies in two ways. The time-dependent FSE of Section 4.3 has $L = d/dt - J_F$ with an initial condition. Then $\ker L^\dagger = \{0\}$ for every Jacobian, so the FSE is a linear initial-value problem with a unique solution, near a bifurcation or not (self-check B-3 works the scalar case). The equilibrium FSE of Theorem 4.1 has $L = D_{\vec{u}}[\vec{f}]$. There SC2 is the assumption that this Jacobian is nonsingular: the FSE then has a unique solution, and the adjoint equation has a unique adjoint variable. That assumption fails at a fold or transcritical bifurcation, where an eigenvalue reaches zero, not near it. Near a fold the Jacobian approaches singularity while $\partial\vec{f}/\partial p$ does not, so the raw derivative grows without bound. Near a transcritical threshold $\partial\vec{f}/\partial p$ vanishes as the Jacobian becomes singular, the raw derivative stays finite, and the normalized index diverges because the equilibrium goes to zero. Section 8.4.1 sorts these cases.

[5]For the square, closed-range operators used here, $\ker L^\dagger = \{0\}$ forces $\ker L = \{0\}$, so existence and uniqueness coincide; for a non-square operator uniqueness would require $\ker L = \{0\}$ as a separate hypothesis.

## B.4 ▪ Connection to Chapters 5 and 6

The following table makes the Chapter 5/6 connection explicit in the language of this appendix:

| Concept | Chapter 5 (matrices) | Chapter 6 (ODEs) |
|---|---|---|
| Space $H$ | $\mathbb{R}^n$ | $L^2([0,T])^n$ |
| Operator $L$ | $A$ (matrix mult.) | $d/dt - J_F$ |
| Adjoint $L^\dagger$ | $A^T$ | $-d/dt - J_F^T$ |
| Adjoint equation | $A^T\vec{v} = \vec{c}$ | $-\dot{\lambda} = J_F^T\lambda + g_u$ |
| Boundary cond. | none (algebraic) | $\lambda(T) = +\nabla h^T$ |
| Sensitivity formula | $\vec{v}^T\vec{r}_j$ | $+\int_0^T \lambda^T F_{p_j}\, dt$ |
| Key operation | matrix transpose | integration by parts |

The formal apparatus of Hilbert spaces and adjoint operators shows that Chapters 5 and 6 are not two separate methods; they are the same method applied to two different Hilbert spaces.

## B.5 ▪ Exercises

*Self-check.* Let $A = \begin{pmatrix} 3 & 1 \\ 0 & 2 \end{pmatrix}$ and response $\vec{c} = (1,0)^T$. (a) Write the adjoint equation $A^T\vec{v} = \vec{c}$ and solve for $\vec{v}$. (b) For the right-hand side $\vec{b} = (b_1, b_2)^T$ and solution $\vec{u} = A^{-1}\vec{b}$, compute $J = \vec{c}^T\vec{u}$ directly. (c) Verify using the adjoint that $\partial J/\partial b_i = v_i$, consistent with the sensitivity formula $dJ/d\vec{b} = \vec{v}$.

*Answer:* (a) $A^T = \begin{pmatrix} 3 & 0 \\ 1 & 2 \end{pmatrix}$. Solving $A^T\vec{v} = (1,0)^T$: $3v_1 = 1$, $v_1 + 2v_2 = 0$, giving $\vec{v} = (1/3, -1/6)^T$. (b) $A^{-1} = \frac{1}{6}\begin{pmatrix} 2 & -1 \\ 0 & 3 \end{pmatrix}$. $\vec{u} = A^{-1}\vec{b} = \frac{1}{6}(2b_1 - b_2,\, 3b_2)^T$. $J = u_1 = (2b_1 - b_2)/6$. (c) $\partial J/\partial b_1 = 2/6 = 1/3 = v_1$. ✓$\partial J/\partial b_2 = -1/6 = v_2$. ✓The adjoint gives all sensitivities with one solve.

*Self-check.* The Lagrange identity (B.2) for the operator $L = d/dt - F_u$ (the ODE linearization) states: $\frac{d}{dt}[\lambda^T w] = \lambda^T(Lw) - (L^\dagger\lambda)^T w$. (a) For the scalar case $L = d/dt - a(t)$ (constant $a$), write out both sides of the Lagrange identity explicitly and verify it holds. (b) If $w$ satisfies $Lw = r(t)$ (the FSE with forcing $r$) and $\lambda$ satisfies $L^\dagger\lambda = 0$, integrate the Lagrange identity over

$[0,T]$ to obtain a formula for $\lambda(T)w(T) - \lambda(0)w(0)$ in terms of $\int_0^T \lambda r\,dt$.

*Answer:* (a) $L = d/dt - a$, so $L^\dagger = -d/dt - a$ (from integration by parts). LHS: $d(\lambda w)/dt = \dot\lambda w + \lambda \dot w$. RHS: $\lambda(Lw) - (L^\dagger\lambda)w = \lambda(\dot w - aw) - (-\dot\lambda - a\lambda)w = \lambda\dot w - a\lambda w + \dot\lambda w + a\lambda w = \dot\lambda w + \lambda\dot w$. LHS = RHS. ✓(b) Integrating: $[\lambda w]_0^T = \int_0^T \lambda^T(Lw)\,dt - \int_0^T (L^\dagger\lambda)^T w\,dt = \int_0^T \lambda r\,dt - 0$. So $\lambda(T)w(T) - \lambda(0)w(0) = \int_0^T \lambda(t)r(t)\,dt$.

*Self-check.* The Fredholm alternative (Section B.3) states that $Lu = f$ has a solution if and only if $f \perp \ker(L^\dagger)$. For $L = d/dt - a$ on $[0,T]$ with $u(0) = 0$ (zero initial condition), find $\ker(L^\dagger)$ and state the solvability condition for $Lu = f$ in terms of an integral of $f$, or explain why none is needed.

*Answer:* $L^\dagger = -d/dt - a$ with terminal condition $v(T) = 0$. $L^\dagger v = 0$ means $-\dot v = av$, so $v(t) = Ce^{-a(t-T)}$. Applying $v(T) = 0$: $Ce^0 = 0$, so $C = 0$ and $\ker(L^\dagger) = \{0\}$. Since the kernel is trivial, the Fredholm alternative says $Lu = f$ has a unique solution for every $f$ (with $u(0) = 0$), namely $u(t) = \int_0^t e^{a(t-s)} f(s)\,ds$, and no integral condition on $f$ is needed. This holds for every $a$, $a = 0$ included: a zero Jacobian, the scalar analogue of a bifurcation, does not make the kernel nontrivial. $L$ has no eigenvalues at all, since $\dot u - au = \mu u$ with $u(0) = 0$ forces $u \equiv 0$ for every $\mu$. The time-dependent FSE therefore always has a unique solution; why SA can still fail near a bifurcation is the subject of §8.4.1.

Draft

# Appendix C

# Eigenvalue Sensitivity and Singular Value Decomposition

This appendix covers two topics moved from Chapter 5 to keep the main text focused: sensitivity of eigenvalues and eigenvectors (important for stability analysis), and the singular value decomposition (important for understanding ill-conditioning).

## C.1 ▪ Sensitivity of Eigenvalues

Let $A(\varepsilon) = A_0 + \varepsilon B$ be a matrix depending on a scalar parameter $\varepsilon$. The eigenvalue $\lambda_k(\varepsilon)$ and eigenvector $\vec{x}_k(\varepsilon)$ satisfy $A(\varepsilon)\vec{x}_k = \lambda_k \vec{x}_k$.

### C.1.1 ▪ Symmetric Matrices

For symmetric $A_0$ with distinct eigenvalues, differentiating $A\vec{x}_k = \lambda_k \vec{x}_k$ at $\varepsilon = 0$:

$$\left.\frac{d\lambda_k}{d\varepsilon}\right|_{\varepsilon=0} = \vec{x}_k^T B \vec{x}_k. \tag{C.1}$$

The eigenvalue sensitivity is a quadratic form in the perturbation matrix and the normalized eigenvector.

### C.1.2 ▪ Non-Symmetric Matrices

For non-symmetric $A_0$, both left eigenvectors $\vec{y}_k$ (satisfying $\vec{y}_k^T A = \lambda_k \vec{y}_k^T$) and right eigenvectors $\vec{x}_k$ are needed:

$$\left.\frac{d\lambda_k}{d\varepsilon}\right|_{\varepsilon=0} = \frac{\vec{y}_k^T B \vec{x}_k}{\vec{y}_k^T \vec{x}_k}. \tag{C.2}$$

The formula is well-defined when $\lambda_k$ is a simple (distinct) eigenvalue, because in that case $\vec{y}_k^T \vec{x}_k \neq 0$.

**Proposition C.1.** *If $A_0$ has distinct eigenvalues $\lambda_k \neq \lambda_j$, then the corresponding left and right eigenvectors satisfy $\vec{y}_k^T \vec{x}_j = 0$ for $k \neq j$ (*biorthogonality*) and $\vec{y}_k^T \vec{x}_k \neq 0$.*

*Proof.* Let $A\vec{x}_j = \lambda_j \vec{x}_j$ and $\vec{y}_k^T A = \lambda_k \vec{y}_k^T$. Multiply the first equation on the left by $\vec{y}_k^T$ and the second on the right by $\vec{x}_j$:

$$\vec{y}_k^T A\vec{x}_j = \lambda_j(\vec{y}_k^T \vec{x}_j) \qquad \text{and} \qquad \vec{y}_k^T A\vec{x}_j = \lambda_k(\vec{y}_k^T \vec{x}_j).$$

Subtracting: $(\lambda_k - \lambda_j)(\vec{y}_k^T \vec{x}_j) = 0$. Since $\lambda_k \neq \lambda_j$, we conclude $\vec{y}_k^T \vec{x}_j = 0$.

To see that $\vec{y}_k^T \vec{x}_k \neq 0$: if it were zero, then $\vec{x}_k$ would be orthogonal to every $\vec{y}_j$ (via biorthogonality for $j \neq k$ and the assumed zero for $j = k$). But the left eigenvectors $\{\vec{y}_j\}$ span $\mathbb{R}^n$ when eigenvalues are distinct [22, Theorem 1.4.5], so no nonzero vector can be orthogonal to all of them. This contradiction shows $\vec{y}_k^T \vec{x}_k \neq 0$. □

For a real matrix with complex eigenvalues the same argument applies with the conjugate transpose $\vec{y}_k^*$ in place of $\vec{y}_k^T$ and $\mathbb{C}^n$ in place of $\mathbb{R}^n$.

Repetition alone does not break the formula; defectiveness does. If $\lambda_k$ is repeated but *semisimple*, its algebraic and geometric multiplicities agree, bases can still be chosen with $\vec{y}_k^T \vec{x}_k \neq 0$, and what fails is uniqueness: the individual eigenvectors are no longer determined, so the eigenvalues are in general only directionally differentiable. In this case formula (C.2) must be read with care. Evaluated on an arbitrary basis of the eigenspace it need not return a branch derivative at all: for $A(0) = \mathbf{0}$ with $A'(0) = \left[\begin{smallmatrix} 0 & 1 \\ 1 & 0 \end{smallmatrix}\right]$ the two branch derivatives are $\pm 1$, yet the quotient evaluated at $\vec{x} = \vec{e}_1$ returns 0. The branch derivatives are recovered only by restricting the perturbation to the eigenspace and taking the eigenvalues of that restriction; the single-vector quotient applies unambiguously only when $\lambda_k$ is simple. If $\lambda_k$ is *defective*, its geometric multiplicity falls below its algebraic multiplicity, $\vec{y}_k^T \vec{x}_k = 0$, and formula (C.2) fails outright: for a Jordan block of size $m$ the eigenvalues respond to a perturbation of size $\epsilon$ like $\epsilon^{1/m}$ rather than linearly. In applications it is defectiveness, not repetition, that marks the breakdown.

When $\vec{y}_k^T \vec{x}_k$ is small but nonzero (for unit vectors), the eigenvalue is highly sensitive to perturbations. This occurs near a defective eigenvalue of a non-normal matrix, where two eigenvalues are also close. Closeness alone does not produce it: a symmetric matrix keeps $\vec{y}_k^T \vec{x}_k = 1$ however close its eigenvalues are.

For the SIR Jacobian at the endemic equilibrium near $\mathcal{R}_0 = 1$, one eigenvalue stays at $-\nu_m$ and the other two form a complex pair over the range of Chapter 8's laboratory. As $\mathcal{R}_0 \to 1$ the pair shrinks toward zero and its left and right eigenvectors turn toward orthogonal; the pair coalesces,

defectively, a little above $\mathcal{R}_0 = 1$. Both effects shrink the smallest singular value (Section C.2), and together they produce the condition-number spike observed in Chapter 8. The spike is a property of $J_F$; why the indices themselves diverge at the threshold is a separate matter (§8.4.1).

## C.2 ▪ Singular Value Decomposition

Every matrix $A \in \mathbb{R}^{m\times n}$ has the *singular value decomposition* (SVD):

$$A = U\Sigma V^T, \tag{C.3}$$

where $U \in \mathbb{R}^{m\times m}$ and $V \in \mathbb{R}^{n\times n}$ are orthogonal and $\Sigma \in \mathbb{R}^{m\times n}$ is diagonal with nonnegative entries $\sigma_1 \geq \sigma_2 \geq \cdots \geq \sigma_r > 0 = \sigma_{r+1} = \cdots$ (the singular values).

### C.2.1 ▪ SVD and the Condition Number

For a square matrix of full rank, $r = n$ and the condition number from Chapter 8 is $\kappa_2(A) = \sigma_1/\sigma_n$. If $A$ is rank deficient then $\sigma_n = 0$ and $\kappa_2(A) = \infty$; the finite ratio $\sigma_1/\sigma_r$ over the nonzero singular values conditions the pseudoinverse problem on the range of $A$, not $A$ itself. The SVD makes clear why: the smallest singular value $\sigma_n$ measures how close $A$ is to being rank-deficient. When $\sigma_n \approx 0$, $A$ is nearly singular and tiny perturbations to the right-hand side $\vec{b}$ can produce large changes in the solution $\vec{u} = A^{-1}\vec{b}$.

### C.2.2 ▪ SVD and SA

The sensitivity Jacobian $\mathbf{S}$ from Chapter 3 can be analyzed using its SVD. For $\mathbf{S}$ shaped as (QOIs) $\times$ (POIs), the right singular vectors of $\mathbf{S}$ identify the combinations of parameters (POIs) that most strongly affect the model output, and the left singular vectors identify the combinations of outputs (QOIs) most strongly driven by those parameter combinations.

This connects to Principal Component Analysis (PCA): the right singular vectors of the design matrix $X$ in a regression are the principal components. The analogous object in SA is the sensitivity Jacobian $\mathbf{S}$ above, not the sampled parameter matrix. PCA of an LHS input sample (Chapter 9) describes the sampling design, which is close to isotropic by construction, and says nothing about which parameter directions drive the output.

### C.2.3 ▪ SVD and the Pseudoinverse

When $A$ is not square or is rank-deficient, the Moore-Penrose pseudoinverse $A^+ = V\Sigma^+U^T$ provides the minimum-norm least-squares solution: $\vec{u} = A^+\vec{b}$.

The pseudoinverse appears in parameter estimation from sensitivity data and in regularized inversion methods.

## C.3 ▪ Exercises

*Self-check.* The sensitivity Jacobian $\mathbf{S}$ for the SIR model at $T = 60$ days (from the Chapter 3 computational laboratory) is a $3 \times 4$ matrix. (a) Explain what the SVD of $\mathbf{S}$ reveals about the SA problem that the individual sensitivity indices do not. (b) If the largest singular value of $\mathbf{S}$ is $\sigma_1 = 2.3$ and the smallest *nonzero* singular value is $\sigma_3 = 0.02$ (recall a $3 \times 4$ matrix has at most three nonzero singular values), what is the condition number $\kappa_2(\mathbf{S})$? Interpret this number in plain language for the SA context. (c) The right singular vector $\vec{v}_1$ corresponding to $\sigma_1$ identifies the most "influential" linear combination of POIs. If $\vec{v}_1 \approx (0.7, 0.7, 0.1, 0.0)^T$ for POIs $(c, \beta, \tau_R, \nu_m)$, what does this tell you about the SA of the SIR model?

> *Answer:* (a) The SVD reveals which *combinations* of parameters drive the largest output changes. Individual SIs only measure one-at-a-time effects; the SVD captures the directions in parameter space that most amplify perturbations to the model output. (b) $\kappa_2 = 2.3/0.02 = 115$. This means that a perturbation in the direction $\vec{v}_3$ (least influential nonzero parameter combination) has only $1/115$ the output effect of a perturbation in direction $\vec{v}_1$. The SA result is dominated by a small subspace of parameter space. (c) The most influential combination weights $c$ and $\beta$ almost equally ($0.7 \approx 0.7$) with little contribution from $\tau_R$ and essentially none from $\nu_m$. This confirms the structural correlation identified in Chapter 3: $c$ and $\beta$ appear only as the product $c\beta$ in the SIR model, so they always have equal influence on any output.

*Self-check.* For a symmetric $2 \times 2$ matrix $A = \begin{pmatrix} a & b \\ b & d \end{pmatrix}$ with $a > d > 0$ and $b$ small, the eigenvalue sensitivity formula (C.1) gives $d\lambda_1/d\varepsilon = \vec{x}_1^T B \vec{x}_1$ where $B$ is the perturbation matrix. (a) Find the eigenvectors of $A$ for $b = 0$. (b) For the perturbation $B = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$ (an off-diagonal perturbation), compute $d\lambda_1/d\varepsilon$ at $b = 0$. (c) Near the threshold $\mathcal{R}_0 = 1$ of the SIR model, eigenvalues of the (non-symmetric) Jacobian come close together. Does

closeness alone make formula (C.2) unreliable? If not, what does?

*Answer:* (a) For $b = 0$: eigenvectors are $\vec{e}_1 = (1,0)^T$ ($\lambda_1 = a$) and $\vec{e}_2 = (0,1)^T$ ($\lambda_2 = d$). (b) $d\lambda_1/d\varepsilon = \vec{x}_1^T B \vec{x}_1 = (1,0)\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}(1,0)^T = 0$. The leading eigenvalue has zero first-order sensitivity to this off-diagonal perturbation (by symmetry; it appears at second order). (c) Closeness alone does not. For a symmetric matrix $\vec{y}_k = \vec{x}_k$, so $\vec{y}_k^T \vec{x}_k = 1$ however close the eigenvalues are; $\mathrm{diag}(1, 1+\delta)$ is an example. Formula (C.2) fails as an eigenvalue becomes defective, where $\vec{y}_k^T \vec{x}_k \to 0$ for unit vectors (Section C.1). That needs a non-normal matrix: $\left[\begin{smallmatrix} 0 & 1 \\ \delta & 0 \end{smallmatrix}\right]$ has eigenvalues $\pm\sqrt{\delta}$ with $|\vec{y}_k^T \vec{x}_k| = 2\sqrt{\delta}/(1+\delta)$, which is 0.063 at $\delta = 10^{-3}$. The SIR Jacobian is non-symmetric, and its complex pair coalesces defectively a little above $\mathcal{R}_0 = 1$; the formula fails there. That point is not the bifurcation. At $\mathcal{R}_0 = 1$ a single simple eigenvalue reaches zero, and the eigenvalue $-\nu_m$ becomes double without becoming defective. The divergence of the indices at the threshold has another cause (§8.4.1).

Draft

Draft

# Appendix D
# Periodic Solutions and Floquet Theory

For models with periodic forcing or limit-cycle behavior, SA of the periodic orbit requires tools beyond those developed in Chapter 4. This appendix introduces Floquet theory and the sensitivity of periodic solutions, based on the 2009 monograph [3].

## D.1 ▪ Floquet Theory

Consider the linear, $T$-periodic ODE $\dot{\vec{u}} = A(t)\vec{u}$ where $A(t+T) = A(t)$. The *fundamental matrix solution* $\Phi(t)$ satisfies $\dot{\Phi} = A(t)\Phi$, $\Phi(0) = I$.

The *monodromy matrix* is $M = \Phi(T)$: the fundamental matrix evaluated at the end of one period. Its eigenvalues $\mu_1, \ldots, \mu_n$ are the *Floquet multipliers*.

**Floquet's Theorem.** The fundamental matrix has the form $\Phi(t) = P(t)e^{Bt}$ where $P(t+T) = P(t)$ is periodic and $B$ is a constant matrix satisfying $e^{BT} = M$.

The Floquet multipliers determine stability, and the statement takes two forms that are easily conflated.

For the linear $T$-periodic system above, the zero solution is asymptotically stable when every $|\mu_k| < 1$, and unstable when some $|\mu_k| > 1$. Nothing forces any multiplier to equal one.

For a limit cycle of an *autonomous* system $\dot{\vec{u}} = \vec{F}(\vec{u})$, the variational equation about the orbit always carries the multiplier $\mu_1 = 1$, with eigenvector $\vec{F}(\vec{u}^*(0))$ tangent to the orbit: sliding the orbit along itself maps it to itself. That multiplier reflects time translation and carries no stability information. The orbit is orbitally asymptotically stable when the remaining $n-1$ multipliers satisfy $|\mu_k| < 1$, and unstable when any of them exceeds one in modulus.

## D.2 ▪ Sensitivity of the Period

For a nonlinear system $\dot{\vec{u}} = \vec{F}(\vec{u};\vec{p})$ with a periodic orbit $\vec{u}^*(t;\vec{p})$ of period $T(\vec{p})$, the period itself is a QOI. Its sensitivity is the Chapter 6 adjoint construction carried out on a closed orbit instead of a fixed interval.

Because the orbit closes on itself, there is no terminal condition to impose. The adjoint is instead the $T$-periodic solution $\vec{z}(t)$ of

$$\dot{\vec{z}} = -J_F^T(\vec{u}^*(t))\,\vec{z}, \qquad \vec{z}(t+T) = \vec{z}(t), \tag{D.1}$$

made unique by the phase normalization $\vec{z}(t)^T\vec{F}(\vec{u}^*(t);\vec{p}) = 1$. This product is constant along the orbit for any solution of (D.1), so fixing it at one instant fixes it at every instant. The period sensitivity is then

$$\frac{dT}{dp_j} = -\int_0^T \vec{z}(t)^T \frac{\partial \vec{F}}{\partial p_j}\big(\vec{u}^*(t);\vec{p}\big)\,dt. \tag{D.2}$$

The integration is essential. Perturbing $p_j$ changes the vector field at every point of the orbit, and the period responds to the phase shift accumulated over the whole circuit. The sign is the expected one: where the perturbation pushes the state along the direction of travel, the orbit closes sooner and $T$ decreases. Periodicity also forces $M^T\vec{z}(0) = \vec{z}(0)$, so $\vec{z}(0)$ is the left eigenvector of the monodromy matrix belonging to the unit multiplier, scaled by the normalization. That eigenvector fixes the direction in which the orbit responds; only the integral supplies the magnitude.

*Unifying Idea 2.* Equation (D.1) is the adjoint ODE (6.16) with $g_u = 0$, and (D.2) pairs that adjoint against $\partial\vec{F}/\partial p_j$ just as the sensitivity formula (6.18) does. Only the boundary condition is new: periodicity and a phase normalization replace the terminal condition (6.17), and the pairing enters with the opposite sign because $T$ is defined implicitly by the return condition rather than read off a cost functional.

## D.3 ▪ The van der Pol Example

The van der Pol oscillator $\ddot{u} - \epsilon(1-u^2)\dot{u} + u = 0$ (written as a system: $\dot{u}_1 = u_2$, $\dot{u}_2 = \epsilon(1-u_1^2)u_2 - u_1$) has a stable limit cycle for $\epsilon > 0$.

For small $\epsilon$, the period is approximately $T \approx 2\pi(1+\epsilon^2/16)$. Differentiating: $dT/d\epsilon = \pi\epsilon/4$. The normalized period sensitivity is therefore

$$\mathcal{S}_\epsilon^T = \frac{\epsilon}{T}\frac{dT}{d\epsilon} = \frac{\epsilon}{2\pi(1+\epsilon^2/16)}\cdot\frac{\pi\epsilon}{4} = \frac{\epsilon^2}{8(1+\epsilon^2/16)},$$

which is small for small $\epsilon$ (approximately $\epsilon^2/8$ for $\epsilon \ll 1$) but grows with increasing nonlinearity.

For larger $\epsilon$, the limit cycle becomes increasingly asymmetric, the asymptotic period expansion fails, and $dT/d\epsilon$ must be computed numerically from (D.2): integrate the orbit over one period, solve (D.1) for the periodic adjoint, apply the phase normalization, and evaluate the integral.

## D.4 ▪ Sensitivity of Floquet Multipliers

The sensitivity of the Floquet multiplier $\mu_k$ with respect to parameter $p_j$ uses the eigenvalue sensitivity formula from Appendix C.1 applied to the monodromy matrix:

$$\frac{d\mu_k}{dp_j} = \frac{\vec{y}_k^T \frac{\partial M}{\partial p_j} \vec{x}_k}{\vec{y}_k^T \vec{x}_k}, \tag{D.3}$$

where $\vec{x}_k$ and $\vec{y}_k$ are the right and left eigenvectors of $M$ for eigenvalue $\mu_k$, and $\partial M/\partial p_j$ is computed by integrating the variational equation for $\partial\Phi/\partial p_j$ over one period.

When $|\mu_k| \to 1$ for some $k \neq 1$, the periodic orbit approaches a bifurcation, where its stability changes. A real multiplier can pass through $-1$ (a period-doubling: an orbit of twice the period appears) or through $+1$. A small orbit can also shrink onto an equilibrium (a Hopf point), its multiplier $\mu_k$ tending to 1. At a period-doubling, and as an orbit shrinks to a Hopf point, the multiplier typically crosses at a finite rate, so (D.3) stays finite. By (D.3), $d\mu_k/dp_j$ can diverge only if $\vec{y}_k^T \vec{x}_k \to 0$ or $\partial M/\partial p_j$ grows without bound. That happens at a fold of cycles, where two periodic orbits merge and vanish as a multiplier reaches $+1$. None of this is the mechanism of Chapter 8's transcritical threshold, where the index diverges because the equilibrium goes to zero. A multiplier tends to a point on the unit circle, not to zero. Section 8.4.1 sorts these cases.

*Connection to Chapter 5 and Chapter 6.* The computation of $\partial M/\partial p_j$ requires integrating the variational equation $\dot{\partial\Phi}/\partial p_j = J_F(t)\,\partial\Phi/\partial p_j + \partial J_F/\partial p_j \cdot \Phi$ over one period, the same forward sensitivity equation developed in Chapter 4 and analyzed in Chapter 5. The left eigenvector $\vec{y}_k$ in equation (D.3) plays the role of the adjoint variable: it weights the state perturbation $\partial\Phi/\partial p_j \cdot \vec{x}_k$ to extract the component that matters for eigenvalue $\mu_k$, mirroring exactly the role $\vec{\lambda}$ plays in Chapter 6 for extracting the gradient of $J$ with respect to $p$. Students who have worked through Chapters 5–6 will recognize this as another instance of the same duality: the forward computation propagates perturbations forward in time; the left eigenvector (or adjoint variable) weights them by their downstream impact.

## D.5 ▪ Exercises

*Self-check.* The van der Pol oscillator with $\epsilon = 0.5$ has a stable limit cycle. (a) Using the approximation $T \approx 2\pi(1 + \epsilon^2/16)$, compute the period and the normalized sensitivity $\mathcal{S}_\epsilon^T$. (b) Is the period more or less sensitive to $\epsilon$ than a linear oscillator's period is to its natural frequency? Explain. (c) As $\epsilon$ increases from 0.5 to 2.0, does the period sensitivity increase or decrease? What does this imply about using local SA at large $\epsilon$?

*Answer:* (a) $T \approx 2\pi(1 + 0.25/16) = 2\pi \times 1.01563 \approx 6.381$. $dT/d\epsilon = 2\pi(\epsilon/8) = 2\pi(0.0625) \approx 0.393$. $\mathcal{S}_\epsilon^T = (\epsilon/T)(dT/d\epsilon) = (0.5/6.381)(0.393) \approx 0.031$. (b) A linear oscillator's period is independent of amplitude (and $\epsilon$ corresponds to the nonlinear damping coefficient, not the natural frequency). For $\epsilon = 0.5$, the period sensitivity $\approx 0.031$ is small: the period is nearly insensitive to $\epsilon$ in the weakly nonlinear regime. (c) The sensitivity increases. Computed from (D.2) with the periodic adjoint, the index is 0.030 at $\epsilon = 0.5$ (part (a)'s estimate gave 0.031) and 0.298 at $\epsilon = 2$, about tenfold. Beyond the weakly nonlinear regime the small-$\epsilon$ estimate overstates both the period and its slope: at $\epsilon = 2$ the computed values are $T = 7.630$ and $dT/d\epsilon = 1.138$, against 7.854 and $\pi\epsilon/4 = 1.571$. The formula's index there, 0.400, overstates the true 0.298. What fails at large $\epsilon$ is the expansion, not local SA: (D.2) still gives the exact local derivative. Global SA (Chapter 9) would add something only if $\epsilon$ were uncertain over a range.

*Self-check.* A compartmental epidemic model exhibits a periodic outbreak pattern with period $T^* = 365$ days (annual epidemic) under seasonal forcing. The Floquet multiplier associated with the periodic orbit is $\mu = 0.85$. (a) Is this periodic orbit stable or unstable? How do you know? (b) If a parameter perturbation changes $\mu$ to 1.02, what happens to the epidemic dynamics? (c) The sensitivity of $\mu$ to the transmission rate $\beta$ is $d\mu/d\beta = 3.2$ (computed via (D.3)), and the nominal transmission rate is $\beta^* = 0.3$. If $\beta$ increases by 5%, estimate the new value of $\mu$. Is the periodic outbreak still stable?

*Answer:* (a) Stable: $|\mu| = 0.85 < 1$, so perturbations from the periodic orbit decay over successive periods. The orbit is an attractor. (b) If $\mu = 1.02 > 1$, the periodic orbit becomes unstable: nearby trajectories drift away from the annual cycle toward some other attractor, which the multiplier alone does not identify. Two outcomes are ruled out. A multiplier passing through $+1$ does not produce period-doubling, which needs a multiplier through $-1$. A seasonally forced model has no endemic equilibrium to settle on: a constant state with $I > 0$ cannot keep $\dot{I} = 0$ while $\beta$ varies. (c) $\Delta\mu \approx (d\mu/d\beta)\,\Delta\beta = 3.2 \times (0.05 \times \beta)$. With $\beta^* = 0.3$, $\Delta\beta = 0.015$ and $\Delta\mu \approx 3.2 \times 0.015 = 0.048$. New $\mu \approx 0.85 + 0.048 = 0.898 < 1$: still stable. The first-order approximation suggests stability is preserved for this perturbation size.

# Appendix E
# The Adjoint in Optimization

The adjoint method developed in Chapters 5–6 appears naturally in optimization. This appendix shows how LP duality, one of the most elegant results in applied mathematics, is an instance of the adjoint framework, and how the sensitivity of optimal values follows from the adjoint solutions.

## E.1 ▪ Linear Programming Duality

The *primal linear program* is:

$$\min_{\vec{x}} \ \vec{c}^T\vec{x} \quad \text{subject to} \quad A\vec{x} \geq \vec{b},\ \vec{x} \geq 0. \tag{E.1}$$

Its *dual* is:

$$\max_{\vec{y}} \ \vec{b}^T\vec{y} \quad \text{subject to} \quad A^T\vec{y} \leq \vec{c},\ \vec{y} \geq 0. \tag{E.2}$$

The dual constraint $A^T\vec{y} \leq \vec{c}$ involves the transposed matrix $A^T$, exactly the structure of the adjoint equation $A^T\vec{v} = \vec{c}$ from Chapter 5. The dual variables $\vec{y}$ play the role of the adjoint vector $\vec{v}$.

**Strong Duality Theorem.** If both the primal (E.1) and dual (E.2) are feasible, their optimal values are equal:

$$\min_{\text{primal}} \vec{c}^T\vec{x}^* = \max_{\text{dual}} \vec{b}^T\vec{y}^*.$$

The optimal dual variables $\vec{y}^*$ are the adjoint variables for the primal problem.

## E.2 ▪ Sensitivity of the Optimal Value

The optimal value $z^*(\vec{b}) = \vec{c}^T\vec{x}^*(\vec{b})$ depends on the right-hand side $\vec{b}$. The *shadow price* (or *dual price*) of constraint $i$ is:

$$\frac{\partial z^*}{\partial b_i} = y_i^*, \tag{E.3}$$

where $y_i^*$ is the $i$-th component of the optimal dual vector. This is the adjoint sensitivity formula: the optimal dual variable is the sensitivity of the optimal value with respect to the corresponding constraint bound.

The normalized version gives a sensitivity index: $\mathcal{S}_{b_i}^{z^*} = (b_i/z^*)y_i^*$, directly analogous to the $\mathcal{S}_{p_j}^J = (p_j/J)(dJ/dp_j)$ formula from Chapter 2.

**Example E.1.** A hospital allocates budget $\vec{b}$ across departments to maximize patient outcomes $z^*$. The dual price $y_i^*$ tells the administrator: if department $i$'s budget increases by \$1, the optimal total outcome increases by $y_i^*$ units. This is SA of the optimal value with respect to the budget constraints.

## E.3 ▪ Quadratic Programming

For the equality-constrained quadratic program $\min_{\vec{x}} \frac{1}{2}\vec{x}^T H\vec{x} + \vec{c}^T\vec{x}$ subject to $A\vec{x} = \vec{b}$, the Lagrangian is $\mathcal{L} = \frac{1}{2}\vec{x}^T H\vec{x} + \vec{c}^T\vec{x} - \vec{\lambda}^T(A\vec{x} - \vec{b})$, and stationarity in $\vec{x}$ gives the adjoint equation $H\vec{x} + \vec{c} = A^T\vec{\lambda}$. The envelope theorem then yields

$$\frac{\partial z^*}{\partial b_i} = \lambda_i^*, \tag{E.4}$$

so a unit increase in $b_i$ changes the optimal value by $\lambda_i^*$, improving or worsening it according to the sign of $\lambda_i^*$. This is the LP shadow price (E.3) again, written in the same sign convention: $\vec{\lambda}^* = \vec{y}^*$, one object with one sign throughout. Bound constraints $\vec{x} \geq 0$ can be added; they carry their own multipliers and complementarity conditions, and under strict complementarity they leave (E.4) unchanged.

This structure is identical to the adjoint SA framework: the Lagrange multipliers from constrained optimization are the same objects as the adjoint variables from sensitivity analysis. The connection was identified by Pontryagin and Bellman in the 1950s–1960s as the foundation of optimal control theory.

Draft

## E.4 ▪ Connection to Optimal Control

The adjoint ODE from Chapter 6 is not unique to SA; it is the fundamental object in Pontryagin's minimum principle for optimal control.

Given a controlled dynamical system $\dot{\vec{u}} = \vec{F}(\vec{u}, \vec{p}, t)$ and an objective $J = \int_0^T g(\vec{u}, \vec{p})\, dt + h(\vec{u}(T))$, the Pontryagin minimum principle states that, along an optimal trajectory, the costate $\vec{\lambda}(t)$ satisfies the same adjoint ODE (6.16) derived in Chapter 6, while the optimal control $\vec{p}^*(t)$ minimizes the Hamiltonian $\mathcal{H} = g + \vec{\lambda}^T \vec{F}$ at each time.

In SA, we compute the adjoint to find how $J$ changes with parameters. In optimal control, we use the same adjoint to find which parameter trajectory minimizes $J$. The mathematical structure is identical.

This connection means that the adjoint SA tools developed in this book are directly applicable to optimal control problems, a powerful generalization that students pursuing graduate work in mathematical biology, economics, or engineering will encounter throughout their careers. Smith [59] develops parameter identifiability, influence, and selection in Ch. 7 §7.4–7.5, using the sensitivity Jacobian from our Chapter 4 to identify the measurements that most efficiently reduce uncertainty in model predictions.

## E.5 ▪ Exercises

*Self-check.* A hospital network allocates budget across three departments: emergency ($b_1$ = \$400K), ICU ($b_2$ = \$300K), surgery ($b_3$ = \$250K). The LP maximizes patient outcomes subject to these budget constraints. The optimal dual variables are $\vec{y}^* = (1.8, 2.4, 1.1)$ (outcomes per \$1000 additional budget). (a) Compute the normalized sensitivity index $\mathcal{S}_{b_i}^{z^*}$ for each department, assuming $z^* = 1715$ outcomes. (b) Which department has the highest shadow price? What does this mean for hospital policy? (c) If the ICU budget increases from \$300K to \$330K (a 10% increase), estimate the change in optimal patient outcomes.

*Answer:* (a) $\mathcal{S}_{b_i}^{z^*} = (b_i/z^*)\, y_i^*$: $\mathcal{S}_{b_1}^{z^*} = (400/1715) \times 1.8 = 0.42$. $\mathcal{S}_{b_2}^{z^*} = (300/1715) \times 2.4 = 0.42$. $\mathcal{S}_{b_3}^{z^*} = (250/1715) \times 1.1 = 0.16$. The three indices sum to 1, as they must when the budgets are the only constraints: then $z^* = \vec{b}^T \vec{y}^* = 720 + 720 + 275 = 1715$ by the Strong Duality Theorem. (b) The ICU has the highest shadow price ($y_2^* = 2.4$): each additional \$1000 in ICU budget yields 2.4 more patient outcomes: the highest marginal return. Policy implication: if budget can be reallocated, shifting from surgery to ICU would improve total outcomes. (c) $\Delta z^* \approx y_2^*\, \Delta b_2 = 2.4 \times 30 = 72$ additional outcomes. The 10% budget increase in ICU produces a $72/1715 \approx 4.2\%$ improvement in total outcomes: $\mathcal{S}_{b_2}^{z^*} = 0.42$, so $10\% \times 0.42 = 4.2\%$. ✓

*Self-check.* Consider the primal LP $\min\{2x_1 + 3x_2 : x_1 + x_2 \geq 4,\ 2x_1 + x_2 \geq$

6, $x_1, x_2 \geq 0\}$. (a) Find the optimal primal solution $\vec{x}^*$ and optimal value $z^*$ by checking the corner points of the feasible region. (b) Write the dual LP and find $\vec{y}^*$ (the dual optimal solution) using complementary slackness. (c) Verify strong duality: $\vec{c}^T\vec{x}^* = \vec{b}^T\vec{y}^*$. (d) Use the shadow price $y_1^*$ to estimate the new optimal value if the first constraint's right-hand side changes from $b_1 = 4$ to $b_1 = 5$. Verify by re-solving.

*Answer:* (a) The feasible region is defined by $x_1 + x_2 \geq 4$, $2x_1 + x_2 \geq 6$, $x_1, x_2 \geq 0$. The corner points of the feasible region are: $(4, 0)$ (intersection of $x_1 + x_2 = 4$ and $x_2 = 0$), $(0, 6)$ (intersection of $2x_1 + x_2 = 6$ and $x_1 = 0$), and $(2, 2)$ (intersection of the two constraints). Objective values: $2(4) + 3(0) = 8$; $2(0) + 3(6) = 18$; $2(2) + 3(2) = 10$. The minimum is $z^* = 8$ at $\vec{x}^* = (4,\, 0)^T$.
(b) The dual of $\min\{\vec{c}^T\vec{x} : A\vec{x} \geq \vec{b},\ \vec{x} \geq 0\}$ is $\max\{\vec{b}^T\vec{y} : A^T\vec{y} \leq \vec{c},\ \vec{y} \geq 0\}$. Here $A = \begin{pmatrix} 1 & 1 \\ 2 & 1 \end{pmatrix}$, $\vec{b} = (4, 6)^T$, $\vec{c} = (2, 3)^T$, giving:

$$\max\{4y_1 + 6y_2\} \ \text{ s.t. } \ y_1 + 2y_2 \leq 2, \ \ y_1 + y_2 \leq 3, \ \ y_1, y_2 \geq 0.$$

At $\vec{x}^* = (4, 0)$: constraint 1 is active ($x_1 + x_2 = 4$) and constraint 2 is inactive ($2x_1 + x_2 = 8 > 6$). By complementary slackness, $y_2^* = 0$ (since constraint 2 is not active). The dual constraint corresponding to $x_1 > 0$ must hold with equality: $y_1^* + 2(0) = 2$, so $y_1^* = 2$. Thus $\vec{y}^* = (2,\, 0)^T$.
(c) Strong duality: $\vec{c}^T\vec{x}^* = 2(4) + 3(0) = 8$ and $\vec{b}^T\vec{y}^* = 4(2) + 6(0) = 8$. Equal. ✓
(d) The shadow price $y_1^* = 2$ means each unit increase in $b_1$ increases $z^*$ by 2: for a minimization problem with $\geq$ constraints, raising $b_1$ *tightens* the constraint and shrinks the feasible region, so the optimal value can only rise. Estimate: $z^* \approx 8 + 2 \times (5 - 4) = 10$. Verification: with $b_1 = 5$ the relevant corners are $(5, 0)$ with $z = 10$, $(1, 4)$ with $z = 14$, and $(0, 6)$ with $z = 18$, so $(5, 0)$ is optimal. Note that $(5, 0)$ lies on $x_1 + x_2 = 5$ and $x_2 = 0$; the two constraint boundaries $x_1 + x_2 = 5$ and $2x_1 + x_2 = 6$ meet at $(1, 4)$. Exact value: $z^* = 10$. ✓The shadow price approximation is exact here because the optimal basis (constraint 1 active, $x_2 = 0$) is unchanged by this perturbation.

*Self-check.* The optimal control connection (Section E.4) states that the adjoint ODE from Chapter 6 is identical to the costate equation in Pontryagin's minimum principle. (a) For the simple epidemic control problem: minimize total infections $J = \int_0^T I(t)\, dt$ for the SIR model (4.15)–(4.17) with a control $u(t)$ that scales the incidence $c\beta SI/N$ by $1 - u$ in both $\dot{S}$ and $\dot{I}$, so that $\dot{I} = (c\beta(1-u)S/N - \gamma_R - \nu_m)I$, write the Hamiltonian $\mathcal{H}(\vec{u}, \vec{p}, \lambda, t)$. (b) Write the costate (adjoint) equations $-\dot{\lambda}_i = \partial\mathcal{H}/\partial u_i$, one for each *state* $u_i \in \{S, I, R\}$. Note that the scalar control $u(t)$ in part (a) is a different object from the state vector $\vec{u}$; stationarity in the control, $\partial\mathcal{H}/\partial u = 0$, is a separate condition. (c) Compare these costate equations with the adjoint

ODE derived in Chapter 6 for $J = \int_0^T I\,dt$. Are they the same?

*Answer:* (a) Hamiltonian: $\mathcal{H} = I + \lambda_S \dot{S} + \lambda_I \dot{I} + \lambda_R \dot{R} = I + \lambda_S(\nu_m N - c\beta(1-u)SI/N - \nu_m S) + \lambda_I(c\beta(1-u)SI/N - (\gamma_R + \nu_m)I) + \lambda_R(\gamma_R I - \nu_m R)$. (b) Costate equations: $-\dot{\lambda}_S = \partial\mathcal{H}/\partial S = c\beta(1-u)I/N(\lambda_I - \lambda_S) - \nu_m \lambda_S$, etc. (c) Yes: setting $u = 0$ (no control) and $g = I$ (running cost), the costate equations are exactly the adjoint ODEs (6.21)–(6.23) from Chapter 6. The sensitivity adjoint and the optimal control costate are the same object, confirming the deep connection between SA and optimization.

Draft

Draft

# Bibliography


[1] R. M. Anderson and R. M. May, *Infectious Diseases of Humans: Dynamics and Control*, Oxford University Press, Oxford, 1991.

[2] L. M. Arriola and J. M. Hyman, *Being sensitive to uncertainty*, Computing in Science & Engineering, 9 (2007), pp. 10–20.

[3] ———, *Sensitivity analysis for uncertainty quantification in mathematical models*, in Mathematical and Statistical Estimation Approaches in Epidemiology, G. Chowell, J. M. Hyman, L. Bettencourt, and C. Castillo-Chavez, eds., Springer, New York, 2009.

[4] A. G. Baydin, B. A. Pearlmutter, A. A. Radul, and J. M. Siskind, *Automatic differentiation in machine learning: a survey*, Journal of Machine Learning Research, 18 (2018), pp. 1–43.

[5] S. M. Blower and H. Dowlatabadi, *Sensitivity and uncertainty analysis of complex models of disease transmission: An HIV model, as an example*, International Statistical Review, 62 (1994), pp. 229–243.

[6] E. Borgonovo, *A new uncertainty importance measure*, Reliability Engineering & System Safety, 92 (2007), pp. 771–784.

[7] E. Borgonovo, *Sensitivity Analysis: An Introduction for the Management Scientist*, vol. 251 of International Series in Operations Research & Management Science, Springer, Cham, 2017.

[8] G. E. P. Box and N. R. Draper, *Empirical Model-Building and Response Surfaces*, John Wiley & Sons, New York, 1987. The remark that all models are wrong but some are useful appears on p. 424.

[9] D. G. Cacuci, *Sensitivity theory for nonlinear systems. I. nonlinear functional analysis approach*, Journal of Mathematical Physics, 22 (1981), pp. 2794–2802.

[10] D. G. Cacuci, *Sensitivity and Uncertainty Analysis, Volume I: Theory*, Chapman and Hall/CRC, Boca Raton, 2003.

[11] F. Campolongo, J. Cariboni, and A. Saltelli, *An effective screening design for sensitivity analysis of large models*, Environmental Modelling & Software, 22 (2007), pp. 1509–1518.

[12] Y. Cao, S. Li, and L. Petzold, *Adjoint sensitivity analysis for differential-algebraic equations: algorithms and software*, Journal of Computational and Applied Mathematics, 149 (2002), pp. 171–191.


Draft

[13] R. T. Q. Chen, Y. Rubanova, J. Bettencourt, and D. K. Duvenaud, *Neural ordinary differential equations*, Advances in Neural Information Processing Systems, 31 (2018).

[14] N. Chitnis, J. M. Hyman, and J. M. Cushing, *Determining important parameters in the spread of malaria through the sensitivity analysis of a mathematical model*, Bulletin of Mathematical Biology, 70 (2008), pp. 1272–1296.

[15] O. Diekmann, J. A. P. Heesterbeek, and J. A. J. Metz, *On the definition and the computation of the basic reproduction ratio $R_0$ in models for infectious diseases in heterogeneous populations*, Journal of Mathematical Biology, 28 (1990), pp. 365–382.

[16] M. B. Giles and N. A. Pierce, *An introduction to the adjoint approach to design*, Flow, Turbulence and Combustion, 65 (2000), pp. 393–415.

[17] A. Griewank and A. Walther, *Evaluating Derivatives: Principles and Techniques of Algorithmic Differentiation*, SIAM, Philadelphia, 2nd ed., 2008.

[18] J. C. Helton and F. J. Davis, *Latin hypercube sampling and the propagation of uncertainty in analyses of complex systems*, Reliability Engineering & System Safety, 81 (2003), pp. 23–69.

[19] J. Herman and W. Usher, *SALib: An open-source Python library for sensitivity analysis*, Journal of Open Source Software, 2 (2017), p. 97.

[20] N. J. Higham, *Accuracy and Stability of Numerical Algorithms*, Society for Industrial and Applied Mathematics, Philadelphia, PA, 2nd ed., 2002.

[21] T. Homma and A. Saltelli, *Importance measures in global sensitivity analysis of nonlinear models*, Reliability Engineering & System Safety, 52 (1996), pp. 1–17.

[22] R. A. Horn and C. R. Johnson, *Matrix Analysis*, Cambridge University Press, Cambridge, 2nd ed., 2012.

[23] J. M. Hyman, Z. Qu, and L. Xue, *Avoiding pitfalls in epidemic modeling: An introduction to formulating, analyzing, and fitting models.* Manuscript under review at Springer; a decision is not expected before 2027, 2026.

[24] R. L. Iman and W. J. Conover, *A distribution-free approach to inducing rank correlation among input variables*, Communications in Statistics — Simulation and Computation, 11 (1982), pp. 311–334.

[25] M. J. W. Jansen, *Analysis of variance designs for model output*, Computer Physics Communications, 117 (1999), pp. 35–43.

[26] W. O. Kermack and A. G. McKendrick, *A contribution to the mathematical theory of epidemics*, Proceedings of the Royal Society of London A, 115 (1927), pp. 700–721.

[27] S. Kucherenko, S. Tarantola, and P. Annoni, *Estimation of global sensitivity indices for models with dependent variables*, Computer Physics Communications, 183 (2012), pp. 937–946.

[28] Y. A. Kuznetsov, *Elements of Applied Bifurcation Theory*, Springer, New York, 3rd ed., 2004.

[29] X. Lu and E. Borgonovo, *Global sensitivity analysis in epidemiological modeling*, European Journal of Operational Research, 304 (2023), pp. 9–24.

Draft

[30] C. C. Margossian, *A review of automatic differentiation and its efficient implementation*, WIREs Data Mining and Knowledge Discovery, 9 (2019), p. e1305.

[31] S. Marino, I. B. Hogue, C. J. Ray, and D. E. Kirschner, *A methodology for performing global uncertainty and sensitivity analysis in systems biology*, Journal of Theoretical Biology, 254 (2008), pp. 178–196.

[32] M. D. McKay, R. J. Beckman, and W. J. Conover, *A comparison of three methods for selecting values of input variables in the analysis of output from a computer code*, Technometrics, 21 (1979), pp. 239–245.

[33] R. Mester, A. Landeros, C. Rackauckas, and K. Lange, *Differential methods for assessing sensitivity in biological models*, PLOS Computational Biology, 18 (2022), p. e1009598.

[34] M. D. Morris, *Factorial sampling plans for preliminary computational experiments*, Technometrics, 33 (1991), pp. 161–174.

[35] J. D. Murray, *Mathematical Biology I: An Introduction*, Springer, Berlin, 3rd ed., 2007.

[36] U. Naumann, *The Art of Differentiating Computer Programs: An Introduction to Algorithmic Differentiation*, SIAM, Philadelphia, 2012.

[37] I. M. Navon, *Data assimilation for numerical weather prediction: A review*, in Data Assimilation for Atmospheric, Oceanic and Hydrologic Applications, S. K. Park and L. Xu, eds., Springer, Berlin, 2009, pp. 21–65.

[38] A. B. Owen, *Scrambled net variance for integrals of smooth functions*, The Annals of Statistics, 25 (1997), pp. 1541–1562.

[39] ———, *Scrambling Sobol' and Niederreiter–Xing points*, Journal of Complexity, 14 (1998), pp. 466–489.

[40] ———, *Monte Carlo theory, methods and examples.* Self-published, available online, 2013. `https://artowen.su.domains/mc/`.

[41] ———, *Sobol' indices and Shapley value*, SIAM/ASA Journal on Uncertainty Quantification, 2 (2014), pp. 245–251.

[42] F. Pianosi, K. Beven, J. Freer, J. W. Hall, J. Rougier, D. B. Stephenson, and T. Wagener, *Sensitivity analysis of environmental models: a systematic review with practical workflow*, Environmental Modelling & Software, 79 (2016), pp. 214–232.

[43] L. S. Pontryagin, V. G. Boltyanskii, R. V. Gamkrelidze, and E. F. Mishchenko, *The Mathematical Theory of Optimal Processes*, Interscience Publishers, John Wiley & Sons, New York, 1962. Translated from the Russian by K. N. Trirogoff; edited by L. W. Neustadt.

[44] A. Puy, W. Becker, S. Lo Piano, and A. Saltelli, *A comprehensive comparison of total-order estimators for global sensitivity analysis*, International Journal for Uncertainty Quantification, 12 (2022), pp. 1–18.

[45] Z. Qu, L. Xue, and J. M. Hyman, *Modeling the transmission of Wolbachia in mosquitoes for controlling mosquito-borne diseases*, SIAM Journal on Applied Mathematics, 78 (2018), pp. 826–852.

Draft

[46] S. Razavi, A. Jakeman, A. Saltelli, C. Prieur, B. Iooss, E. Borgonovo, E. Plischke, S. Lo Piano, T. Iwanaga, W. Becker, S. Tarantola, J. H. A. Guillaume, J. Jakeman, H. Gupta, N. Melillo, G. Rabitti, V. Chabridon, Q. Duan, X. Sun, S. Smith, R. Sheikholeslami, N. Hosseini, M. Asadzadeh, A. Puy, S. Kucherenko, and H. R. Maier, *The future of sensitivity analysis: An essential discipline for systems modeling and policy support*, Environmental Modelling & Software, 137 (2021), p. 104954.

[47] A. Saltelli, *Making best use of model evaluations to compute sensitivity indices*, Computer Physics Communications, 145 (2002), pp. 280–297.

[48] A. Saltelli, K. Aleksankina, W. Becker, P. Fennell, F. Ferretti, N. Holst, S. Li, and Q. Wu, *Why so many published sensitivity analyses are false: a systematic review of sensitivity analysis practices*, Environmental Modelling & Software, 114 (2019), pp. 29–39.

[49] A. Saltelli, P. Annoni, I. Azzini, F. Campolongo, M. Ratto, and S. Tarantola, *Variance based sensitivity analysis of model output. design and estimator for the total sensitivity index*, Computer Physics Communications, 181 (2010), pp. 259–270.

[50] A. Saltelli, K. Chan, and E. M. Scott, *Sensitivity Analysis*, John Wiley & Sons, Chichester, 2000.

[51] A. Saltelli, A. Jakeman, S. Razavi, and Q. Wu, *Sensitivity analysis: a discipline coming of age*, Environmental Modelling & Software, 146 (2021), p. 105226.

[52] A. Saltelli, M. Ratto, T. Andres, F. Campolongo, J. Cariboni, D. Gatelli, M. Saisana, and S. Tarantola, *Global Sensitivity Analysis: The Primer*, John Wiley & Sons, Chichester, 2008.

[53] A. Saltelli, M. Ratto, S. Tarantola, and F. Campolongo, *Sensitivity analysis practices: strategies for model-based inference*, Reliability Engineering & System Safety, 91 (2006), pp. 1109–1125.

[54] A. Saltelli, S. Tarantola, and F. Campolongo, *Sensitivity analysis as an ingredient of modeling*, Statistical Science, 15 (2000), pp. 377–395.

[55] A. Saltelli, S. Tarantola, F. Campolongo, and M. Ratto, *Sensitivity Analysis in Practice: A Guide to Assessing Scientific Models*, Wiley, Chichester, 2004.

[56] A. Saltelli, S. Tarantola, and K. P.-S. Chan, *A quantitative model-independent method for global sensitivity analysis of model output*, Technometrics, 41 (1999), pp. 39–56.

[57] Z. Sirkes and E. Tziperman, *Finite difference of adjoint or adjoint of finite difference?*, Monthly Weather Review, 125 (1997), pp. 3373–3378.

[58] R. C. Smith, *Uncertainty Quantification: Theory, Implementation, and Applications*, vol. 12 of Computational Science and Engineering, SIAM, Philadelphia, 2014.

[59] ——, *Uncertainty Quantification: Theory, Implementation, and Applications, Second Edition*, Computational Science and Engineering, SIAM, Philadelphia, 2024.

[60] I. M. Sobol', *Global sensitivity indices for nonlinear mathematical models and their Monte Carlo estimates*, Mathematics and Computers in Simulation, 55 (2001), pp. 271–280.

[61] S. H. Strogatz, *Nonlinear Dynamics and Chaos*, Westview Press, Boulder, CO, 2nd ed., 2018.

[62] S. Tarantola, F. Ferretti, S. Lo Piano, M. Kozlova, A. Lachi, R. Rosati, A. Puy, P. Roy, G. Vannucci, M. Kuc-Czarnecka, and A. Saltelli, *An annotated timeline of sensitivity analysis*, Environmental Modelling & Software, 174 (2024), p. 105977.

[63] P. van den Driessche and J. Watmough, *Reproduction numbers and sub-threshold endemic equilibria for compartmental models of disease transmission*, Mathematical Biosciences, 180 (2002), pp. 29–48.

[64] J. Wu, R. Dhingra, M. Gambhir, and J. V. Remais, *Sensitivity analysis of infectious disease models: methods, advances and their application*, Journal of the Royal Society Interface, 10 (2013), p. 20121018.

Draft

Draft

# Glossary

This glossary is a finding aid for the reader who meets a term away from where it was first defined. Each entry states what the term means and points to the section that develops it. Symbols are collected separately in the Notation and Conventions section; the choice among the sensitivity indices is laid out in Table 9.4.

**Adjoint method** A technique for computing all parameter sensitivities of a scalar response at a cost independent of the number of parameters, by solving one auxiliary (adjoint) system backward in time. The adjoint is the transpose of the forward sensitivity problem. See Chapter 5.

**Adjoint operator** The generalization of the matrix transpose to a linear operator $L$ on an inner product space: the operator $L^\dagger$ defined by $\langle Lu, v\rangle = \langle u, L^\dagger v\rangle$ for all $u, v$. The adjoint method rests on this identity. See Appendix B.

**Algorithmic (automatic) differentiation** The evaluation of exact derivatives of a function specified by a program, by applying the chain rule to its elementary operations. Reverse-mode algorithmic differentiation is the adjoint method applied to a computation. See Chapter 5.

**ANOVA decomposition** The decomposition of a function's variance into contributions from each parameter and each combination of parameters; the basis of the Sobol indices. See Chapter 9.

**Backpropagation** The name used in machine learning for reverse-mode algorithmic differentiation, the method by which neural networks are trained. It is the adjoint method under another name: a reader who knows backpropagation already knows the book's central computational tool. See Chapter 5.

**Basic reproduction number ($\mathcal{R}_0$)** The expected number of secondary infections produced by one infectious individual in a fully susceptible population; for the running SIR model, $\mathcal{R}_0 = c\beta\tau_R$. See Chapter 2.

Draft

**Bifurcation** A parameter value at which the qualitative behavior of a model changes: equilibria merge, or an equilibrium changes stability. How local sensitivity analysis fails at a bifurcation depends on what is differentiated. The book distinguishes two kinds: at a *fold* bifurcation two equilibria collide and annihilate, and the raw derivative of the equilibrium diverges; at a *transcritical* bifurcation two equilibria exchange stability, the raw derivative stays finite, and only the normalized index diverges. The epidemic threshold is transcritical. See Chapter 8.

**Bootstrap** A resampling procedure for placing confidence intervals on estimated sensitivity indices without assuming a distributional form. See Chapter 9.

**Checkpointing** A strategy for the adjoint method that trades computation for memory, storing the forward solution at selected times and recomputing the rest, rather than storing the entire trajectory. See Chapter 7.

**Condition number** A measure, $\kappa(A) = \|A\| \, \|A^{-1}\|$, of how much a problem's output can change relative to a change in its input; large $\kappa$ signals ill-conditioning. See Chapter 8 and Appendix C.

**Delta index** A moment-independent global sensitivity measure (Borgonovo's $\delta$) based on the shift a parameter induces in the entire output distribution, not only its variance. See Chapter 9.

**Discretize-then-adjoint /adjoint-then-discretize** The two orderings of the operations that produce a numerical adjoint. Discretize-then-adjoint forms the adjoint of the discretized problem; adjoint-then-discretize discretizes the continuous adjoint. The two need not agree, and the distinction matters for gradient accuracy. See Chapter 7.

**eFAST** The extended Fourier Amplitude Sensitivity Test, a spectral method for estimating first-order and total Sobol indices. See Chapter 9.

**Eigenvalue sensitivity** The derivative of an eigenvalue of a matrix $A(\varepsilon)$ with respect to a parameter, given for a simple eigenvalue by a ratio of left and right eigenvectors. The formula requires care when the eigenvalue is repeated (see *semisimple / defective eigenvalue*). See Appendix C.

**Elasticity** A synonym for the normalized sensitivity index, borrowed from economics: the fractional change in the response per fractional change in the parameter. An elasticity of one means a one-percent change in

the parameter produces a one-percent change in the response, not that the relationship is linear. See Chapter 2.

**Factor prioritization, fixing, and mapping** The three settings in which a global sensitivity analysis is read: which parameters to measure more precisely (prioritization), which can be fixed without loss (fixing), and which drive the output into a region of interest (mapping). See Chapter 9.

**Floquet multipliers** The eigenvalues of the monodromy matrix of a periodic system, governing the stability of a periodic orbit and the sensitivity of its period. See Appendix D.

**Forward sensitivity equation (FSE)** The differential equation obtained by differentiating the model with respect to a parameter and solving alongside it; it yields the sensitivity of the full trajectory to one parameter. See Chapter 4.

**Fredholm alternative** The statement that a linear equation $Lu = f$ is solvable exactly when $f$ is orthogonal to the kernel of the adjoint operator $L^\dagger$. See Appendix B.

**Gaussian process emulator (surrogate model)** A cheap statistical approximation to an expensive model, fitted to a set of runs and used in its place for sensitivity analysis. See Chapter 9.

**Global sensitivity analysis** The study of how a model's output responds to parameters varied over their whole plausible ranges simultaneously, as opposed to local analysis at a single nominal point. See Chapter 9.

**Hilbert space** A complete inner product space, one in which every Cauchy sequence converges within the space. The function space $L^2([a, b])$ is the example the book uses, and completeness is what lets the Sobol variance decomposition be stated. See Appendix A.

**Inner product** A function $\langle \cdot, \cdot \rangle$ on a vector space that is symmetric, linear in each argument, and positive-definite, generalizing the dot product; on a function space it is an integral, $\langle f, g \rangle = \int fg$. Its *induced norm* is $\|u\| = \sqrt{\langle u, u \rangle}$. Orthogonality, the adjoint, and the Fredholm alternative are all stated through it. See Appendix A.

**Jansen estimator** An efficient variance-based estimator of Sobol indices that reuses model evaluations across the first-order and total-effect calculations. See Chapter 9.

**Latin hypercube sampling (LHS)** A stratified sampling scheme that spreads

points evenly across each parameter's range, improving coverage over independent uniform sampling. See Chapter 9.

**Low-discrepancy sequence** A deterministic sequence (such as a Sobolśequence) that fills space more evenly than random points, reducing the variance of a sensitivity estimate at a given sample size. See Chapter 9.

**Morris screening (elementary effects)** A low-cost method for ranking many parameters by importance, using a small number of one-at-a-time perturbations to estimate the mean and spread of each parameter's effect. See Chapter 9.

**Parameter of interest (POI) /Quantity of interest (QOI)** The input under study (POI) and the output being analyzed (QOI). A sensitivity index always relates one QOI to one POI. See Chapter 1.

**Polynomial chaos expansion** A representation of a model's output as a series in orthogonal polynomials of its random inputs, from which sensitivity indices can be read off analytically. See Chapter 9.

**PRCC (partial rank correlation coefficient)** A sampling-based sensitivity measure that ranks parameters by their monotonic influence on the output after removing the effect of the others. See Chapter 9.

**Pseudoinverse** The generalization of the matrix inverse to non-square or rank-deficient matrices, built from the singular value decomposition; it solves a least-squares problem on the range of the matrix. A near-zero smallest singular value signals ill-conditioning. See Appendix C.

**Saltelli construction** A sampling design that arranges model evaluations so that first-order and total Sobol indices can be estimated together with minimal cost. See Chapter 9.

**Semisimple / defective eigenvalue** A repeated eigenvalue is *semisimple* when its geometric and algebraic multiplicities agree and *defective* when the geometric multiplicity is smaller. The simple eigenvalue-sensitivity formula must be read with care in the semisimple case and fails outright in the defective case, where eigenvalues respond to a perturbation like a fractional power rather than linearly. See Appendix C.

**Sensitivity index** The central object of the book: the normalized (relative) derivative $\mathcal{S}_p^Q = (p/Q)\,\partial Q/\partial p$, the fractional change in a response $Q$ per fractional change in a parameter $p$. A sensitivity index describes a *model*, not the system the model represents; it inherits every assumption built into that model. See Definition in Chapter 2.

**Shapley effects** A global sensitivity measure that allocates the output variance among parameters by averaging each parameter's marginal contribution over all orderings, giving a fair share even under correlation. See Chapter 9.

**Sobol indices** Variance-based global sensitivity measures. The first-order index is the fraction of output variance explained by a parameter alone; the total-effect index adds every interaction that parameter participates in. See Chapter 9.

**Standardized (rank) regression coefficient** A sensitivity measure from a linear (or rank-linear) fit of output to inputs, valid when that fit captures most of the output variance. See Chapter 9.

**Structural identifiability** Whether a model's parameters can, even in principle, be determined from perfect output data; parameters that appear only in a fixed combination are not separately identifiable. See Chapter 3.

Draft

Draft

# Index

absolute change, 22
adjoint method, 91
  backward in time, 108
  ODE derivation, 109
adjoint operator, 213
adjoint sensitivity analysis, 7
adjoint vector, 90
adjoint-then-discretize, 138
algorithmic differentiation
  discontinuities, 139
  forward mode, 97
  reverse mode, 98
  software, 135
ANOVA decomposition, 179

backpropagation, 98
basic reproduction number ($\mathcal{R}_0$), 32
bifurcation
  fold, 153
  SA failure near, 151
  transcritical, 152
bootstrap, 185

checkpointing, 138
  adjoint, 138
commutative diagram, 93
condition number, 157
  SVD definition, 219
correlated POIs
  constraint, 57
  empirical, 57
  structural, 56
CVODES, 85, 127

delta index, 191
discretize-then-adjoint, 138

eFAST, 184
eigenvalue sensitivity, 217
elasticity, 25, 43, *see also* sensitivity index
elementary effect, 187
eta-squared ($\eta^2$), 181

factor fixing, 171
factor mapping, 171
factor prioritization, 171
Floquet multipliers, 223
Floquet theory, 223
force of infection, 31
forward sensitivity analysis, 7
forward sensitivity equation (FSE), 69
Fredholm Alternative, 214
Fredholm alternative, 214

Gaussian process emulator, 189
global sensitivity analysis, 11
  when needed, 161

Halton sequence, 199
Hilbert space, 210

ill-conditioning, 157

Draft

implicit function theorem, 71
inner product, 209
  $L^2$, 210

Jansen estimator, 183

Lagrange identity, 214
Lagrange multiplier, 109
Latin hypercube sampling, 66, 175
LHS, *see* Latin hypercube sampling
linear operator, 213
linear programming
  dual, 227
  primal, 227
low-discrepancy sequences, 175, 176
  Halton, 199
  Sobol, 199

MATCONT, 166
method-question mismatch, 150
moment-independent sensitivity, 191
monodromy matrix, 223
Morris screening, 187

norm
  induced, 209

orthogonality, 210

parameter of interest (POI), 3
partial rank correlation coefficient, *see* PRCC
polynomial chaos expansion, 189
Pontryagin maximum principle, 228
Pontryagin minimum principle, 229
PRCC, 177
  as nonparametric correlation, 177
proliferation problem, 77
pseudoinverse, 219

quantity of interest (QOI), 5

regularized change, 23
relative change, 23
rounding error, 51
Runge–Kutta method, 139

SALib, 66, 176
Saltelli construction, 174
sensitivity index, 25
  dimensionlessness, 26, 172
  logarithmic form, 26
  normalized, 25
  ratio form, 25
sensitivity Jacobian, 47
  normalized, 47
separatrix, 159
shadow price, 228
Shapley effects, 191
sigma-normalized index, 172
sign rule, 24
Sobol indices, 180
  correlated POIs, 190
  first-order, 180
  total, 180
Sobol sequence, 199
Sobol' sequence
  scrambled, 175
standardized regression coefficient, 163
strong duality theorem, 227
structural identifiability, 56, 57
SUNDIALS, 127
surrogate model, 188

time-varying sensitivity, 173
tornado plot, 36
trajectory sensitivity, 159
truncation error, 51

U-curve, 52

Draft

unifying ideas, 12

variance inflation factor (VIF), 157

Draft