


Overview of the Motor System (Part 2)  
Start of spinal reflexes


Professor Malcolm MacIver

FIGURE 16.7 Motor neuron recruitment in the cat medial gastrocnemius muscle under different behavioral conditions



After Walmsley et al. (1978) *J. Neurophys.* 41: 1203–1216.

FIGURE 16.9 The number of active motor units and their rate of firing both increase with voluntary force




After Monster and Chan (1977).

# Features of $\alpha$ motor neurons

	That innervate small motor units		That innervate large motor units
Axonal diameter	Smaller	C O N T I N U U M	Larger
Input resistance	Larger		Smaller
Excitability	Greater		Lower
EPSP amplitude	Larger		Smaller
Duration of after-hyperpolarization	Longer		Shorter
	<ul style="list-style-type: none"> <li>-Easily depolarized</li> <li>-Slow rate of firing</li> <li>-Control slow muscle fibers needed for posture</li> </ul>		<ul style="list-style-type: none"> <li>-Difficult to depolarize</li> <li>-Fast rate of firing</li> <li>-Control fibers that produce max force used for jumping</li> </ul>

## Increase motor unit size, alpha motor neuron exhibits:

Increased	Decreased
Cell body size	Input resistance
Dendritic Complexity	Excitability
Short-term EPSP potentiation with repeated activation	Ia EPSP amplitude
Axonal diameter (i.e. faster conduction)	PSP decay constant
Number of axonal branches (i.e., more muscle fibers innervated)	Duration of after-hyperpolarization



# Nervous system is extremely plastic

- Growth and development
- Injury
- Training

Upper motor centers vs. motor unit plasticity

Aids in understanding of motor neuron and muscle physiology

Great implications in training and rehabilitation

# Motor system plasticity

Modification of strength and endurance of motor behavior through experience

- Depends partly on changes in motor unit phenotype
- But MAINLY on changes in supraspinal motor centers



# Nervous system is extremely plastic


- Growth and development
- Injury
- Training

Upper motor centers vs. **motor unit plasticity**

Aids in understanding of motor neuron and muscle physiology

Great implications in training and rehabilitation

# Types of Motor Unit



3 types of alpha motor neuron

- Excitability
- Discharge frequency

3 types of muscle fibers

- Contraction speed
- Fatigability




Motor units are important site of plasticity in the nervous system

Results of studies with cross-innervation and electrical stimulation show properties of motor neurons determine muscle fiber behavior

Motor unit behavior is modified by training

Important implications for exercise and rehabilitation programs

# Cross-Innervation



Soleus- slow muscle with constant, low frequency innervation

TA- fast muscle with intermittent, high frequency innervation

With cross-innervation:

- Soleus become fast
- TA became slow


Shows motorneuron innervation determines muscle fiber phenotype

# Electrical Stimulation

Electrical stimulation can change behavior of muscle fiber

- Frequency dependence

Electrical stimulation at low frequencies can prevent slow muscle from becoming fast muscle with disuse




# Cross-innervation experiments


Alk mATPase

Before

Control



56 days  
stim



- Fast fatigable fiber
- Fast fatigue resistant fiber
- ★ Slow

Stained muscle for myosin ATPase activity

Before

Fast fatigable and fast fatigue resistant fibers are darker

Slow fibers are lighter

After chronic electrical nerve stimulation similar to the innervation that slow fibers receive

Almost all fibers show phenotype of slow fibers (lighter)

Ability of motor neurons and the muscles they innervate to change physically and functionally

- Acquire new motor skills
- Modify the strength and endurance of motor behavior

## In general exercise:

- slows the contractile properties of motor units
- Increases endurance of fibers

The impact of exercise is proportional to the order of recruitment

- Exercise has a greater impact on slow motor units at low exertion levels
- Only has an impact on fast fibers if high intensity exercise is performed





Most commonly, exercise “slows” contractile properties of motor units while increasing endurance and strength of muscle fibers

S motor units affected most at low exertion levels

FR and FF motor units affected only if recruited by higher intensities of exercise

Appreciable gains shown in non-exercised limb

- Recruitment and adaptation of central neural circuits that have access to contralateral motor units

Documented gains in muscle strength with imagined exercise

Appreciable gains shown in non-exercised limb

- Recruitment and adaptation of central neural circuits that have access to contralateral motor units

Documented gains in muscle strength with  
**imagined** exercise

Appreciable gains shown in non-exercised limb

- Recruitment and adaptation of central neural circuits that have access to contralateral motor units


Documented gains in muscle strength with  
**imagined** exercise

BMD ENG 301  
Quantitative Systems Physiology  
(Nervous System)

## Spinal Reflex Circuits

Malcolm MacIver

# Organization of neural structures involved in the control of movement




# **Reflexes**

- 1. A simple motor action that is stereotypical and repeatable**
- 2. Elicited by a sensory stimulus, the strength of the motor action being graded with the intensity of the stimulus**

Dr. Charles Sherrington 1857-1952  
(1932 Nobel Prize for discoveries  
regarding functions of neurons)  
– Published “The integrative  
Action of the Nervous  
System” in 1906








### Speed of Signal transmission

If a signal needs to propagate to the brain and back, a typical reaction time is **0.25** seconds, whereas a reflex will typically take closer to **0.15** seconds.

# Fixed Action Patterns *versus* Reflexes




**REFLEX**

**FIXED ACTION PATTERN**


## Fixed Action Patterns *versus* Reflexes

Reflex	Fixed Action Pattern
A simple stereotyped and repeatable motor action	A complex motor act, involving a specific temporal sequence of component motor acts
Elicited by a sensory stimulus. Magnitude of the motor response increases with the strength of the sensory drive.	Generated internally or initiated by a sensory stimulus. The stimulus triggers release of the coordinated motor act. The action may be all-or-none or graded in intensity, and it may be dependent on the type of sensory stimulus or internal state, but it maintains its basic pattern.

## Simplest Circuit




**Extensor**




# Stretch reflex circuitry


(A) Muscle spindle




(B)



(C)








## Structure of Spindle

Sensory axons coil around intrafusal fibers

Stretch in fibers activates mechanically gated ion channels in sensory axons

Ia sensory around bag fibers

II sensory around bag and chain




Eight to ten intrafusal fibers

Contained in collagen sheath

Two types of fibers

- Nuclear bag fibers
  - Excite Ia and II sensory afferents
  - Sensitive to dynamic changes (rapidly adapting)
  - Non-contractile
- Nuclear chain fibers
  - Half the size of bag fibers
  - Excite II sensory afferents only
  - Sensitive to static levels of stretch
  - Contractile



## Primary afferents (group Ia)

- Sense changes in muscle length
  - Limb dynamics (velocity and direction)

## Secondary afferents (group II)

- Sense sustained muscle stretch
  - Static limb position



