

Northern Pikeminnow Abundance in Deadwater Slough, Salmon River, Idaho, and Potential Impacts to Local Chinook Salmon

--Manuscript Draft--

Manuscript Number:	
Article Type:	Research Article
Full Title:	Northern Pikeminnow Abundance in Deadwater Slough, Salmon River, Idaho, and Potential Impacts to Local Chinook Salmon
Short Title:	Northern Pikeminnow in Deadwater Slough
Corresponding Author:	Michael Ackerman, M.S. Mount Hood Environmental McCall, Idaho UNITED STATES
Keywords:	Northern Pikeminnow; Chinook salmon; predation; mark-recapture; bioenergetics
Abstract:	<p>Predation on emigrating juvenile salmonids by piscivorous fishes is a widely-studied source of mortality within reservoirs and dam tailraces in the Columbia River basin. Native Northern Pikeminnow have been estimated to consume ~8% of the approximately 200 million juvenile salmonids emigrating through the lower Snake and Columbia rivers, annually. Less is known about the interaction between upstream, river-dwelling Northern Pikeminnow and their impacts on salmonid recovery above the reservoir systems. In this study, we examine the abundance of Northern Pikeminnow in a slow-water reach of the Salmon River, Idaho, known as Deadwater Slough, and consider their potential impacts on local Chinook salmon populations. The Deadwater Slough is within a migration corridor for several Chinook salmon populations as well as other Endangered Species Act-listings including steelhead and Sockeye salmon. We estimated the abundance of Northern Pikeminnow in Deadwater Slough, an approximately 1.9 kilometer reach of the Salmon River, to be 19,499 in the fall and 10,352 in the spring, corresponding with the peak emigration windows of juvenile Chinook salmon. Using these abundance values, we estimated Northern Pikeminnow consumption of juvenile Chinook salmon. Assuming 60% of the Northern Pikeminnow diet is fish, of which 50% is juvenile Chinook salmon, we estimated the Northern Pikeminnow population can consume 61,409 juvenile Chinook salmon, annually. After performing a sensitivity analysis, we estimated Northern Pikeminnow predation on juvenile Chinook salmon was equivalent to 377 (95% CI: 161 - 935) returning adults. Given the relatively small size of the Salmon River, the high densities and potential consumption rates of Northern Pikeminnow in Deadwater Slough suggest predation likely has a consequential impact on Chinook salmon recovery in the Upper Salmon River.</p>
Order of Authors:	Michael W Ackerman, M.S. Nicholas J Porter, M.S. Kevin E See, Ph.D. Tara E Blackman, M.S. Michael S Hall Richard A Carmichael
Opposed Reviewers:	
Additional Information:	
Question	Response
Financial Disclosure Enter a financial disclosure statement that describes the sources of funding for the	The funders had no role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript.

work included in this submission. Review the [submission guidelines](#) for detailed requirements. View published research articles from [PLOS ONE](#) for specific examples.

This statement is required for submission and **will appear in the published article** if the submission is accepted. Please make sure it is accurate.

Unfunded studies

Enter: *The author(s) received no specific funding for this work.*

Funded studies

Enter a statement with the following details:

- Initials of the authors who received each award
- Grant numbers awarded to each author
- The full name of each funder
- URL of each funder website
- Did the sponsors or funders play any role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript?
- **NO** - Include this sentence at the end of your statement: *The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.*
- **YES** - Specify the role(s) played.

* typeset

Competing Interests

Use the instructions below to enter a competing interest statement for this submission. On behalf of all authors, disclose any [competing interests](#) that could be perceived to bias this work—acknowledging all financial support and any other relevant financial or non-financial competing interests.

This statement is **required** for submission and **will appear in the published article** if the submission is accepted. Please make sure it is accurate and that any funding sources listed in your Funding Information

The authors have declared that no competing interests exist.

later in the submission form are also declared in your Financial Disclosure statement.

View published research articles from [PLOS ONE](#) for specific examples.

NO authors have competing interests

Enter: *The authors have declared that no competing interests exist.*

Authors with competing interests

Enter competing interest details beginning with this statement:

I have read the journal's policy and the authors of this manuscript have the following competing interests: [insert competing interests here]

* typeset

Ethics Statement

Enter an ethics statement for this submission. This statement is required if the study involved:

- Human participants
- Human specimens or tissue
- Vertebrate animals or cephalopods
- Vertebrate embryos or tissues
- Field research

Write "N/A" if the submission does not require an ethics statement.

General guidance is provided below. Consult the [submission guidelines](#) for detailed instructions. **Make sure that all information entered here is included in the Methods section of the manuscript.**

N/A

Format for specific study types

Human Subject Research (involving human participants and/or tissue)

- Give the name of the institutional review board or ethics committee that approved the study
- Include the approval number and/or a statement indicating approval of this research
- Indicate the form of consent obtained (written/oral) or the reason that consent was not obtained (e.g. the data were analyzed anonymously)

Animal Research (involving vertebrate animals, embryos or tissues)

- Provide the name of the Institutional Animal Care and Use Committee (IACUC) or other relevant ethics board that reviewed the study protocol, and indicate whether they approved this research or granted a formal waiver of ethical approval
- Include an approval number if one was obtained
- If the study involved *non-human primates*, add *additional details* about animal welfare and steps taken to ameliorate suffering
- If anesthesia, euthanasia, or any kind of animal sacrifice is part of the study, include briefly which substances and/or methods were applied

Field Research

Include the following details if this study involves the collection of plant, animal, or other materials from a natural setting:

- Field permit number
- Name of the institution or relevant body that granted permission

Data Availability

Authors are required to make all data underlying the findings described fully available, without restriction, and from the time of publication. PLOS allows rare exceptions to address legal and ethical concerns. See the [PLOS Data Policy](#) and [FAQ](#) for detailed information.

Yes - all data are fully available without restriction

A Data Availability Statement describing where the data can be found is required at submission. Your answers to this question constitute the Data Availability Statement and **will be published in the article**, if accepted.

Important: Stating 'data available on request from the author' is not sufficient. If your data are only available upon request, select 'No' for the first question and explain your exceptional situation in the text box.

Do the authors confirm that all data underlying the findings described in their manuscript are fully available without restriction?

Describe where the data may be found in full sentences. If you are copying our sample text, replace any instances of XXX with the appropriate details.

- If the data are **held or will be held in a public repository**, include URLs, accession numbers or DOIs. If this information will only be available after acceptance, indicate this by ticking the box below. For example: *All XXX files are available from the XXX database (accession number(s) XXX, XXX).*
- If the data are all contained **within the manuscript and/or Supporting Information files**, enter the following:
All relevant data are within the manuscript and its Supporting Information files.
- If neither of these applies but you are able to provide **details of access elsewhere**, with or without limitations, please do so. For example:

Data cannot be shared publicly because of [XXX]. Data are available from the XXX Institutional Data Access / Ethics Committee (contact via XXX) for researchers who meet the criteria for access to confidential data.

The data underlying the results presented in the study are available from (include the name of the third party

All data and code for the analyses presented here can be found in a GitHub repository release (<https://doi.org/10.5281/zenodo.6458124>).

<p><i>and contact information or URL).</i></p> <ul style="list-style-type: none">• This text is appropriate if the data are owned by a third party and authors do not have permission to share the data. <p>* typeset</p>	
Additional data availability information:	

Dr. Chenette
Editor-in-Chief
PLOS ONE

June 12, 2022

Dear Dr. Chenette:

I would like to submit our manuscript titled “Northern Pikeminnow Abundance in Deadwater Slough, Salmon River, Idaho, and Potential Impacts to Local Chinook Salmon” to be considered for publication as a research article in PLOS ONE.

The research paper uses a mark-recapture survey to estimate the abundance of Northern Pikeminnow, a piscine predator, in an uncharacteristically slow reach of the Salmon River, Idaho, that has at least been partially formed due to human activities. We estimate a large population size of Northern Pikeminnow and use a bioenergetics approach to consider their impacts to local ESA-listed Chinook salmon populations. We identify that predation by Northern Pikeminnow on local Chinook salmon should be considered among limiting factors hindering population recovery efforts in the region. We believe these findings will be of interest to readers interested in salmon recovery and will be a valuable contribution to PLOS ONE.

We declare that this manuscript and content is original, has not been published previously, and is not currently being considered for publication elsewhere. As the corresponding author, I confirm that the manuscript has been read and approved for submission by all the co-authors.

We hope you find our research article suitable for publication and a valuable addition to PLOS ONE. We look forward to hearing from you on your decision. In the meantime, please feel free to contact me with any questions or concerns.

Sincerely,

Michael Ackerman
Senior Scientist
Mount Hood Environmental
PO Box 4282
McCall, Idaho 83638
Cell: 208-469-0896
Email: mike.ackerman@mthoodenvironmental.com

Northern Pikeminnow Abundance in Deadwater Slough, Salmon River, Idaho, and Potential Impacts to Local Chinook Salmon

Michael W. Ackerman^{1,2,✉}, Nicholas J. Porter², Kevin E. See^{2,3}, Tara E. Blackman⁴, Michael S.
Hall², and Richard A. Carmichael²

12 April, 2022

¹ Mount Hood Environmental, PO Box 4282, McCall, Idaho, 83638, USA

² Biomark, Inc., 705 South 8th Street, Boise, Idaho, 83702, USA

³ Washington Department of Fish and Wildlife, Fish Program, Science Division, 1111
Washington Street NE, Olympia, Washington, 98501, USA

⁴ Mount Hood Environmental, 39085 Pioneer Boulevard #100 Mezzanine, Sandy, Oregon,
97055, USA

✉Correspondence: [Michael W. Ackerman <mike.ackerman@mthoodenvironmental.com>](mailto:mike.ackerman@mthoodenvironmental.com)

Running Head: Northern Pikeminnow in Deadwater Slough

Keywords: Northern Pikeminnow; Chinook salmon; predation; mark-recapture; bioenergetics

Abstract

Predation on emigrating juvenile salmonids by piscivorous fishes is a widely-studied source of mortality within reservoirs and dam tailraces in the Columbia River basin. Native Northern Pikeminnow have been estimated to consume ~8% of the approximately 200 million juvenile salmonids emigrating through the lower Snake and Columbia rivers, annually. Less is known about the interaction between upstream, river-dwelling Northern Pikeminnow and their impacts on salmonid recovery above the reservoir systems. In this study, we examine the abundance of Northern Pikeminnow in a slow-water reach of the Salmon River, Idaho, known as Deadwater Slough, and consider their potential impacts on local Chinook salmon populations. The Deadwater Slough is within a migration corridor for several Chinook salmon populations as well as other Endangered Species Act-listings including steelhead and Sockeye salmon. We estimated the abundance of Northern Pikeminnow in Deadwater Slough, an approximately 1.9 kilometer reach of the Salmon River, to be 19,499 in the fall and 10,352 in the spring, corresponding with the peak emigration windows of juvenile Chinook salmon. Using these abundance values, we estimated Northern Pikeminnow consumption of juvenile Chinook salmon. Assuming 60% of the Northern Pikeminnow diet is fish, of which 50% is juvenile Chinook salmon, we estimated the Northern Pikeminnow population can consume 61,409 juvenile Chinook salmon, annually. After performing a sensitivity analysis, we estimated Northern Pikeminnow predation on juvenile Chinook salmon was equivalent to 377 (95% CI: 161 - 935) returning adults. Given the relatively small size of the Salmon River, the high densities and potential consumption rates of Northern Pikeminnow in Deadwater Slough suggest predation likely has a consequential impact on Chinook salmon recovery in the Upper Salmon River.

INTRODUCTION

The Snake and Columbia rivers drain a network of inland tributaries that provide essential spawning and rearing habitat for anadromous Pacific salmon *Oncorhynchus* spp. and steelhead *O. mykiss*. These fishes were historically abundant throughout the basin and consequently have tremendous ecological, cultural, and economic value (Lewis et al. 2019; Atlas et al. 2021). Over the last century, salmonid stocks throughout the Snake and Columbia rivers have undergone significant declines related to habitat modification in the freshwater environment (e.g., removal of beavers, river channel simplification, water withdrawals, hydrosystem development) and poor marine conditions, affecting their survival and recruitment at multiple life stages (Justice et al. 2017; Clark et al. 2020; Crozier et al. 2020). Consequently, action agencies have made considerable investments in the rehabilitation of tributary ecosystems and evaluation of factors attributed to the species decline (Roni et al. 2018; White et al. 2021).

Predation on emigrating juveniles by piscivorous fishes is another important, and potentially under-estimated, source of salmonid mortality. In the lower mainstem Snake and Columbia rivers it is estimated that predation on out-migrating salmonids during peak emigration has a significant negative impact on the overall population and success of recovery efforts (Tabor et al. 1993; Beamesderfer et al. 1996; Friesen and Ward 1999). Dams and reservoirs in the Columbia River are the primary locations associated with high rates of piscine predation on salmonids (Petersen 1994; Ward et al. 1995). There are generally two mechanisms that explain these high predation zones. First, migration (movement) rates of juvenile salmon are reduced during reservoir passage (Venditti et al. 2000), thereby increasing the time migrating smolts are vulnerable to predation. Second, reservoirs and downstream tailraces associated with dams on

the Snake and Columbia rivers create favorable slow-water habitat for predatory fishes known to consume juvenile salmonids, such as the Northern Pikeminnow *Ptychocheilus oregonsis*. Northern Pikeminnow tolerate and thrive in relatively warm, slow-water habitats (Wydoski and Whitney 2003) and thus have benefited from dams on the Columbia River, becoming abundant predators of salmonid outmigrants (Knutsen and Ward 1999). Northern Pikeminnow are estimated to consume ~8% (16.4 million) of the approximately 200 million juvenile salmonids emigrating through the lower Snake and Columbia rivers, annually (Beamesderfer et al. 1996).

Most predation studies have focused on reservoirs (Murphy et al. 2021) and mainstem reaches (Tabor et al. 1993; Ward et al. 1995; Shively et al. 1996; Zimmerman and Ward 1999) of the Columbia River basin whereas considerably less is known about the interaction between upstream, river-dwelling piscivorous fishes and their impacts on salmonid recovery above the reservoir systems (Rubenson et al. 2020). Upstream habitats containing slower water velocities and other attributes that support piscivorous predators may overlap with essential habitat for some salmonid species and life stages. One such example is Deadwater Slough, an approximately 1.9 km long reach of uncharacteristically slow and deep water in the Salmon River, Idaho. The Deadwater Slough is within a section of the Salmon River containing historically important overwinter rearing habitat for juvenile Chinook salmon *O. tshawytscha* and is part of the migratory pathway for upstream Chinook salmon populations, the endangered Snake River Sockeye salmon *O. nerka* population (Axel et al. 2015), and several populations of threatened Snake River steelhead. The slough is also inhabited by piscivorous predators, including Northern Pikeminnow and Smallmouth Bass *Micropterus dolomieu*, that are potential sources of mortality for rearing and migrating salmonids.

The Salmon River was historically the most productive tributary for Chinook salmon in the Columbia River basin (Nemeth and Kiefer 1999), but those populations have become depleted in recent decades. Therefore, quantifying the mortality of spring/summer-run Chinook salmon (hereafter Chinook salmon) in the Upper Salmon River associated with piscine predation is of particular interest. Importantly, the Salmon River supports eight extant populations of Chinook salmon upstream of Deadwater Slough (National Oceanic and Atmospheric Administration 2017). This includes the Lemhi River, which was historically the largest population, and is a prioritized candidate for restoration of natural processes to increase production of juvenile Chinook salmon (Zimmerman et al. 2012).

Chinook salmon in the Upper Salmon River are stream-type and exhibit two distinct migration tactics; downstream rearing (DSR) and natal reach rearing (NRR) (Copeland et al. 2014). The DSR migrants leave their natal area as subyearlings between June and November and typically overwinter in downstream, mainstem habitats until the following spring when they emigrate to the ocean as smolts. Alternatively, NRR migrants remain in their natal areas for approximately one year after emergence until emigration to the ocean as smolts. Diversity of migration tactics provides a mechanism for coping with adverse conditions in freshwater rearing and migration environments and buffers against catastrophic events, thereby increasing population resiliency (Dodson et al. 2013). Deadwater Slough represents an important habitat for Chinook salmon in the Upper Salmon River as it supports rearing and migration of juveniles from all upstream populations including fall DSR and spring NRR migrants.

Recent studies have examined the downstream movement, distribution, and apparent survival (hereafter ‘survival’) of juvenile salmonid emigrants through the Salmon River, including the Deadwater Slough reach. Sockeye salmon migrating through the mainstem Salmon River during

spring were estimated to have 32% lower survival in the Deadwater Slough reach compared to directly adjacent reaches (Axel et al. 2015). Similarly, emigrating DSR Chinook salmon had an approximate 10% reduction in transition probability through the Deadwater Slough compared to surrounding reaches during fall and early winter months (Ackerman et al. 2018; Porter et al. 2019). Low survival was attributed to lack of fish cover and low-velocity water delaying movement rates, thereby increasing predation risk.

In this study, we estimated the abundance of Northern Pikeminnow in Deadwater Slough and evaluated potential impacts to juvenile salmon emigrants, focusing on DSR and NRR Chinook salmon. We hypothesized that high densities of piscivorous predators in the Deadwater Slough might explain the reduced survival observed for juvenile Chinook salmon. To test this, our objectives were four-fold:

1. Estimate the abundance of Northern Pikeminnow in Deadwater Slough during the peaks of fall DSR and spring NRR juvenile emigrations;
2. Document predation on juvenile Chinook salmon during the DSR and NRR emigration periods using gastric lavage;
3. Use an established bioenergetics approach to estimate consumption potential (grams) of the Northern Pikeminnow population on DSR and NRR Chinook salmon emigrants at Deadwater Slough;
4. Estimate how consumption of juvenile Chinook salmon emigrants at Deadwater Slough by Northern Pikeminnow might impact adult returns to the Upper Salmon River.

METHODS

Study Site

The Deadwater Slough is an approximately 1.9 km section of the mainstem Salmon River located roughly 6 river kilometers (rkm) downstream from the town of North Fork, Idaho (Figure 1). The downstream end of the slough is located at the confluence of Dump Creek and the Salmon River. A large alluvial fan at the mouth of Dump Creek has, at least partially, created a hydraulic control in the Salmon River resulting in the formation of Deadwater Slough. Although the origin and timing of the alluvial fan and Deadwater Slough is somewhat ambiguous (Reichmuth et al. 1985; USACE 1986), some believe they coincided with the failure of a small mining reservoir in the Dump Creek drainage around 1897 that deposited substantial amounts of sediment at its confluence with the Salmon River (Emerson 1973). The slough is currently a slow, deep (approximately 6 m maximum depth) section in the river, spanning 12 hectares, averaging 68 m wide, and has characteristics resembling a small reservoir.

Northern Pikeminnow Demographics

We estimated the population size of piscine predators in the Deadwater Slough using a mark-recapture survey design and a catch per unit effort (CPUE) approach. Predators were sampled near the peaks of the fall DSR and spring NRR emigrations. During our initial survey in 2019, Northern Pikeminnow were the most prevalent piscine predator in Deadwater Slough and consequently became our focal taxa. Our intent was to survey during the emigration of the 2018 Chinook salmon brood year that occurred fall 2019 and spring 2020, however logistical constraints during the onset of the COVID-19 pandemic delayed the spring 2020 survey until

2021. Over this period, an additional fall survey occurred during the 2020 DSR emigration. Fall surveys were constrained to two weeks to minimize the effect of Northern Pikeminnow immigration and emigration on our population size estimate.

Multiple capture methods were employed during the fall 2019 survey to reduce selectivity and bias for species and size classes. Methods included raft electrofishing, fyke netting, snorkeling, and angling. After evaluating all methods, angling was the most effective method for capturing Northern Pikeminnow while also minimizing potential impacts to Endangered Species Act (ESA)-listed adult steelhead that were present during our surveys. Less effective methods were abandoned in subsequent years and the following analyses will focus on fish captured by angling, unless otherwise noted.

Our study relied heavily on volunteer anglers who were permitted to fish anywhere within Deadwater Slough. Each survey day, anglers boated or hiked their catch (periodically or upon filling a livewell) to a processing station at the boat ramp, approximately 500 m downstream from the top of the slough. For each fish, we recorded the date of capture, species, total length (TL; mm), and whether the fish was previously marked. Unmarked fish were given a physical mark (e.g., hole punch of lower caudal, upper caudal, left pelvic, right pelvic) unique to each day and then were released. Fish release sites were distributed throughout Deadwater Slough to facilitate mixing back into the population. Fish that died prior to release were included in that day's total capture (or recapture) count, but not included in the number of marked fish available for recapture after that date. Finally, we recorded the angling start and end time for each crew (person or combinations of persons) to calculate CPUE as the number of Northern Pikeminnow caught per angler hour.

Two types of mark-recapture estimators were used to explore variation in abundance estimates: a single census and a multiple census estimator. For the single census estimator, we treated the first week of the survey as the mark event and the following week as the recapture event, pooling data within each week. Alternatively, the multiple census estimator treated each day as a sampling event and used information about the total number of marked fish from all previous events to calculate the total abundance. For the single census estimator, we used the Chapman-modified Lincoln-Peterson estimator:

$$\hat{N} = \frac{(M + 1)(n + 1)}{(m + 1)} - 1$$

where M is the total number of fish marked and returned to the population during the first event (week), n is the total number of fish caught in the second event (week), and m is the number of marked fish caught during the second week. For a multiple census estimator, we used the Chapman-modified Schnabel estimator:

$$\hat{N} = \frac{\sum_{i=1}^k n_i M_i}{(\sum_{i=1}^k m_i) + 1}$$

where M , n , and m are indexed by each survey (day), i . The 95% confidence intervals of the Schnabel estimator were calculated using a Poisson approximation (Krebs 1999).

Because fish capture was dependent on volunteer anglers, there was concern that sampling effort was biased to locations preferred by anglers. If release sites for marked fish were not proximal to preferred angling locales, recently released fish may have been less susceptible to immediate capture than unmarked fish. To account for this potential bias, we included an adjusted “delayed-

mixing” Schnabel estimator that assumed fish marked and released on a given survey day were not available for recapture for 48 hours. The delayed-mixing Schnabel estimator is given by:

$$\hat{N} = \frac{\sum_{i=1}^k n_i M_{i-1}}{(\sum_{i=1}^k m_i) + 1}$$

All mark-recapture abundance estimators include a Chapman correction for small sample sizes (Chapman 1951) and assumed: (1) the population is closed (no immigration, emigration, births or deaths during the survey period), (2) all fish have equal chance of being caught in subsequent survey days, (3) marking a fish does not affect its chances of recapture, (4) no loss of marks, and (5) marks are not missed or misidentified.

Surveys conducted in spring 2021 did not follow a mark-recapture study design so Northern Pikeminnow abundance was estimated as the ratio of mean total CPUE in the fall surveys to the total CPUE in the spring survey, then multiplied that ratio by the mean abundance from fall surveys:

$$\hat{N}_{spring} = \bar{N}_{fall} * \frac{CPUE_{spring}}{CPUE_{fall}}$$

This approach assumes equal capture probabilities for fall and spring surveys. Fall and spring population estimates were converted into linear and areal densities by dividing by the length and area of Deadwater Slough, respectively. Length and area were measured in QGIS software (QGIS Development Team 2022) and using drone-generated orthoimagery.

Anglers reported that Northern Pikeminnow in Deadwater Slough were large-bodied. To describe the population size structure, we calculated the proportional stock density (PSD) for

208 Northern Pikeminnow in Deadwater Slough. PSD is the percentage of all fish that meet criteria
209 for “quality-length” (larger) individuals and “stock-length” individuals:

$$210 \quad PSD = 100 * \frac{FQ}{FS}$$

211 where FQ is the number of fish \geq quality-length, and FS is the number of fish \geq stock-length.

212 For Northern Pikeminnow in Deadwater Slough, we used 380 mm TL for quality-length and 250
213 mm TL for stock-length (Winther et al. 2020).

214 **Stomach Contents**

215 Gastric lavage (Foster 1977) was used to examine the stomach contents of Northern Pikeminnow
216 for the presence of juvenile Chinook salmon and other fishes occupying Deadwater Slough.

217 Immediately following lavage, stomach contents of individuals were preserved with 99%
218 isopropyl alcohol in whirl-paks to be analyzed in a controlled environment. For each stomach
219 sample, total wet weight (grams) was recorded, including all fish and non-fish items (e.g.,

220 macroinvertebrates, organic matter). Fish and fish remnants were identified to the lowest
221 taxonomic unit using diagnostic bones (Hansel et al. 1988; Frost 2000) or were categorized as
222 unknown. Approximately 5% of Northern Pikeminnow were euthanized for dissection ($n = 75$)
223 after gastric lavage to validate the efficacy of the methodology. Contents of livewells were
224 examined during fish processing to ensure food items were not regurgitated.

225 **Fish Consumption Potential**

226 To estimate the total consumption potential of Northern Pikeminnow in Deadwater Slough
227 during the peaks of fall DSR and spring NRR emigrations, we used the Fish Bioenergetics v4.0

application developed by Deslauriers et al. (2017) applied in R statistical software (R Core Team 2021). The daily rate of consumption in grams for an individual Northern Pikeminnow was estimated based on predator and prey energy densities, predator start and end weights, and water temperatures. Separate models were run for the fall period (September 15 - November 30) and the spring period (March 1 - May 31) to coincide with peak emigrations of DSR and NRR juveniles from the Lemhi River, the largest Chinook salmon population in the Upper Salmon River.

Predator energy density for Northern Pikeminnow was fixed at 6,703 Joules (J)/g (Deslauriers et al. 2017). Prey energy densities were fixed at 3,000 J/g for invertebrates and 21,500 J/g for juvenile Chinook salmon (Moss et al. 2016). Because we were unable to differentiate juvenile Chinook salmon from other fish prey in model runs, we assume all fish prey have the same energy densities as juvenile Chinook salmon. The average TL of Northern Pikeminnow caught in Deadwater Slough during our study was converted to fork length (FL) and then to weight (grams) using a weight-length formula from Parker et al. (1995).

$$\log_{10}W_s = -4.886 + 2.986[\log_{10}(FL)]$$

This resulted in average fall DSR and spring NRR starting weights of 598.7 and 430.7 g for Northern Pikeminnow with average TLs of 394.1 and 352.9 mm, respectively. We assumed no growth in individual Northern Pikeminnow, resulting in equal start and end weights. Mean daily water temperatures were summarized from 15-minute interval temperature readings between March 3, 2013 and June 14, 2021 from USGS gage station 13307000, approximately 22 rkm downstream of Deadwater Slough.

The proportion of fish in the Northern Pikeminnow's diet relative to non-fish prey items is unknown in Deadwater Slough. Therefore, we conducted a series of model runs with varying proportions of fish in the diet ranging from 30 - 90%, in 10% increments. These values were supported by observations elsewhere in the Columbia River basin (54-86%: Shively et al. (1996); 48-86%: Zimmerman and Ward (1999); 37%: Gray and Dauble (2001)) that suggest fish are the majority of Northern Pikeminnow prey by volume. A model run was conducted for each combination of diet scenario and season, resulting in fourteen estimates of the total grams of fish consumed by an individual Northern Pikeminnow. To estimate the total biomass of fish consumed by Northern Pikeminnow in Deadwater Slough, we multiplied the grams of fish consumed by an individual Northern Pikeminnow by the estimated Northern Pikeminnow population sizes during fall DSR and spring NRR emigrations.

Impacts to Chinook Salmon Populations

The proportion of Chinook salmon prey relative to other fish prey in the Northern Pikeminnow diet is also unknown. However, there is some evidence that juvenile salmonids are by far the most consumed fish prey (Shively et al. 1996; Zimmerman and Ward 1999). Moreover, it is estimated that Chinook salmon make up 64.2%, 29.3%, and 49.3% of the fish prey consumed by Northern Pikeminnow in the Columbia River below Bonneville Dam, in Columbia River reservoirs, and in the lower Snake River, respectively (Zimmerman and Ward 1999). Because the bioenergetics model does not differentiate juvenile Chinook salmon from other fish prey items, we performed a sensitivity analysis to assess potential impacts of Northern Pikeminnow predation on local Chinook salmon populations. Using values similar to Zimmerman and Ward (1999), we modeled three diet scenarios where Chinook salmon comprised 30%, 50%, and 65%

of the fish prey, by weight, in the Northern Pikeminnow diet during the fall DSR and spring NRR emigration periods. In this analysis, we assume 60% of the Northern Pikeminnow's total diet is comprised of fish, representing the median scenario in our range of modeled diets.

We obtained an estimate of the number of juvenile Chinook salmon consumed by dividing total biomass consumed by the average weight of DSR (10.3 g) and NRR (10.9 g) emigrants in the Upper Salmon River. The average weights of DSR and NRR emigrants were calculated from fish captured at seven rotary screw traps upstream of Deadwater Slough during the fall and spring periods. Although the primary impact to Chinook salmon in Deadwater Slough by Northern Pikeminnow is juvenile predation, the common metric to evaluate salmon recovery is adult returns. We quantified the potential impact of Northern Pikeminnow predation on adult returns by estimating the number of equivalent adults expected to return to Lower Granite Dam if predation by Pikeminnow was eliminated in Deadwater Slough. To accomplish this, we multiplied the estimated total juvenile Chinook salmon consumed by the median Granite-to-Granite smolt-to-adult return rate (SAR) of 0.00614 (SD = 0.00051) from McCann et al. (2019) for Chinook salmon in the Upper Salmon River.

All data and code for the analyses presented here can be found in a GitHub repository release (<https://doi.org/10.5281/zenodo.6458124>).

RESULTS

Northern Pikeminnow Demographics

Using hook-and-line angling, we caught a total of 1,663 Northern Pikeminnow over the course of the study including 14 recaptures; the overall CPUE was 1.32 Northern Pikeminnow per angler

hour (Table 1). Mark-recapture abundance estimates of Northern Pikeminnow in Deadwater Slough ranged from 12,480 to 18,732 in fall 2019 and from 24,381 to 37,016 in fall 2020 (Table 3). We estimated larger populations of Northern Pikeminnow using the unadjusted multiple census estimator compared to the single census and adjusted delayed-mixing multiple census estimators (Figure 2). Our sampling design most closely matched a multiple census estimator; therefore, the Schnabel estimates were considered most appropriate. Accordingly, the mean Northern Pikeminnow abundance for the two fall sampling events (2019 and 2020) was 27,874 (95% CI: 14,244 - 59,388) using the unadjusted Schnabel estimator. Using the delayed-mixing Schnabel estimator the mean fall abundance estimate was 19,499 (95% CI: 9,952 - 41,597). All subsequent analyses use results from the adjusted delayed-mixing Schnabel estimator. For spring 2021, we estimated 10,352 (95% CI: 5,284 - 22,084) Northern Pikeminnow in Deadwater Slough. Those estimates translate to linear densities of 10,422 and 5,533 Northern Pikeminnow per rkm and areal densities (fish per 100 m²) of 16.3 and 8.7 for fall and spring, respectively. Lengths of Northern Pikeminnow ranged from 176 to 639 mm TL with an average of 389 mm (Figure 3). The PSD for Northern Pikeminnow in Deadwater Slough across all three surveys was 50%.

Stomach Contents

We found that gastric lavage successfully removed food items in all dissected individuals, indicating the method was effective. Of the 1,558 Northern Pikeminnow sampled using gastric lavage, we found contents in 350 stomachs and confirmed fish or fish parts in 44 of those. Northern Pikeminnow captured during spring 2021 had a higher proportion of fish content in their diet compared to individuals captured in the fall surveys (Table 4). The mean wet weight of

total contents for an individual was 0.98 g (median = 0.25 g; SD = 2.16 g). Overall, fish or fish remnants made up 11.7% of all stomach contents examined.

Fish Consumption Potential

During the fall DSR emigration, we estimated an average-size Northern Pikeminnow (394.1 TL mm) to consume 43.55 g of fish across the season to maintain their body size, assuming 60% of their diet consisted of fish prey (Figure 4). Fall consumption ranged from 35.74 g for a diet of 30% fish to 46.91 g for a diet of 90% fish. During the spring NRR emigration, we estimated an average-size Northern Pikeminnow (352.9 TL mm) to consume 42.51 g of fish, assuming 60% of their diet was fish prey (Figure 4). Spring consumption ranged from 34.01 for a diet of 30% fish to 44.33 for a diet of 90% fish. Daily consumption rates were higher early in the fall and late in the spring and corresponded with higher water temperatures, especially above 8 degrees Celsius (Figure 4). Given estimated mean Northern Pikeminnow population sizes of 19,499 in the fall and 10,352 in the spring 2021, and a diet consisting of 60% fish, the Northern Pikeminnow population was estimated to consume 0.8 metric tons of fish during the fall DSR emigration and 0.4 metric tons during the spring NRR emigration in Deadwater Slough.

Impacts to Chinook Salmon Populations

Given yearly fish consumption estimates during the fall DSR and spring NRR periods, we produced a sensitivity analysis to calculate the consumption of juvenile Chinook salmon where 30%, 50%, and 65% of total fish prey consumed were juvenile Chinook salmon (Figure 5). At the median value of 50%, we estimated that 61,409 (95% CI: 31,342 - 131,004) juvenile Chinook salmon would be consumed. Using the median Granite-to-Granite SAR for Chinook

salmon in the Upper Salmon River, we estimated the “adult equivalents” of juveniles consumed to be 377 (95% CI: 161 - 935) adults (Figure 6).

DISCUSSION

All mark-recapture abundance estimators suggest a large population of Northern Pikeminnow occupies Deadwater Slough. The mean population size of Northern Pikeminnow was estimated to be 19,499 during the fall DSR emigration and 10,352 during the spring NRR emigration even after accounting for the potential of delayed mixing of marked and released fish. Those estimates translate to linear densities (fish per rkm) that are 15 and 8 times greater than densities reported by Beamesderfer and Rieman (1991) for the John Day Reservoir and approximately 4 and 2 times greater than projections for the lower Columbia River (Beamesderfer et al. 1996). Beamesderfer and Rieman (1991) acknowledged that their methods were unsuitable for sampling offshore, noting that water velocity, depth, and irregular bottom contours, and barge traffic made sampling offshore ineffective; therefore, it is possible that their estimates may have been low. Conversely, the maximum depth at Deadwater Slough was approximately 6 m with a relatively homogenous bottom contour, making angling an effective method throughout the entire reach. Our estimated densities of Northern Pikeminnow suggest that slow-water reaches outside of reservoir complexes on the Snake and Columbia rivers may support exceptionally high predator densities, consistent with findings of Harnish et al. (2014), Gray and Dauble (2001), and Zimmerman and Ward (1999). Given the dearth of current data available throughout the Columbia River basin, it is unclear how Northern Pikeminnow abundance estimates in this study compare to elsewhere in the basin; additional information on contemporary piscine predator abundances may be needed (Widener et al. 2021).

357 Several assumptions in our estimators may have influenced the magnitude of our abundance
358 results. First, all models assumed a closed population; however, it is possible that immigration
359 and emigration occurred. Nevertheless, our estimators still provide unbiased estimates of
360 abundance assuming the immigration and emigration rates were equal between marked and
361 unmarked fish. Emigration of marked individuals could reduce the marking fraction in the
362 population leading to an upward bias in abundance estimates, but emigration rates would have
363 needed to be substantial. We also captured two fish with marks from previous years, indicating
364 seasonal residency of some fish. The multiple census estimators, which we report, are more
365 robust to this assumption because the marking fraction was estimated daily. We additionally
366 have little reason to believe that emigration occurs during the two-week survey window owing to
367 favorable habitat for Northern Pikeminnow in Deadwater Slough relative to adjacent reaches.
368 Given the size of the sampling area and the short duration of our surveys, the closed population
369 assumption was likely met. This assumption could likewise have affected results if mortality
370 occurred for some marked fish released back to the population. During field processing, no
371 mortalities or injuries post release were observed and fish injured prior to release were retained.
372 Nevertheless, the potential for mortality amongst marked and released fish cannot be discounted.
373 Next, our abundance estimates assumed that capture and recapture events are random samples of
374 the population. Angling methods often have a size selection bias, thereby limiting abundance
375 estimates to a size range susceptible to angling, resulting in a conservative estimate of the total
376 population. The estimators also assumed equal catchability of individual fish between sampling
377 events; two fish were recaptured during multiple days within a survey, indicating that marked
378 fish were continually susceptible to angling. It is also possible that marked and released fish may
379 not have immediately mixed thoroughly back into the population. Although we attempted to

release marked fish evenly throughout Deadwater Slough, we anecdotally observed anglers congregating in particular areas of the slough to socialize or exploit “good” fishing locales. Violations of the assumption of equal catchability between marked and unmarked fish may have led to an overestimate of abundance of Northern Pikeminnow in our study if marked fish have lower catchability. This bias will be proportional to the difference in catchability of marked and unmarked fish. As an example, if marked fish were only 70% as likely to be caught as unmarked fish, the true abundance is closer to 70% of our abundance estimate. This would additionally account for marked and released fish being “hook shy” for a period of time. The adjusted delayed-mixing Schnabel estimator may account for these biases; however, the magnitude of bias is unknown.

The spring abundance estimate was smaller than fall abundance estimates due to the spring CPUE being approximately half of the mean fall CPUE (Table 1). Although our approach assumed equal capture probabilities between the fall and spring surveys, we believe that higher spring flows may result in lower capture probabilities of Northern Pikeminnow. Therefore, we believe our spring abundance estimate to be conservative.

In addition to the large population abundance, the observed (50%) PSD in this study was greater than observations of 41% below Bonneville Dam and 18% in Bonneville Reservoir (Winther et al. 2020). This suggests that a larger fraction of Northern Pikeminnow in Deadwater Slough are of a quality size relative to populations reported elsewhere in the Columbia River. Notably, the Idaho state catch-and-release record Northern Pikeminnow, measuring 639 mm TL, was caught in Deadwater Slough during the fall 2020 survey alluding to the favorable conditions for Northern Pikeminnow.

402 We performed gastric lavage on nearly all Northern Pikeminnow collected during this study.
403 Observed prey species included juvenile Chinook salmon as well as Redside Shiner
404 *Richardsonius balteatus*, Largescale Sucker *Catostomus macrocheilus*, sculpin *Cottus* spp., and
405 Mountain Whitefish *Prosopium williamsoni*, though decomposition from digestion rendered
406 most fish contents unidentifiable. Although our approach provided useful information on prey
407 species, the use of angling and gastric lavage to quantify diet composition in Northern
408 Pikeminnow can be problematic. First, diets of angled fish can differ from those collected by
409 other methods such as electrofishing (Hodgson and Cochran 1988). For instance, angled fish are
410 more likely to have empty stomachs (Jurajda et al. 2016); possibly because they are captured
411 while actively searching for food, suggesting they are hungry. Second, unlike other predators
412 such as Smallmouth Bass and Walleye, that are commonly sampled using gastric lavage (Kamler
413 and Pope 2001), Northern Pikeminnow are cyprinids and therefore lack a true stomach.
414 Consequently, performing gastric lavage on cyprinid species can be ineffective for accurately
415 quantifying diet composition (Hartleb and Moring 1995), or alternately, requires a modification
416 in methodology to flush food items out the vent of the fish (Wasowicz and Valdez 1994).
417 Therefore, we consider our diet composition estimates from gastric lavage to be conservative.
418 Our bioenergetics approach assumed that all available prey consumed by Northern Pikeminnow
419 have energy densities equal to juvenile Chinook salmon. Generalizing energy densities using a
420 single species is a common approach (Petersen and Ward 1999). Other salmonids including
421 juvenile steelhead and Sockeye salmon are likely also available prey in Deadwater Slough,
422 especially during the spring migration, and so generalizing energy densities among salmonids
423 may not be problematic. Additionally, hatchery smolt releases (Chinook salmon, steelhead, and
424 Sockeye salmon) are also prevalent in the mainstem Salmon River during the spring migration.

To address the prevalence of other fish in the Northern Pikeminnow's diet, we considered scenarios where DSR and NRR Chinook salmon represent only 30%, 50%, and 65% of total fish prey consumed. These values may be considered conservative during the peak emigration periods when Chinook salmon are abundant and the Northern Pikeminnow diet shifts almost entirely to piscivory, presumably on juvenile Chinook salmon (Poe et al. 1991; Shively et al. 1996). This is likely the case during fall months when DSR Chinook salmon are the dominant prey species available within the Salmon River and Deadwater Slough as 1) few to no hatchery releases are present in the river and 2) fall emigrations of steelhead and Sockeye salmon are less prominent than Chinook salmon. During spring when natural-origin steelhead and Sockeye salmon are also actively emigrating from the Upper Salmon River and hatchery-origin releases of all three species (Chinook salmon, steelhead, Sockeye salmon) are present in the river, less than 50% Chinook salmon in the Northern Pikeminnow's diet may be more likely. Future work to quantify and identify juvenile salmonids in Deadwater Slough during their seasonal migration would be useful to validate the diet composition assumptions used in our model and to understand impacts to local populations.

Chinook salmon populations above Deadwater Slough are within the Upper Salmon major population group (MPG) which supports eight independent, extant populations including Salmon River (above Redfish Lake Creek), Valley Creek, Yankee Fork Salmon River, East Fork Salmon River, Salmon River (mainstem below Redfish Lake Creek), Pahsimeroi River, Lemhi River, and North Fork Salmon River (National Oceanic and Atmospheric Administration 2017). Recovery of the MPG is desired to support local fisheries and economies. At least five of the eight populations must meet criteria set forth by McElhany et al. (2000) and the Interior Columbia Technical Recovery Team (2007) for the MPG to be considered viable and for recovery of the

Snake River Evolutionary Significant Unit. We estimated that 61,409 juvenile Chinook salmon may be consumed by Northern Pikeminnow assuming that 60% of their diet is fish and 50% of fish prey are Chinook salmon. For context, this is approximately 1.35 times the average annual combined DSR and NRR emigration estimates for brood years 2011 - 2018 at a rotary screw trap located in the lower Lemhi River (Poole et al. 2019; Feeken et al. 2020; McClure et al. 2021), the largest population in the Upper Salmon MPG. Even under the most conservative scenario where 30% of the Northern Pikeminnow diet is fish and 30% of fish prey are Chinook salmon, we estimate that 29,988 juvenile Chinook salmon may be consumed which is 66% of the total DSR and NRR emigration for brood years 2011 - 2018. Considering that diet scenarios used in our study are likely conservative assumptions, especially during the fall DSR emigration, the estimated number of Chinook salmon consumed is substantial.

Spawner abundance is perhaps the most important metric considered in determining a population's viability and productivity. We estimated that consumption of juvenile Chinook salmon by Northern Pikeminnow in the Deadwater Slough is equivalent to 377 adults, annually, which is 68% of the total mean spawner escapement for the Upper Salmon MPG in 2017 – 2019 (Kinzer et al. 2020). In the most conservative scenario, we estimated juvenile consumption equivalent to 184 adults, which is 33% of recent natural-origin adult Chinook salmon returns. In the least conservative scenarios, adult equivalents approached recent escapements for the entire Upper Salmon MPG. The sensitivity analysis used some simplifying assumptions; for example, it assumed no juvenile mortality between Deadwater Slough and Lower Granite Dam which is an unrealistic assumption. Nevertheless, we found converting juveniles consumed to “adult equivalents” a useful exercise to place results in context with a metric commonly used for recovery.

471 In addition to juvenile Chinook salmon, juvenile steelhead and Sockeye salmon (both natural-
472 origin and hatchery releases) are also likely prey items for Northern Pikeminnow in Deadwater
473 Slough. Accordingly, we surmise that juvenile emigration and adult returns of steelhead and
474 Sockeye salmon are also affected by predation, including hatchery populations, which provide
475 for recreational fishing opportunities in the Upper Salmon River. Consequently, reducing
476 predation mortality in Deadwater Slough could potentially benefit multiple upriver natural and
477 hatchery populations, including other ESA-listed species. Because Deadwater Slough is part of
478 the migratory pathway for multiple species and populations of emigrating salmonids, the positive
479 impact could be greater than individual tributary rehabilitation actions which typically benefit a
480 single population. Predation on juveniles from any of the ESA-listed salmonid species is likely
481 detrimental to their recovery.

482 Deadwater Slough is a favorable candidate for management or restoration actions to benefit local
483 Chinook salmon populations. Three potential management actions could reduce predation at
484 Deadwater Slough: 1) removing or reducing the Dump Creek alluvial fan, 2) a local Northern
485 Pikeminnow bounty program to encourage harvest in Deadwater Slough aimed at reducing the
486 predator population size, and 3) adding structure or cover within Deadwater Slough to provide
487 refuge for juvenile salmonids to reduce predation rates. Reducing or removing the Dump Creek
488 alluvial fan has the benefit of restoring natural fluvial processes in the Salmon River that likely
489 existed in the reach prior to the formation or increase in size of the alluvial fan. Restoring natural
490 processes could reduce Northern Pikeminnow densities to levels similar to upstream and
491 downstream reaches where higher survival and transition probabilities for juvenile salmon have
492 been observed relative to Deadwater Slough (Axel et al. 2015; Ackerman et al. 2018; Porter et al.
493 2019). Managers ought also to consider the feasibility and net benefit of restoring fluvial

processes relative to potential losses for recreational fishing and bird watching opportunities (<https://www.audobon.org/important-bird-areas/deadwater-slough>).

Northern Pikeminnow bounty programs have proven successful at reducing population sizes and impacts on emigrating salmonids in the Columbia River (Winther et al. 2020). A local Northern Pikeminnow bounty program could provide monetary incentive for sportfisherman, in addition to boosting the local economy from lost revenues. Bounty programs could be conducted year-round or seasonally to coincide with peak juvenile outmigration(s). Each approach would require continual/annual effort and harvest to suppress the population size and as such, appropriate funding would be needed to support such a program. Lastly, a more passive approach could be to improve cover or structure within Deadwater Slough that provides refugia for juvenile salmonids during their emigration. Cover would need to be appropriately sized to provide concealment to juvenile fishes while reducing access by larger fishes like Northern Pikeminnow. Adding cover could be a cost-effective approach; however, its potential effectiveness is unclear.

Although the origins of Deadwater Slough are somewhat ambiguous (Reichmuth et al. 1985; USACE 1986), it appears likely that human activities in Dump Creek have either exacerbated or contributed to its formation, and in turn, created favorable conditions for Northern Pikeminnow. Among the three surveys, our lowest population abundance estimate was greater than 10,000 Northern Pikeminnow occupying Deadwater Slough, suggesting a remarkably high density given the size of the area. The slow water velocity and lack of cover for fish also create conditions where juvenile salmonids that are rearing in or emigrating through Deadwater Slough, including Chinook salmon, are susceptible to predation. We estimate that, at a minimum, Northern Pikeminnow in Deadwater Slough consume greater than 15,000 juvenile Chinook salmon (the lower endpoint of the confidence interval for the most conservative scenario); however, it is

517 likely that this value is closer to 61,000, annually. The magnitude of predation by Northern
518 Pikeminnow on adult returns is equivalent to 33% and 94% of recent adult escapements.
519 Deadwater Slough predates the ESA-listing of Chinook salmon populations in the Upper Salmon
520 MPG in the 1990s and is therefore unlikely to be the primary cause for the population's decline.
521 However, predation by Northern Pikeminnow in Deadwater Slough and elsewhere should be
522 considered among limiting factors hindering recovery efforts of ESA-listed Chinook salmon
523 populations in the Upper Salmon MPG.

524

ACKNOWLEDGEMENTS

The authors extend much appreciation to the many volunteers that assisted with field efforts including collaborators from the Bureau of Reclamation, Idaho Governor's Office of Species Conservation, Idaho Department of Fish and Game, and the Lemhi Regional Land Trust, among others. Special thanks to Braden Lott, Jared Barker, Brian Hamilton, Tulley Mackey, and Chelsea Welke for their help in the field. We further appreciate the administrative support and guidance from staff at Inter-Fluve. This manuscript benefited from reviews by Sean Gibbs, Mark Roes, and Ian Courter. Funding for this study was provided by the Bureau of Reclamation, Pacific Northwest Regional Office (contract No. 140R1021F0018) with assistance from the Idaho Governor's Office of Species Conservation.

LITERATURE CITED

- Ackerman, M. W., G. A. Axel, R. A. Carmichael, and K. See. 2018. Movement and distribution of sp/sum Chinook salmon presmolts in the mainstem Salmon River, pilot study. Technical report prepared for Idaho Governor's Office of Species Conservation.
- Atlas, W. I., N. C. Ban, J. W. Moore, A. M. Tuohy, S. Greening, A. J. Reid, N. Morven, E. White, W. G. Housty, J. A. Housty, C. N. Service, L. Greba, S. Harrison, C. Sharpe, K. I. R. Butts, W. M. Shepert, E. Sweeney-Bergen, D. Macintyre, M. R. Sloat, and K. Connors. 2021. Indigenous systems of management for culturally and ecologically resilient Pacific salmon (*Oncorhynchus* spp.) fisheries. *BioScience* 71(2):186–204.
- Axel, G. A., M. Peterson, C. C. Kozfkay, B. P. Sandford, M. G. Nesbit, B. J. Burke, K. E. Frick, and J. J. Lamb. 2015. Characterizing migration and survival between the upper Salmon River basin and Lower Granite Dam for juvenile Snake River sockeye salmon, 2014. Report of research by Fish Ecology Division, Northwest Fisheries Science Center, National Marine Fisheries Service, National Oceanic and Atmospheric Administration and Idaho Department of Fish and Game. Prepared for Division of Fish and Wildlife, Bonneville Power Administration. 36 pp.
- Beamesderfer, C. P., D. L. Ward, and A. A. Nigro. 1996. Evaluation of the biological basis for a predator control program on Northern Pikeminnow (*Ptychocheilus Oregonensis*) in the Columbia and Snake rivers. *Canadian Journal of Fisheries and Aquatic Sciences* 53(12):2898–2908.
- Beamesderfer, R. C., and B. E. Rieman. 1991. Abundance and distribution of Northern Squawfish, Walleyes, and Smallmouth Bass in John Day Reservoir, Columbia River. *Transactions of the American Fisheries Society* 120:439–447.

559 Chapman, D. G. 1951. Some Properties of the Hypergeometric Distribution with Applications to
560 Zoological Sample Censuses. University of California Publications Statistics 1:131–160.

561 Clark, C., P. Roni, J. Keeton, and G. Pess. 2020. Evaluation of the removal of impassable
562 barriers on anadromous salmon and steelhead in the Columbia River Basin. Fisheries
563 Management and Ecology 27(1):102–110.

564 Copeland, T., M. W. Ackerman, M. P. Corsi, P. Kennedy, K. K. Wright, M. R. Campbell, and
565 W. C. Schrader. 2014. Wild juvenile steelhead and Chinook salmon abundance and
566 composition at Lower Granite Dam, migratory years 2010 and 2011. Idaho Department
567 of Fish and Game.

568 Crozier, L. G., J. E. Siegel, L. E. Wiesebron, E. M. Trujillo, B. J. Burke, B. P. Sandford, and D.
569 L. Widener. 2020. Snake River Sockeye and Chinook salmon in a changing climate:
570 implications for upstream migration survival during recent extreme and future climates.
571 PLOS ONE 15(9):e0238886.

572 Deslauriers, D., S. R. Chipps, J. E. Breck, J. A. Rice, and C. P. Madenjian. 2017. Fish
573 Bioenergetics 4.0: An R-based modeling application. Fisheries 42(11):586–596.

574 Dodson, J. J., N. Aubin-Horth, V. Thériault, and D. J. Páez. 2013. The evolutionary ecology of
575 alternative migratory tactics in salmonid fishes: alternative migratory tactics as threshold
576 traits. Biological Reviews 88(3):602–625.

577 Emerson, J. L. 1973. A history of the Salmon National Forest. United States Forest Service. 194
578 pp.

579 Feeken, S. F., B. Barnett, E. Felts, E. J. Stark, M. Davison, J. R. Poole, C. McClure, B. A. Knoth,
580 and M. E. Dobos. 2020. Idaho Anadromous Emigrant Monitoring, 2019 Annual Report.
581 IDFG Report Number 20-09:67.

582 Foster, J. R. 1977. Pulsed gastric lavage: An efficient method of removing the stomach contents
583 of live fish. *The Progressive Fish-Culturist* 39(4):166–169.

584 Friesen, T. A., and D. L. Ward. 1999. Management of Northern Pikeminnow and implications
585 for juvenile salmonid survival in the lower Columbia and Snake rivers. *North American*
586 *Journal of Fisheries Management* 19(2):406–420.

587 Frost, C. N. 2000. A key for identifying preyfish in the Columbia River based on diagnostic
588 bones. U.S. Geological Survey, Western Fisheries Research Center, Columbia River
589 Research Laboratory.

590 Gray, R. H., and D. D. Dauble. 2001. Some life history characteristics of cyprinids in the
591 Hanford Reach, Mid-Columbia River. *Northwest Science* 75(2):122–136.

592 Hansel, H. C., S. D. Duke, P. T. Lofy, and G. A. Gray. 1988. Use of diagnostic bones to identify
593 and estimate original lengths of ingested prey fishes. *Transactions of the American*
594 *Fisheries Society* 117(1):55–62.

595 Harnish, R. A., E. D. Green, K. A. Deters, K. D. Ham, Z. Deng, H. Li, B. Rayamajhi, K. W.
596 Jung, and G. A. McMichael. 2014. Survival of wild Hanford Reach and Priest Rapids
597 Hatchery fall Chinook salmon juveniles in the Columbia River: predation implications.
598 Pacific Northwest National Laboratory.

599 Hartleb, C. F., and J. R. Moring. 1995. An improved gastric lavage device for removing stomach
600 contents from live fish. *Fisheries Research* 24(3):261–265.

601 Hodgson, J. R., and P. A. Cochran. 1988. The effect of sampling methodology on diet analysis in
602 largemouth bass (*Micropterus Salmoides*). *SIL Proceedings, 1922-2010* 23(3):1670–
603 1675.

604 Interior Columbia Technical Recovery Team. 2007. Viability criteria for application to interior
 605 Columbia basin salmonid ESUs. National Marine Fisheries Service, Northwest Fisheries
 606 Science Center, National Oceanographic and Atmospheric Administration.

607 Jurajda, P., K. Roche, I. Sedlacek, and L. Vsetickova. 2016. Assemblage characteristics and diet
 608 of fish in the shallow coastal waters of James Ross Island, Antarctica. *Polar Biology*
 609 39(12):2299–2309.

610 Justice, C., S. M. White, D. A. McCullough, D. S. Graves, and M. R. Blanchard. 2017. Can
 611 stream and riparian restoration offset climate change impacts to salmon populations?
 612 *Journal of Environmental Management* 188(2017):212–227.

613 Kamler, J. F., and K. L. Pope. 2001. Nonlethal methods of examining fish stomach contents.
 614 *Reviews in Fisheries Science* 9(1):1–11.

615 Kinzer, R., R. Orme, M. Campbell, J. Hargrove, and K. See. 2020. Report to NOAA Fisheries
 616 for 5-Year ESA Status Review: Snake River Basin Steelhead and Chinook salmon
 617 population abundance, life history, and diversity metrics calculated from in-stream PIT-
 618 tag observations (SY2010-SY2019). IPTDSW (In-stream PIT-tag detection systems
 619 workgroup). 118 pp.

620 Knutsen, C. J., and D. L. Ward. 1999. Biological characteristics of Northern Pikeminnow in the
 621 lower Columbia and Snake rivers before and after sustained exploitation. *Transactions of*
 622 *the American Fisheries Society* 128(6):1008–1019.

623 Krebs, C. J. 1999. *Ecological methodology*. Addison Welsey Educational Publishers, Inc., Menlo
 624 Park, CA.

625 Lewis, D. J., S. J. Dundas, D. M. Kling, D. K. Lew, and S. D. Hacker. 2019. The non-market
 626 benefits of early and partial gains in managing threatened salmon. PLOS ONE
 627 14(8):e0220260.

628 McCann, J., B. Chockley, E. Cooper, B. Hsu, G. Scheer, S. Haeseker, R. Lessard, T. Copeland,
 629 E. Tinus, A. Storch, D. Rawding, and M. DeHart. 2019. Comparative survival study of
 630 PIT-tagged spring/summer/fall Chinook, summer steelhead, and Sockeye. Draft 2019
 631 annual report. BPA Project #19960200. Available at:
 632 https://www.fpc.org/documents/CSS/2019CSS_FullDRAFT.pdf.

633 McClure, C., B. Barnett, E. A. Felts, M. Davison, N. Smith, B. A. Knoth, J. R. Poole, and S. F.
 634 Feeken. 2021. Idaho Anadromous Emigrant Monitoring, 2020 Annual Report. IDFG
 635 Report Number 21-11:65.

636 McElhany, P., M. H. Ruckelshaus, M. J. Ford, T. C. Wainwright, and E. P. Bjorkstedt. 2000.
 637 Viable salmonid populations and the recovery of evolutionarily significant units. U.S.
 638 Department of Commerce, National Oceanic and Atmospheric Administration, Technical
 639 Memorandum. NMFS-NWFSC-42.:156 pp.

640 Moss, J., J. Murphy, E. Fergusson, and R. Heintz. 2016. Allometric relationships between body
 641 size and energy density of juvenile Chinook (*Oncorhynchus Tshawytscha*) and Chum (*O.*
 642 *Keta*) salmon across a latitudinal gradient. North Pacific Anadromous Fish Commission
 643 Bulletin 6(1):161–168.

644 Murphy, C. A., J. D. Romer, K. Stertz, I. Arismendi, R. Emig, F. Monzyk, and S. L. Johnson.
 645 2021. Damming salmon fry: evidence for predation by non-native warmwater fishes in
 646 reservoirs. Ecosphere 12(9):e03757.

647 National Oceanic and Atmospheric Administration. 2017. ESA recovery plan for Snake River
648 spring/summer Chinook salmon (*Oncorhynchus Tshawytscha*) and Snake River basin
649 steelhead (*Oncorhynchus Mykiss*).

650 Nemeth, D. J., and R. B. Kiefer. 1999. Snake River spring and summer Chinook salmon choice
651 for recovery. *Fisheries* 24(10):16–23.

652 Parker, R. M., M. P. Zimmerman, and D. L. Ward. 1995. Variability in biological characteristics
653 of Northern Squawfish in the lower Columbia and Snake rivers 124:335–346.

654 Petersen, J. H. 1994. Importance of spatial pattern in estimating predation on juvenile salmonids
655 in the Columbia River. *Transactions of the American Fisheries Society* 123(6):924–930.

656 Petersen, J. H., and D. L. Ward. 1999. Development and corroboration of a bioenergetics model
657 for Northern Pikeminnow feeding on juvenile salmonids in the Columbia River.
658 *Transactions of the American Fisheries Society* 128(5):784–801.

659 Poe, T. E., H. C. Hansel, S. Vigg, D. E. Palmer, and L. A. Prendergast. 1991. Feeding of
660 predaceous fishes on out-migrating juvenile salmonids in John Day Reservoir, Columbia
661 River. *Transactions of the American Fisheries Society* 120(4):405–420.

662 Poole, J. R., E. Felts, M. Dobos, B. Barnett, M. Davison, C. J. Roth, B. A. Knoth, and E. J. Stark.
663 2019. Idaho Anadromous Emigrant Monitoring, 2018 Annual Report. IDFG Report
664 Number 19-11:71.

665 Porter, N. J., M. W. Ackerman, T. Mackey, G. A. Axel, and K. E. See. 2019. Movement and
666 distribution of Chinook salmon presmolts in the mainstem Salmon River, 2018/2019
667 Annual Report. Prepared by Biomark, Inc. - Applied Biological Services and Fish
668 Ecology Division, Northwest Fisheries Science Center, National Marine Fisheries
669 Service, National Oceanic and Atmospheric Administration. 47 pp.

670 QGIS Development Team. 2022. QGIS Geographic Information System. Open Source
671 Geospatial Foundation.

672 R Core Team. 2021. R: A language and environment for statistical computing. R Foundation for
673 Statistical Computing, Vienna, Austria.

674 Reichmuth, D., A. Potter, M. Knops, and M. Leaverton. 1985. Reducing ice jams along the
675 Salmon River Deadwater Reach: problems, proposals and prospects. Preliminary review,
676 evaluation and proposal. Page 69. Geomax. Prepared for Idaho Department of Fish and
677 Game.

678 Roni, P., P. J. Anders, T. J. Beechie, and D. J. Kaplowe. 2018. Review of tools for identifying,
679 planning, and implementing habitat restoration for Pacific salmon and steelhead. North
680 American Journal of Fisheries Management 38(2):355–376.

681 Rubenson, E. S., D. J. Lawrence, and J. D. Olden. 2020. Threats to rearing juvenile Chinook
682 salmon from nonnative Smallmouth Bass inferred from stable isotope and fatty acid
683 biomarkers. Transactions of the American Fisheries Society 149(3):350–363.

684 Shively, R. S., T. P. Poe, M. B. Sheer, and R. Peters. 1996. Criteria for reducing predation by
685 Northern Squawfish near juvenile salmonid bypass outfalls at Columbia River dams.
686 Regulated Rivers: Research & Management 12(4-5):493–500.

687 Tabor, R. A., R. S. Shively, and T. P. Poe. 1993. Predation of juvenile salmonids by Smallmouth
688 Bass and Northern Pikeminnow in the Columbia River near Richland, Washington. North
689 American Journal of Fisheries Management 13(4):831–838.

690 USACE. 1986. Salmon River flood damage reduction. Lemhi County, Idaho. Detailed project
691 report and Environmental Impact Statement. Page 705. US Army Corps of Engineers.

692 Venditti, D. A., D. W. Rondorf, and J. M. Kraut. 2000. Migratory behavior and forebay delay of
693 radio-tagged juvenile fall Chinook salmon in a lower Snake River impoundment. *North*
694 *American Journal of Fisheries Management* 20(1):41–52.

695 Ward, D. L., J. H. Petersen, and J. J. Loch. 1995. Index of predation on juvenile salmonids by
696 Northern Squawfish in the lower and middle Columbia River and in the lower Snake
697 River. *Transactions of the American Fisheries Society* 124:321–334.

698 Wasowicz, A., and R. A. Valdez. 1994. A nonlethal technique to recover gut contents of
699 Roundtail Chub. *North American Journal of Fisheries Management* 14(3):656–658.

700 White, S. M., S. Brandy, C. Justice, K. A. Morinaga, L. Naylor, J. Ruzycki, E. R. Sedell, J.
701 Steele, A. Towne, J. G. Webster, and I. Wilson. 2021. Progress towards a comprehensive
702 approach for habitat restoration in the Columbia Basin: case study in the Grande Ronde
703 River. *Fisheries* 46(5):229–243.

704 Widener, D. L., J. R. Faulkner, S. G. Smith, T. M. Marsh, and R. W. Zabel. 2021. Survival
705 estimates for the passage of spring-migrating juvenile salmonids through Snake and
706 Columbia River dams and reservoirs, 2020. U.S. Department of Commerce, NOAA
707 Contract Report NMFS-NWFSC-CR-2021-05.

708 Winther, E., C. M. Barr, C. Miller, and C. Wheaton. 2020. Report on the predation index,
709 predator control fisheries, and program evaluation for the Columbia River Basin Northern
710 Pikeminnow Sport Reward Program, 2020 Annual Report. 155 pp. Pacific States Marine
711 Fisheries Commission, Washington Dept. of Fish and Wildlife, Oregon Dept. of Fish and
712 Wildlife.

713 Wydoski, R. S., and R. R. Whitney. 2003. *Inland fishes of Washington: second edition, revised*
714 *and expanded*. University of Washington Press.

715 Zimmerman, M. P., and D. L. Ward. 1999. Index of predation on juvenile salmonids by Northern
716 Pikeminnow in the lower Columbia River basin from 1994-96. Transactions of the
717 American Fisheries Society 128(6):995–1007.

718 Zimmerman, M., K. Krueger, P. Roni, B. Bilby, J. Walters, and T. Quinn. 2012. Intensively
719 Monitored Watersheds Program: an updated plan to monitor fish and habitat responses to
720 restoration actions in the lower Columbia watersheds. Washington Department of Fish
721 and Wildlife. Olympia, Washington.

722

723 **TABLES**

724 TABLE 1. Summary of Northern Pike minnow captures from angling surveys in Deadwater
 725 Slough. Fall surveys included mark and recapture weeks for the single census estimate. Spring
 726 surveys included only a single capture week to estimate CPUE.

Survey Week	Date	Fish Caught	Marked Fish Caught	Marked Fish Returned	CPUE
Fall 2019					
1	Nov 12	29	0	28	1.83
	Nov 13	146	0	146	
	Nov 14	93	1	93	
2	Nov 19	149	2	132	
	Nov 20	104	1	77	
	Nov 21	143	4	118	
Fall 2020					
1	Oct 20	173	0	170	1.23
	Oct 21	188	1	187	
	Oct 22	104	0	102	
	Oct 23	41	0	41	
2	Oct 27	42	0	41	
	Oct 28	47	1	46	
	Oct 29	157	4	156	
	Oct 30	45	0	45	
Spring 2021					
1	May 18	85	-	-	0.81
	May 19	64	-	-	
	May 20	41	-	-	
	May 21	12	-	-	

727

728 TABLE 2. Estimated fish consumed by an individual Northern Pikeminnow to maintain its body
 729 weight during the fall DSR and spring NRR Chinook salmon emigrations. Bioenergetic model
 730 results for each season include seven model runs that assume different percentages of fish in the
 731 diet. Northern Pikeminnow start and end weights were 598.7 g for the fall season and 430.7 g for
 732 the spring season.

Date Range	Fish in Diet (%)	g Fish Consumed (g)
Fall		
Sep 15 - Nov 30	30	35.74
	40	39.37
	50	41.75
	60	43.55
	70	44.88
	80	45.98
	90	46.91
Spring		
Mar 1 - May 31	30	34.01
	40	39.54
	50	39.54
	60	42.51
	70	42.51
	80	43.52
	90	44.33

733

734 TABLE 3. Estimated Northern Pikeminnow in Deadwater Slough and the associated standard
 735 errors and 95% confidence intervals from mark-recapture estimators calculated for the fall
 736 surveys. Standard errors are not available for the multiple census estimators.

Estimator	N	SE	95% CI
Fall 2019			
Chapman	13,298	4,322	6,898 - 27,893
Schnabel	18,732	NA	10,057 - 37,851
Schnabel - Delayed Mixing	12,480	NA	6,701 - 25,219
Fall 2020			
Chapman	24,381	9,066	11,547 - 55,761
Schnabel	37,016	NA	18,430 - 80,924
Schnabel - Delayed Mixing	26,518	NA	13,203 - 57,975

737

738 TABLE 4. Total number of Northern Pikeminnow gastric lavaged and the number of those
 739 individuals with contents (including fish) and with contents that include fish.

Survey	Lavaged	Stomach Contents	Fish Contents
Fall 2019	660	57	12
Fall 2020	793	188	25
Spring 2021	105	105	7
Total	1,558	350	44

740

FIGURES

FIGURE 1. Map of the Deadwater Slough study area within the Upper Salmon River MPG. The Dump Creek alluvial fan is located at the downstream end of the study reach.

FIGURE 2. Estimated abundance of Northern Pikeminnow in Deadwater Slough from three mark-recapture estimators for the fall surveys. Error bars indicate 95% confidence intervals.

FIGURE 3. Length frequency histogram of Northern Pikeminnow caught using hook-and-line angling during the study.

FIGURE 4. The cumulative fish prey consumed (g) by an individual Northern Pikeminnow during the peak Chinook salmon emigrations for fall DSR (top left) and spring NRR (top right) and corresponding daily mean water temperatures (bottom panels). The black line shows the consumption by Northern Pikeminnow when the diet consists of 60% fish and the gray area shows the range between all model runs in the series, from 30% to 90% fish prey. Daily mean water temperatures were summarized from six years of data available from the Shoup gage station.

FIGURE 5. Estimated number of juvenile Chinook salmon consumed by Northern Pikeminnow from a sensitivity analysis of diet scenarios made up of variable percentages of fish. Diet scenarios include 30%, 50%, and 65% Chinook salmon in the total fish prey consumed by Northern Pikeminnow, similar to those reported by Zimmerman and Ward 1999.

FIGURE 6. Estimated “adult equivalents” of the total juvenile Chinook salmon consumed by Northern Pikeminnow in Deadwater Slough. Error bars indicate 95% confidence intervals.

762 Colophon

763 This report was generated on 2022-04-12 15:23:58 using the following computational
764 environment and dependencies:

```
765 #> - Session info -----  
766 ----  
767 #> setting value  
768 #> version R version 4.1.1 (2021-08-10)  
769 #> os Windows 10 x64 (build 22000)  
770 #> system x86_64, mingw32  
771 #> ui RTerm  
772 #> language (EN)  
773 #> collate English_United States.1252  
774 #> ctype English_United States.1252  
775 #> tz America/Denver  
776 #> date 2022-04-12  
777 #> pandoc 2.14.0.3 @ C:/Program Files/RStudio/bin/pandoc/ (via rmarkdown)  
778 #>  
779 #> - Packages -----  
780 ----  
781 #> package * version date (UTC) lib source  
782 #> assertthat 0.2.1 2019-03-21 [1] CRAN (R 4.1.1)  
783 #> backports 1.4.1 2021-12-13 [1] CRAN (R 4.1.2)  
784 #> bit 4.0.4 2020-08-04 [1] CRAN (R 4.1.1)  
785 #> bit64 4.0.5 2020-08-30 [1] CRAN (R 4.1.1)
```

786	#>	bookdown	0.24	2021-09-02	[1]	CRAN	(R 4.1.1)
787	#>	broom	0.7.11	2022-01-03	[1]	CRAN	(R 4.1.2)
788	#>	cachem	1.0.6	2021-08-19	[1]	CRAN	(R 4.1.1)
789	#>	callr	3.7.0	2021-04-20	[1]	CRAN	(R 4.1.1)
790	#>	cellranger	1.1.0	2016-07-27	[1]	CRAN	(R 4.1.1)
791	#>	cli	3.1.1	2022-01-20	[1]	CRAN	(R 4.1.2)
792	#>	colorspace	2.0-2	2021-06-24	[1]	CRAN	(R 4.1.2)
793	#>	crayon	1.5.0	2022-02-14	[1]	CRAN	(R 4.1.2)
794	#>	data.table	1.14.2	2021-09-27	[1]	CRAN	(R 4.1.1)
795	#>	DBI	1.1.2	2021-12-20	[1]	CRAN	(R 4.1.2)
796	#>	dbplyr	2.1.1	2021-04-06	[1]	CRAN	(R 4.1.1)
797	#>	desc	1.4.0	2021-09-28	[1]	CRAN	(R 4.1.1)
798	#>	devtools	2.4.2	2021-06-07	[1]	CRAN	(R 4.1.1)
799	#>	digest	0.6.28	2021-09-23	[1]	CRAN	(R 4.1.1)
800	#>	dplyr	* 1.0.7	2021-06-18	[1]	CRAN	(R 4.1.1)
801	#>	ellipsis	0.3.2	2021-04-29	[1]	CRAN	(R 4.1.1)
802	#>	english	* 1.2-6	2021-08-21	[1]	CRAN	(R 4.1.2)
803	#>	evaluate	0.14	2019-05-28	[1]	CRAN	(R 4.1.1)
804	#>	fansi	1.0.2	2022-01-14	[1]	CRAN	(R 4.1.2)
805	#>	farver	2.1.0	2021-02-28	[1]	CRAN	(R 4.1.1)
806	#>	fastmap	1.1.0	2021-01-25	[1]	CRAN	(R 4.1.1)
807	#>	forcats	* 0.5.1	2021-01-27	[1]	CRAN	(R 4.1.1)
808	#>	fs	1.5.0	2020-07-31	[1]	CRAN	(R 4.1.1)
809	#>	FSA	* 0.9.1	2021-07-17	[1]	CRAN	(R 4.1.2)
810	#>	generics	0.1.2	2022-01-31	[1]	CRAN	(R 4.1.2)

811	#> ggplot2	* 3.3.5	2021-06-25	[1]	CRAN (R 4.1.1)
812	#> glue	1.6.1	2022-01-22	[1]	CRAN (R 4.1.2)
813	#> gridExtra	* 2.3	2017-09-09	[1]	CRAN (R 4.1.2)
814	#> gtable	0.3.0	2019-03-25	[1]	CRAN (R 4.1.1)
815	#> haven	2.4.3	2021-08-04	[1]	CRAN (R 4.1.1)
816	#> here	* 1.0.1	2020-12-13	[1]	CRAN (R 4.1.1)
817	#> highr	0.9	2021-04-16	[1]	CRAN (R 4.1.1)
818	#> hms	1.1.1	2021-09-26	[1]	CRAN (R 4.1.1)
819	#> htmltools	0.5.2	2021-08-25	[1]	CRAN (R 4.1.1)
820	#> http	1.4.2	2020-07-20	[1]	CRAN (R 4.1.1)
821	#> janitor	* 2.1.0	2021-01-05	[1]	CRAN (R 4.1.1)
822	#> jsonlite	1.7.3	2022-01-17	[1]	CRAN (R 4.1.2)
823	#> kableExtra	* 1.3.4	2021-02-20	[1]	CRAN (R 4.1.1)
824	#> knitr	1.37	2021-12-16	[1]	CRAN (R 4.1.2)
825	#> labeling	0.4.2	2020-10-20	[1]	CRAN (R 4.1.1)
826	#> lattice	0.20-44	2021-05-02	[2]	CRAN (R 4.1.1)
827	#> lemon	* 0.4.5	2020-06-08	[1]	CRAN (R 4.1.2)
828	#> lifecycle	1.0.1	2021-09-24	[1]	CRAN (R 4.1.1)
829	#> lubridate	* 1.8.0	2021-10-07	[1]	CRAN (R 4.1.1)
830	#> magrittr	* 2.0.1	2020-11-17	[1]	CRAN (R 4.1.1)
831	#> memoise	2.0.1	2021-11-26	[1]	CRAN (R 4.1.2)
832	#> modelr	0.1.8	2020-05-19	[1]	CRAN (R 4.1.1)
833	#> munsell	0.5.0	2018-06-12	[1]	CRAN (R 4.1.1)
834	#> pillar	1.7.0	2022-02-01	[1]	CRAN (R 4.1.2)
835	#> pkgbuild	1.3.1	2021-12-20	[1]	CRAN (R 4.1.2)

836	#>	pkgconfig	2.0.3	2019-09-22	[1]	CRAN	(R 4.1.1)
837	#>	pkgload	1.2.4	2021-11-30	[1]	CRAN	(R 4.1.2)
838	#>	plyr	1.8.6	2020-03-03	[1]	CRAN	(R 4.1.1)
839	#>	prettyunits	1.1.1	2020-01-24	[1]	CRAN	(R 4.1.1)
840	#>	processx	3.5.2	2021-04-30	[1]	CRAN	(R 4.1.1)
841	#>	ps	1.6.0	2021-02-28	[1]	CRAN	(R 4.1.1)
842	#>	purrr	* 0.3.4	2020-04-17	[1]	CRAN	(R 4.1.1)
843	#>	R6	2.5.1	2021-08-19	[1]	CRAN	(R 4.1.1)
844	#>	Rcpp	1.0.7	2021-07-07	[1]	CRAN	(R 4.1.1)
845	#>	readr	* 2.1.2	2022-01-30	[1]	CRAN	(R 4.1.2)
846	#>	readxl	1.3.1	2019-03-13	[1]	CRAN	(R 4.1.1)
847	#>	remotes	2.4.2	2021-11-30	[1]	CRAN	(R 4.1.2)
848	#>	reprex	2.0.1	2021-08-05	[1]	CRAN	(R 4.1.1)
849	#>	rlang	0.4.12	2021-10-18	[1]	CRAN	(R 4.1.1)
850	#>	rmarkdown	2.11	2021-09-14	[1]	CRAN	(R 4.1.1)
851	#>	rprojroot	2.0.2	2020-11-15	[1]	CRAN	(R 4.1.1)
852	#>	rstudioapi	0.13	2020-11-12	[1]	CRAN	(R 4.1.1)
853	#>	rvest	1.0.2	2021-10-16	[1]	CRAN	(R 4.1.1)
854	#>	scales	1.1.1	2020-05-11	[1]	CRAN	(R 4.1.1)
855	#>	sessioninfo	1.2.2	2021-12-06	[1]	CRAN	(R 4.1.2)
856	#>	snakecase	0.11.0	2019-05-25	[1]	CRAN	(R 4.1.1)
857	#>	stringi	1.7.6	2021-11-29	[1]	CRAN	(R 4.1.2)
858	#>	stringr	* 1.4.0	2019-02-10	[1]	CRAN	(R 4.1.1)
859	#>	svglite	2.0.0	2021-02-20	[1]	CRAN	(R 4.1.1)
860	#>	systemfonts	1.0.3	2021-10-13	[1]	CRAN	(R 4.1.1)

```

861 #> testthat      3.1.0    2021-10-04 [1] CRAN (R 4.1.1)
862 #> tibble        * 3.1.6    2021-11-07 [1] CRAN (R 4.1.2)
863 #> tidyr         * 1.2.0    2022-02-01 [1] CRAN (R 4.1.2)
864 #> tidyselect    1.1.1    2021-04-30 [1] CRAN (R 4.1.2)
865 #> tidytable     * 0.6.5    2021-09-03 [1] CRAN (R 4.1.1)
866 #> tidyverse     * 1.3.1    2021-04-15 [1] CRAN (R 4.1.1)
867 #> tzdb          0.2.0    2021-10-27 [1] CRAN (R 4.1.1)
868 #> usethis       2.1.3    2021-10-27 [1] CRAN (R 4.1.1)
869 #> utf8          1.2.2    2021-07-24 [1] CRAN (R 4.1.1)
870 #> vctrs         0.3.8    2021-04-29 [1] CRAN (R 4.1.1)
871 #> viridisLite   0.4.0    2021-04-13 [1] CRAN (R 4.1.1)
872 #> vroom         1.5.7    2021-11-30 [1] CRAN (R 4.1.2)
873 #> webshot       0.5.2    2019-11-22 [1] CRAN (R 4.1.1)
874 #> withr         2.4.3    2021-11-30 [1] CRAN (R 4.1.2)
875 #> xfun          0.27     2021-10-18 [1] CRAN (R 4.1.1)
876 #> xml2          1.3.3    2021-11-30 [1] CRAN (R 4.1.2)
877 #> yaml          2.2.1    2020-02-01 [1] CRAN (R 4.1.1)

```

```
878 #>
```

```
879 #> [1] C:/Users/mikea/Documents/R/win-library/4.1
```

```
880 #> [2] C:/Program Files/R/R-4.1.1/library
```

```
881 #>
```

```
882 #> -----
```

```
883 ----
```

```
884
```

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5





Figure 6





[Click here to access/download](#)

Supporting Information

Supplement - Avian Predation in Deadwater
Slough.docx

