

i. Significance

Dehydration, whether caused by exposure to extreme environmental conditions, water deprivation, or by infection (e.g. diarrheal illnesses) represents a significant threat to human life. When death is avoided, dehydration may lead to chronic health conditions like renal failure. While the mechanisms underlying physiological compromise are well characterized [ref], some animals possess the ability, much unlike humans, to osmoregulate despite extreme heat and a complete lack of extrinsic water intake [ref]. Specifically, highly adapted desert mice may never drink water [ref], produce an extremely viscous urine (or no urine at all) [ref], and excrete urea in the form of uric acid crystals in the feces [ref]. Focusing on osmoregulation, patterns of renal gene expression have been shown to be highly derived in some desert adapted rodents (e.g. *Dipodomys* [ref]), but not in others like (*Notomys*), and therefore the extent to which differences in gene expression underlies phenotype remains unknown. In addition to its' generality, both studies of non-model organisms focus on a limited number of genes, and therefore may not appropriately assay the complexity of the genetics of extreme renal osmoregulation. In contrast with this work, several studies attempting to understand the effects of dehydration in model organisms (e.g. *Mus* and *Rattus*) exist, though while these works benefit from an extensive genomic and physiologic toolbox, that they lack the appropriate phenotype (extreme osmoregulatory abilities) limits insight. In summary, the study of extreme renal osmoregulation is hindered on one hand by a lack of tools, and on the other by a lack of an appropriate phenotype. This limitation may undercut our ability to efficiently develop novel therapies directed at mitigating the untoward effects of dehydration.

The proposed research uses a novel approach integrating physiology, evolutionary genomics, and computational biology to better understand how animals survive in what appear to be non-survivable conditions. I develop the genomic and physiologic tools in a novel model for the study of extreme osmoregulation, *Peromyscus eremicus*. It's contribution is to effectively leverage the power of a sophisticated toolset of a model organism against a uniquely adapted rodent, which will allow for a synthetic understanding of extreme osmoregulation, and ultimately novel insights into the cause of - and cure for - dehydration related mortality and morbidity.


ii. Innovation

In spite of modern medicine, millions of people die from dehydration. Millions more are affected by chronic kidney disease, a condition whose underlying causes are diverse, though many resemble the pathophysiology of dehydration (e.g. low renal perfusion pressure). The proposed work recognizes that successful treatment requires an appropriate model, and while traditional models are wonderful, they lack the biology (extreme osmoregulation) upon which more successful interventions may be modeled. The desert-adapted rodent *P. eremicus* retains many of the beneficial characteristics of model organisms, while enhancing opportunity to assay interesting biological phenomenon. In addition to this fundamental innovation, the project is innovative in a number of other ways.

- Leverage an unprecedented level of control over the experimental environment by use of a desert chamber where natural conditions can be replicated while simultaneously preserving the ability to manipulate water availability.
- Link detailed information on physiology and metabolism to genomic information using a novel analytical tools under active development in the lab.

Aim 1: Using captive desert-adapted mice, I will correlate multiple physiological variables, as well as their genomic underpinnings to differences in temperature, relative humidity, and water availability. I hypothesize that, as a result of unique mechanisms related to solute and water balance, that average serum electrolyte concentrations will remain relatively constant throughout various experimental manipulations, but the variance in measured levels between individuals will increase in the most extreme conditions. These differences will be echoed in differences in urine electrolytes and concentration.

To better understand the physiological effects of desert conditions on rodents, I will relate multiple physiological variables to differences in temperature, relative humidity, and water availability. These experiments are fundamentally linked to a series of environmental manipulations, described in Figure 1. The experimental design is fully factorial -- meaning that the focal experimental parameter (e.g. temperature) will be tested in the context of the full range of other conditions (e.g. humidity, water availability). Twenty individuals per treatment will be included -- power analyses suggest this sample size will allow me to garner statistical support for patterns with small to medium effect sizes. Together, this design will allow me to tease-apart the physiologic and genomic response to the various conditions.



For each experiment in Aim 1, I will collect information on metabolism, urine concentration and electrolyte content, and serum electrolyte content. In the context of limited water intake, how animals achieve electrolyte balance is unknown. Electrolytes are both easy to assay and are critical to physiological well being. Indeed, proper electrolyte balance is fundamental to all other physiological processes like neuronal signal transduction and muscle (including cardiac) contractility. Here, electrolytes will be measured using the VetScan VS2 critical care panel which includes ALT, BUN, Cl, CRE, GLU, K, Na, bicarbonate ion in a 100uL sample volume. In addition to assaying electrolytes themselves, I will conduct measures of urine electrolytes and specific gravity, as the urinary system represents that major pathway through which these chemicals are lost. These parameters will be measured using an Atago UG- α urine refractometer

and tests conducted at the IDEXX reference lab. Lastly, I will collect key metabolic parameters such as carbon dioxide production and oxygen consumption that may influence water consumption. These tests will be measured during a twenty four hour period at the end of the experimental manipulation, just prior to euthanasia, using a metabolic chamber (Sable Inc.) modified for use in the desert chamber. Together, these data will represent a uniquely rich characterization of the physiological state of a desert rodent held in captivity but importantly, exposed to conditions typical of the natural environment.

Aim 1a-1: Determine the physiologic response to water deprivation in desert adapted rodents.

Background: Water availability is thought to be one of the key attributes that determines the hostility of a given environment. For instance, ecosystems like deserts present unique challenges to the animals that inhabit them -- the ways in which animals overcome these challenges may allow researchers unique insights with biomedical importance.

Research Plan: To accomplish this aim, I will analyze physiological data from animals help with and without water, factorial with respect to the other conditions. Unique patterns present in (for instance)

Outcome: These data will be compiled into a large data matrix, with statistical analyses focusing on the identifying patterns related to differences in heat, water availability, or humidity. In addition, because behavior related to desert survival will be characterized as part of a separate project, changes in electrolytes will be linked to individual level differences in behavior. Ultimately, these physiology data will be combined with the genomic data to understand the relationship between physiology and genomics.

Preliminary data: I have characterized the electrolyte profile of 2 individuals housed at 70F, 50% RH, water *ad lib* and two individuals housed in identical conditions except that drinking water was withheld. Despite being housed in typical laboratory conditions, these animals have remarkably unusual electrolyte panel. For instance, mean serum potassium in an un-hemolyzed sample is unusually high at 8.1mg/dL, while Creatinine is low, with a mean measurement of 0.25mg/dL. mean blood urea nitrogen (BUN) is 47mg/dL. In contrast, animals without *ad lib* water were essentially...

Aim 1a-2: Define the relationship between temperature and physiological state.

Aim 1a-3: Define the relationship between relative humidity and physiological state.

- Describe role of serum electrolytes.
- How are these things regulated in the kidney (link to genes)
- typical dehydration panel
- What I'm going to measure
- expected outcome

Aim 1b: Define patterns of gene expression | isoform use | methylation given differences environmental condition. I will understand the genetic response to extreme heat and aridity via a series of bisulfite sequencing and RNAseq experiments, and will link these patterns to individual physiologic state as defined

in Aim 1. I hypothesize that genes responsible for water and solute transport will be particularly active in the most extreme conditions.

Background: The genomic processes related to desert survival have yet to be characterized. The few studies of genetics that have been done have focused on the role of single members of the Aquaporin gene family (but see Bartolo and Donald (2007)), which are large membrane-bound proteins that are critically involved in renal water transport (Kwon et al., 2009; Verkman, 2002; Brown et al., 1995; Nielsen and Agre, 1995). These studies have shown that changes in Aquaporin (AQP) protein abundance and expression may be related to water availability (Böselt et al., 2009; Gallardo et al., 2005; Bozinovic et al., 2003). In addition to changes in expression, another study showed that the AQP4 pathway was completely lost in the desert rodent *Dipodomys merriami merriami* (Huang et al., 2001). Despite these studies, we have a limited understanding of the genomics of renal water and solute regulation in desert animals. While AQPs are functionally important, water and solute balance is extraordinarily complex, and therefore single-gene studies are necessarily limited in their purview. A more complete understanding of this phenotype and its mechanistic underpinnings will require a sophisticated genome-level approach, which will be the outcome of the proposed research.

Research Plan: For analysis of the bisulfite sequence data, an accurate genome assembly is required. Using the existing draft genome (sequenced using startup funds) as a starting point, I will complete the genome assembly of the primary study organism, *Peromyscus eremicus*. This process will be significantly aided by leveraging the existing genome sequence data available from several other *Peromyscus* species. Specifically, an additional 60X coverage Illumina dataset using 150bp paired end and mate-pair sequencing will supplement the current 40X dataset. Simultaneous with this, I will sequence up to 10X coverage using long-read PacBio sequencing. A short-read assembly of Illumina data will be done using AllPaths-LG (Maccallum et al., 2009). Illumina contigs will be assembled with the corrected PacBio data using the software package SGA (Simpson and Durbin, 2012) or an overlap-layout-consensus assembler (e.g. Celera, Miller et al. (2008)) to generate a primary *de novo* genome assembly. Assemblies will be validated by use of the CEGMA (Parra et al., 2007) and REAPR (Hunt et al., 2013) algorithms. In general, this approach has been shown to be successful in producing high quality draft assemblies, though multiple assembly methods will be evaluated as per the findings of my previous work (Bradnam et al., 2013).

To annotate the genome, RNA and smRNA sequence data from several tissues, including liver, kidney, and brain will be generated. Sequence data will consist of Illumina strand-specific small-insert 150bp paired end sequencing. Each tissue type will be sequenced to a depth of 100 million reads. Assembly will be accomplished using the Trinity software package (Haas et al., 2013; Grabherr et al., 2011) after error correction (MacManes and Eisen, 2013) and quality trimming (MacManes, 2013). Annotation will be completed using the programs Trinotate (<http://trinotate.sourceforge.net/>) and Maker (Cantarel et al., 2008).

To identify patterns of differential isoform use, I will generate normalized cDNA libraries using X, and sequence using the single molecule Pacific Biosystems technology. This approach offers significant advantage over short read high-throughput sequencing technologies (e.g. Illumina, Ion Torrent) where accurate assembly of isoforms is extremely problematic (?).

In addition to the assembly and annotation of the *P. eremicus* genome, a secondary result of this

work is methods development (**enhancing infrastructure**). To this end, I have already already released a transcriptome assembly pipeline (<http://sourceforge.net/projects/tamrs/>) and automated quality control software (<http://sourceforge.net/projects/qcpro/>). In addition to this, I am an active developer of the transcriptome assembly program Trinity and annotation software Trinotate. Given the popularity of high-throughput sequencing, the demand for these types of tool development programs will likely increase.

Outcome: This work will allow me to look deeply into the genomic processes that underlie adaptation.

Preliminary Data: To date, I have generated a RNAseq dataset that consists of approximately 30M 150nt SE Illumina reads from the same 5 animals housed in the 'cold/simulated rain' treatment group from which I collected physiology data

- Describe typical renal gene expression profile
- link to electrolytes
- what I'm going to measure and how
- BiSeq, RNAseq, PacBio transcriptome to get at isoforms
- expected outcome

Aim 1c: Correlate physiology (=phenotype) with genomic patterns.

- Diff gene expression
- coexpression networks
- Patterns of isoform use
- diff methylation

Aim 2: Given the transition from the obligate intake of fluids as infants, to it's complete absence later in life, the ontogeny of physiologic water conservation will be elucidated.

ii. Background

The study of adaptation, or the process through which animals become fitted to their environment has intrigued researchers for decades (Darwin, 1859; Fisher, 1930), though only recently have we had the ability to study the underlying genomic mechanisms. Interestingly, researchers interested in understanding the genetics of adaptation have the ability to ground modern studies of genetics on decades of work aimed at understanding the ecological context within which adaptation occurs. One particularly salient example of the connection between studies of ecology and natural history and modern genomics can be found in the study of physiologic adaptation to desert conditions. Here, remarkable physiologic, morphologic (Dickinson et al., 2007; Huntley et al., 1984; Schmidt-Nielsen and Schmidt-Nielsen, 1950; Schmidt-Nielsen, 1952) and behavioral (Nagy, 1994) adaptation has been studied in the context of desert ecology. These studies provide a rich context for the current work, which aims to define the relationship between physiology and genomics in rodents able to thrive in amongst the most harsh of conditions on Earth.

Though classic research relating morphology (especially renal ultrastructure) and physiology to desert adaptation has been done, we know virtually nothing about how extreme heat and aridity may affect other core physiological and metabolic processes. For instance, the maintenance of normal serum electrolyte concentration is challenging in the context of desert conditions. Indeed, electrolyte derangement is often the ultimate cause dehydration-related death, yet for the vast majority of desert animals, we know nothing of electrolyte balance - not even typical values! Understanding these critical processes is fundamental to our understanding of physiological adaptation, and will be accomplished as part of this project.

Lastly,

Specific Aim 1:

Specific Aim 2:

Preliminary Data: .

Specific Aim 3: Given that desert adapted mice, capable of surviving without water are as neonates dependent on liquid intake, the study of the ontogeny of physiologic water conservation is extremely interesting and relevant to the current work. This phenomenon will be explored using neonate mice in the treatments listed in Aim 1, and methods described in Aim 2. Five neonate mice will be culled per treatment at 3 different timepoints (immediately after birth, mid-lactation, 1 day after weaning). I hypothesize that patterns of gene expression and methylation will resemble those common in conditions where water is available *ab lib*.

iv. Broad Impacts

How desert animals survive without water is an intrinsically interesting example of extreme physiology and adaptation. Indeed, telling people of the remarkable story of a rodent that may never drink water is immediately captivating. Having told hundreds of people, including children, adults, scientifically literate and not, the most common response is something like "Wow! That is amazing. How do they do that?" I intend to **broaden dissemination** by capitalizing on this common response, using traditional and non-traditional methods. While lecture has been the mainstay of scientific communication, its audience tends towards the educated/affluent subset of the population. Given one of my primary goals as a scientists is to broadly share my work, I will use non-traditional modes of communication which will include social media and a blog. All publications are hosted on a public preprint server, and published using open access options. All data and code is freely available using public repositories like Github, Dryad, or Figshare. In

addition to this, I will develop an outreach program that aims to present research findings to school-aged children. These program will be multi-dimensional, with programs geared towards several different age groups.

Broadening participation is an issue about which I care deeply. As a scientist of Native American descent, I strive to be a role model and mentor for underrepresented groups. To this end, I am developing an internship program- in collaboration with the American Indian Higher Education Consortium (<http://aihec.org/>), where qualified Native students may spend a summer working in my lab either in directed or independent research programs. It is my hope that students introduced to my lab via internship may ultimately elect to join the lab. This method of recruitment supplements my general interest in recruiting students from underrepresented groups.

Lastly, my efforts aimed at broadening dissemination and participation are all within the context of a research program with substantial **benefits to society**. First, Earth is becoming both warmer and drier, and accurate prediction of animals' response to a changing climate requires knowledge of how animals currently living in dry/hot climates survive - this project will deepen the requisite knowledge. Next, kidney disease effects millions of Americans (http://www.cdc.gov/diabetes/projects/pdfs/ckd_factsheet.pdf). Though the causes are diverse, the pathophysiology of many resemble dehydration (e.g. low renal perfusion pressure). Thus, having a better understanding of how desert rodents endure water stress seemingly without complication may enhance our ability to explain why humans and most other mammals cannot.

v. Summary & Future Directions

The maintenance of water balance in animals is one of the most important physiologic processes, and is critical to survival. Osmoregulation in animals living in desert environments is particularly challenging, as extreme heat and aridity is common. Gaining a deeper understanding of the physiological adaptations that allow for desert survival is important and has obvious implications for climate change science, conservation, and human health and medicine. The proposed research aims to generate a uniquely rich dataset, leveraging cutting edge genomic techniques against careful characterization of physiology, all within an ecological context of desert life. A carefully constructed plan for broadening dissemination and participation ensures that the scientific process as well as its results will be available to all interested parties.

This proposal represents the foundational steps toward developing *P. eremicus* as a model system for the study of desert adaptation. Indeed, this model offers the scientific community a unique opportunity to gain a deep understanding into the physiology and genomics of desert adaptation- an insight which is impossible to achieve using traditional model system. While not a part of this proposal, this work lays the groundwork for future studies where causal links are made between phenotype and genotype, using technologies like CRISPR-Cas9 and RNAi.

References

- Bartolo, R.C., Donald, J.A., 2007. The distribution of renal hyaluronan and the expression of hyaluronan synthases during water deprivation in the Spinifex hopping mouse, *Notomys alexis*. *Comparative biochemistry and physiology Part A, Molecular & integrative physiology* 148, 853--860.
- Böselt, I., Römpler, H., Hermsdorf, T., Thor, D., Busch, W., Schulz, A., Schöneberg, T., 2009. Involvement of the V2 vasopressin receptor in adaptation to limited water supply. *PLOS ONE* 4, e5573.
- Bozinovic, F., Gallardo, P.A., Visser, G.H., Cortés, A., 2003. Seasonal acclimatization in water flux rate, urine osmolality and kidney water channels in free-living degus: molecular mechanisms, physiological processes and ecological implications. *Journal of Experimental Biology* 206, 2959--2966.
- Bradnam, K.R., Fass, J.N., Alexandrov, A., Baranay, P., Bechner, M., Birol, I., Boisvert, S., Chapman, J.A., Chapuis, G., Chikhi, R., Chitsaz, H., Chou, W.C., Corbeil, J., Del Fabbro, C., Docking, T.R., Durbin, R., Earl, D., Emrich, S., Fedotov, P., Fonseca, N.A., Ganapathy, G., Gibbs, R.A., Gnerre, S., Godzaridis, E., Goldstein, S., Haimel, M., Hall, G., Haussler, D., Hiatt, J.B., Ho, I.Y., Howard, J., Hunt, M., Jackman, S.D., Jaffe, D.B., Jarvis, E., Jiang, H., Kazakov, S., Kersey, P.J., Kitzman, J.O., Knight, J.R., Koren, S., Lam, T.W., Lavenier, D., Laviolette, F., Li, Y., Li, Z., Liu, B., Liu, Y., Luo, R., Maccallum, I., **MacManes**, M.D., Maillet, N., Melnikov, S., Naquin, D., Ning, Z., Otto, T.D., Paten, B., Paulo, O.S., Phillippy, A.M., Pina-Martins, F., Place, M., Przybylski, D., Qin, X., Qu, C., Ribeiro, F.J., Richards, S., Rokhsar, D.S., Ruby, J.G., Scalabrin, S., Schatz, M.C., Schwartz, D.C., Sergushichev, A., Sharpe, T., Shaw, T.I., Shendure, J., Shi, Y., Simpson, J.T., Song, H., Tsarev, F., Vezzi, F., Vicedomini, R., Vieira, B.M., Wang, J., Worley, K.C., Yin, S., Yiu, S.M., Yuan, J., Zhang, G., Zhang, H., Zhou, S., Korf, I.F., 2013. Assemblathon 2: evaluating *de novo* methods of genome assembly in three vertebrate species. *GigaScience* 2, 10.
- Brown, D., Katsura, T., Kawashima, M., Verkman, A.S., Sabolic, I., 1995. Cellular distribution of the aquaporins: A family of water channel proteins. *Histochemistry and Cell Biology* 104, 1--9.
- Cantarel, B.L., Korf, I., Robb, S.M.C., Parra, G., Ross, E., Moore, B., Holt, C., Sánchez Alvarado, A., Yandell, M., 2008. MAKER: an easy-to-use annotation pipeline designed for emerging model organism genomes. *Genome Research* 18, 188--196.
- Darwin, C., 1859. *On the Origin of Species by Means of Natural Selection or the Preservation of Favored Races in the Struggle for Life*. 1st ed., J. Murray, London.
- Dickinson, H., Moritz, K., Wintour, E.M., Walker, D.W., Kett, M.M., 2007. A comparative study of renal function in the desert-adapted spiny mouse and the laboratory-adapted C57BL/6 mouse: response to dietary salt load. *American Journal of Physiology-Renal Physiology* 293, F1093--8.
- Fisher, R., 1930. *The Genetical Theory of Natural Selection*. Oxford University, London.
- Gallardo, P.A., Cortés, A., Bozinovic, F., 2005. Phenotypic flexibility at the molecular and organismal level allows desert-dwelling rodents to cope with seasonal water availability. *Physiological and Biochemical Zoology* 78, 145--152.
- Grabherr, M.G., Haas, B.J., Yassour, M., Levin, J.Z., Thompson, D.A., Amit, I., Adiconis, X., Fan, L., Raychowdhury, R., Zeng, Q., Chen, Z., Mauceli, E., Hacohen, N., Gnirke, a., Rhind, N., di Palma, F., Birren, B.W., Nusbaum, C., Lindblad-Toh, K., Friedman, N., Regev, A., 2011. Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nature Biotechnology* 29, 644--652.

- Haas, B.J., Papanicolaou, A., Yassour, M., Grabherr, M., Blood, P.D., Bowden, J., Couger, M.B., Eccles, D., Li, B., Lieber, M., **MacManes**, M.D., Ott, M., Orvis, J., Pochet, N., Strozzi, F., Weeks, N., Westerman, R., William, T., Dewey, C.N., Henschel, R., Leduc, R.D., Friedman, N., Regev, A., 2013. *De novo* transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. *Nature protocols* 8, 1494--1512.
- Huang, Y., Tracy, R., Walsberg, G.E., Makkinje, A., Fang, P., Brown, D., Van Hoek, A.N., 2001. Absence of aquaporin-4 water channels from kidneys of the desert rodent *Dipodomys merriami merriami*. *American Journal of Physiology-Renal Physiology* 280, F794--F802.
- Hunt, M., Kikuchi, T., Sanders, M., Newbold, C., Berriman, M., Otto, T.D., 2013. REAPR: a universal tool for genome assembly evaluation. *Genome Biology* 14, R47.
- Huntley, A.C., Costa, D.P., Rubin, R.D., 1984. The contribution of nasal countercurrent heat exchange to water balance in the northern elephant seal, *Mirounga angustirostris*. *The Journal of experimental biology* 113, 447--454.
- Kwon, T.H., Nielsen, J., Nielsen, J., Møller, H.B., Møller, H.B., Fenton, R.A., Fenton, R.A., Nielsen, S., Frøkiaer, J., 2009. Aquaporins in the Kidney, in: *Aquaporins*. Springer Berlin Heidelberg, Berlin, Heidelberg, pp. 95--132.
- Maccallum, I., Przybylski, D., Gnerre, S., Burton, J., Shlyakhter, I., Gnirke, a., Malek, J., McKernan, K., Ranade, S., Shea, T.P., Williams, L., Young, S., Nusbaum, C., Jaffe, D.B., 2009. ALLPATHS 2: small genomes assembled accurately and with high continuity from short paired reads. *Genome Biology* 10, R103.
- MacManes, M.D., 2013. On the optimal trimming of high-throughput mRNA sequence data. Technical Report.
- MacManes, M.D., Eisen, M.B., 2013. Improving transcriptome assembly through error correction of high-throughput sequence reads. *PeerJ* 1, e113.
- Miller, J.R., Delcher, A.L., Koren, S., Venter, E., Walenz, B.P., Brownley, A., Johnson, J., Li, K., Mobarry, C., Sutton, G., 2008. Aggressive assembly of pyrosequencing reads with mates. *Bioinformatics (Oxford, England)* 24, 2818--2824.
- Nagy, K.A., 1994. Seasonal Water, Energy and Food Use by Free-Living, Arid-Habitat Mammals. *Australian Journal of Zoology* 42, 55--63.
- Nielsen, S., Agre, P., 1995. The aquaporin family of water channels in kidney. *Kidney international* .
- Parra, G., Bradnam, K., Korf, I., 2007. CEGMA: a pipeline to accurately annotate core genes in eukaryotic genomes. *Bioinformatics (Oxford, England)* 23, 1061--1067.
- Schmidt-Nielsen, B., 1952. Renal Tubular Excretion of Urea in Kangaroo Rats. *American Journal of Physiology* 170, 45--56.
- Schmidt-Nielsen, B., Schmidt-Nielsen, K., 1950. Pulmonary Water Loss in Desert Rodents. *American Journal of Physiology* 162, 31--36.
- Simpson, J.T., Durbin, R., 2012. Efficient *de novo* assembly of large genomes using compressed data structures. *Genome Research* 22, 549--556.
- Verkman, A.S., 2002. Physiological importance of aquaporin water channels. *Annals of medicine* 34, 192--200.