

The maintenance of water balance is critical for survival. Humans are exquisitely sensitive to changes in osmolality, with slight derangement eliciting physiologic compromise. When the loss of water exceeds dietary intake, dehydration - and in extreme cases, death - can occur. Far from uncommon, millions of people die every year as a direct result of dehydration. In contrast to humans, animals living in desert habitats thrive without water and endure extreme heat and intense drought, as a direct result of unique adaptations. These adaptation all them to survive conditions fatal to humans and most other animals. Despite being a well-known ecological phenomenon with obvious implications for human health, we know very little of the underlying mechanisms that allow for survival in desert environments. **The proposed research uses a novel approach integrating physiology, evolutionary genomics, and computational biology to better understand how animals survive in what appear to be non-survivable conditions.** This proposal represents the foundational steps toward developing the cactus mouse (*Peromyscus eremicus*) as a model system for the study of physiologic water conservation. Indeed, this model offers the scientific community a unique opportunity to gain a deep understanding into the physiology and genomics of osmoregulation in extreme environments – a critically important insight that is impossible using traditional model system like *Mus*, that like humans, die with subjected to these conditions. While not a part of this proposal, this project lays the groundwork for my long-term research goal – to identify the causal links between phenotype and genotype, using emerging technologies like the CRISPR-Cas9 system. Ultimately, understanding the mechanisms underlying extreme osmoregulation may suggest novel treatment strategies for conditions (e.g. diarrhea) resulting in acute dehydration in humans.

**SPECIFIC AIM 1:** To characterize the the physiology and adaptive response (differential gene expression, patterns of methylation or isoform use) in desert-adapted mice in response to extreme heat and aridity.

The working hypothesis is that while desert-adapted mice may demonstrate genome wide expression patterns suggestive of stress (e.g. heat shock proteins) during dehydration, these responses function to preserve normal physiology and thus serum electrolytes will be similar to mice with unrestricted access to water.

**SPECIFIC AIM 2:** To determine the ontogeny of extreme osmoregulatory ability, from the neonatal period during which fluid (milk) intake is obligate through weaning, when oral fluid intake is exceptionally rare.

I hypothesize that patterns of renal gene expression during fetal development through weaning will resemble patterns of gene expression, isoform use, and methylation typical of adult mice when water is freely available.

The proposed project aims to integrate studies of physiology, genomics, and computational biology to gain a deep understanding of a fundamental physiological problem – how to conserve water when intake is limited. Although dehydration is both common and dangerous, a large swath of the biology underlying its physiological effects is currently invisible to researchers using traditional mammalian models of disease that lack the eco-evolutionary history present in desert-adapted mice. This project will fill a critically important gap in our understanding, which is in support of the specific research aims of the National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK).

### i. Significance

Dehydration, whether caused by exposure to extreme environmental conditions, water deprivation, or by infection (e.g. diarrheal illnesses) represents a significant threat to human life. In spite of modern medicine, millions of people die every year from dehydration. Compounding issues of exposure and illness, are public health issues regarding the delivery of safe drinking water. With global climate change, these challenges are thought to become only more severe and as a result, research providing insight into the mechanisms underlying physiologic resistance to acute dehydration is urgently needed. The response to acute dehydration in humans and traditional mammalian models is generally maladaptive and may include death - this response limits our ability to develop novel insights into this important cause of human mortality. As such, the study of dehydration-tolerant mammalian models will significantly enhance our understanding, and will provide fodder for novel treatments. **The proposed work aims study extreme osmoregulation in a uniquely suited novel desert-adapted model organism.**

While the mechanisms underlying physiological compromise in dehydration are well characterized (Roberts, Pope, Newson, Lolait, O'Carroll, 2010), some animals possess the ability, much unlike humans, to osmoregulate despite extreme heat and a complete lack of extrinsic water intake (Nagy Gruchacz, 1994). Specifically, highly adapted desert mice may never drink water, produce an extremely viscous urine, or no urine at all, and excrete urea in the form of uric acid crystals in the feces (K. Schmidt-Nielsen Schmidt-Nielsen, 1952). This phenotype results in an animal that is very resistant to dehydration-related physiologic compromise, and is in stark contrast to the phenotype of humans and traditional model organisms (e.g. *Mus* and *Rattus*). Although model organisms are attractive targets for study, they lack the requisite biology which may limit insight. In contrast with traditional model organisms, non-model desert-adapted organisms may provide a unique opportunity to study dehydration tolerance, though they typically lack many of the genomic and physiologic tools characteristic of model organisms. Despite this, renal gene expression has been characterized for several genes in desert animals, and was shown to be highly derived in some (e.g. *Dipodomys* (Huang et al., 2001)), but not in others (e.g. *Notomys* (Weaver, Walker, Alcorn, Skinner, 1994)). No studies characterizing genome-wide patterns of gene expression, methylation or isoform in desert-adapted water stressed animals has been done and therefore the extent to which differences in these parameters underlie phenotype remains unknown. The proposed work effectively integrates the power of a model organism with the unique biology of a desert-adapted rodent, the cactus mouse (*Peromyscus eremicus*), to generate insights into extreme osmoregulation not current possible.


### ii. Innovation

The proposed work recognizes that successful treatment requires an appropriate model, and while traditional models are powerful, they lack the biology (extreme osmoregulation) upon which more successful interventions may be modeled. The desert-adapted rodent *P. eremicus* retains many of the beneficial characteristics of model organisms, while enhancing opportunity to assay interesting biological phenomenon. In addition to this fundamental innovation, the project is innovative in a number of other ways.

- Experimental, conceptual, theoretical, and technical innovation: The proposed project leverages unprecedented control over environmental conditions using an ideally suited novel model organism and unique analytical methods to understand basic physiology in dehydration resistant organisms.

**Aim 1:** To characterize the the physiology and adaptive response (differential gene expression, patterns of methylation or isoform use) in desert-adapted mice in response to extreme heat and aridity.

To better understand the physiological effects of desert conditions on rodents, I will relate multiple physiological variables to differences in temperature, relative humidity, and water availability. These experiments are fundamentally linked to a series of environmental manipulations, described in Figure 1. The experimental design is fully factorial -- meaning that the focal experimental parameter (e.g. water availability) will be tested in the context of the full range of other conditions (e.g. humidity, temperature). Animal care is standardized between experiments and includes measures to reduce the water content of food and bedding materials. Both of these will be dried in a standard desiccation oven to less than 1% water/volume. Twenty individuals per treatment will be included -- power analyses suggest this sample size will allow me to garner statistical support for patterns with small to medium effect sizes. Together, this design will allow me to tease-apart the physiologic and genomic response to the various conditions.



For each experiment in Aim 1, I will collect information including metabolism, urine concentration and electrolyte content, serum electrolyte content, as well as more basic measures like body temperature and body weight. In the context of limited water intake, how animals achieve electrolyte balance is unknown. Electrolytes are both easy to assay and are critical to physiological well being. Indeed, proper electrolyte balance is fundamental to all other physiological processes like neuronal signal transduction and muscle (including cardiac) contractility. Here, electrolytes will be measured using the VetScan VS2 critical care panel which includes ALT, BUN, Cl, CRE, GLU, K, Na, bicarbonate ion in a 100uL sample volume.

In addition to assaying electrolytes themselves, I will conduct measures of urine electrolytes and specific gravity, as the urinary system represents that major pathway through which these chemicals are lost. These parameters will be measured using an Atago UG- $\alpha$  urine refractometer and tests conducted at the IDEXX reference lab. Lastly, I will weigh animals to the nearest 0.1gm every other day, including the

day of sacrifice. Body temperature will be assayed with weighing using a digital thermometer and probe designed by World Precision Instruments (Sarasota, FL). In connection with this, feces will be collected and water content will be assayed using standard methods.

I will collect key metabolic parameters such as carbon dioxide production and oxygen consumption that may influence water consumption. In addition, I will measure the change in relative humidity within the metabolic chamber, which will allow me to understand the rate of pulmonary water loss (or gain). These tests will be measured during a twenty four hour period at the end of the experimental manipulation, just prior to euthanasia, using a metabolic chamber (Sable Inc.) modified for use in the desert chamber. Together, these data will represent a uniquely rich characterization of the physiological state of a desert rodent held in captivity but importantly, exposed to conditions typical of the natural environment.

**Aim 1a:** Determine the physiologic response to drinking-water deprivation, extreme temperature, and humidity in the desert-adapted rodent *P. eremicus*. I hypothesize that, as a result of unique mechanisms related to solute and water balance, that average serum electrolyte concentrations will remain relatively constant throughout various experimental manipulations, but the variance in measured levels between individuals will increase in the most extreme conditions. These differences will be echoed in differences in urine electrolytes and concentration. Predictions regarding other parameters are detailed in Table 1.

Background: The human body consist of 60% water (Jéquier Constant, 2009). Far from a static reservoir, proper physiologic function requires water for countless processes including nutrient transport (Haussinger, 1996), signal transduction, pH balance, thermal regulation (Montain, Latzka, Sawka, 1999) and the removal of metabolic waste. To accomplish these things, approximately 2 liters of fluid are used daily - these fluids are lost mainly via the GI and GI system, and by evaporative loss, which is accelerated greatly in extremes of heat and aridity (Cheuvront, Kenefick, Montain, Sawka, 2010). These losses must be matched by intake (Jéquier Constant, 2009), mainly in the form of oral fluid intake. Though the body possesses limited reserves, when loss exceeds intake over even a period of time, dehydration and in extreme cases, death can occur. Humans and most other animals are exquisitely sensitive to dehydration, and possess limited compensatory mechanisms. In contrast, desert rodents survive in extreme environmental conditions, often without fluid intake. Understanding the mechanisms underlying this remarkable phenotype requires we understand the physiology that accompanies it. The work described here aims to characterize the physiology of dehydration resistance in desert adapted rodents.

While the prolonged absence of drinking water is invariably fatal for humans and many other animals, one potentially mitigating effect may be the acquisition of water (or limitation of loss) via the pulmonary vasculature, which is known to be variably permeable to water (Berger et al., 2011; Goralski, Boucher, Button, 2010). While pulmonary water acquisition has not been quantified in humans or in mammalian models, the pulmonary vasculature is ideally positioned to retain water from inspired air. Following this, relative humidity - the amount of extractable water present in respired air may be important to overall hydration status. The design described above incorporates two different levels of humidity to begin to disentangle the effects of drinking water from water acquisition via the pulmonary system.

Although water stress is obviously important to the survival of desert rodents - a phenotype which

is relevant to human health and wellness, extreme temperatures represent another way in which physiological processes may be challenged. While desert animals may thrive in extreme heat, humans cannot. The physiological response is characterized in model organisms, but not in other animals adapted to these conditions. Genes like the heat-shock proteins are protective in humans, but no record of their activity on desert rodents is known.

**Research Plan:** To accomplish this aim, I will analyze physiologic data from animals held with and without drinking water, factorial with respect to the other conditions (e.g. temperature and humidity). The specific experiments described in Figure 1 will allow me to tease apart the effects of water deprivation from other parameters. Though the data I propose to collect is described above, in brief, I plan to collect blood and urine electrolytes and urine specific gravity. I will collect data on fecal water content, animal weight and temperature, as well as a battery of metabolic parameters. The specific predictions regarding several of these parameters are described in Table 1.

The statistical treatment of the data will include a multivariate regression (either linear or non-linear) to establish the relationships between the data. Many of these analyses will be conducted with non-parametric tests, as data are often non-normally distributed nor independent. One of the most interesting comparisons will be to understand the relationship between serum sodium and urine sodium, urine concentration, fecal water content, and changes in body weight. Ultimately (e.g. Aim 1b) I will link these data with patterns of gene expression, methylation, and isoform use to gain a synthetic understanding of dehydration resistance.

**Preliminary data:** I have characterized the electrolyte profile of 2 individuals housed at 70F, 50% RH, water *ad lib* and two individuals housed in identical conditions except that drinking water was withheld. Despite being housed in typical laboratory conditions, these animals have remarkably unusual electrolyte panel. For instance, mean serum potassium in an un-hemolyzed sample is unusually high at 8.1mg/dL, while Creatinine is low, with a mean measurement of 0.25mg/dL. mean blood urea nitrogen (BUN) is 47mg/dL. In contrast, animals without *ad lib* water were essentially...

		Serum E-lytes	Urine Conc.	Fecal Water	Weight
Hot/Dry Hot/Wet		Mean Na, Cl, K ↑ to =, Var ↑↑	↑↑↑↑↑	↓↓↓↓↓	↓↓↓↓↓
	Rain	Mean Na, Cl, K ↑ to =	↑↑	↓↓↓	↓↓↓
	Humidity	Mean Na, Cl, K ↑ to =	↑↑↑	↓↓↓	↓↓↓
Cold/Dry Cold		Mean Na, Cl, K ↑ to =, Var ↑↑	↑↑↑↑↑	↓↓↓↓↓	↓↓↓↓↓
	Rain	Mean Na, Cl, K ↑ to =	↑	↓	↓
	Humidity	Mean Na, Cl, K ↑ to =	↑↑	↓↓	↓↓

Table 1: Say something about predictions here.

**Aim 1b:** Define patterns of gene expression | isoform use | methylation given differences environmental condition. I will understand the genetic response to extreme heat and aridity via a series of bisulfite and RNA and PacBio sequencing experiments, and will link these patterns to individual physiologic state as defined in Aim 1. I hypothesize that genes responsible for water and solute transport will be particularly active in the most extreme conditions in renal and pulmonary tissues, while genes involved in the activation of the hypothalamic-neurohypophyseal system will be upregulated in the hypothalamus.

**Background:** Broadly speaking, genes underlie the vast majority of observable phenotypes. Whether this relationship is mediated by patterns of expression (e.g. (Teets et al., 2012)), which itself may be mediated by differences in methylation (Brenet et al., 2011), or by use alternative splice isoforms (Yukutake Yasui, 2010), linking genotype to phenotype is extremely difficult. In addition to these mechanisms, function (=phenotype) may be determined by post-translational modifications like phosphorylation of specific sites (Moeller, MacAulay, Knepper, Fenton, 2009). The identification of these mechanisms is important, not only because in doing so we gain a deeper understanding of evolution, but also because these molecular mechanisms may be later used as targets for drug development or other therapeutic intervention. With regards to resistance to dehydration, the development of novel therapies is critical, as millions of people die yearly as a consequence.

In model organisms, dehydration precipitates a physiological response that is largely driven by the neuroendocrine system. Very much simplified, the cascade begins with the stimulation of osmoreceptors (Arsenijevic Baertschi, 1985), which in turn stimulates neurons located in the paraventricular and supraoptic nuclei of the hypothalamus to release ADH (Zingg, Lefebvre, Almazan, 1986). ADH then binds to vasopressin-responsive receptors located in the renal medulla, resulting in Aquaporin movement to the surface of the collecting duct (Nielsen et al., 1995) which encourages water re-uptake. In addition to the aquaporins, the renin-angiotensin-aldosterone system (Gubler Antignac, 2009), natriuretic peptides (Totsune et al., 1994), the SLC and mTOR families (Ortells et al., 2012), and potentially other yet to be discovered pathways are important to water balance. Far from canonical, each stage in these cascades is dynamic and therefore pathways revealed in *Mus* and humans may not be equivalent to pathways in uniquely adapted desert animals, particularly given radically different phenotype.

The genomic processes related to desert survival have yet to be characterized. The few studies of genetics that have been done have focused on the role of expression of single members of the Aquaporin gene family (but see (Bartolo Donald, 2007)), which are large membrane-bound proteins that are critically involved in renal water transport (D. Brown, Katsura, Kawashima, Verkman, Sabolic, 1995; Kwon et al., 2009; Nielsen Agre, 1995; Verkman, 2002). These studies have shown that changes in Aquaporin (AQP) protein abundance and expression may be related to water availability (Bozinovic, Gallardo, Visser, Cortés, 2003; Bösel et al., 2009; Gallardo, Cortés, Bozinovic, 2005). In addition to changes in expression, another study showed that the AQP4 pathway was completely lost in the desert rodent *Dipodomys merriami merriami* (Huang et al., 2001). Despite these studies, we have a limited understanding of the genomics of renal water and solute regulation in desert animals. While AQPs are functionally important, water and solute balance is extraordinarily complex, and therefore single-gene studies are necessarily limited in their purview. A more complete understanding of this phenotype and its mechanistic underpinnings will require a sophisticated genome-level approach, which will be the outcome of the proposed research. In contrast to the limited amount known about patterns of renal gene expression, we know much less about gene expression in other tissues, and absolutely nothing about differential methylation or isoform use, even though we know that these complexities are functionally important to this specific function (Silberstein et al., 2004; Yukutake Yasui, 2010).

**Research Plan:** The analysis of the genome wide patterns of response to dehydration will be con-

ducted using the same individuals for which I collected physiology data. To accomplish this goal, I will map RNAseq reads derived from kidney, lung, and hypothalamus to the existing annotated draft genome, which was sequenced using startup funds. This phase of the project will be accomplished using the short read aligned BWA (Li, 2013) and best practices previously established (MacManes, 2014). Differential expression will be evaluated via the Cufflinks package (Trapnell et al., 2012), while evidence for coordinated changes in large numbers of genes will be done using the software wgcna (Langfelder Horvath, 2008).

Accurate isoform reconstruction is notoriously difficult using high-throughout short read data data like Illumina sequencing (Hiller, Jiang, Xu, Wong, 2009; Pyrkosz, Cheng, Brown, 2013), despite the advent of longer read lengths and newer analytical techniques (Jiang Wong, 2009; LeGault Dewey, 2013). In projects like this, where differential isoform use may be critical to phenotype, a different approach may be warranted. For instance, the sequencing technology available from Pacific Biosystems (PacBio) is suggested to provide a resolution to the isoform reconstruction problems (Au et al., 2013), specifically because because it involves a long-read single molecule sequencing strategy (Eid et al., 2009). To identify patterns of differential isoform use, I will sequence poly-A selected mRNA samples using PacBio technology. Because throughput is relatively low, which may limit the precision with which quantitation can be achieved, I will explore ways to accurately estimate isoform specific expression. One previously unexplored approach involves estimating expression in the program eXpress using only those reads that map uniquely and unambiguously to a specific isoform. Because this approach is uncharacterized, I will also develop a set of isoform specific PCR primers that will allow me to estimate expression using qPCR.

Lastly, aside from differences in expression of isoform use, patterns of methylation could be important in the development of extreme osmoregulation - indeed, methylation has been shown to be important to many other complex phenotypes including behavior (Keller, 2010) and metabolism (Foret et al., 2012). To understand patterns of methylation, I will generate a large bisulfite sequence dataset, which will contain information from ever individual included in the mRNAseq experiments, described above. The

To annotate the genome, RNA and smRNA sequence data from several tissues, including liver, kidney, and brain will be generated. Sequence data will consist of Illumina strand-specific small-insert 150bp paired end sequencing. Each tissue type will be sequenced to a depth of 100 million reads. Assembly will be accomplished using the Trinity software package (Grabherr et al., 2011; Haas et al., 2013) after error correction (MacManes Eisen, 2013) and quality trimming (MacManes, 2013). Annotation will be completed using the programs Trinotate (<http://trinotate.sourceforge.net/>) and Maker (Cantarel et al., 2008).

To identify patterns of differential isoform use, I will generate normalized cDNA libraries using X, and sequence using the single molecule Pacific Biosystems technology. This approach offers significant advantage over short read high-throughput sequencing technologies (e.g. Illumina, Ion Torrent) where accurate assembly of isoforms is extremely problematic (Pyrkosz et al., 2013).

In addition to the assembly and annotation of the *P. eremicus* genome, a secondary result of this work is methods development (**enhancing infrastructure**). To this end, I have already already released a transcriptome assembly pipeline (<http://sourceforge.net/projects/tamrs/>) and automated quality control software (<http://sourceforge.net/projects/qcpro/>). In addition to this, I am an active developer of the transcriptome assembly program Trinity and annotation software Trinotate. Given the popularity

of high-throughput sequencing, the demand for these types of tool development programs will likely increase.

**Preliminary Data:** To date, I have generated a RNAseq dataset that consists of approximately 30M 150nt SE Illumina reads from the same 5 animals housed in the 'cold/simulated rain' treatment group from which I collected physiology data. I have DDDDDDDDDDDDD

**Expected Outcome:** Upon completion of Aim 1, I will have a synthetic understanding of the physiologic and genomics patterns associated with extreme osmoregulation. These data will allow me to generate a list of genes and genomic regions putatively linked to the phenotype of interest. This list is critical, and will form the basis for my first R01 submission, which will propose the development of a system where manipulation of specific genes is possible (e.g. the CRISPR/CAS9 transgenic system), thus moving the work from correlation to causation. This grant will be developed and submitted during the second year of the COBRE tenure. In addition this, the completion of Aim 1 will allow me to become more proficient in the collection and bioinformatic analysis of physiology data. Lastly, part of Aim1B involved the development of a novel pipeline for the identification of differential isoform use using PacBio RNA sequence data. This skill will be useful to my broader scientific goals, as well as to the broader scientific community.

Regarding dissemination, the work will be published in open access journals, after rapid release using preprint servers. I envision several papers that are a direct result of this work, include papers describing the physiological and metabolic response to water deprivation as well as their genomic responses. In addition, I aim to publish a more methods-oriented paper surrounding the study of isoform using PacBio data. Aside from peer-reviewed publication, results will be disseminated via social media, my blog, and at the annual meeting of the Society for the Study of Evolution.

**Aim 2: Given the transition from the obligate intake of fluids as infants, to it's complete absence later in life, the ontogeny of physiologic water conservation will be elucidated.**

**Background:** Given that desert adapted mice, capable of surviving without water are as neonates dependent on liquid intake, the study of the ontogeny of physiologic water conservation is extremely interesting and relevant to the current work.

**Research Plan:** This phenomenon will be explored using fetal and neonatal mice whose mothers are exposed to treatments and an abbreviated set of methods listed in Aim 1. Many of the physiological measurements (e.g. blood and urine analyses) will be impossible to collect in very young animals secondary to sample volume requirements, though a full battery of genomic tests should be possible. To evaluate the ontogeny, five fetal and neonatal mice will be culled per treatment at four different timepoints (immediately prior to birth, 2 hours after birth, mid-lactation (approximately 10 days after birth), 1 day after weaning). I hypothesize that patterns of gene expression, methylation, and isoform use will resemble those common in conditions where water is available *ab lib*.

**Expected Outcome:** Upon completion of Aim 2, I will have a synthetic understanding of the genomics patterns associated with the ontogeny of extreme osmoregulation. These data will allow me to generate a list of genes and genomic regions putatively linked to the phenotype of interest. This list is critical, and



will form the basis for my first R01 submission, which will propose the development of a system where manipulation of specific genes is possible (e.g. the CRISPR/CAS9 transgenic system), thus moving the work from correlation to causation. This grant will be developed and submitted during the second year of the COBRE tenure. In addition this, the completion of Aim 1 will allow me to become more proficient in the collection and bioinformatic analysis of physiology data.

Regarding dissemination, the work will be published in open access journals, after rapid release using preprint servers. I envision several papers that are a direct result of this work, include papers describing the physiological and metabolic response to water deprivation. In addition to peer-reviewed publication, results will be disseminated via social media, my blog, and at the annual meeting of the Society for the Study of Evolution.

Activity	FY2015	FY2016	FY2017
Recruit PDF, grad students, undergraduates	X		
Increase Colony Size & ID animals for experiments	X		
Conduct Physiology Experiments -- AIM 1A	X		
Collect & Analyze expression data -- AIM 1B	X		
Analyze Bisulfite and PacBio data -- AIM 1B		X	
Animal breeding in prep for Aim 2		X	
Collect & Analyze genomic data -- AIM 2		X	X
Write papers & submit		X	X
Present results at international conference		X	X
Prepare R01 & and resubmit as needed		X	X
Train Undergrad, Grad students, & PDF	X	X	X
Disseminate info	X	X	X

Arsenijevic, Y., Baertschi, A. J. (1985). Activation of the hypothalamo-neurohypophyseal system by hypertonic superfusion of the rat mesentery. *Brain Research*, 347(1), 169–172. doi:10.1016/0006-8993(85)90907-2

Au, K. F., Sebastiano, V., Afshar, P. T., Durruthy, J. D., Lee, L., Williams, B. A., et al. (2013). Characterization of the human ESC transcriptome by hybrid sequencing. *Pnas*, 201320101. doi:10.1073/pnas.1320101110

Bartolo, R. C., Donald, J. A. (2007). The distribution of renal hyaluronan and the expression of hyaluronan synthases during water deprivation in the *Spinifex* hopping mouse, *Notomys alexis*. *Comparative Biochemistry and Physiology Part a, Molecular Integrative Physiology*, 148(4), 853–860. doi:10.1016/j.cbpa.2007.08.027

Berger, G., Guetta, J., Klorin, G., Badarneh, R., Braun, E., Brod, V., et al. (2011). Sepsis impairs alveolar epithelial function by downregulating Na-K-ATPase pump. *AJP: Lung Cellular and Molecular Physiology*, 301(1), L23–L30. doi:10.1152/ajplung.00010.2010

Bozinovic, F., Gallardo, P. A., Visser, G. H., Cortés, A. (2003). Seasonal acclimatization in water flux rate, urine osmolality and kidney water channels in free-living degus: molecular mechanisms, physiological processes and ecological implications. *Journal of Experimental Biology*, 206(17), 2959–2966. doi:10.1242/jeb.00509

Böselt, I., Römpler, H., Hermsdorf, T., Thor, D., Busch, W., Schulz, A., Schöneberg, T. (2009). Involvement of the V2 vasopressin receptor in adaptation to limited water supply. *PloS One*, 4(5), e5573. doi:10.1371/journal.pone.0005573

Brenet, F., Moh, M., Funk, P., Feierstein, E., Viale, A. J., Socci, N. D., Scandura, J. M. (2011). DNA Methylation of the First Exon Is Tightly Linked to Transcriptional Silencing. *PloS One*, 6(1), e14524. doi:10.1371/journal.pone.0014524

Brown, D., Katsura, T., Kawashima, M., Verkman, A. S., Sabolic, I. (1995). Cellular distribution of the aquaporins: A family of water channel proteins. *Histochemistry and Cell Biology*, 104(1), 1–9. doi:10.1007/BF01464780

Cantarel, B. L., Korf, I., Robb, S. M. C., Parra, G., Ross, E., Moore, B., et al. (2008). MAKER: an easy-to-use annotation pipeline designed for emerging model organism genomes. *Genome Research*, 18(1), 188–196. doi:10.1101/gr.6743907

Cheuvront, S. N., Kenefick, R. W., Montain, S. J., Sawka, M. N. (2010). Mechanisms of aerobic performance impairment with heat stress and dehydration. *Journal of Applied Physiology*, 109(6), 1989–1995. doi:10.1152/jappphysiol.00367.2010

Eid, J., Fehr, A., Gray, J., Luong, K., Lyle, J., Otto, G., et al. (2009). Real-time DNA sequencing from single polymerase molecules. *Science (New York, NY)*, 323(5910), 133–138. doi:10.1126/science.1162986

Foret, S., Kucharski, R., Pellegrini, M., Pellegrini, M., Feng, S., Feng, S., et al. (2012). DNA methylation dynamics, metabolic fluxes, gene splicing, and alternative phenotypes in honey bees. *Proceedings of the National Academy of Sciences*, 109(13), 4968–4973. doi:10.1073/pnas.1202392109

Gallardo, P. A., Cortés, A., Bozinovic, F. (2005). Phenotypic flexibility at the molecular and organismal level allows desert-dwelling rodents to cope with seasonal water availability. *Physiological and Biochemical Zoology*, 78(2), 145–152. doi:10.1086/425203

Goralski, J. L., Boucher, R. C., Button, B. (2010). Osmolytes and ion transport modulators: new strategies for airway surface rehydration. *Current Opinion in Pharmacology*, 10(3), 294–299. doi:10.1016/j.coph.2010.04.003

Grabherr, M. G., Haas, B. J., Yassour, M., Levin, J. Z., Thompson, D. A., Amit, I., et al. (2011). Full-length transcriptome assembly from RNA-Seq data without a reference genome. *Nature Biotechnology*, 29(7), 644–652. doi:10.1038/nbt.1883

Gubler, M. C., Antignac, C. (2009). Renin–angiotensin system in kidney development: renal tubular dysgenesis. *Kidney International*, 77(5), 400–406. doi:10.1038/ki.2009.423

Haas, B. J., Papanicolaou, A., Yassour, M., Grabherr, M., Blood, P. D., Bowden, J., et al. (2013). De novo transcript sequence reconstruction from RNA-seq using the Trinity platform for reference generation and analysis. *Nature Protocols*, 8(8), 1494–1512. doi:10.1038/nprot.2013.084

Haussinger, D. (1996). The role of cellular hydration in the regulation of cell function. *Biochem J*. Hiller, D., Jiang, H., Xu, W., Wong, W. H. (2009). Identifiability of isoform deconvolution from junction arrays and RNA-Seq. *Bioinformatics (Oxford, England)*, 25(23), 3056–3059. doi:10.1093/bioinformatics/btp544

Huang, Y., Tracy, R., Walsberg, G. E., Makkinje, A., Fang, P., Brown, D., Van Hoek, A. N. (2001). Absence of aquaporin-4 water channels from kidneys of the desert rodent *Dipodomys merriami merriami*. *American Journal of Physiology-Renal Physiology*, 280(5), F794–F802.

Jéquier, E., Constant, F. (2009). Water as an essential nutrient: the physiological basis of hydration. *European Journal of Clinical Nutrition*, 64(2), 115–123.

doi:10.1038/ejcn.2009.111 Jiang, H., Wong, W. H. (2009). Statistical inferences for isoform expression in RNA-Seq. *Bioinformatics* (Oxford, England), 25(8), 1026–1032. doi:10.1093/bioinformatics/btp113 Keller, L. (Ed.). (2010). *The Honey Bee Epigenomes: Differential Methylation of Brain DNA in Queens and Workers*. *PLoS Biology*, 8(11), e1000506. doi:10.1371/journal.pbio.1000506 Kwon, T.-H., Nielsen, J., Nielsen, J., Møller, H. B., Møller, H. B., Fenton, R. A., et al. (2009). Aquaporins in the Kidney. In *Aquaporins* (Vol. 190, pp. 95–132). Berlin, Heidelberg: Springer Berlin Heidelberg. doi:10.1007/978-3-540-79885-9\_5 Langfelder, P., Horvath, S. (2008). *WGCNA: an R package for weighted correlation network analysis*. *BMC Bioinformatics*, 9, 11. doi:10.1186/1471-2105-9-559 LeGault, L. H., Dewey, C. N. (2013). *Inference of alternative splicing from RNA-Seq data with probabilistic splice graphs*. *Bioinformatics* (Oxford, England). doi: 10.1093/bioinformatics/btt396 /DC1 Li, H. (2013, March 16). *Aligning sequence reads, clone sequences and assembly contigs with BWA-MEM*. *MacManes, M. D. (2014). On the optimal trimming of high-throughput RNA-Seq data*. *Frontiers in Genetics*, 5. doi: 10.3389/fgene.2014.00013/abstract MacManes, M. D., throughput sequence reads. *PeerJ*, 1, e113. doi: 10.7717/peerj.113 Moeller, H. B., MacAulay, N., Knepper, M. A., Fenton, R. A., et al. (2009). *Terminal tail of aquaporin-2 for water transport: evidence against channel gating*. *American Journal of Physiology—Renal Physiology*, 296(3), F649–57. doi: 10.1152/ajprenal.90682.2008 Montain, S., Latzka, W., Sawka, N. (1999). *Fluid balance and metabolism of the Desert-Dwelling Kangaroo Rat (Dipodomys merriami)*. *Physiological Zoology*, 67(6), 1461–1478. doi:10.1038/ki.1995.389 Nielsen, S., Chou, C. L., Marples, D., Christensen, E. I., Kishore, B. K., Knepper, M. A. (1995). *Vasopressin increases water channel stoplasmamembrane*. *Pnas*, 92(4), 1013–1017. Ortells, M. C., Moranco, B., Drews-Elger, K., Violin, J. D., Rodriguez, C., Aramburu, J. (2012). *Transcriptional regulation of gene expression during osmotic stress responses by the* *10.1093/nar/gks038* Pyrkosz, A. B., Cheng, H., Brown, C. T. (2013, March 10). *RNA-Seq Mapping Errors When Using Induced Stress*. *Journal of Neuroendocrinology*, 23(1), 12–19. doi: 10.1111/j.1365-2826.2010.02074.x Schmidt-Nielsen, K., Schmidt-Nielsen, B. (1952). *Water metabolism of desert mammals*. *Physiological Reviews*, 32(2), 135–167. Soler, N., Brown, D., VanHoek, A. N. (2004). *Membrane organization and function of M1 and M23 isoforms of aquaporin 4 in epithelial cells*. *American Journal of Physiology—Renal Physiology*, 287(3), F501–11. doi: 10.1152/ajprenal.00439.2003 10.1073/pnas.1218661109 Totsune, K., Totsune, K., Takahashi, K., Murakami, O., Murakami, O., Satoh, F., et al. (1999). *HY P.24.6.758* Trapnell, C., Roberts, A., Goff, L., Pertea, G., Kim, D., Kelley, D. R., et al. (2012). *Differential gene expression experiments with TopHat and Cufflinks*. *Nature Protocols*, 7(3), 562–578. doi: 10.1038/nprot.2012.016 Verkman, A. S. (2009). *10.1080/ann.34.3.192.200* Weaver, D., Walker, L., Alcorn, D., Skinner, S. (1994). *The contribution of renin and vasopressin to the regulation of water permeability through aquaporin* *10.1016/0300-9629(94)90061-2* Yukutake, Y., Yasui, M. (2010). *Regulation of Water Permeability Through Aquaporin 4*. *Nsc*, 168(4), 885–891. doi: 10.1016/j.neuroscience.2009.10.029 Zingg, H. H., Lefebvre, D., Almazan, G. (1986). *Regulation of water permeability through aquaporin 4*