

# Characterization of the transcriptome, nucleotide sequence polymorphism and selection in the desert adapted mouse *Peromyscus eremicus*

Matthew D. MacManes<sup>1</sup>, Michael B. Eisen<sup>2</sup>,

**1** Department of Molecular, Cellular and Biomedical Sciences, University of New Hampshire  
Institution Name, Durham NH, USA

**2** HHMI and University of California, Berkeley, Berkeley, CA, USA

\* E-mail: macmanes@gmail.com, @PeroMHC

## Abstract

## Introduction

For biologists interested in understanding the relationship between fitness, genotype, and phenotype, modern sequencing technologies provide for an unprecedented opportunity to gain a deep understanding of genome level processes that together, underlie adaptation. One interesting example of adaptation lies in animals ability to survive desert conditions. Here, heat *and* drought provide for powerful selective forces, testing animals' ability to osmoregulate and thus to survive.

Specifically, the maintenance of water balance in animals is one of the most important physiologic processes, and is critical to desert survival. Indeed, mammals are exquisitely sensitive to changes in osmolality, with slight derangement eliciting physiologic compromise. When the loss of water exceeds dietary intake, dehydration - and in extreme cases, death - can occur. Unlike most mammals, animals living in desert habitats are subjected to long periods of extreme heat and intense drought. As a result, desert animals have evolved mechanisms through which physiologic homeostasis is maintained despite severe and prolonged dehydration. While the elucidation of these mechanisms is beyond the scope of this manuscript, we aim here to lay the groundwork by characterizing the transcriptome from four distinct tissues (brain, liver, kidney, testes). Further, via sequencing the renal tissue of a total of 15 additional animals, we characterize nucleotide polymorphism and genome wide patterns of natural selection.

## Results

### Subsection 1

### Subsection 2

## Discussion

## Materials and Methods

### RNA extraction and Sequencing

Total RNA was extracted from each tissue using a TRIzol extraction (Invitrogen) following the manufacturers instructions. Because preparation of an RNA library suitable for sequencing is dependent on having high quality, intact RNA, a small aliquot of each total RNA extract was analyzed on a Bioanalyzer 2100 (Agilent, Palo Alto, CA, USA). Following confirmation of sample quality, the reference sequencing libraries were made using the TruSeq stranded RNA prep kit (Illumina), while an unstranded TruSeq kit was used to construct the other sequencing libraries. A unique index was ligated to each sample to allow

for multiplexed sequencing. Libraries were then pooled to contain equimolar quantities of each individual library and submitted for Illumina sequencing on a HiSeq 2000 sequencer at The Vincent Coates Genome Sequencing Lab at UC Berkeley.

## **Acknowledgments**

## **Figure Legends**

## **Tables**