

Characterization of the transcriptome, nucleotide sequence polymorphism, and natural selection in the desert adapted mouse *Peromyscus eremicus*

Matthew D. MacManes¹, Michael B. Eisen²,

¹ Department of Molecular, Cellular and Biomedical Sciences, University of New Hampshire, Durham NH, USA

² HHMI and University of California, Berkeley, Berkeley, CA, USA

* E-mail: macmanes@gmail.com, @PeroMHC

1 Abstract

2 Introduction

3 Deserts are widely considered one of Earth's harshest environments. Animals living in desert
4 environments are forced to endure intense heat and drought, and in turn, species having evolved
5 in these environments are likely to have evolved specialised mechanisms that may enhance
6 fitness. While living in deserts likely involves a large number of adaptive phenotypes, the abil-
7 ity to osmoregulate – to maintain the proper water and electrolyte balance – appears to be
8 paramount [1]. Indeed, the maintenance of water balance in animals is one of the most impor-
9 tant physiologic processes, and is critical to desert survival. Mammals are exquisitely sensitive
10 to changes in osmolality, with slight derangement eliciting physiologic compromise. When the
11 loss of water exceeds dietary intake, dehydration - and in extreme cases, death - can occur. Un-
12 derstanding these mechanisms will significantly enhance our understanding of the physiologic
13 processes underlying osmoregulation in extreme environments, having implications for studies
14 of human health, conservation, and climate change.

15

16 The genes and structures responsible for the maintenance of water and electrolyte balance
17 are well characterized in model organisms such as mice [2], rats [3–5], and humans [6–8]. These
18 studies, many of which have been enabled by newer sequencing technologies, serve as a founda-
19 tion for studies of renal genomics in non-model organisms. In particular, because researchers
20 have long been interested in desert adaptation, a number of studies have looked at the mor-
21 phology or expression of single genes in the renal tissues of desert adapted rodents *Phyllotis*
22 *darwini* [9], *Psammomys obesus* [10], and *Perognathus penicillatus* [11]. More recently, full re-
23 nal transcriptomes have been generated for *Dipodomys spectabilis* and *Chaetodipus baileyi* [12]
24 as well as *Abrothrix olivacea* [13].

25

26 These studies provide a rich context for the current and future work, aimed at developing

a synthetic understanding of the the genetic and genomic underpinnings of desert adaptation in rodents. As a first step, we have sequenced, assembled, and characterized the transcriptome (using four tissue types - liver, kidney, testes, brain), of a desert adapted cricetid rodent endemic to the Southwest United States [14], *Peromyscus eremicus*. These animals have a lifespan typical of small mammals, and therefore an individual may live it's entire life without ever drinking water. These rodents have distinct advantage over other desert animals (e.g. *Dipodomys*) in that they breed readily in captivity, which enables future laboratory studies of the phenotype of interest. In addition the focal species is positioned in a clade of well known animals (e.g. *P. californicus*, *P. maniculatus* and *P. polionotus*) [15] with growing genetic and genomic resources [16–18] which together suggest that future comparative studies are possible.

While the elucidation of the mechanisms underlying adaptation to desert survival is beyond the scope of this manuscript, we aim here to lay the groundwork by characterizing the transcriptome from four distinct tissues (brain, liver, kidney, testes). These data will be included in current larger effort aimed at sequencing the entire genome. Further, via sequencing the renal tissue of a total of 15 additional animals, we characterize nucleotide polymorphism and genome wide patterns of natural selection. Together, these investigations will aid in our overarching goal – to understand the genetic bases of adaptation in *P. eremicus*.

Materials and Methods

Animal Collection and Study Design

To begin to understand how genes may underlie desert adaptation, I collected 15 individuals from a single population *P. eremicus* over a two year time period (2012-2013). These individuals were captured in live traps, then euthanized using isoflurane overdose and decapitation. Immediately post-mortem, the abdominal and pelvic organs were removed, cut in half (in the case of kidney), placed in RNAlater and flash frozen in liquid Nitrogen. Removal of the brain, with similar preservation techniques, followed that. Time from euthanasia to removal of all organs never exceeded five minutes. Samples were transferred to a -80C freezer at a later date. These procedures were approved by the University of California Berkeley Animal Care and Use Committee and follow guidelines established by the American Society of Mammalogy for the use of wild animals in research [19].

RNA extraction and Sequencing

Total RNA was extracted from each tissue using a TRIzol extraction (Invitrogen) following the manufacturers instructions. Because preparation of an RNA library suitable for sequencing is

dependent on having high quality, intact RNA, a small aliquot of each total RNA extract was analyzed on a Bioanalyzer 2100 (Agilent, Palo Alto, CA, USA). Following confirmation of sample quality, the reference sequencing libraries were made using the TruSeq stranded RNA prep kit (Illumina), while an unstranded TruSeq kit was used to construct the other sequencing libraries. A unique index was ligated to each sample to allow for multiplexed sequencing. Reference libraries (n=4 tissue types) were then pooled to contain equimolar quantities of each individual library and submitted for Illumina sequencing using two lanes of 150nt paired end sequencing using the rapid-mode of the HiSeq 2500 sequencer at The Hubbard Center for Genome Sciences (University of New Hampshire). The remaining 14 libraries were similarly multiplexed and sequenced in a mixture of 100nt paired and single end across two lanes of an Illumina HiSeq 2000 at the Vincent G. Coates Gnome Center (University of California, Berkeley).

Sequence Data Preprocessing and Assembly

The raw sequence reads were error corrected using the software `bleed` [20], using `kmer=25`, based on the developers recommendations. The error corrected adapter and quality trimmed following recommendations from MacManes [21] and Mbandi [22]. Specifically, adapter sequence contamination was removed, and low quality nucleotides (defined as `PHRED < 2`) were removed using the program `Trimmomatic` version 0.32 [23]. Reads from each tissue were assembled using `Trinity` version released 17 July 2014 [24]. We used flags indicating the stranded nature of sequencing reads and set maximum allowable physical distance between read pairs to 999nt. The assembly was conducted on a linux workstation with 64 cores and 512Gb RAM. To filter the raw sequence assembly, I downloaded *Mus musculus* cDNA and ncRNA datasets from Ensembl (ftp://ftp.ensembl.org/pub/release-75/fasta/mus_musculus/), and the *Peromyscus maniculatus* reference transcriptome from NCBI (ftp://ftp.ncbi.nlm.nih.gov/genomes/Peromyscus_maniculatus_bairdii/RNA/). I used a relaxed blastN procedure (evalue set to 10^{-10}) to identify contigs in the *P. eremicus* dataset that are likely biological in origin. I then concatenated the filtered assemblies from each tissue into a single file, reducing redundancy using the software `cd-hit-est` [27] using default setting except that only identical sequences were removed. The resulting assembly file was characterized using the software package `transrate` (<https://github.com/Blahah/transrate>).

Assembled Sequence Annotation

From the filtered assembly, I extracted putative coding sequences using `Transdecoder` version 16Jan2014 (<http://transdecoder.sourceforge.net/>). These putative protein coding sequences were annotated using default settings of the `blastx` algorithm [28] against the Swis-

sProt database downloaded on 1 March 2014. Because transcriptome assemblies typically contain non-coding elements (e.g. ncRNA) in addition to protein coding sequence, we annotated the entire filtered dataset using the NCBI nt dataset, downloaded on 1 March 2014. Lastly, because rapidly evolving nucleotide sequences may evade detection by blast algorithms, we used HMMER3 [29] to search for conserved protein domains contained in the Pfam database [30].

To identify sequences unique to each tissue type, I mapped sequence reads from each tissue type to the reference assembly using bwa-mem. We estimated expression individually for the four tissues. Interesting patterns of expression, including instances where expression was limited to a single tissue type were identified.

Population Genomics

In addition to the reference individual sequenced at four different tissue types, we sequenced 15 other conspecific individuals from the same population, located in Palm Desert, California. Sequence data were mapped to the reference transcriptome using bwa-mem. The alignments were sorted and converted to BAM format, then passed to the program ANGSD, which was used for calculating the folded site frequency spectrum (SFS) and Tajima's D [31].

Natural Selection

To characterize natural selection on several genes related to water and ion homeostasis, we identified several of the transcripts identified as experiencing positive selection in a recent work on desert-adapted *Dipodomys* rodents. The coding sequence corresponding to these genes, Solute Carrier family 2 member 9 (Slc2a9) and the Vitamin D3 receptor (Vdr), were extracted from the dataset, aligned using the software MACSE [32] to homologous sequences identified by the conditional reciprocal best blast procedure (CRBB, [33]) implemented in transrate. These alignments were inputted into the software codeABC version 1.6.0 [34].

Results

RNA extraction, Sequencing, Assembly, Mapping

RNA was extracted from the hypothalamus, renal medulla, testes, or liver from each individual using sterile technique. TRIzol extraction resulted in a large amount of high quality ($RIN \geq 8$) total RNA, which was used as input. Libraries were constructed as per the standard Illumina

protocol, and were sequenced as described above. The number of reads per library varied from 56 million strand-specific paired-end reads in Peer360 kidney, to 9 million single-end reads in Peer321. Adapter sequence contamination and low-quality nucleotides were eliminated, which resulted in a loss of <2% of reads.

129

Transcriptome assembly was accomplished using the program Trinity. The raw assembly contained 743314 assembled sequences measuring 418Mb. This assembly was filtered using TMP >1 as a threshold. The filtered assembly contained 130764 sequences measuring 149Mb. From this filtered dataset, I extracted 64355 putative coding sequences (60Mb). Of these 64355 sequences, 37960 were complete exons (containing both start and stop codons), while others were either truncated at the 5-prime end (16880 sequences), 3-prime end (4203 sequences), or were internal (5312 sequences having neither stop nor start codon).

137 Subsection 2

138 Discussion

139 Acknowledgments

140 References

- 141 1. Walsberg G (2000) Small mammals in hot deserts: Some generalizations revisited. *Bio-*
142 *science* 50: 109–120.
- 143 2. Tatum R, Zhang Y, Salleng K, Lu Z, Lin JJ, et al. (2009) Renal salt wasting and chronic
144 dehydration in claudin-7-deficient mice. *AJP: Renal Physiology* 298: F24–F34.
- 145 3. Romero DG, Plonczynski MW, Welsh BL, Gomez-Sanchez CE, Zhou MY, et al. (2007)
146 Gene expression profile in rat adrenal zona glomerulosa cells stimulated with aldosterone
147 secretagogues. *Physiological Genomics* 32: 117–127.
- 148 4. Rojek A, Rojek A, Fuchtbauer E, Fuchtbauer E, Kwon T, et al. (2006) Severe urinary
149 concentrating defect in renal collecting duct-selective AQP2 conditional-knockout mice.
150 *Proceedings of The National Academy of Sciences of The United States of America* 103:
151 6037–6042.
- 152 5. Nielsen S, CHOU C, MARPLES D, CHRISTENSEN E, KISHORE B, et al. (1995) Vaso-
153 pressin Increases Water Permeability of Kidney Collecting Duct by Inducing Transloca-

- tion of Aquaporin-Cd Water Channels to Plasma-Membrane. Proceedings of The National Academy of Sciences of The United States of America 92: 1013–1017.
6. Mobasheri A, Marples D, Young IS, Floyd RV, Moskaluk CA, et al. (2007) Distribution of the AQP4 Water Channel in Normal Human Tissues: Protein and Tissue Microarrays Reveal Expression in Several New Anatomical Locations, including the Prostate Gland Seminal Vesicles. Channels 1: 30–39.
 7. Bedford JJ, Bedford JJ, Leader JP, Leader JP, Walker RJ, et al. (2003) Aquaporin expression in normal human kidney and in renal disease. Journal of the American Society of Nephrology : JASN 14: 2581–2587.
 8. Nielsen S, KWON T (1999) Physiology and Pathophysiology of Renal Aquaporins. Journal of the
 9. Gallardo PA, Cortés A, Bozinovic F (2005) Phenotypic flexibility at the molecular and organismal level allows desert-dwelling rodents to cope with seasonal water availability. Physiological and Biochemical Zoology 78: 145–152.
 10. Kaissling B, De Rouffignac C, Barrett JM, Kriz W (1975) The structural organization of the kidney of the desert rodent *Psammomys obesus*. Anatomy and embryology 148: 121–143.
 11. Altschuler EM, Altschuler EM, Nagle RB, Nagle RB, Braun EJ, et al. (1979) Morphological study of the desert heteromyid kidney with emphasis on the genus *Perognathus*. The Anatomical record 194: 461–468.
 12. Marra NJ, Romero a, DeWoody Ja (2014) Natural selection and the genetic basis of osmoregulation in heteromyid rodents as revealed by RNA-seq. Molecular Ecology 23: 2699–2711.
 13. Giorello FM, Feijoo M, a GD, Valdez L, Opazo JC, et al. (2014) Characterization of the kidney transcriptome of the South American olive mouse *Abrothrix olivacea* 15: 1–10.
 14. Veal R, Caire W (2001) *Peromyscus eremicus*. Mammalian Species 118: 1–6.
 15. Feng BJ, Sun LD, Soltani-Arabshahi R, Bowcock AM, Nair RP, et al. (2007) Toward a Molecular Phylogeny for *Peromyscus*: Evidence from Mitochondrial Cytochrome- b Sequences. Journal of Mammalogy 88: 1146–1159.
 16. Shorter KR, Owen A, anderson V, Hall-South AC, Hayford S, et al. (2014) Natural Genetic Variation Underlying Differences in *Peromyscus* Repetitive and Social/Aggressive Behaviors. Behavior genetics .

- 186 17. Panhuis TM, Panhuis TM, Broitman-Maduro G, Broitman-Maduro G, Uhrig J, et al.
 187 (2011) Analysis of Expressed Sequence Tags from the Placenta of the Live-Bearing Fish
 188 *Poeciliopsis* (Poeciliidae). *Journal of Heredity* 102: 352–361.
- 189 18. Shorter KR, Crossland JP, Webb D, Szalai G, Felder MR, et al. (2012) *Peromyscus* as a
 190 Mammalian Epigenetic Model. *Genetics Research International* 2012: 1–11.
- 191 19. Sikes RS, Gannon WL, Animal Care and Use Committee of the American Society of
 192 Mammalogists (2011) Guidelines of the American Society of Mammalogists for the use of
 193 wild mammals in research. *Journal of Mammalogy* 92: 235–253.
- 194 20. Heo Y, Wu XL, Chen D, Ma J, Hwu WM (2014) BLESS: bloom filter-based error cor-
 195 rection solution for high-throughput sequencing reads. *Bioinformatics* (Oxford, England)
 196 30: 1354–1362.
- 197 21. MacManes MD (2014) On the optimal trimming of high-throughput mRNA sequence
 198 data. *Frontiers in Genetics* 5.
- 199 22. Christoffels A (2014) A glance at quality score: implication for *de novo* transcriptome
 200 reconstruction of Illumina reads : 1–5.
- 201 23. Lohse M, Bolger AM, Nagel A, Fernie AR, Lunn JE, et al. (2012) RobiNA: a user-friendly,
 202 integrated software solution for RNA-Seq-based transcriptomics. *Nucleic Acids Research*
 203 40: W622–7.
- 204 24. Haas BJ, Papanicolaou A, Yassour M, Grabherr M, Blood PD, et al. (2013) *De novo*
 205 transcript sequence reconstruction from RNA-seq using the Trinity platform for reference
 206 generation and analysis. *Nature protocols* 8: 1494–1512.
- 207 25. Li H (2013) Aligning sequence reads, clone sequences and assembly contigs with BWA-
 208 MEM .
- 209 26. Roberts A, Pachter L (2013) Streaming fragment assignment for real-time analysis of
 210 sequencing experiments. *Nature Methods* 10: 71–73.
- 211 27. Li W, Li W, Godzik A, Godzik A (2006) Cd-hit: a fast program for clustering and
 212 comparing large sets of protein or nucleotide sequences. *Bioinformatics* (Oxford, England)
 213 22: 1658–1659.
- 214 28. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, et al. (2009) BLAST+:
 215 architecture and applications. *BMC Bioinformatics* 10: 421.

- 216 29. Wheeler TJ, Wheeler TJ, Eddy SR, Eddy SR (2013) nhmmer: DNA homology search
217 with profile HMMs. *Bioinformatics (Oxford, England)* 29: 2487–2489.
- 218 30. Punta M, Coggill PC, Eberhardt RY, Mistry J, Tate J, et al. (2012) The Pfam protein
219 families database. *Nucleic Acids Research* 40: D290–301.
- 220 31. Korneliussen TS, Moltke I, Albrechtsen A (2013) Calculation of Tajima’s D and other
221 neutrality test statistics from low depth next-generation sequencing data. *BMC ...*
- 222 32. Ranwez V, Harispe S, Delsuc F, Douzery EJP (2011) MACSE: Multiple Alignment of
223 Coding SEquences Accounting for Frameshifts and Stop Codons. *PLOS ONE* 6: e22594.
- 224 33. Aubry S, Kelly S, Kämpers BMC, Smith-Unna RD, Hibberd JM (2014) Deep Evolutionary
225 Comparison of Gene Expression Identifies Parallel Recruitment of Trans-Factors in Two
226 Independent Origins of C4 Photosynthesis. *PLOS Genetics* 10: e1004365.
- 227 34. Lopes JS, Arenas M, Posada D, Beaumont MA (2013) Coestimation of recombination,
228 substitution and molecular adaptation rates by approximate Bayesian computation 112:
229 255–264.

230 **Figure Legends**

231 **Tables**

232 **Table 1**

233

DATASET	NUM. RAW READS
PEER360 TESTES	32M PE
PEER360 LIVER	53M PE
PEER360 KIDNEY	56M PE
PEER360 BRAIN	23M PE
PEER305	19M PE
PEER308	15M PE
PEER319	14M PE
PEER321	9M SE
PEER340	16M PE
PEER352	14M PE
PEER354	9M SE
PEER359	14M PE
PEER365	16M PE
PEER366	16M PE
PEER368	14M PE
PEER369	14M PE
PEER372	17M SE
PEER373	23M SE
PEER380	16M SE
PEER382	14M SE

234

235

Table 1. The number of sequencing reads per sample