
Cardioembolic estimate points the same way in all 12 analyses; the 95% CI excludes 1 in 7
Shaded band: range of IVW estimates over 35 leave−one−locus−out analyses (OR 0.836−0.871; P 0.005−0.031 in every one)
Estimator (all 41 SNPs)
IVW, random effects (primary)
Weighted median
MR−Egger
MR−RAPS (robust to outliers)
0.853 (0.754−0.966) n=41
0.862 (0.717−1.036) n=41
0.907 (0.729−1.129) n=41
0.856 (0.753−0.972) n=41
Removing loci / restricting SNP set
Without HP locus (chr16, 4 SNPs)
Without LPA locus (chr6, 3 SNPs)
Without HP, LPA and MHC loci
31 SNPs shared by all five outcomes
0.843 (0.727−0.977) n=37
0.871 (0.767−0.987) n=38
0.867 (0.744−1.009) n=34
0.833 (0.729−0.952) n=31
Removing AF−related instruments
Without SNPs with AF p < 0.001
Without SNPs near AF lead SNPs
Without either AF−related group
0.884 (0.779−1.003) n=38
0.845 (0.746−0.958) n=39
0.876 (0.771−0.996) n=36
Earlier release (partly overlapping cases)
MEGASTROKE 2018 CES, IVW
0.907 (0.794−1.036) n=51
0.7
0.8
0.9
1.0
1.2
Odds ratio for cardioembolic stroke per 1−SD higher GlycA (95% CI)