

Section 1. Clinical Findings

Patients in cohort 1 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 2. Clinical Findings

Patients in cohort 2 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 3. Clinical Findings

Patients in cohort 3 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 4. Clinical Findings

Patients in cohort 4 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 5. Clinical Findings

Patients in cohort 5 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 6. Clinical Findings

Patients in cohort 6 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 7. Clinical Findings

Patients in cohort 7 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 8. Clinical Findings

Patients in cohort 8 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 9. Clinical Findings

Patients in cohort 9 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 10. Clinical Findings

Patients in cohort 10 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 11. Clinical Findings

Patients in cohort 11 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 12. Clinical Findings

Patients in cohort 12 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 13. Clinical Findings

Patients in cohort 13 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 14. Clinical Findings

Patients in cohort 14 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 15. Clinical Findings

Patients in cohort 15 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 16. Clinical Findings

Patients in cohort 16 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 17. Clinical Findings

Patients in cohort 17 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 18. Clinical Findings

Patients in cohort 18 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 19. Clinical Findings

Patients in cohort 19 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 20. Clinical Findings

Patients in cohort 20 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 21. Clinical Findings

Patients in cohort 21 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 22. Clinical Findings

Patients in cohort 22 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 23. Clinical Findings

Patients in cohort 23 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 24. Clinical Findings

Patients in cohort 24 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 25. Clinical Findings

Patients in cohort 25 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 26. Clinical Findings

Patients in cohort 26 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 27. Clinical Findings

Patients in cohort 27 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 28. Clinical Findings

Patients in cohort 28 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 29. Clinical Findings

Patients in cohort 29 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 30. Clinical Findings

Patients in cohort 30 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 31. Clinical Findings

Patients in cohort 31 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 32. Clinical Findings

Patients in cohort 32 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 33. Clinical Findings

Patients in cohort 33 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 34. Clinical Findings

Patients in cohort 34 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 35. Clinical Findings

Patients in cohort 35 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 36. Clinical Findings

Patients in cohort 36 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 37. Clinical Findings

Patients in cohort 37 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 38. Clinical Findings

Patients in cohort 38 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 39. Clinical Findings

Patients in cohort 39 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 40. Clinical Findings

Patients in cohort 40 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 41. Clinical Findings

Patients in cohort 41 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 42. Clinical Findings

Patients in cohort 42 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 43. Clinical Findings

Patients in cohort 43 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 44. Clinical Findings

Patients in cohort 44 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 45. Clinical Findings

Patients in cohort 45 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 46. Clinical Findings

Patients in cohort 46 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 47. Clinical Findings

Patients in cohort 47 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 48. Clinical Findings

Patients in cohort 48 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 49. Clinical Findings

Patients in cohort 49 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.

Section 50. Clinical Findings

Patients in cohort 50 demonstrated a measurable change in estimated glomerular filtration rate over the twelve month observation window. The magnitude of the change varied with baseline proteinuria.

Secondary endpoints included hospitalization for volume overload and the initiation of renal replacement therapy. Neither endpoint reached the prespecified threshold for statistical significance.