



Package leaflet: Information for the user

Amlodipine 5 mg tablets
Amlodipine 10 mg tablets
amlodipine

Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.

- Keep this leaflet. You may need to read it again.
- If you have any further questions, ask your doctor or pharmacist.
- This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any side effects, talk to your doctor or pharmacist. This includes any possible side effects not listed in this leaflet. See section 4.

What is in this leaflet:

1. What Amlodipine is and what it is used for
2. What you need to know before you take Amlodipine
3. How to take Amlodipine
4. Possible side effects
5. How to store Amlodipine
6. Contents of the pack and other information

1. What Amlodipine is and what it is used for

Amlodipine contains the active substance amlodipine which belongs to a group of medicines called calcium antagonists.

Amlodipine is used to treat high blood pressure (hypertension) or a certain type of chest pain called angina, a rare form of which is Prinzmetal's or variant angina.

In patients with high blood pressure this medicine works by relaxing blood vessels, so that blood passes through them more easily. In patients with angina Amlodipine works by improving blood supply to the heart muscle which then receives more oxygen and as a result chest pain is prevented. This medicine does not provide immediate relief of chest pain from angina.

2. What you need to know before you take Amlodipine

Do not take Amlodipine

- If you are allergic (hypersensitive) to amlodipine, or any of the other ingredients of this medicine listed in section 6, or to any other calcium antagonists. This may be itching, reddening of the skin or difficulty in breathing.
- If you have severe low blood pressure (hypotension)
- If you have narrowing of the aortic heart valve (aortic stenosis) or cardiogenic shock (a condition where your heart is unable to supply enough blood to the body).
- If you suffer from heart failure after a heart attack

Warnings and precautions

Talk to your doctor or pharmacist before taking Amlodipine
You should inform your doctor if you have or have had any of the following conditions:

- Recent heart attack
- Heart failure
- Severe increase in blood pressure (Hypertensive crisis)
- Liver disease
- You are elderly and your dose needs to be increased

Children and adolescents

Amlodipine has not been studied in children under the age of 6 years. Amlodipine should only used for hypertension in children and adolescents from 6 years to 17 years of age (see section 3).
For more information, talk to your doctor.

Other medicines and Amlodipine

Tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines, including medicines obtained without a prescription.

Amlodipine may affect or be affected by other medicines, such as:

- ketoconazole, itraconazole (anti-fungal medicines)

- ritonavir, indinavir, nelfinavir (so called protease inhibitors used to treat HIV)
- rifampicin, erythromycin, clarithromycin (antibiotics)
- hypericum perforatum (St. John's Wort)
- verapamil, diltiazem (heart medicines)
- dantrolene (infusion for severe body temperature abnormalities)
- tacrolimus, sirolimus, temsirolimus, and everolimus (medicines used to alter the way your immune system works)
- simvastatin (cholesterol lowering medicine)
- cyclosporine (an immunosuppressant)

Amlodipine may lower your blood pressure even more if you are already taking other medicines to treat your high blood pressure.

Amlodipine with food and drink

Grapefruit juice and grapefruit should not be consumed by people who are taking Amlodipine. This is because grapefruit and grapefruit juice can lead to an increase in the blood levels of the active ingredient amlodipine, which can cause an unpredictable increase in the blood pressure lowering effect of Amlodipine.

Pregnancy and breast-feeding
Pregnancy

The safety of amlodipine in human pregnancy has not been established. If you think you might be pregnant, or are planning to get pregnant, you must tell your doctor before you take Amlodipine.

Breast-feeding

Amlodipine has been shown to pass into breast milk in small amounts. If you are breast-feeding or about to start breast-feeding you must tell your doctor before taking Amlodipine.

Ask your doctor or pharmacist for advice before taking any medicine.

Driving and using machines

Amlodipine may affect your ability to drive or use machines. If the tablets make you feel sick, dizzy or tired, or give you a headache, do not drive or use machines and contact your doctor immediately.

Amlodipine contains sodium

This medicine contains less than 1 mmol sodium (23 mg) per tablet, which means it is essentially 'sodium-free'.

3. How to take Amlodipine

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Adults

The recommended initial dose is Amlodipine 5 mg once daily. The dose can be increased to Amlodipine 10 mg once daily.

This medicine can be used before or after food and drinks. You should take this medicine at the same time each day with a drink of water. Do not take Amlodipine with grapefruit juice.

Use in children and adolescents

For children and adolescents (6-17 years old), the recommended usual starting dose is 2.5 mg a day. The maximum recommended dose is 5 mg a day. Amlodipine 2.5 mg is not currently available and the 2.5 mg dose cannot be obtained with Amlodipine 5 mg tablets as these tablets are not manufactured to break into two equal halves.

It is important to keep taking the tablets. Do not wait until your tablets are finished before seeing your doctor.

If you take more Amlodipine than you should

Taking too many tablets may cause your blood pressure to become low or even dangerously low. You may feel dizzy, lightheaded, faint or weak. If blood pressure drop is severe enough shock can occur. Your skin could feel cool and clammy and you could lose consciousness. Seek immediate medical attention if you take too many Amlodipine tablets.

Excess fluid may accumulate in your lungs (pulmonary oedema) causing shortness of breath that may develop up to 24-48 hours after intake.

If you forget to take Amlodipine

Do not worry. If you forget to take a tablet, leave out that dose completely. Take your next dose at the right time. Do not take a double dose to make up for a forgotten dose.

Black A/s: 165 x 350 mm

 AUROBINDO Packaging Development		Product Name	Component	Item Code	Date & Time
		Amlodipine	Leaflet	P1540047	06.10.2025 & 9.00 AM
Team Leader		Customer / Country	Version No.	Reason Of Issue	Reviewed / Approved by APPROVED By RA Milpharm at 10:37 am, Oct 11, 2025
		Milpharm_Unit-3 & Unit 15	01	Revision	
Initiator		Dimensions	No. of Colours : 01		
Artist:		165 x 350 mm			
 vision Graphic Designers		Pharmacode			
		40047			
Additional Information : Supersede Item Code: P1533156					
Sign / Date					

P1540047

If you stop taking Amlodipine
Your doctor will advise you how long to take this medicine. Your condition may return if you stop using this medicine before you are advised.
If you have any further questions on the use of this medicine, ask your doctor or pharmacist.

4. Possible side effects

Like all medicines, this medicine can cause side effects, although not everybody gets them.

Visit your doctor **immediately** if you experience any of the following side effects after taking this medicine.

- Sudden wheeziness, chest pain, shortness of breath or difficulty in breathing
- Swelling of eyelids, face or lips
- Swelling of the tongue and throat which causes great difficulty breathing
- Severe skin reactions including intense skin rash, hives, reddening of the skin over your whole body, severe itching, blistering, peeling and swelling of the skin, inflammation of mucous membranes (Stevens Johnson Syndrome, toxic epidermal necrolysis) or other allergic reactions
- Heart attack, abnormal heart beat
- Inflamed pancreas which may cause severe abdominal and back pain accompanied with feeling very unwell

The following **very common side effect** has been reported. If this causes you problems or if it **lasts for more than one week**, you should contact your doctor.

Very common (may affect more than 1 in 10 people)
• Oedema (fluid retention)

The following **common side-effects** have been reported. If any of these cause you problems or if they **last for more than one week**, you should **contact your doctor**.

Common (may affect up to 1 in 10 people)
• Headache, dizziness, sleepiness (especially at the beginning of treatment)
• Palpitations (awareness of your heart beat), flushing
• Abdominal pain, feeling sick (nausea)
• Altered bowel habits, diarrhoea, constipation, indigestion,
• Tiredness, weakness
• Visual disturbances, double vision
• Muscle cramps
• Ankle swelling

Other side-effects that have been reported include the following list. If any of these get serious, or if you notice any side-effects not listed in this leaflet, please tell your doctor or pharmacist.

Uncommon (may affect up to 1 in 100 people)
• Mood changes, anxiety, depression, sleeplessness
• Trembling, taste abnormalities, fainting,
• Numbness or tingling sensation in your limbs; loss of pain sensation
• Ringing in the ears
• Low blood pressure
• Sneezing/running nose caused by inflammation of the lining of the nose (rhinitis)
• Cough
• Dry mouth, vomiting (being sick)
• Hair loss, increased sweating, itchy skin, red patches on skin, skin discolouration
• Disorder in passing urine, increased need to urinate at night, increased number of times of passing urine
• Inability to obtain an erection; discomfort or enlargement of the breasts in men
• Pain, feeling unwell
• Joint or muscle pain, back pain
• Weight increase or decrease

Rare (may affect up to 1 in 1,000 people)
• Confusion

Very rare (may affect up to 1 in 10,000 people)
• Decreased numbers of white blood cells, decrease in blood platelets which may result in unusual bruising or easy bleeding
• Excess sugar in blood (hyperglycaemia)
• A disorder of the nerves which can cause muscular weakness, tingling or numbness
• Swelling of the gums, bleeding gums
• Abdominal bloating (gastritis)

- Abnormal liver function, inflammation of the liver (hepatitis), yellowing of the skin (jaundice), liver enzyme increase which may have an effect on some medical tests
- Increased muscle tension
- Inflammation of blood vessels, often with skin rash
- Sensitivity to light
- Disorders combining rigidity, tremor, and/or movement disorders

Not known (cannot be estimated from the available data)
• Trembling, rigid posture, mask-like face, slow movements and a shuffling, unbalanced walk.

Reporting of side effects
If you get any side effects, talk to your doctor or pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via Yellow Card Scheme Website: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store. By reporting side effects you can help provide more information on the safety of this medicine.

5. How to store Amlodipine

Keep this medicine out of the sight and reach of children.

This medicinal product does not require any special storage conditions.

Do not use this medicine after the expiry date which is stated on the carton after EXP. The expiry date refers to the last day of that month.

Do not throw away any medicines via wastewater or household waste. Ask your pharmacist how to throw away medicines you no longer use. These measures will help protect the environment.

6. Contents of the pack and other information

What Amlodipine contains
- The active substance is amlodipine as amlodipine besilate.
Each tablet contains amlodipine besilate equivalent to 5 mg of amlodipine.
Each tablet contains amlodipine besilate equivalent to 10 mg of amlodipine.
- The other ingredients are microcrystalline cellulose [E 460a], calcium hydrogen phosphate, sodium starch glycolate, magnesium stearate [E 470B].

What Amlodipine looks like and contents of the pack
Tablet.

Amlodipine 5 mg tablets
White or almost white, flat, bevelled edges, barrel-shaped tablet debossed with "C" on one side and "58" on the other side. The size is 8 mm x 6 mm.

Amlodipine 10 mg tablets
White or almost white, flat, bevelled edges, round [diameter 10 mm] tablet debossed with "C" on one side and "59" on the other side.

Amlodipine tablets are available in PVC/ PVdC-Aluminium blister.

Pack sizes:
Blister pack:
7, 10, 14, 15, 20, 28, 30, 50, 56, 60, 84, 90, 98, 100, 120, 200, 250, 300 and 500 tablets.

Not all pack sizes may be marketed.

Marketing Authorisation Holder
Milpharm Limited
1 Roundwood Avenue,
Stockley Park, Uxbridge,
UB11 1AF
United Kingdom

Manufacturer
APL Swift Services (Malta) Limited
HF 26, Hal Far Industrial Estate, Hal Far
Brizebugia BBG 3000
Malta

or
Milpharm Limited
1 Roundwood Avenue,
Stockley Park, Uxbridge,
UB11 1AF
United Kingdom

This leaflet was last approved in 10/2025.