KAHLER-EINSTEIN SUBMANIFOLDS OF COMPLEX SPACE FORMS

MATTHEW DRURY

Note. This is an incomplete PhD thesis in Mathematics. It was written between 2010 and 2011 at Indiana University under the advisement of Ji-Ping Sha.

The thesis contains an exposition on the basics of complex differential geometry, focusing on the method of moving frames. It then uses that framework to derive Calabi's and Chern's curvature restrictions on Kahler-Einstein submanifolds of complex space forms. It includes a novel formula for the third covariant derivative of the second fundamental form.

1. Introduction

Let $\tilde{M}^n(c)$ be an n dimensional Kähler space form with constant holomorphic curvature c. As is the case in Riemannian geometry, such an $\tilde{M}$ is locally holomorphically isometric to a standard model space: $\mathbb{C}^n$ with its standard flat metric if $c = 0$, $\mathbb{C}P^n(c)$ with its Fubini-Study metric if $c > 0$, or $\mathbb{C}H^n(c)$ with its hyperbolic metric if $c < 0$. Let M be a complex submanifold of $\tilde{M}$. As is well known, this submanifold inherits a Kähler structure from that of $\tilde{M}$. We call M a Kahler-Einstein submanifold if the induced metric on M is Einstein. In this paper we address the problem of classifying what submanifolds of complex space forms are Einstein.

The first person to obtain results in this direction was Calabi in his thesis [C1]. This remarkable work contains a wealth of innovative mathematics, methodically studying isometric embedding in flat and positively curved space forms. As his main tool Calabi introduces the diastasis D_M, a real analytic function, determined by the metric, which is defined on a neighborhood of the diagonal in $M \times M$. The most important property of D_M is its simple behavior with respect to submanifolds: if $M \to \tilde{M}$ is a holomorphic embedding then the diastasis of $\tilde{M}$ restricts to the diastasis of M. This makes the diastasis a tool perfectly suited to the study of complex

submanifolds, and using it Calabi is able to deduce necessary and sufficient conditions for a Kahler manifold to isometrically embed into $\mathbb{C}^n$ or $\mathbb{C}P^n(c)$.

We will elaborate on these results enough to see how Calabi's work settles the problem of classifying Einstein curves isometrically embedded in $\mathbb{C}^n$ or $\mathbb{C}P^n(c)$. Before beginning we first remark that a Kähler-Einstein curve is simply a space form of a single complex dimension.

Let p be a point in some Kähler manifold M, and let $z : U \to \mathbb{C}$ be a complex coordinate system with $z(p) = 0$. Then the diastasis has a power series expansion near (p, p):

$$D_M(0, z(q)) = \sum_{IJ} b_{IJ} z(q)^I \overline{z(q)}^J$$

where I, J are multi-indices. The matrix (b_{IJ}) is hermitian, and one can define its rank. Then, according to Calabi, some neighbourhood of p in M can be isometrically embedded into $\mathbb{C}^n$ if and only if the rank of b_{IJ} is less than or equal to n. Furthermore, if this rank is equal to n, then no neighbourhood can be embedded in $\mathbb{C}^m$ for any $m < n$, and any two embeddings into $\mathbb{C}^n$ differ by a unitary transformation. By examining the diastatic functions of $\mathbb{C}P^1(c)$ and $\mathbb{C}H^1(c)$ one sees that they do not satisfy the criterion above, and hence there is no local isometric embedding of these manifolds into $\mathbb{C}^n$. Therefore the only isometrically embedded Einstein complex curves in $\mathbb{C}^n$ are complex lines.

Similar considerations to those above yield a necessary and sufficient condition for a Kähler manifold to isometrically embed in $\mathbb{C}P^n(c)$. This criterion immediately implies that $\mathbb{C}$ and $\mathbb{C}H^1$ cannot be isometrically embedded into $\mathbb{C}P^n$ even locally. As for $\mathbb{C}P^1$, in the last theorem of his paper Calabi applies his criterion to prove:

Theorem (Calabi): The space $\mathbb{C}P^m(c)$ can be globally embedded into $\mathbb{C}P^n(\tilde{c})$ if and only if the following conditions are satisfied:

(1) $c \leq \tilde{c}$
(2) $\tilde{c} = kc$ for some positive integer k
(3) $n \geq \begin{pmatrix} m+k \\ k \end{pmatrix} - 1$

Taking $\tilde{c} = 1$ and $m = 1$, we find that $\mathbb{C}P^1(c)$ embeds into $\mathbb{C}P^n(1)$ if and only if $c = \frac{1}{k}$ for some k, and $n \geq k$. Looking at this another way, we see that any Einstein curve isometrically embedded in $\mathbb{C}P^n$ must have constant sectional curvature equal to one of the numbers $1, \frac{1}{2}, \ldots, \frac{1}{n}$.

Examples of such curves are easy to construct. Let $[z_1, z_2]$ be the homogeneous coordinates on $\mathbb{C}P^1$. Then the map:

$$[z_1, z_2] \mapsto \left[\sqrt{\binom{n}{0}} z_1^n, \sqrt{\binom{n}{1}} z_1^{n-1} z_2, \cdots, \sqrt{\binom{n}{n}} z_2^n\right]$$

isometrically embeds $\mathbb{C}P^1(\frac{1}{n})$ into $\mathbb{C}P^n(1)$.

The next person to make progress on the problem was Smyth in his thesis [S], who addressed the case of a complete Einstein hypersurface embedded in a space form. Let Q^n denote the smooth algebraic variety in $\mathbb{C}P^{n+1}$ which is defined by the homogeneous polynomial equation:

$$z_0^2 + z_1^2 + \cdots + z_{n+1}^2 = 0$$

which is called a hypersphere. Then Smyth's result is:

Theorem (Smyth): If $n \geq 2$ then:

- $\mathbb{C}P^n$ and the complex hypersphere Q^n are the only complex hypersurfaces in $\mathbb{C}P^{n+1}$ which are complete and Einstein.
- $\mathbb{C}^n$ is the only simply connected complex hypersurface in $\mathbb{C}^{n+1}$.
- $\mathbb{C}H^n$ is the only simply connected complex hypersurface in $\mathbb{C}H^{n+1}$.

Smyth proves his theorem with an in depth analysis of the second fundamental form of a potential Einstein hypersurface. His work is heavily connection theoretic, playing with the curvature tensors of the space form and hypersurface until he can show that any Einstein hypersurface is a symmetric space. Then he calls upon Cartan's classification of Hermitian symmetric spaces to narrow down the possibilities for the submanifold to those in the theorem.

Smyth's work was soon generalized by Chern [Ch], who proved the corresponding local result. Chern used his complete mastery of Cartan's connection form formalism to prove that any Einstein hypersurface in a space form of curvature c must have Einstein constant λ equal to either $\frac{(n+1)c}{2}$ or $\frac{nc}{2}$. In case $\lambda = \frac{(n+1)c}{2}$ it is easy to prove that M must be totally geodesic. It follows from the Gauss equation that when $c < 0$ the $\lambda = \frac{nc}{2}$ case cannot occur. On the other hand, in the case $\lambda = \frac{nc}{2}$ and $c > 0$, Chern gives a geometric proof that M is locally isometric to the hypersphere Q^{n-1}.

More progress was made by Tsukada [T] who generalized Chern's work to submanifolds of codimension 2. Tsukada's theorem is:

Theorem (Tsukada) : Let M be an isometrically embedded Kähler Einstein submanifold of codimension 2 in a complex space form $\tilde{M}(c)$. Then:

- If $c \leq 0$, then M is totally geodesic.

- If $c > 0$, then M is either totally geodesic, or M is a hypersphere Q^{n-2} inside of some totally geodesic hypersurface $\mathbb{C}P^{n-1}(c)$.

Tsukada's argument involves a detailed study of the second fundamental form of M and its first two covariant derivatives. By carefully investigating the linear algebraic properties of these tensors he obtains the result above.

Finally we have the work of Umehara, which is based on the ideas of Calabi. In a sequence of papers [Um1, Um2] Umehara investigates in detail the properties of Calabi's diastatic function.

In his first paper, Umehara studies the structure of the set of real analytic function on a Kähler manifold. More precisely, he introduces the algebra:

$$\Lambda(M) = \text{span}_{\mathbb{R}}\{h\bar{k} + \bar{h}k : h, k \text{ are holomorphic functions on } M\}$$

If M is compact, then $\Lambda(M)$ coincides with the algebra of all real analytic functions on M, and in any case, the structure of $\Lambda(M)$ is intimately connected with the complex geometry of M. While studying $\Lambda(M)$ Umehara is able to put the results of Calabi in a convenient framework, which greatly clarifies ones understanding of the situation. Next, in [Um2], Umehara addresses the general case of an Einstein submanifold of a complex space form. It is easy to write down a condition, involving only the diastasis, for M to be embeddable as an Einstein submanifold of $\tilde{M}(c)$. Using this condition, and his understanding of the structure of $\Lambda(M)$, Umehara succeeds in completely solving the classification problem in the case $c \leq 0$, obtaining the following result:

Theorem (Umehara) : Any Einstein submanifold of $\mathbb{C}^n$ or $\mathbb{C}H^n$ is totally geodesic.

Unfortunately Umehara does not make any progress in the positive curvature case, but even with this drawback, his work is both profound and beautiful.

Reflecting on these results, one immediately sees that there are currently two approaches to the classification problem. The approach of Calabi and Umehara is analytic, and studies the diastatic function of a space form to draw conclusions about the geometry of submanifolds of $\tilde{M}$. The work of Smyth, Chern and Tsukada is purely differential geometric, and uses classical connection theoretic tools to draw conclusions about submanifolds with careful linear algebraic arguments. One wonders what the connection is between these two approaches, and whether results proved with one technique are approachable with the other. Also, one wonders whether either technique has exhausted what it can tell us about the problem - can pure differential geometry be more carefully applied to draw further conclusions, and can the diastasis (and the structure of $\Lambda(M)$) be further investigated

to draw conclusions in the positive curvature case? These are the type of questions we wish to address in this paper.

2. Background on Kahler Geometry

In this section we outline the basic properties of Kahler geometry that we will be using throughout this paper.

2.1. **Invariant Metrics and the Associated 2-form.**

Let M be a complex manifold with complex structure J.

Definition 2.2. A Riemannian metric on M is called J-invariant if $g(JX, JY) = g(X, Y)$ for all $X, Y \in TM$.

Note that any complex manifold can be given a J-invariant Riemannian metric, for we can locally use the standard euclidean metric on $\mathbb{C}^n$, and then assemble one globally using a partition of unity. We denote also by g the complex bilinear extension of the given metric to the complexified tangent bundle $T_{\mathbb{C}}M$.

As usual we consider the local holomorphic coordinate frame fields:

$$Z_i = \frac{\partial}{\partial z_i} = \frac{1}{2}\left(\frac{\partial}{\partial x_i} - i\frac{\partial}{\partial y_i}\right) = \frac{1}{2}(X_i - iY_i)$$

and:

$$\bar{Z}_i = \frac{\partial}{\partial \bar{z}_i} = \frac{1}{2}\left(\frac{\partial}{\partial x_i} + i\frac{\partial}{\partial y_i}\right) = \frac{1}{2}(X_i + iY_i)$$

Then, since g is J-invariant and $Y_i = JX_i$, we have:

$$\begin{aligned} g(Z_i, Z_j) &= g(X_i - iY_i, X_j - iY_j) \\ &= g(X_i, X_j) - ig(X_i, Y_j) - ig(Y_i, X_j) - g(Y_i, Y_j) \\ &= g(X_i, X_j) - ig(X_i, Y_j) + ig(Y_j, X_i) - g(X_i, X_j) \\ &= 0 \end{aligned}$$

Similarly, $g(\bar{Z}_i, \bar{Z}_j) = 0$. We define the symbols of the Riemannian metric g by::

$$g_{ij} = g(Z_i, \bar{Z}_j)$$

then:

$$g(\bar{Z}_i, Z_j) = \overline{g(Z_i, \bar{Z}_j)} = \bar{g}_{ij}$$

and we have the following expression for the metric in a holomorphic coordinate frame field:

$$g = g_{ij} dz_i \otimes d\bar{z}_j + \bar{g}_{ij} d\bar{z}_i \otimes dz_j$$

Remark 2.3. It is easy to now derive an expression for the metric in terms of the original real coordinates. Indeed, it is easy to check that:

$$dz = dx + idy \quad \text{and} \quad d\bar{z} = dx - idy$$

so we get:

$$\begin{aligned} g &= g_{ij} dz_i \otimes d\bar{z}_j + \bar{g}_{ij} d\bar{z}_i \otimes dz_j \\ &= g_{ij}(dx_i + idy_i) \otimes (dx_j - idy_j) + \bar{g}_i j(dx_i - idy_i) \otimes (dx_j + idy_j) \\ &= (g_{ij} + \bar{g}_{ij}) dx_i \otimes dx_j + (-ig_{ij} + i\bar{g}_{ij}) dx_i \otimes dy_j \\ &\quad + (ig_{ij} - i\bar{g}_{ij}) dy_i \otimes dx_j + (g_{ij} + \bar{g}_{ij}) dy_i \otimes dy_j \\ &= 2\mathrm{Re}(g_{ij}) dx_i \otimes dx_j + 2\mathrm{Im}(g_{ij}) dx_i \otimes dy_j \\ &\quad - 2\mathrm{Im}(g_{ij}) dy_i \otimes dx_j + 2\mathrm{Re}(g_{ij}) dy_i \otimes dy_j \end{aligned}$$

which gives an expression for the metric in real coordinates.

Observe that if g is a J-invariant metric, then the equation:

$$\omega(X, Y) = g(JX, Y)$$

defines a real valued 2-form on M. Indeed, by the J-invariance of g, we have:

$$\omega(Y, X) = g(JY, X) = g(J^2 Y, JX) = -g(Y, JX) = -g(JX, Y) = \omega(X, Y)$$

Clearly we can recover g from ω with the formula:

$$g(X, Y) = \omega(X, JY)$$

Again, we also denote by ω the complex bilinear linear extension of this form to the complexified tangent bundle. We then have:

$$\omega(Z_i, Z_j) = g(JZ_i, Z_j) = ig(Z_i, Z_j) = 0$$

and similarly $\omega(\bar{Z}_i, \bar{Z}_j) = 0$. Finally:

$$\omega(Z_i, \bar{Z}_j) = g(JZ_i, \bar{Z}_j) = ig(Z_i, \bar{Z}_j) = ig_{ij}$$

so we get the following local expression for ω in a holomorphic coordinate frame field:

$$\omega = ig_{ij} dz_i \wedge d\bar{z}_j$$

In particular, ω is a $(1,1)$-form. We refer to ω as the 2-form associated to the metric g.

Remark 2.4. While we used a holomorphic coordinate frame field in our calculations, by reviewing the argument above the reader may verify that the results hold in any frame field for the bundle TM^+.

2.5. **The Connection and the Kahler Condition.** Let ∇ denote the Riemannian connection of the metric g, and extend it complex linearly in both components to the complexified tangent bundle. Recall that this connection canonically induces connections on all bundles of tensors associated to $T_{\mathbb{C}}M$, and in particular we can write ∇J since J is a tensor of type $(1,1)$ on $T_{\mathbb{C}}M$. As usual, $\nabla_X J$ is again a $(1,1)$ tensor for any vector field X, and is defined by the equation:

$$(\nabla_X J)(Y) = \nabla_X(JY) - J(\nabla_X Y)$$

With this setup we have the following proposition relating ∇J and $d\omega$:

Proposition 2.6. *Let M be a complex manifold, g a J-invariant Riemannian metric, ω the 2-form associated to g, and let ∇ be the Riemannian connection of g. Then we have:*

$$d\omega(X,Y,Z) = g((\nabla_X J)Y, Z) + g((\nabla_Y J)Z, X) + g((\nabla_Z J)X, Y)$$

$$2g((\nabla_X J)Y, Z) = d\omega(X,Y,Z) - d\omega(X, JY, JZ)$$

Proof. We start by verifying the first formula. It suffices to compute $d\omega(X,Y,Z)$ for X, Y, Z coordinate vectors, and if we use a holomorphic coordinate system, then X, Y, Z, JX, JY and JZ are mutually commuting vector fields (i.e. the Lie bracket of any two of these vector fields vanishes). Then:

$$\begin{aligned}
d\omega(X,Y,Z) &= X\omega(Y,Z) + Y\omega(Z,X) + Z\omega(X,Y) \\
&= Xg(JY,Z) + Yg(JZ,X) + Zg(JX,Y) \\
&= g(\nabla_X(JY), Z) + g(JY, \nabla_X Z) + g(\nabla_Y(JZ), X) \\
&\quad + g(JZ, \nabla_Y X) + g(\nabla_Z(JX), Y) + g(JX, \nabla_Z Y) \\
&= g((\nabla_X J)Y, Z) + g(J(\nabla_X Y), Z) + g(JY, \nabla_X Z) \\
&\quad + g((\nabla_Y J)Z, X) + g(J(\nabla_Y Z), X) + g(JZ, \nabla_Y X) \\
&\quad + g((\nabla_Z J)X, Y) + g(J(\nabla_Z X), Y) + g(JX, \nabla_Z Y)
\end{aligned}$$

Now we simply observe that, since $[X.Z] = 0$ we have:

$$g(JY, \nabla_X Z) = g(JY, \nabla_Z X) = -g(Y, J(\nabla_Z X))$$

and similarly for the terms above ending the two subsequent lines. In summary, six of the terms in the above expression cancel in pairs, and we are left with:

$$d\omega(X,Y,Z) = g((\nabla_X J)Y, Z) + g((\nabla_Y J)Z, X) + g((\nabla_Z J)X, Y)$$

which is what we wanted to show.

To prove the second formula, we write out:

$$\begin{aligned} d\omega(X, JY, JZ) &= X\omega(JY, JZ) + JY\omega(JZ, X) + JZ\omega(X, JY) \\ &= -Xg(Y, JZ) - JYg(Z, X) + JZg(X, Y) \end{aligned}$$

and so the right hand side becomes:

$$\begin{aligned} &d\omega(X, Y, Z) - d\omega(X, JY, JZ) \\ &\qquad = Yg(JZ, X) + Zg(JX, Y) + JYg(Z, X) - JZg(X, Z) \end{aligned}$$

Moving on to the left hand side we first write:

$$\begin{aligned} 2g((\nabla_X J)Y, Z) &= 2g(\nabla_X(JY), Z) - 2g(J(\nabla_X Y), Z) \\ &= 2g(\nabla_X(JY), Z) + 2g(\nabla_X Y, JZ) \end{aligned}$$

and now use the Koszul formula ([CE] page 2). The results are:

$$\begin{aligned} 2g(\nabla_X(JY), Z) &= Xg(JY, Z) + JYg(X, Z) - Zg(X, JY) \\ &= Xg(JY, Z) + JYg(X, Z) + Zg(JX, Y) \end{aligned}$$

and:

$$\begin{aligned} 2g(\nabla_X Y, JZ) &= Xg(Y, JZ) + Yg(X, JZ) - JZg(X, Y) \\ &= -Xg(JY, Z) + Yg(X, JZ) - JZg(X, Y) \end{aligned}$$

Adding these gives us:

$$2g((\nabla_X J)Y, Z) = Yg(JZ, X) + Zg(JX, Y) + JYg(Z, X) - JZg(X, Z)$$

which is what we wanted to show. □

The next proposition is a cornerstone of Kahler geometry:

Proposition 2.7. *Let M be a complex manifold with g a J-invariant Riemannian metric, let ω be the 2-form associated to g, and let the ∇ be Riemannian connection of g. Then the following conditions are equivalent:*

1. $\nabla J = 0$
2. $d\omega = 0$
3. *For every point $p \in M$ there is a smooth real valued function f, defined in a neighbourhood of p, so that $\omega = i\partial\bar{\partial}f$*

Proof. The previous proposition immediately implies that 1) and 2) are equivalent. It is also clear that 3) $\Rightarrow$ 2), since $d = \partial + \bar{\partial}$.

We now show that 2) $\Rightarrow$ 3). Assume that $d\omega = 0$. Now ω is a real valued 2-form on TM, so by the Poincaré lemma there is a real valued 1-form α, defined in a neighbourhood of any given point p, so that $\omega = d\alpha$. Extending

these forms complex linearly to $T_{\mathbb{C}}M$, we can write $\alpha = \beta + \bar{\beta}$, where β is a $(1,0)$-form. Now we have:

$$\omega = d\alpha = \partial\beta + \bar{\partial}\beta + \partial\bar{\beta} + \bar{\partial}\bar{\beta}$$

and since ω is a $(1,1)$-form, it follows that $\partial\beta = \bar{\partial}\bar{\beta} = 0$ (since $\partial\beta$ is a $(2,0)$-form and $\bar{\partial}\bar{\beta}$ is a $(0,2)$-form). It follows, from the Poincaré-Grothendieck lemma, that $\beta = \partial\phi$ for some smooth complex function ϕ defined near p. Assembling all of this information gives us:

$$\omega = \bar{\partial}\beta + \partial\bar{\beta} = \bar{\partial}\partial\phi + \partial\bar{\partial}\bar{\phi} = \partial\bar{\partial}(\bar{\phi} - \phi)$$

So setting $f = i(\bar{\phi} - \phi)$ does the trick (note that f so defined is real valued because $\bar{\phi} - \phi$ is imaginary valued). □

Now we are prepared to make the following important definition:

Definition 2.8. Let M be a complex manifold with a J-invariant Riemannian metric. If any of the three equivalent conditions in the previous proposition are satisfied, then M is called a Kahler manifold, g a Kahler metric, and ω the Kahler form associated to the metric. Further, any function f which locally satisfies the condition 3) is called a Kahler potential for the metric.

One of the great advantages of Kahler geometry over Riemannian geometry is the simplification that comes about during computations. We now investigate how this simplification arises, and derive formulas for the Christoffel symbols of a Kahler manifold.

Let:

$$Z_i = \frac{\partial}{\partial z_i} = \frac{1}{2}\left(\frac{\partial}{\partial x_i} - i\frac{\partial}{\partial y_i}\right) = \frac{1}{2}\left(X_i - iY_i\right)$$

be some local holomorphic coordinate frame field for the bundle TM^+, with $\bar{Z}_i$ the corresponding frame field for the bundle TM^-. Then, using the symmetry of the connection, the fact that all of the relevant brackets vanish, and the fact that J is parallel, we have:

$$\begin{aligned}
\nabla_{Z_i}\bar{Z}_j &= \frac{1}{4}\nabla_{(X_i - iJX_i)}(X_j + iJX_j) \\
&= \frac{1}{4}(\nabla_{X_i}X_j + i\nabla_{X_i}(JX_j) - i\nabla_{JX_i}X_j + \nabla_{JX_i}(JX_j)) \\
&= \frac{1}{4}(\nabla_{X_i}X_j + iJ\nabla_{X_i}X_j - iJ\nabla_{X_j}X_i - \nabla_{X_j}X_i) \\
&= 0
\end{aligned}$$

and consequently $\nabla_{\bar{Z}_i}Z_j = \overline{\nabla_{Z_i}\bar{Z}_j} = 0$ as well. Finally, we note that, for any complex vector field X on M:

$$J\nabla_X Z_i = \nabla_X(JZ_i) = \nabla_X(iZ_i) = i\nabla_X Z_i$$

so that $\nabla_X Z_j \in TM^+$. Saying this in another way: on a Kahler manifold M the subbundle TM^+ of $T_{\mathbb{C}}M$ is parallel with respect to the Riemannian connection of a Kahler metric. In exactly the same way, TM^- is also a parallel subbundle of $T_{\mathbb{C}}M$.

Now an important question must be addressed. There is a canonical bundle isomorphism $\Phi : TM \to TM^+$ which sends the complex structure J on TM to multiplication by i on TM^+. This isomorphism is given by:

$$X \mapsto \frac{1}{2}(X - iJX)$$

which we abbreviate as $X \mapsto Z$. Extending the Riemannian connection ∇ on TM to $T_{\mathbb{C}}M$, and then restricting it to the parallel subbundle TM^+, we get a connection on TM^+, and it is important for us to be aware of the relationship between these two objects. Thankfully, the best of all possible results holds:

Proposition 2.9. *Let X and Y be any two vector fields on TM, then $\Phi(\nabla_X Y) = \nabla_{\Phi X}\Phi Y$. That is, the connection induced on TM^+ by the isomorphism Φ agrees with the one obtained by first extending ∇ to be complex bilinear and then restricting it to TM^+.*

Proof. It suffices to verify this in a holomorphic coordinate frame field, so let $Z_i = \frac{1}{2}(X_i - iY_i)$ be such a field. Note that the inverse of the isomorphism Φ is given by $Z \mapsto Z + \bar{Z}$. With this in mind, we compute:

$$\begin{aligned}\nabla_{X_i} X_j &= \nabla_{Z_i + \bar{Z}_i}(Z_j + \bar{Z}_j) \\ &= \nabla_{Z_i} Z_j + \overline{\nabla_{Z_i} Z_j} \\ &= \Phi^{-1}(\nabla_{Z_i} Z_j)\end{aligned}$$

so:

$$\Phi(\nabla_{X_i} X_j) = \nabla_{Z_i} Z_j = \nabla_{\Phi X_i}\Phi X_j$$

Continuing in this manner:

$$\begin{aligned}\nabla_{Y_i} Y_j &= \nabla_{i(\bar{Z}_i - Z_i)}(i(\bar{Z}_j - Z_j)) \\ &= -(\nabla_{Z_i} Z_j + \overline{\nabla_{Z_i} Z_j}) \\ &= -\Phi^{-1}(\nabla_{Z_i} Z_j)\end{aligned}$$

so:

$$\Phi(\nabla_{Y_i} Y_j) = -\nabla_{Z_i} Z_j = \nabla_{iZ_i}(iZ_j) = \nabla_{\Phi Y_i}\Phi Y_j$$

For the mixed terms we use the fact, alluded to earlier, that Φ sends J to multiplication by i. For:

$$\begin{aligned}\Phi(\nabla_{X_i} Y_j) = \Phi(\nabla_{X_i} JX_j) = i\Phi(\nabla_{X_i} X_j) = i\nabla_{\Phi X_i}\Phi X_j \\ = \nabla_{\Phi X_i} i\Phi X_j = \nabla_{\Phi X_i}\Phi Y_j\end{aligned}$$

The computation involving the other mixed term is entirely analogous. □

Now we investigate the local expression of ∇ on a Kahler manifold. Again, let Z_i denote and holomorphic coordinate frame field for M. Since TM^+ is a parallel subbundle of $T_{\mathbb{C}}M$, and $\nabla_{\bar{Z}_i} Z_j = 0$, we define the (complex) Christoffel symbols Γ_{ij}^k of ∇ by:[1]

$$\nabla_{Z_i} Z_j = \Gamma_{ij}^k Z_k$$

and observe that we then also have:

$$\nabla_{\bar{Z}_i} \bar{Z}_j = \bar{\Gamma}_{ij}^k \bar{Z}_k$$

We would like to derive an expression for these symbols in terms of the symbols of the Riemannian metric, and this is very easy to do. Notice that:

$$\Gamma_{ij}^k g_{kl} = g(\nabla_{Z_i} Z_j, \bar{Z}_l) = Z_i g(Z_j, \bar{Z}_l) = \frac{\partial g_{jl}}{\partial z_i}$$

and therefore:

$$\Gamma_{ij}^k = \Gamma_{ij}^r g_{rl} g^{lk} = \frac{\partial g_{jl}}{\partial z_i} g^{lk}$$

This equation expresses the Christoffel symbols (and hence the connection) in a holomorphic coordinate system. Comparing this to the analogous formula from Riemannian geometry will convince the reader that a great simplification has occurred upon narrowing our focus to Kahler manifolds.

2.10. **The Curvature.** Now let's move on to the Riemannian curvature. We remind the reader that the Riemannian curvature is a $(3,1)$-tensor that is defined, for $X, Y, Z \in TM$, by:

$$R(X,Y)Z = \nabla_X \nabla_Y Z - \nabla_Y \nabla_X Z - \nabla_{[X,Y]} Z$$

As has become commonplace, we extend this to be $\mathbb{C}$ bilinear (more honestly, R inherits its complex bi-linearity from ∇). Then, if Z_i is a holomorphic coordinate frame field for TM^+, we have (by arguments entirely analogous to those used for ∇):

$$R(Z_i, Z_j)Z_k = 0$$

and:

$$R(\bar{Z}_i, \bar{Z}_j)\bar{Z}_k = \overline{R(Z_i, Z_j)Z_k} = 0$$

Note also that $JR = RJ$, since J is parallel, and hence we have:

$$JR(\bar{Z}_i, Z_j)Z_k = R(\bar{Z}_i, Z_j)(JZ_k) = R(\bar{Z}_i, Z_j)(iZ_k) = iR(\bar{Z}_i, Z_j)Z_k$$

that is, $R(\bar{Z}_i, Z_j)Z_k \in TM^+$.

[1]The summation convention on repeated indices will always be in effect

Of course, R satisfies all of the usual symmetries of a curvature tensor, and in Kahler case additional symmetries appear. Let X, Y, Z, W be any complex vector fields on a Kahler manifold. Then:

$$\begin{aligned} g(R(X,Y)JZ, JW) &= g(JR(X,Y)Z, JW) \\ &= g(R(X,Y)Z, W) \end{aligned}$$

and consequently we also have:

$$\begin{aligned} g(R(JX,JY)Z, W) &= g(R(Z,W)JX, JY) \\ &= g(R(Z,W)X, Y) \\ &= g(R(X,Y)Z, W) \end{aligned}$$

Now let's take a look at the local expression for the curvature tensor on a Kahler manifold. As is the case for the connection, the resulting formulas are drastically more simple than their Riemannian counterparts. Let Z_i be any holomorphic coordinate frame field for TM^+, and define the complex curvature symbols by:

$$R(\bar{Z}_i, Z_j)Z_k = R^l_{ijk} Z_l$$

It is easy to derive a local expression for these curvature symbols. Indeed:

$$\begin{aligned} R(\bar{Z}_i, Z_j)Z_k &= \nabla_{\bar{Z}_i} \nabla_{Z_j} Z_k - \nabla_{Z_j} \nabla_{\bar{Z}_i} Z_k \\ &= \nabla_{\bar{Z}_i} \left(\Gamma^l_{jk} Z_l \right) \\ &= \frac{\partial \Gamma^l_{jk}}{\partial \bar{z}_i} Z_l + \Gamma^l_{jk} \nabla_{\bar{Z}_i} Z_l \\ &= \frac{\partial \Gamma^l_{jk}}{\partial \bar{z}_i} Z_l \end{aligned}$$

so we conclude that:

$$R^l_{ijk} = \frac{\partial \Gamma^l_{jk}}{\partial \bar{z}_i}$$

Definition 2.11. Let M be a Kahler manifold, and let $X \in TM$. The holomorphic sectional curvature of M at X is defined to be:

$$K(X) = \frac{g(R(X, JX)JX, X)}{g(X, X)^2}$$

that is, the sectional curvature of the 2-plane spanned by X and JX.

In a later section we will need to know how to compute the holomorphic sectional curvature using a holomorphic coordinate frame. With this in mind we offer:

Lemma 2.12. *Given $X \in TM$ write $Z = \frac{1}{2}(X - iJX)$. Then:*

$$K(X) = \frac{g(R(X, JX)JX, X)}{g(X, X)^2} = -\frac{g(R(Z, \bar{Z})\bar{Z}, Z)}{g(Z, \bar{Z})^2}$$

Proof. Observe that $X = Z + \bar{Z}$ and $JX = i(\bar{Z} - Z)$. Now we may simply compute:

$$R(X, JX) = R(Z + \bar{Z}, i(\bar{Z} - Z)) = -i(R(\bar{Z}, Z) - R(Z, \bar{Z})) = -2iR(\bar{Z}, Z)$$

Therefore:

$$g(R(X, JX)JX, X) = (-2i)^2 g(R(\bar{Z}, Z)Z, \bar{Z}) = -4g(R(Z, \bar{Z})\bar{Z}, Z)$$

On the other hand:

$$g(X, X) = g(Z + \bar{Z}, Z + \bar{Z}) = 2g(Z, \bar{Z})$$

Putting these two computations together yields the result we are after. □

Definition 2.13. A Kahler manifold is said to have constant holomorphic sectional curvature c when $K(X) = c$ for all $X \in TM$.

In the next section we will give examples of Kahler manifolds with constant holomorphic sectional curvature. Until then, we offer the reader the following characterization.

Proposition 2.14. *A Kahler manifold has constant holomorphic sectional curvature equal to c if and only if its curvature tensor is given by:*

$$g(R(X, Y)Z, W) = \frac{c}{4}(g(X, W)g(Y, Z) - g(X, Z)g(Y, W) + g(X, JW)g(Y, JZ) - g(X, JZ)g(Y, JW) + 2g(X, JY)g(W, JZ))$$

Proof. Let's use the notation:

$$R(X, Y, Z, W) = g(R(X, Y)Z, W)$$

for the curvature tensor, and write $\mathcal{R}(X, Y, Z, W)$ for the tensor defined by the right hand side of the equation we are trying to verify. The reader may directly check that $\mathcal{R}$ satisfies all of the symmetry properties of the Riemannian curvature tensor[2]. That is:

$$\mathcal{R}(X, Y, Z, W) = -\mathcal{R}(Y, X, Z, W)$$
$$\mathcal{R}(X, Y, Z, W) = -\mathcal{R}(X, Y, W, Z)$$
$$\mathcal{R}(X, Y, Z, W) = \mathcal{R}(Z, W, X, Y)$$

and the Bianchi identity holds as well:

$$\mathcal{R}(X, Y, Z, W) + \mathcal{R}(Y, Z, X, W) + \mathcal{R}(Z, X, Y, W) = 0$$

[2]we call a tensor with these symmetries an algebraic curvature tensor

Furthermore, $\mathcal{R}$ satisfies the additional Kahler symmetry properties:

$$\mathcal{R}(JX, JY, Z, W) = \mathcal{R}(X, Y, JZ, JW) = \mathcal{R}(X, Y, Z, W)$$

Let's now consider the tensor $Q = R - \mathcal{R}$, which is also an algebraic curvature tensor, and additionally satisfies the Kahler symmetry properties we highlighted earlier. Upon considering the following calculation:

$$\begin{aligned}\mathcal{R}(X, JX, JX, X) &= \frac{c}{4}(g(X, X)g(JX, JX) - g(X, JX)^2 + g(X, JX)^2 \\ &\quad + g(X, X)g(JX, JX) + 2g(X, X)^2) \\ &= cg(X, X)^2\end{aligned}$$

we see that M has constant holomorphic sectional curvature c if and only if $Q(X, JX, JX, X) = 0$ for all $X \in TM$. Consequently, to prove the proposition, we need to show that $Q = 0$ if and only if $Q(X, JX, JX, X) = 0$ for all X. We begin by polarizing the equation $Q(X, JX, JX, X) = 0$:

$$\begin{aligned}0 &= Q(X + Y, JX + JY, JX + JY, X + Y) \\ &= 4Q(X, JX, Y, JX) + 2Q(X, Y, X, Y) + 6Q(X, JY, X, JY) \\ &\quad + 4Q(Y, JY, Y, JX)\end{aligned}$$

Although we have suppressed much of the computation to preserve readability, the symmetry properties of an algebraic curvature tensor and the Kahler symmetry properties were used to simplify the results of the computation above. Now let's replace X by tX in the equation above, the result is a cubic polynomial in t which must vanish identically, and hence the coefficient of t^2 in this polynomial must be zero. That is:

$$2Q(X, Y, X, Y) + 6Q(X, JY, X, JY) = 0$$

for all $X, Y \in TM$. We now apply this last equation twice:

$$Q(X, Y, X, Y) = -3Q(X, JY, X, JY) = 9Q(X, Y, X, Y)$$

It follows immediately that $Q(X, Y, Y, X) = 0$ for all $X, Y \in TM$.

To complete the argument, we cite the well known fact that an algebraic curvature tensor is completely determined by the values of $Q(X, Y, X, Y)$ (any reader who doubts this is invited to consult any textbook on Riemannian geometry, for example, [dC], page 94). We immediately conclude that $Q = 0$, and hence the argument is completed. □

Corollary 2.15. *A Kahler manifold has constant holomorphic sectional curvature if and only if its curvature tensor is given by:*

$$\begin{aligned}R(X, Y)Z = \frac{c}{4}&(g(Y, Z)X - g(X, Z)Y - g(Y, JZ)JX \\ &+ g(X, JZ)JY + 2g(X, JY)JZ)\end{aligned}$$

Proof. Simply use the J-invariance of g to pass all the J's off of the W's in the formula from the previous proposition. □

The above expression for the curvature tensor in manifolds of constant holomorphic sectional curvature leads to the following important corollary:

Corollary 2.16. *Any two Kahler manifolds with the same constant holomorphic sectional curvature are locally isometric, and if they are both simply connected, then they are globally isometric.*

Proof. This follows immediately from the Cartan-Ambrose-Hicks theorem (see [CE] page 31), using the expression for the curvature tensor in the previous corollary and the fact that g and J are invariant under parallel translation. □

2.17. **The Ricci Curvature and Kahler-Einstein Manifolds.** We now want to introduce one of the main objects of study in this paper, Kahler-Einstein manifolds. To begin, we give the important definition:

Definition 2.18. The Ricci tensor of a Kahler manifold is the complex valued 2-tensor on M defined by:

$$\mathrm{Ric}(X,Y) = \mathrm{tr}(Z \mapsto R(Z,X)Y)$$

for $X, Y \in T_{\mathbb{C}}M$.

As in the Riemannian case, the Ricci tensor is symmetric. Therefore, there is an associated $\mathbb{C}$-linear map $T_{\mathbb{C}}M \to T_{\mathbb{C}}M$, which we also call Ric, defined by:

$$g(\mathrm{Ric}(X), Y) = \mathrm{Ric}(X,Y)$$

The Kahler environment lends itself to a convenient expression for this map.

Proposition 2.19. *Suppose that $\{X_i, JX_i\}$, for $i = 1, \ldots, n$, is an orthonormal frame for the tangent space T_PM. Then, for $X \in T_pM$, we have:*

$$Ric(X) = \sum_i R(X_i, JX_i)JX$$

Proof. We compute, using the usual summation convention to avoid excessive use of Σ notation:

$$\begin{aligned} g(\mathrm{Ric}(X), Y) &= \mathrm{Ric}(X, Y) \\ &= g(R(X_i, X)Y, X_i) + g(R(JX_i, X)Y, JX_i) \\ &= -g(R(X_i, X)JX_i, JY) - g(R(JX_i, X)JX_i, Y) \\ &= g(R(X, X_i)JX_i, JY) + g(R(JX_i, X)X_i, JY) \\ &= -g(R(X_i, JX_i)X, JY) \\ &= g(R(X_i, JX_i)JX, Y) \end{aligned}$$

We have used the Bianchi identity in the fourth line of the computation above. □

As a consequence of the above proposition we deduce the J-invariance of the Ricci tensor, for:

$$\begin{aligned} \mathrm{Ric}(JX, JY) &= g(\mathrm{Ric}(JX), JY) \\ &= g(R(X_i, JX_i)J^2X, JY) \\ &= g(R(X_i, JX_i)JX, Y) \\ &= \mathrm{Ric}(X, Y) \end{aligned}$$

Consequently, we can define an associated $(1,1)$-form, the Ricci form, by:

$$\mathrm{ric}(X, Y) = \mathrm{Ric}(JX, Y)$$

Let's once again investigate how all of this looks in a holomorphic local coordinate system. Since Ric is $\mathbb{C}$-bilinear and symmetric, we can repeat the proof we gave for g to show that:

$$\mathrm{Ric}(Z_i, Z_j) = \mathrm{Ric}(\bar{Z}_i, \bar{Z}_j) = 0$$

So we set:

$$\mathrm{Ric}_{ij} = \mathrm{Ric}(Z_i, \bar{Z}_j)$$

Before we begin deriving a local expression for these symbols, we make a small preliminary calculation. First note that the trace used in defining the ricci curvature can be calculated by viewing R as a real linear tensor on TM or a complex linear tensor on $T_{\mathbb{C}}M$. Therefore, if we let Z_i be a local holomorphic coordinate frame field, so that $\{Z_i, \bar{Z}_i\}$ is a frame field for $T_{\mathbb{C}}M$, and $\{Z^i, \bar{Z}^i\}$ is a frame field for $T^*_{\mathbb{C}}M$, we have:

$$\begin{aligned} \mathrm{Ric}(Z_i, \bar{Z}_j) &= \mathrm{tr}(Z \to R(Z, Z_i)\bar{Z}_j) \\ &= Z^k(R(Z_k, Z_i)\bar{Z}_j) + \bar{Z}^k(R(\bar{Z}_k, Z_i)\bar{Z}_j) \end{aligned}$$

Because $R(Z_r, Z_i) = 0$, the first term above is zero. As for the other, playing around with the curvature tensor gives us:

$$\begin{aligned} R(\bar{Z}_k, Z_i)\bar{Z}_j &= -\overline{R(\bar{Z}_i, Z_k)Z_j} \\ &= -\overline{R^r_{ikj} Z_r} \\ &= -\bar{R}^r_{ikj}\bar{Z}_r \end{aligned}$$

so $\bar{Z}^k(R(\bar{Z}_k, Z_i), \bar{Z}_j) = -\bar{R}^k_{ikj}$. Therefore:

$$\mathrm{Ric}(Z_i, \bar{Z}_j) = \mathrm{tr}(Z \to (Z, Z_i)\bar{Z}_j) = -\bar{R}^k_{ikj} = -\frac{\partial \bar{\Gamma}^k_{kj}}{\partial z_i}$$

Setting $G = (g_{ij})$, we have:

$$\Gamma^k_{kj} = \Gamma^k_{jk} = \frac{\partial g_{kl}}{\partial z_j} g^{lk} = \mathrm{tr}\left(\frac{\partial G}{\partial z_j} G^{-1}\right) = \frac{\partial}{\partial z_j} \log\det G$$

where in the last equality we have used a classic formula from linear algebra. Now observe that, since $G^t = \bar{G}$, we have:

$$\det G = \det G^t = \det \bar{G} = \overline{\det G}$$

so $\det G$ is real valued. Putting this all together, we obtain:

$$\mathrm{Ric}(Z_i, \bar{Z}_j) = -\frac{\partial \bar{\Gamma}^k_{kj}}{\partial z_i} = -\frac{\partial}{\partial z_i}\frac{\partial}{\partial \bar{z}_j} \log\det G$$

that is:

$$\mathrm{Ric}_{ij} = -\frac{\partial^2}{\partial z_i \partial \bar{z}_j} \log\det G$$

This formula becomes particularly simple in the context of the Ricci form, for:

$$\mathrm{ric} = i\mathrm{Ric}_{ij} dz_j \wedge d\bar{z}_i = -i\partial\bar{\partial} \log\det G$$

from which it follows immediately that the Ricci form is a closed $(1,1)$-form.

Definition 2.20. A Kahler manifold is called Kahler-Einstein if there is a constant λ so that $\mathrm{ric} = \lambda\omega$, or equivalently, $\mathrm{Ric} = \lambda g$. The constant λ is called the Einstein constant of M.

Clearly an Einstein constant is a real number.

Proposition 2.21. *A Kahler manifold, of complex dimension n, with constant holomorphic sectional curvature c, is a Kahler-Einstein manifold with Einstein constant $\frac{(n+1)c}{2}$.*

Proof. We use the characterization of metrics of constant holomorphic sectional curvature from a previous proposition. Let $e_1, e_2, \ldots, e_{2n}$ be an orthonormal frame for some tangent space to M, and let X, Y be any two

tangent vectors at this same point. Then we have:

$$\begin{aligned}\mathrm{Ric}(X,Y) &= R(X,e_i,e_i,Y)\\ &= \frac{c}{4}(g(X,Y)g(e_i,e_i) - g(X,e_i)g(Y,e_i) + g(X,JY)g(e_i,Je_i)\\ &\quad - g(X,Je_i)g(e_i,JY) + 2g(X,Je_i)g(Y,Je_i))\\ &= \frac{c}{4}(2ng(X,Y) - g(X,Y) + g(X,Y) + 2g(X,Y))\\ &= \frac{(n+1)c}{2}g(X,Y)\end{aligned}$$

where we have used that $g(X,e_i)g(e_i,Y) = g(X,Y)$ in the computation. □

3. Kahler Manifolds with Constant Holomorphic Sectional Curvature

In this section we will discuss the various examples of simply connected Kahler manifolds of constant holomorphic sectional curvature.

3.1. **Flat Kahler Manifolds.** Consider the complex coordinate space $\mathbb{C}^n$, given a metric according to:

$$g = dz_i \otimes d\bar{z}_i + d\bar{z}_i \otimes dz_i$$

This is clearly a J-invariant metric, and the associated 2-form is given by:

$$\omega = i dz_i \wedge d\bar{z}_i$$

which can be written as:

$$\omega = i\partial\bar{\partial}|z|^2$$

Hence g is a Kahler metric arising from the global potential function $|z|^2$. We see immediately that in the standard global coordinate system:

$$g_{ij} = \delta^i_j \text{ and } g^{ij} = \delta^i_j$$

so:

$$\Gamma^k_{ij} = 0 \text{ and } R^l_{ijk} = 0$$

from which is follows that g has constant holomorphic sectional curvature zero.

3.2. **The Fubini-Study Metric.** Recall that the complex projective space $\mathbb{C}P^n$ can be viewed as the collection of all complex lines in $\mathbb{C}^{n+1}$. From this point of view there is clearly a natural complex n-plane bundle $E \to \mathbb{C}P^n$ whose total space is:

$$E = \{(p,v) : p \in \mathbb{C}P^n, v \in p\} \subset \mathbb{C}P^n \times \mathbb{C}^{n+1}$$

The reader will find it routine to check that this is a holomorphic vector bundle.

Take a holomorphic local section $S : \mathbb{C}P^n \to E$, and consider the locally defined, closed, $(1,1)$-form on $\mathbb{C}P^n$, given by:

$$\omega = i\partial\bar{\partial}\log(S\bar{S}^t)$$

In fact, the form ω does not depend on the choice of local section S. Any other local section T is related to S by the formula $T = \alpha S$, where $\alpha : \mathbb{C}P^n \to \mathbb{C}$ is some locally defined holomorphic function. Therefore, since α is holomorphic, and $\bar{\alpha}$ is anti-holomorphic:

$$\begin{aligned}
\omega' &= i\partial\bar{\partial}\log\det(T\bar{T}^t) \\
&= i\partial\bar{\partial}\log(\alpha S\bar{S}^t\bar{\alpha}) \\
&= i\partial\bar{\partial}(\log(\alpha) + \log(S\bar{S}^t) + \log(\bar{\alpha})) \\
&= i\partial\bar{\partial}\log\det(\bar{S}^t S) \\
&= \omega
\end{aligned}$$

We conclude that ω is a globally defined $(1,1)$-form on $\mathbb{C}P^n$.

Definition 3.3. The Fubini-Study metric on $\mathbb{C}P^n$ is given by $g(X,Y) = \omega(X, JY)$.

We now want to verify that the Fubini-Study metric is Kahler. Note that $d\omega = 0$ by definition, so we only need to verify that g is positive definite and J-invariant.

To begin, recall that there is a transitive action of the unitary group U_{n+1} on $\mathbb{C}P^n$; this action is induced from the standard action of the unitary group on $\mathbb{C}^{n+1}$. For a given $A \in U_{n+1}$ we write $f_A : \mathbb{C}P^n \to \mathbb{C}P^n$ for this induced map, which is easily verified to be holomorphic. We claim that f_A preserves the Fubini-Study metric, that is, $f_A^*\omega = \omega$. To see this, again let S be a holomorphic local section of E, say over some open set U. Then $T = (S \circ f_A)A^{-1}$ is another such section, this one over the open set $f_A^{-1}(U)$. Within this framework, we can calculate:

$$\begin{aligned}
f_A^*\omega &= i\partial\bar{\partial}f_A^*\log\det(S\bar{S}^t) \\
&= i\partial\bar{\partial}\log\det((S \circ f_A)(\bar{S}^t \circ f_A)) \\
&= i\partial\bar{\partial}\log(((S \circ f_A)A^{-1})(A(\bar{S}^t \circ f_A))) \\
&= i\partial\bar{\partial}\log(T\bar{T}^t) \\
&= \omega
\end{aligned}$$

where we have used the fact that A is a unitary matrix in the third line.

Because the action above is transitive, holomorphic, and preserves the action of J, we only need to check that g is positive definite and J-invariant at one

point. To this end, let's find a local expression for g at the point $[1, 0, \ldots, 0]$ in $\mathbb{C}P^n$. As usual, we consider the open neighbourhood:

$$U = \{[z_0, z_1, \ldots, z_n] : z_0 \neq 0\}$$

which (after normalization) supports the coordinate function:

$$[1, z_1, \ldots, z_n] \mapsto (z_1, \ldots, z_n)$$

Clearly then, a holomorphic section over U is:

$$[1, z_1, \ldots, z_n] \mapsto (1, z_1, \ldots, z_n)$$

and hence we have:

$$\begin{aligned}
\omega &= i\partial\bar{\partial}\log(1 + |z_1|^2 + \cdots + |z_n|^2) \\
&= i\partial\left(\frac{z_i d\bar{z}_i}{1 + |z|^2}\right) \\
&= i\frac{(1 + |z|^2)dz_i \wedge d\bar{z}_i - \bar{z}_j dz_j \wedge z_i d\bar{z}_i}{(1 + |z|^2)^2} \\
&= i\frac{(1 + |z|^2)\delta^i_j - \bar{z}_i z_j}{(1 + |z|^2)^2} dz_i \wedge d\bar{z}_j
\end{aligned}$$

where we have used the abbreviation $|z|^2 = \sum z_i \bar{z}_i$ for convenience. Evaluating this expression at $z = 0$ gives us:

$$\omega|_{z=0} = i\delta^i_j dz_i \wedge d\bar{z}_j = i dz_i \wedge d\bar{z}_i$$

and therefore:

$$g|_{z=0} = dz_i \otimes d\bar{z}_i + d\bar{z}_i \otimes dz_i$$

which shows that g is positive definite and J-invariant at $z = 0$. According to our observation above, it follows that g is J-invariant and positive definite everywhere.

Now we would like to continue the discussion and show that the Fubini-Study metric has constant holomorphic sectional curvature. To begin, we calculate the Christoffel symbols of g in the standard homogeneous coordinate system.

Recall that in a holomorphic coordinate frame field on a Kahler manifold we have:

$$\nabla_{Z_i} Z_j = \Gamma^k_{ij} Z_k \text{ and } \nabla_{\bar{Z}_i} \bar{Z}_j = \bar{\Gamma}^k_{ij} \bar{Z}_k$$

with all the mixed covariant derivatives vanishing. The Christoffel symbols are given by the formula:

$$\Gamma^k_{ij} = \frac{\partial g_{jr}}{\partial z_i} g^{rk}$$

We have seen that the Fubini-Study metric is given in the usual coordinate frame by:

$$g_{ij} = \frac{(1 + |z|^2)\delta^i_j - \bar{z}_i z_j}{(1 + |z|^2)^2}$$

and it is easy to check that:

$$g^{ij} = (1+|z|^2)(\delta^i_j + \bar{z}_i z_j)$$

We begin by calculating the partial derivative that appears in the formula for the Christoffel symbols:

$$\begin{aligned}\frac{\partial g_{jr}}{\partial z_i} &= \frac{\partial}{\partial z_i}\left(\frac{(1+|z|^2)\delta^j_r - \bar{z}_j z_r}{(1+|z|^2)^2}\right) \\ &= \frac{(\delta^j_r \bar{z}_i - \delta^r_i \bar{z}_j)(1+|z|^2)^2 - 2(1+|z|^2)\bar{z}_i((1+|z|^2)\delta^j_r - \bar{z}_j z_r)}{(1+|z|^2)^4} \\ &= \frac{\delta^j_r \bar{z}_i - \delta^r_i \bar{z}_j}{(1+|z|^2)^2} - \frac{2\bar{z}_i \delta^j_r}{(1+|z|^2)^2} + \frac{2\bar{z}_i \bar{z}_j z_r}{(1+|z|^2)^3}\end{aligned}$$

Focusing on these three terms one by one, we have, for the first:

$$\begin{aligned}\frac{\delta^j_r \bar{z}_i - \delta^r_i \bar{z}_j}{(1+|z|^2)^2} g^{rk} &= \frac{\delta^j_r \bar{z}_i - \delta^r_i \bar{z}_j}{(1+|z|^2)^2}(1+|z|^2)(\delta^r_k + \bar{z}_r z_k) \\ &= \frac{1}{1+|z|^2}(\delta^j_r \delta^r_k \bar{z}_i - \delta^r_i \delta^r_k \bar{z}_j + \delta^j_r \bar{z}_r \bar{z}_i z_k - \delta^r_i \bar{z}_r \bar{z}_j z_k) \\ &= \frac{\delta^j_k \bar{z}_i - \delta^i_k \bar{z}_j}{1+|z|^2}\end{aligned}$$

for the second:

$$\begin{aligned}-\frac{2\bar{z}_i \delta^j_r}{(1+|z|^2)^2} g^{rk} &= -\frac{2\bar{z}_i \delta^j_r}{(1+|z|^2)^2}(1+|z|^2)(\delta^r_k + \bar{z}_r z_k) \\ &= -\frac{2\delta^j_k \bar{z}_i + 2\bar{z}_i \bar{z}_j z_k}{1+|z|^2}\end{aligned}$$

and finally, for the third:

$$\begin{aligned}\frac{2\bar{z}_i \bar{z}_j z_r}{(1+|z|^2)^3} g^{rk} &= \frac{2\bar{z}_i \bar{z}_j z_r}{(1+|z|^2)^3}(1+|z|^2)(\delta^r_k + \bar{z}_r z_k) \\ &= \frac{2}{(1+|z|^2)^2}(\delta^r_k \bar{z}_i \bar{z}_j z_r + \bar{z}_i \bar{z}_j z_k z_r \bar{z}_r) \\ &= \frac{2}{(1+|z|^2)^2}(\delta^r_k \bar{z}_i \bar{z}_j z_r + \bar{z}_i \bar{z}_j z_k |z|^2) \\ &= \frac{2}{(1+|z|^2)^2}(1+|z|^2)\bar{z}_i \bar{z}_j z_k \\ &= \frac{2\bar{z}_i \bar{z}_j z_k}{1+|z|^2}\end{aligned}$$

Putting all these ingredients together, we find that the Christoffel symbols are given by:

$$\Gamma^k_{ij} = -\frac{\delta^k_i \bar{z}_j + \delta^k_j \bar{z}_i}{1 + |z|^2}$$

We move on to the curvature tensor. Recall that in a holomorphic coordinate frame we have:

$$R(Z_i, Z_j)Z_l = 0$$

so that the curvature symbols are defined by the equation:

$$R(\bar{Z}_i, Z_j)Z_l = R^r_{ijl} Z_r$$

In the previous chapter we showed that the curvature symbols are given by the following formula:

$$R^r_{ijl} = \frac{\partial \Gamma^r_{jl}}{\partial \bar{z}_i}$$

In the case of the Fubini-Study metric, in the standard homogeneous coordinate frame, then have:

$$\begin{aligned} R^r_{ijl} &= \frac{\partial}{\partial \bar{z}_i}\left(-\frac{\delta^k_i \bar{z}_j + \delta^k_j \bar{z}_i}{1 + |z|^2}\right) \\ &= -\frac{(\delta^r_j \delta^i_l + \delta^r_l \delta^j_i)(1 + |z|^2) - z_i(\delta^r_j \bar{z}_l + \delta^r_l \bar{z}_j)}{(1 + |z|^2)^2} \end{aligned}$$

so:

$$R\left(\frac{\partial}{\partial \bar{z}_i}, \frac{\partial}{\partial z_j}\right)\frac{\partial}{\partial z_l} = -\frac{\delta^i_l(1 + |z|^2) - z_i \bar{z}_l}{(1 + |z|^2)^2}\frac{\partial}{\partial z_j} - \frac{\delta^i_j(1 + |z|^2) - z_i \bar{z}_j}{(1 + |z|^2)^2}\frac{\partial}{\partial z_l}$$

Again, because U_{n+1} is a transitive group of isometries of the Fubini-Study metric, we only need to calculate the holomorphic curvature at a single point. Evaluating the above expression for the curvature at $z = 0$, we get:

$$R\left(\frac{\partial}{\partial \bar{z}_i}, \frac{\partial}{\partial z_j}\right)\frac{\partial}{\partial z_l}\bigg|_{z=0} = -\delta^i_l \frac{\partial}{\partial z_j} - \delta^i_j \frac{\partial}{\partial z_l}$$

Therefore we have:

$$\begin{aligned} g\left(R\left(\frac{\partial}{\partial x_i}, \frac{\partial}{\partial y_i}\right)\frac{\partial}{\partial y_i}, \frac{\partial}{\partial x_i}\right) &= -4g\left(R\left(\frac{\partial}{\partial \bar{z}_i}, \frac{\partial}{\partial z_i}\right)\frac{\partial}{\partial z_i}, \frac{\partial}{\partial \bar{z}_i}\right) \\ &= -4g\left(-2\frac{\partial}{\partial z_i}, \frac{\partial}{\partial \bar{z}_i}\right) \\ &= 8 \end{aligned}$$

so the holomorphic sectional curvature in the z_i direction is:

$$K\left(\frac{\partial}{\partial x_i}\right) = \frac{8}{g\left(\frac{\partial}{\partial x_i}, \frac{\partial}{\partial x_i}\right)} = \frac{8}{4\left(\frac{\partial}{\partial z_i}, \frac{\partial}{\partial \bar{z}_i}\right)} = 2$$

A similar calculation yields that $K\left(\frac{\partial}{\partial y_i}\right) = 2$. We conclude that the Fubini-Study metric has constant holomorphic sectional curvature equal to 2.

3.4. **The Complex Hyperbolic Space.** Consider the unit disk in $\mathbb{C}^n$:

$$\mathbb{D}^n = \{z \in \mathbb{C}^n : |z|^2 < 1\}$$

In analogy to our procedure for the Fubini-Study metric, we define a 2-form on $\mathbb{D}^n$ by:

$$\begin{aligned}\omega &= -i\partial\bar{\partial}\log(1-|z|^2) \\ &= i\frac{(1-|z|^2)\delta^i_j + z_j\bar{z}_i}{(1-|z|^2)^2}dz_i \wedge d\bar{z}_j\end{aligned}$$

Definition 3.5. The (complex) hyperbolic metric on $\mathbb{D}^n$ is given by $g(X,Y) = \omega(JX,Y)$.

In the standard global coordinate frame for $\mathbb{D}^n$ the symbols for this metric are clearly:

$$g_{ij} = \frac{(1-|z|^2)\delta^i_j + z_j\bar{z}_i}{(1-|z|^2)^2}$$

Following closely our procedure for the Fubini-Study metric, we will begin by investigating the isometries of g. Consider the group $U_{1,n}$ of matrices that preserve the standard indefinite bilinear form with signature $(1,n)$, that is:

$$U_{1,n} = \{A \in GL_{n+1}(\mathbb{C}) : AI_{1,n}\bar{A}^t = I_{1,n}\}$$

where $I_{1,n}$ is the matrix:

$$\begin{pmatrix} -1 & 0 \\ 0 & I_n \end{pmatrix}$$

We will produce a transitive action by holomorphic isometries of this group on $\mathbb{D}^n$. To accomplish this, first consider the holomorphic projections $\pi_0 : \mathbb{C}^{n+1} \to \mathbb{C}$ and $\pi : \mathbb{C}^{n+1} \to \mathbb{C}^n$ given by:

$$\pi_0(z_0, z) = z_0$$

and:

$$\pi(z_0, z) = z$$

Now, let $U_{1,n}$ act on $\mathbb{D}^n$ (on the right) by setting:

$$f_A(z) = \frac{\pi((1,z)A)}{\pi_0((1,z)A)}$$

We first need to check that $f_A(z) \in \mathbb{D}^n$ for all $A \in U_{1,n}$ and $z \in \mathbb{D}^n$. To see this, we simply write:

$$\begin{aligned}
|f_A(z)|^2 &= \frac{|\pi((1,z)A)|^2 - |\pi_0((1,z)A)|^2 + |\pi_0((1,z)A)|^2}{|\pi_0((1,z)A)|^2} \\
&= \frac{|\pi((1,z)A)|^2 - |\pi_0((1,z)A)|^2}{|\pi_0((1,z)A)|^2} + 1 \\
&= \frac{(1,z)AI_{1,n}\bar{A}^t(1,\bar{z})^t}{\pi_0((1,z)A)} + 1 \\
&= \frac{(1,z)I_{1,n}(1,\bar{z})^t}{\pi_0((1,z)A)} + 1 \\
&= \frac{|z|^2 - 1}{\pi_0((1,z)A)} + 1 \\
&< 1
\end{aligned}$$

To check that this is actually a group action it is convenient for us to give a slightly different description of the situation. Consider $\mathbb{D}^n$ as a subset of $\mathbb{C}P^n$ defined by:

$$\mathbb{D} = \left\{ [z_0, z] : \frac{|z|^2}{|z_0|^2} < 1 \right\} = \left\{ [1, z] : |z|^2 < 1 \right\}$$

Then $U_{1,n}$ acts on this disk by $[z_0, z] \mapsto [z_0, z]A$, and the map $[z_0, z] \mapsto \frac{z}{z_0}$ sends the new action into the old one. But the new definition clearly results in a group action, and hence the old one does as well. We conclude that each f_A is a homeomorphism, so it remains to see that each is an isometry, and that the overall action is transitive.

To see that we have an action by isometries, it suffices to check that $f_A^* \omega = \omega$. First observe that:

$$\begin{aligned}
(1, f_A(z)) &= \left(1, \frac{\pi((1,z)A)}{\pi_0((1,z)A)}\right) \\
&= \frac{(\ \pi_0((1,z)A), \pi((1,z)A)\ )}{\pi_0((1,z)A)} \\
&= \frac{(1,z)A}{\pi_0((1,z)A)}
\end{aligned}$$

with a similar result holding for $(1, \overline{f_A(z)})$. Next, notice that we can express ω as:

$$\omega = -i\partial\bar{\partial} \log(\ -(1,z)I_{1,n}(1,\bar{z})^t\ )$$

and using this expression we can calculate the pullback:

$$\begin{aligned} f_A^*\omega &= -i\partial\bar{\partial}\log(-(1,f_A(z))I_{1,n}(1,\overline{f_A(z)})^t) \\ &= -i\partial\bar{\partial}\log\left(-\frac{(1,z)AI_{1,n}\bar{A}^t(1,\bar{z})^t}{|\pi_0((1,z)A)|^2}\right) \\ &= -i\partial\bar{\partial}\log(-(1,z)AI_{1,n}\bar{A}^t(1,\bar{z})^t) \\ &\quad + i\partial\bar{\partial}\log(\pi_0((1,z)A)) + i\partial\bar{\partial}\overline{\log(\pi_0((1,z)A))} \\ &= -i\partial\bar{\partial}\log(-(1,z)I_{1,n}(1,\bar{z})^t) \\ &= \omega \end{aligned}$$

It follows that f_A is an isometry for all A.

Finally, we show that the action of $U_{1,n}$ is transitive. To do this, it suffices to show that, given $z \in \mathbb{D}^n$, there is an $A \in U_{1,n}$ such that $f_A(z) = 0$; this is equivalent to solving the equation $\pi((1,z)A) = 0$ for $A \in U_{1,n}$. This is easy enough, take the vector $(1,z)$, normalize it, and then extend it to a basis for $\mathbb{C}^{n+1}$ which is orthonormal with respect to the indefinite bilinear form defined by is $I_{1,n}$. Use this basis as the columns in a matrix A, which is then easily seen to lie in $U_{1,n}$. Since $(1,z)$ is orthogonal to the last n columns of the matrix, clearly $\pi((1,z)A^{-1}) = 0$.

All together, we have shown that there is a transitive action of holomorphic isometries on $\mathbb{D}^n$ by the matrix group $U_{1,n}$. So, as was the case for the Fubini-Study metric, to see that g is positive definite and J-invariant, it suffices to look at a single point. We have:

$$g_{ij}|_{z=0} = \delta^i_j$$

and so, indeed, g is J-invariant, positive definite, and hence a Kahler metric on $\mathbb{D}^n$. This Kahler metric is called the complex hyperbolic metric.

Now we can move on to show that g has constant holomorphic sectional curvature. TO begin, it is easy to parallel our work for the Fubini-Study metric and use the g_{ij} symbols to calculate the Christoffel symbols of the hyperbolic metric:

$$\Gamma^k_{ij} = \frac{\delta^i_k\bar{z}_j + \delta^j_k\bar{z}_i}{1-|z|^2}$$

Continuing in the same manner, for the curvature symbols we get:

$$R^r_{ijl} = \frac{\partial\Gamma^r_{jl}}{\partial\bar{z}_i} = \frac{(1-|z|^2)(\delta^r_j\delta^i_l + \delta^r_l\delta^i_j) + \delta^r_j z_i\bar{z}_l + \delta^r_l z_i\bar{z}_j}{(1-|z|^2)^2}$$

so that:

$$R\left(\frac{\partial}{\partial\bar{z}_i}, \frac{\partial}{\partial z_j}\right)\frac{\partial}{\partial z_l} = \frac{(1-|z|^2)\delta^i_l + z_i\bar{z}_l}{(1-|z|^2)^2}\frac{\partial}{\partial z_j} + \frac{(1-|z|^2)\delta^i_j + z_i\bar{z}_j}{(1-|z|^2)^2}\frac{\partial}{\partial z_l}$$

Since we have already shown g to be homogeneous, we may focus our attention at the point $z = 0$, obtaining:

$$R\left(\frac{\partial}{\partial \bar{z}_i}, \frac{\partial}{\partial z_j}\right)\frac{\partial}{\partial z_l}\bigg|_{z=0} = \delta^i_l \frac{\partial}{\partial z_j} + \delta^i_j \frac{\partial}{\partial z_l}$$

which is exactly the negation of what we obtained for the Fubini-Study metric. It is now immediate that the complex hyperbolic metric has constant holomorphic sectional curvature equal to -2.

3.6. **Summary.** Above, we have constructed simply connected Kahler manifolds with constant holomorphic sectional curvatures equal to $0, 2$ and -2. It is easy to now produce metrics of any constant holomorphic sectional curvature. Indeed, suppose we take a Kahler metric g and scale it to get another metric λg. Following through the various definitions and formulas, we see that the Christoffel symbols and curvature symbols in any coordinate system are left unchanged. The only difference occurs in the last step, when calculating the holomorphic sectional curvature; upon evaluating the curvature tensor inside the metric, the numerator is scaled by λ, and the denominator by λ^2, and hence the final result is scaled by $\frac{1}{\lambda}$.

Consequently, if $c > 0$, then the 2-form:

$$\omega = i\frac{2}{c}\partial\bar{\partial}\log(1 + |z|^2)$$

(in an appropriate coordinate system) defines a metric of constant holomorphic sectional curvature c on $\mathbb{C}P^n$. This metric is called the Fubini-Study metric of holomorphic curvature c, and we denote the resulting Kahler manifold by $\mathbb{C}P^n(c)$. Similarly, the 2-form:

$$\omega = -i\frac{2}{c}\partial\bar{\partial}\log(1 - |z|^2)$$

defines a metric of constant holomorphic sectional curvature $-c$ on $\mathbb{D}^n$, which we call the complex hyperbolic metric of holomorphic curvature $-c$, and denote the resulting Kahler manifold by $\mathbb{C}H^n(c)$.

We now have, for every real number c, a simply connected complex manifold with a Kahler metric of constant holomorphic sectional curvature c. Recalling from the previous chapter that such a space is unique up to isometry, we have completed the proof of the following theorem:

Theorem 3.7. *Let M be a simply connected Kahler manifold of constant holomorphic sectional curvature c and complex dimension n, then:*

- *If $c = 0$, then M is isometric to $\mathbb{C}^n$ with its flat Kahler metric.*
- *If $c > 0$, then M is isometric to $\mathbb{C}P^n(c)$.*
- *If $c < 0$, then M is isometric to $\mathbb{C}H^n(c)$.*

4. The Hermitian Metric and the Connection and Curvature Forms

In this section we take a closer look at some of the results we have already obtained, and recast them in the language of Hermitian metrics and the Cartan formalism of differential forms.

4.1. **The Hermitian Metric.** Let M be a Kahler manifold with metric g, associated 2-form ω, and Riemannian connection ∇. Consider the complex valued 2-tensor on TM, defined by:

$$h(X,Y) = g(X,Y) - i\omega(X,Y)$$

Since M is a complex manifold, its tangent bundle is naturally a complex vector bundle, with the complex structure encoded by the automorphism J. In this context h is a hermitian metric, for:

$$\begin{aligned} h(JX,Y) &= g(JX,Y) - i\omega(JX,Y) \\ &= \omega(X,Y) + ig(X,Y) \\ &= ih(X,Y) \end{aligned}$$

and similarly, $h(X,JY) = -ih(X,Y)$. In a holomorphic coordinate frame field $Z_i = \frac{1}{2}(X_i - iY_i)$ we define the symbols of this metric by:

$$h_{ij} = h(X_i, X_j) = h(Y_i, Y_j)$$

These are related to the g symbols by:

$$\begin{aligned} g_{ij} = g(Z_i, \bar{Z}_j) &= \frac{1}{4} g(X_i - iY_i, X_j + iY_j) \\ &= \frac{1}{2}(g(X_i, X_j) - ig(Y_i, X_j)) \\ &= \frac{1}{2}(g(X_i, X_j) - i\omega(X_i, X_j)) \\ &= \frac{1}{2} h(X_i, X_j) \\ &= \frac{1}{2} h_{ij} \end{aligned}$$

Therefore we have the following expressions for g and the associated 2-form in a holomorphic coordinate frame field:

$$g = \frac{h_{ij}}{2} dz_i \otimes d\bar{z}_j + \frac{\bar{h}_{ij}}{2} d\bar{z}_i \otimes dz_i$$

and:

$$\omega = \frac{ih_{ij}}{2} dz_i \wedge d\bar{z}_j$$

Remark 4.2. Note that the above expressions hold for the h and g symbols in any local frame field for TM^+, not just in a holomorphic coordinate frame field. In particular, if the frame $\{e_1, \dots, e_n\}$ is unitary with respect to h, and $\{e^1, \dots, e^n\}$ is its dual frame field, then the 2-form associated to the metric is given by:

$$\omega = \frac{i}{2} e^i \wedge \bar{e}^i$$

We observe now, for later use, an important property of the hermitian metric on a Kahler manifold:

Proposition 4.3. *The Riemannian connection ∇ is compatible with the hermitian metric, that is:*

$$Xh(Y, Z) = h(\nabla_X Y, Z) + h(Y, \nabla_X Z)$$

Proof. We simply calculate using the definition of h, the compatibility of ∇ with g, and the fact that J is parallel:

$$\begin{aligned} Xh(Y, Z) &= X(g(Y, Z) - ig(JY, Z)) \\ &= g(\nabla_X Y, Z) + g(Y, \nabla_X Z) - ig(J\nabla_X Y, Z) - ig(JY, \nabla_X Z) \\ &= h(\nabla_X Y, Z) + h(Y, \nabla_X, Z) \end{aligned}$$

which is what we wanted to show. □

Remark 4.4. Using the bundle isomorphism $\Phi : TM \to TM^+$ we can transfer h to a hermitian metric on TM^+, which we denote by $\tilde{h}$. The induced connection on TM^+ is then compatible with the metric $\tilde{h}$. We remind the reader that this induced connection agrees with what we obtain by extending ∇ to $T_{\mathbb{C}}M$ complex linearly, and then restricting to TM^+.

4.5. **The Connection and Curvature Forms.** We now turn our attention in another direction temporarily, and discuss a different way of organizing the information contained in the connection on a Kahler manifold. Again, we work with a fixed holomorphic coordinate frame field Z_i for TM^+.

Define a collection of complex valued one forms on $T_{\mathbb{C}}M$ by:

$$\nabla_X Z_j = \theta_j^k(X) Z_k$$

Sometimes we choose to think of θ as a matrix of one forms (with the top index for the rows and bottom for the columns) and sometimes as a single $M_n(\mathbb{C})$ valued 1-form, depending on the context. From this perspective the definition of θ can be abbreviated as follows: assemble the frame into a row vector $Z = (Z_i, \dots, Z_n)$, then:

$$\nabla Z = Z\theta$$

Because θ expresses the connection ∇ in a local coordinate frame field, we call it the connection form.

Comparing the definition of θ to the relations $\nabla_{Z_i} Z_j = \Gamma_{ij}^k Z_k$ and $\nabla_{\bar{Z}_i} Z_j = 0$, we see that:

$$\theta_j^k = \Gamma_{ij}^k dz_i$$

So θ is a $(1,0)$-form.

The previous equation lets us easily derive a formula for θ in terms of h, for we have:

$$\Gamma_{ij}^k = \frac{\partial g_{jl}}{\partial z_i} g^{lk} = \frac{\partial h_{jl}}{\partial z_i} h^{lk}$$

and hence:

$$\theta_j^k = \frac{\partial h_{jl}}{\partial z_i} h^{lk} dz_i = \left(\frac{\partial h_{jl}}{\partial z_i} dz_i \right) h^{lk} = \partial h_{jl}\ h^{lk} = (\partial h\ h^{-1})_{jk}$$

which we can write as a single matrix equation:

$$\theta = (\partial h\ h^{-1})^t = (h^t)^{-1} (\partial h)^t = \bar{h}^{-1}\ \partial \bar{h}$$

Now let's turn to the curvature. In analogy with θ we define some complex bilinear objects Ω on $T_{\mathbb{C}} M$ by the formula:

$$R(X,Y) Z_k = \Omega_k^l(X,Y) Z_l$$

since $R(X,Y) = -R(Y,X)$, we see that the Ω's are complex valued 2-forms. To relate them to our earlier work, recall that:

$$R(Z_i, Z_j) = R(\bar{Z}_i, \bar{Z}_j) = 0$$

and:

$$R(\bar{Z}_i, Z_j) Z_k = R_{ijk}^l Z_l$$

so that:

$$\Omega_k^l = R_{ijk}^l d\bar{z}_i \wedge dz_j$$

From which we conclude that Ω is a $(1,1)$-form.

Now recall the formula for the curvature symbols in a holomorphic coordinate frame field:

$$R_{ijk}^l = \frac{\partial \Gamma_{jk}^l}{\partial \bar{z}_i}$$

Substituting this into the previous equation gives us:

$$\Omega_k^l = \frac{\partial \Gamma_{jk}^l}{\partial \bar{z}_i} d\bar{z}_i \wedge dz_j = \bar{\partial}(\Gamma_{jk}^l dz_j) = \bar{\partial} \theta_k^l$$

or more succinctly:

$$\Omega = \bar{\partial} \theta$$

It will be convenient for us to complete the emerging picture and investigate $\partial\theta$. Differentiate the formula $\theta = \bar{h}^{-1}\partial\bar{h}$, and using the linear algebraic formula $\partial\bar{h}^{-1} = -\bar{h}^{-1}\partial\bar{h}\bar{h}^{-1}$ along the way:

$$\begin{aligned}\partial\theta &= \partial(\bar{h}^{-1}\partial\bar{h}) = (\partial\bar{h}^{-1}) \wedge \partial\bar{h} - \bar{h}^{-1}\partial^2\bar{h} \\ &= -\bar{h}^{-1}(\partial\bar{h})\bar{h}^{-1} \wedge \partial\bar{h} = -\bar{h}^{-1}\partial\bar{h} \wedge \bar{h}^{-1}\partial\bar{h} \\ &= -\theta \wedge \theta\end{aligned}$$

So we have derived the two equations:

$$\bar{\partial}\theta = \Omega \quad \text{and} \quad \partial\theta + \theta \wedge \theta = 0$$

Which, since $d = \partial + \bar{\partial}$, can be combined in to the single:

$$\Omega = d\theta + \theta \wedge \theta$$

We would like to remove the dependence on a holomorphic coordinate frame from these results, and in fact, this can be very easily done:

Proposition 4.6. *Let $X_1, X_2, \ldots, X_n$ be any local frame field for TM^+, and assemble it into a row vector $\mathbf{X} = (X_1, \ldots, X_n)$. Define connection and curvature forms with respect to this frame by $\nabla\mathbf{X} = \mathbf{X}\tilde{\theta}$ and $R\mathbf{X} = \mathbf{X}\tilde{\Omega}$. Then:*

$$\tilde{\Omega} = d\tilde{\theta} + \tilde{\theta} \wedge \tilde{\theta}$$

Proof. We first choose a local holomorphic frame field $\mathbf{Z} = (Z_1, \ldots, Z_n)$ to relate the given frame to; that is, we write $\mathbf{X} = \mathbf{Z}A$ where A is a locally defined change of frame function from M to $GL_n(\mathbb{C})$. We write θ and Ω for the connection and curvature forms with respect to $\mathbf{Z}$, which we already know satisfy the equation:

$$\Omega = d\theta + \theta \wedge \theta$$

We have:

$$\begin{aligned}\nabla\mathbf{X} &= \nabla(\mathbf{Z}A) = (\nabla\mathbf{Z})A + \mathbf{Z}dA \\ &= \mathbf{Z}\theta A + \mathbf{Z}dA \\ &= \mathbf{X}A^{-1}\theta A + \mathbf{X}A^{-1}dA \\ &= \mathbf{X}(A^{-1}\theta A + A^{-1}dA)\end{aligned}$$

and

$$\begin{aligned}R\mathbf{X} &= R(\mathbf{Z}A) = (R\mathbf{Z})A \\ &= \mathbf{Z}\Omega A \\ &= \mathbf{X}A^{-1}\Omega A\end{aligned}$$

so we conclude that:

$$\tilde{\theta} = A^{-1}\theta A + A^{-1}dA \quad \text{and} \quad \tilde{\Omega} = A^{-1}\Omega A$$

Therefore:

$$\begin{aligned}
d\tilde{\theta} + \tilde{\theta} \wedge \tilde{\theta} &= d(A^{-1}\theta A + A^{-1}dA) + (A^{-1}\theta A + A^{-1}dA) \wedge (A^{-1}\theta A + A^{-1}dA) \\
&= d(A^{-1}) \wedge \theta A + A^{-1}d\theta A - A^{-1}\theta \wedge dA + d(A^{-1}) \wedge dA + A^{-1}\theta \wedge \theta A \\
&\quad + A^{-1}\theta \wedge dA + A^{-1}dA \wedge A^{-1}\theta A + A^{-1}dA \wedge A^{-1}dA \\
&= -A^{-1}dAA^{-1} \wedge \theta A + A^{-1}d\theta A - A^{-1}\theta \wedge dA - A^{-1}dAA^{-1} \wedge dA \\
&\quad + A^{-1}\theta \wedge \theta A + A^{-1}\theta \wedge dA + A^{-1}dA \wedge A^{-1}\theta A + A^{-1}dA \wedge A^{-1}dA \\
&= A^{-1}d\theta A + A^{-1}\theta \wedge \theta A \\
&= A^{-1}\Omega A \\
&= \tilde{\Omega}
\end{aligned}$$

as we wanted to show. □

We need another formula in our subsequent work, and now is a good time to discuss it.

Proposition 4.7. *Let $\mathbf{X} = (X_1, \ldots, X_n)$ be any frame field for the bundle TM^+, and let $\mathbf{X}^* = (X_1^*, \ldots, X_n^*)^t$ be its dual frame field (note the transpose). Then the equation:*

$$d\mathbf{X}^* + \theta \wedge \mathbf{X}^* = 0$$

is equivalent to the connection ∇ being torsion free.

Proof. Let Y and Z be any two vector fields in TM^+. We can write these in the given frame X as:

$$Y = \mathbf{X}(\mathbf{X}^*(Y)) \quad \text{and} \quad Z = \mathbf{X}(\mathbf{X}^*(Z))$$

Let's calculate the torsion of the connection using these two equations. We have:

$$\begin{aligned}
\nabla_Y Z &= \nabla_Y(\mathbf{X}(\mathbf{X}^*(Z))) \\
&= (\nabla_Y \mathbf{X})(\mathbf{X}^*(Z)) + \mathbf{X}(Y(\mathbf{X}^*(Z))) \\
&= (\mathbf{X}\theta(Y))(\mathbf{X}^*(Z)) + \mathbf{X}(Y(\mathbf{X}^*(Z))) \\
&= \mathbf{X}(\ \theta(Y)\mathbf{X}^*(Z) + Y(\mathbf{X}^*(Z))\ )
\end{aligned}$$

and by switching Y with Z:

$$\nabla_Z Y = \mathbf{X}(\ \theta(Z)\mathbf{X}^*(Y) + Z(\mathbf{X}^*(Y))\ )$$

Now, using that $[Y, Z] = \mathbf{X}(\mathbf{X}^*([Y, Z]))$, we can express the torsion tensor as:

$$\begin{aligned}
T(Y, Z) &= \nabla_Y Z - \nabla_Z Y - [Y, Z] \\
&= \mathbf{X}(\ Y(\mathbf{X}^*(Z)) - Z(\mathbf{X}^*(Y)) - \mathbf{X}^*([Y, Z]) + \theta(Y)\mathbf{X}^*(Z) - \theta(Z)\mathbf{X}^*(Y)\ ) \\
&= \mathbf{X}(\ d\mathbf{X}^*(Y, Z) + \theta \wedge \mathbf{X}^*(Y, Z)\ )
\end{aligned}$$

which clearly completes the proof. □

We have defined our connection forms relative to a chosen frame for TM^+, and different choices of frames can be convenient in different situations. We have already seen that a holomorphic coordinate frame brings about certain simplifications in the theory, for example, in such a frame ω is a $(1,0)$-form. Let's now take a look at how things go if we use a unitary frame $e_1, e_2, \ldots, e_n$ for TM^+, in the sense that:

$$\tilde{h}(e_i, e_j) = \delta_j^i$$

We want to look at an additional property of the connection form in such a frame; to do so, we differentiate the above equation:

$$\begin{aligned} 0 &= X\tilde{h}(e_i, e_j) \\ &= \tilde{h}(\nabla_X e_i, e_j) + \tilde{h}(e_i, \nabla_X e_j) \\ &= \tilde{h}(\theta_i^l(X)e_l, e_j) + \tilde{h}(e_i, \theta_j^l(X)e_l) \\ &= \theta_i^l(X)\delta_j^l + \bar{\theta}_j^l(X)\delta_l^i \\ &= \theta_i^j(X) + \bar{\theta}_j^i(X) \end{aligned}$$

Which verifies the following result:

Proposition 4.8. *If $e_1, \ldots, e_n$ is a unitary frame field for the bundle TM^+ with respect to the hermitian metric $\tilde{h}$, then the connection form with respect to this frame is skew-hermitian, that is $\theta^t = -\bar{\theta}$.*

We note at this point that propositions 3 and 4 actually characterize the connection forms completely, since, as we know from Riemannian geometry, there is a unique connection which is both compatible with h and torsion free (this could also be verified directly).

We need one more result about the curvature form of a manifold of constant holomorphic sectional curvature, which is a restatement of our previous determination of the curvature tensor of these spaces.

Proposition 4.9. *Let $\lambda = (\epsilon_1, \ldots, \epsilon_n)$ be any local frame field for the bundle TM^+, let $\lambda^* = (\epsilon^1, \ldots, \epsilon^n)^t$ be the dual frame field, and let Ω be the curvature form with respect to this frame. Then M has constant holomorphic sectional curvature c if and only if:*

$$\Omega = \frac{c}{2}\left(-i\omega I + \frac{1}{2}\lambda^* \wedge (\bar{\lambda}^*)^t \bar{h}\right)$$

Proof. Recall the definition of Ω:

$$R(X,Y)\epsilon_i = \Omega_i^l(X,Y)\epsilon_l$$

so that:

$$\Omega_i^j(X,Y) = \epsilon^j(R(X,Y)\epsilon_i)$$

Now, M has constant holomorphic sectional curvature c if and only if its curvature tensor is given by:

$$\begin{aligned} R(X,Y)Z = \frac{c}{4}(&g(Y,Z)X - g(X,Z)Y - g(Y,JZ)JX \\ &+ g(X,JZ)JY + 2g(X,JY)JZ) \end{aligned}$$

so we get:

$$\begin{aligned} \Omega_i^j(X,Y) &= \epsilon^j(R(X,Y)\epsilon_i) \\ &= \frac{c}{4}(g(Y,\epsilon_i)\epsilon^j(X) - g(X,\epsilon_i)\epsilon^j(Y) - g(Y,J\epsilon_i)\epsilon^j(JX) \\ &\quad + g(X,J\epsilon_i)\epsilon^j(JY) + 2g(X,JY)\epsilon^j(J\epsilon_i)) \end{aligned}$$

Let's take a look at the terms in this expression one by one, starting with the last. Since $\epsilon_i \in TM^+$, we have:

$$2g(X,JY)\epsilon^j(J\epsilon_i) = -2\omega(X,Y)\epsilon^j(i\epsilon_i) = -2i\delta_j^i\omega(X,Y)$$

which gives us the first term in our target expression for Ω.

Now let's examine the term $g(Y,\epsilon_i)\epsilon^j(X)$. First, we write Y in the given frame as:

$$Y = \epsilon^l(Y)\epsilon_l + \bar{\epsilon}^l(Y)\bar{\epsilon}_l$$

and therefore:

$$g(Y,\epsilon_i) = g(\epsilon^l(Y)\epsilon_l + \bar{\epsilon}^l(Y)\bar{\epsilon}_l, \epsilon_i) = g(\bar{\epsilon}_l,\epsilon_i)\bar{\epsilon}^l(Y)$$

so the term becomes:

$$g(Y,\epsilon_i)\epsilon^j(X) = g(\bar{\epsilon}_l,\epsilon_i)\bar{\epsilon}^l(Y)\epsilon^j(X)$$

Similarly we acquire:

$$g(X,\epsilon_i)\epsilon^j(Y) = g(\bar{\epsilon}_l,\epsilon_i)\bar{\epsilon}^l(X)\epsilon^j(Y)$$

Moving on, let's now concentrate on the term $g(Y,J\epsilon_i)\epsilon^j(JX)$. First:

$$g(Y,J\epsilon_i) = -\omega(Y,\epsilon_i) = -\omega(\bar{\epsilon}_l,\epsilon_i)\bar{\epsilon}_l(Y)$$

so we need to examine $\epsilon^j(JX)$. To simplify this, recall that we the complex tangent bundle splits as a direct sum of J-invariant subspaces: $T_{\mathbb{C}}M = TM^+ \oplus TM^-$; so write $X = X^+ + X^-$ where $X^+ \in TM^+$ and $X^- \in TM^-$. Then we have:

$$\epsilon^j(JX) = \epsilon^j(JX^+ + JX^-) = \epsilon^j(JX^+) = \epsilon^j(iX^+) = i\epsilon^j(X^+) = i\epsilon^j(X)$$

So altogether:

$$g(Y,J\epsilon_i)\epsilon^j(JX) = -i\omega(\bar{\epsilon}_l,\epsilon_i)\bar{\epsilon}_l(Y)\epsilon^j(X)$$

Similarly:

$$g(X, J\epsilon_i)\epsilon^j(JY) = -i\omega(\bar{\epsilon}_l, \epsilon_i)\bar{\epsilon}_l(X)\epsilon^j(Y)$$

Putting this all together, we get:

$$\begin{aligned}
\Omega_i^j(X,Y) &= \epsilon^j(R(X,Y)\epsilon_i) \\
&= \frac{c}{4}(g(\bar{\epsilon}_l, \epsilon_i)\bar{\epsilon}^l(Y)\epsilon^j(X) - g(\bar{\epsilon}_l, \epsilon_i)\bar{\epsilon}^l(X)\epsilon^j(Y) \\
&\quad + i\omega(\bar{\epsilon}_l, \epsilon_i)\bar{\epsilon}_l(Y)\epsilon^j(X) - i\omega(\bar{\epsilon}_l, \epsilon_i)\bar{\epsilon}_l(X)\epsilon^j(Y) - 2i\delta_j^i\omega(X,Y)) \\
&= \frac{c}{4}(\ (g(\epsilon_i, \bar{\epsilon}_l) - i\omega(\epsilon_i, \bar{\epsilon}_l))(\bar{\epsilon}^l(Y)\epsilon^j(X) - \bar{\epsilon}^l(X)\epsilon^j(Y)) \\
&\quad - 2i\delta_j^i\omega(X,Y))
\end{aligned}$$

Now:

$$g(\epsilon_i, \bar{\epsilon}_l) - i\omega(\epsilon_i, \bar{\epsilon}_l) = g_{il} - i(ig_{il}) = 2g_{il} = h_{il}$$

and:

$$\bar{\epsilon}^l(Y)\epsilon^j(X) - \bar{\epsilon}^l(X)\epsilon^j(Y) = \epsilon^j \wedge \bar{\epsilon}^l(X,Y)$$

so we have shown that:

$$\begin{aligned}
\Omega_i^j(X,Y) &= \frac{c}{4}(\ h_{il}\epsilon^j \wedge \bar{\epsilon}^l(X,Y) - 2i\delta_j^i\omega(X,Y)\ ) \\
&= \frac{c}{4}(\ \epsilon^j \wedge \bar{\epsilon}^l(X,Y)\bar{h}_{li} - 2i\delta_j^i\omega(X,Y)\ )
\end{aligned}$$

Which is exactly what we wanted. □

Finally, we prove a result characterising Kahler-Einstein manifolds in terms of their curvature forms.

Proposition 4.10. *A manifold is Kahler Einstein of constant λ if and only if its curvature form in any local frame for TM^+ satisfies $tr\ \Omega = -i\lambda\omega$.*

Proof. We begin by proving the result in a local holomorphic coordinate frame field for TM^+. Let Z_i be such a frame field, and let Z^i be its dual. Recall that the Kahler metric is Einstein if and only if $\mathrm{ric} = \lambda\omega$, and in this case we have:

$$\begin{aligned}
\lambda\omega(Z_i, \bar{Z}_j) &= \mathrm{ric}(Z_i, \bar{Z}_j) = i\mathrm{Ric}(Z_i, \bar{Z}_j) \\
&= i\mathrm{tr}(Z \mapsto R(Z, Z_i)\bar{Z}_j) \\
&= i\bar{Z}^k(R(\bar{Z}_k, Z_i)\bar{Z}_j) \\
&= i\bar{Z}^k(-R(\bar{Z}_j, \bar{Z}_k)Z_i - R(Z_i, \bar{Z}_j)\bar{Z}_k) \\
&= -i\bar{Z}^k(\bar{\Omega}_r^k(Z_i, \bar{Z}_j)\bar{Z}_r) \\
&= -i\bar{\Omega}_k^k(Z_i, \bar{Z}_j)
\end{aligned}$$

Taking the conjugate of both sides, and using that ω is real valued, gives the result we want.

Given an arbitrary frame field $\mathbf{X}$, let $\tilde{\Omega}$ denote the curvature form in this frame. Let $A \in GL_n(\mathbb{C})$ be the change of frame matrix from $\mathbf{X}$ to $\mathbf{Z}$, and recall that $\tilde{\Omega} = A^{-1}\Omega A$. We then have:

$$\mathrm{tr}\tilde{\Omega} = \mathrm{tr}(A^{-1}\Omega A) = \mathrm{tr}\Omega$$

which proves the result in the frame $\mathbf{X}$. □

4.11. **Some More Discussion of the Fubini Study Metric.** We will now give another perspective on the Fubini-Study metric, and study its connection and curvature forms.

Consider the normalization map $z_0 : \mathbb{C}^{n+1} - \{0\} \to S^{2n-1}$ defined by:

$$z_0(z) = \frac{z}{|z|}$$

which is smooth on (since we have removed the origin). Using the symbol $\cdot$ to denote the usual Hermitian inner product on $\mathbb{C}^{n+1}$, define a hermitian metric on $\mathbb{C}^{n+1} - \{0\}$ by:

$$ds^2 = dz_0 \cdot dz_0 - (dz_0 \cdot z_0)(z_0 \cdot dz_0)$$

Observe that this metric is invariant under the multiplicative action of $\mathbb{C} - \{0\}$ on $\mathbb{C}^{n+1}$, and it therefore induces a metric on the quotient space $\mathbb{C}P^n$. We will prove that this is the Fubini-Study metric of constant holomorphic sectional curvature 4.

Let $p \in \mathbb{C}P^n$, and let Z_0 be a fixed unit vector in the fiber over p. Then it is easy to see that the projection $\pi : \mathbb{C}^{n+1} - \{0\} \to \mathbb{C}P^n$ induces an isometry from the unitary complement of Z_0 equipped with the metric ds^2 to the tangent space $T_p\mathbb{C}P^n$ with the induced metric, since for vectors in the unitary complement we have $dz_0 \cdot z_0 = 0$. Therefore, given a unitary frame $\mathbf{e} = \{e_1, \ldots, e_n\}$ for $T_p\mathbb{C}^n$, there is a unique unitary frame $\{Z_1, \ldots, Z_n\}$ for $Z_0^{\perp}$ mapping isometrically to $\mathbf{e}$. So suppose we have a local section of unit vectors $Z_0 : U \subset \mathbb{C}P^n \to \mathbb{C}^{n+1} - \{0\}$ for the projection map, and a local unitary frame field for TM^+ (where $M = \mathbb{C}P^n$) over U. Denoting the $k \times k$ unitary matrices by U_k, we can lift all this structure to map:

$$\Phi : U \to U_{n+1}$$

which is defined in our notation as:

$$\Phi(p) = (Z_0, Z_1, \ldots, Z_n)$$

Now each $dZ_i : U \to \mathbb{C}^{n+1}$ is a $\mathbb{C}^{n+1}$ valued 1-form, and hence can be expressed as:

$$dZ_i = \phi_i^j Z_j$$

where the ϕ's are complex valued 1-forms on U. We can write this definition succinctly as $d\mathbf{Z} = \mathbf{Z}\phi$, where here we think of ϕ as a $M_n(\mathbb{C})$ valued 1-form. Clearly, these 1-forms can be expressed as:

$$\phi_i^j = dZ_i \cdot Z_j$$

Since dZ_0 is an isometry from $T_p\mathbb{C}P^n$ to its image in $Z_0^\perp$, this immediately implies that:

$$\phi_0^i = dZ_0 \cdot Z_i = e^i$$

Differentiating the equation $Z_i \cdot Z_j = \delta_j^i$ tells us that:

$$\phi_i^j = -\bar{\phi}_j^i$$

that is, the matrix ϕ is skew hermitian.

We can also calculate the exterior derivative of ϕ. Let's write:

$$Z_i = (z_{i0}, z_{i1}, \ldots, z_{in})$$

then:

$$dZ_i \cdot Z_j = \bar{z}_{jr} dz_{ir}$$

and hence, since $\bar{z}_{sr} z_{tr} = \delta_s^t$:

$$\begin{aligned} d(dZ_i \cdot Z_j) &= d\bar{z}_{jr} \wedge dz_{ir} \\ &= (\bar{\phi}_j^s \bar{z}_{sr}) \wedge (\phi_i^t z_{tr}) \\ &= \bar{\phi}_j^s \wedge \phi_i^t \bar{z}_{sr} z_{tr} \\ &= \bar{\phi}_j^s \wedge \phi_i^s \\ &= -\phi_s^j \wedge \phi_i^s \end{aligned}$$

which shows that $d\phi = -\phi \wedge \phi$.

Next, we relate the forms ϕ to the connection forms of the metric ds^2 on $\mathbb{C}P^n$.

Lemma 4.12. *Let $\theta_i^j = \phi_i^j - \delta_i^j \phi_0^0$, then θ is the connection form of the metric ds^2 with respect to the unitary frame e_i.*

Proof. First observe that since ϕ is skew hermitian, so is θ. Therefore we only need to check that the torsion free equation is satisfied:

$$\begin{aligned} de^i = d\phi_0^i &= -\phi_r^i \wedge \phi_0^r \\ &= -\phi_r^i \wedge e^r - \phi_0^i \wedge \phi_0^0 \\ &= (-\phi_r^i + \delta_r^i \phi_0^0) \wedge e^r \\ &= -\theta_r^i \wedge e^r \end{aligned}$$

so $de + \theta \wedge e = 0$, as we wanted to show. □

Proposition 4.13. *The metric induced on $\mathbb{C}P^n$ by ds^2 has constant holomorphic curvature equal to 4, and hence is isometric to a Fubini-Study metric.*

Proof. We calculate the curvature forms using the fundamental equation and the previous lemma:

$$\begin{aligned} \Omega_i^j &= d\theta_i^j + \theta_r^j \wedge \theta_i^r \\ &= d(\phi_i^j - \delta_i^j \phi_0^0) + (\phi_r^j - \delta_r^j \phi_0^0) \wedge (\phi_i^r - \delta_i^r \phi_0^0) \\ &= -\phi_0^j \wedge \phi_i^0 - \phi_r^j \wedge \phi_i^r + \delta_i^j \phi_0^0 \wedge \phi_0^0 + \delta_i^j \phi_r^0 \wedge \phi_0^r \\ &\quad \phi_r^j \wedge \phi_i^r - \delta_i^r \phi_r^j \wedge \phi_0^0 - \delta_r^j \phi_0^0 \wedge \phi_i^r + \delta_r^j \delta_i^r \phi_0^0 \wedge \phi_0^0 \\ &= -\phi_0^j \wedge \phi_i^0 + \delta_i^j \phi_r^0 \wedge \phi_0^r \\ &= e^j \wedge \bar{e}^i - \delta_i^j \bar{e}^r \wedge e^r \\ &= \delta_i^j e^r \wedge \bar{e}^r + e^j \wedge \bar{e}^i \end{aligned}$$

Now note that since we are using a unitary frame, $h_{ij} = \delta_j^i$, and consequently:

$$\omega = \frac{i}{2} e^r \wedge \bar{e}^r$$

and therefore we have:

$$\Omega_i^j = -2i\delta_j^i \omega + e^j \wedge \bar{e}^i$$

which, from an earlier proposition, implies that the metric has constant holomorphic sectional curvature 4. □

5. Kahler Submanifolds

In this section we study complex submanifolds of an ambient Kahler manifold.

Let M and $\tilde{M}$ be complex manifolds with complex structures J and $\tilde{J}$ respectively. Recall that a smooth map $f : M \to \tilde{M}$ is holomorphic if and only if $df \circ J = \tilde{J} \circ df$.

Definition 5.1. We say M is an immersed (or embedded) complex submanifold of $\tilde{M}$ if it is a real submanifold, and the immersion (or embedding) map is holomorphic.

As usual in this context, we will suppress the notation for the immersion map, and consider M as a subset of $\tilde{M}$. Notice that, according to the comment preceding the definition, if M is a complex submanifold of $\tilde{M}$, then TM is a subbundle of $T\tilde{M}|_M$, and the complex structure of $\tilde{M}$ restricts to that of M, that is, $\tilde{J}|_{TM} = J$.

Now suppose that $\tilde{M}$ is a Kahler manifold with Kahler metric $\tilde{g}$. Then $g = \tilde{g}|_M$ is a Riemannian metric on M, which, according to the previous comment, is J-invariant. The associated 2-form for this metric is then $\tilde{\omega} = \omega|_M$, and we have:

$$d\omega = d(\tilde{\omega}|_M) = (d\tilde{\omega})|_M = 0$$

so g is a Kahler metric. In this situation we call M a Kahler submanifold of $\tilde{M}$.

Recall that a locally defined real valued function on $\tilde{M}$ is called a potential for the Kahler metric if:

$$\tilde{\omega} = i\partial\bar{\partial}f$$

The relation between potentials for an ambient space and a Kahler submanifold is as clear-cut as possible:

Proposition 5.2. *If f is a Kahler potential for $\tilde{g}$ then $f|_M$ is a Kahler potential for g.*

Proof. This follows from the simple calculation:

$$i\partial\bar{\partial}(f|_M) = (i\partial\bar{\partial}f)|_M = \tilde{\omega}|_M = \omega$$

□

Another important property of Kahler submanifolds is:

Proposition 5.3. *If M is a Kahler submanifold of $\tilde{M}$, then the normal bundle of M in $\tilde{M}$ is $\tilde{J}$-invariant.*

Proof. Let X be tangent to M and Y normal to M, both at the same point. It suffices to show that $\tilde{g}(X, \tilde{J}Y) = 0$. But:

$$\tilde{g}(X, \tilde{J}Y) = -\tilde{g}(\tilde{J}X, Y) = 0$$

since $\tilde{J}X$ is also tangent to M. □

Now we introduce, as usual, the second fundamental form $B : TM \otimes TM \to NM$, which is defined as:

$$B(X,Y) = (\tilde{\nabla}_X Y)^\perp = \tilde{\nabla}_X Y - \nabla_X Y$$

where $\tilde{\nabla}$ is the Riemannian connection associated to the metric $\tilde{g}$, and ∇ the induced connection on M. As is well known, the second fundamental form is symmetric. In the Kahler case, we have the following additional property:

Proposition 5.4. *If M is a Kahler submanifold of $\tilde{M}$, and B is its second fundamental form, then:*

$$B(X,JY) = B(JX,Y) = \tilde{J}B(X,Y)$$

Proof. Again, we simply calculate:

$$B(X,JY) = \tilde{\nabla}_X JY - \nabla_X JY = \tilde{J}(\tilde{\nabla}_X Y - \nabla_X Y) = \tilde{J}B(X,Y)$$

and consequently:

$$B(JX,Y) = B(Y,JX) = \tilde{J}B(Y,X) = \tilde{J}B(X,Y)$$

which is what we wanted to show. □

From here on we will drop the tedious distinction between structures on the ambient space, and the corresponding structures on the submanifold. We will resume using tildes only when confusion would otherwise result, since all of the previous propositions show that in most cases the distinction is immaterial.

Remark 5.5. Suppose we now extend B complex linearly in both components to the bundle $T_{\mathbb{C}}M$. Using the usual arguments we get that:

$$B(X,Y) = 0 \quad \text{for} \quad X \in TM^+, Y \in TM^-$$

and that:

$$B(\bar{X},\bar{Y}) = \overline{B(X,Y)}$$

Let $N_{\mathbb{C}}M$ denote the unitary complement of $T_{\mathbb{C}}M$ inside $T_{\mathbb{C}}\tilde{M}$, and let NM^+ and NM^- be the i and $-i$ eigenspaces of J in $N_{\mathbb{C}}M$ respectively. Then for $X, Y \in TM^+$ we have:

$$JB(X,Y) = B(JX,Y) = B(iX,Y) = iB(X,Y)$$

so $B(X,Y) \in NM^+$. In the same way, for $X, Y \in TM^-$ we have $B(X,Y) \in NM^-$.

Now recall the most important fact about the second fundamental form, the Gauss Equation:

Gauss Equation : If X, Y, Z and W are tangent to M at some point p, then:

$$R(X,Y,Z,W) = \tilde{R}(X,Y,Z,W) + g(B(X,W),B(Y,Z)) \\ - g(B(X,Z),B(Y,W))$$

In the Kahler setting, we deduce the following relation between the holomorphic sectional curvatures:

$$\begin{aligned} K(X) &= R(X,JX,JX,X) \\ &= \tilde{R}(X,JX,JX,X) + g(B(X,X),B(JX,JX)) \\ &\quad - g(B(X,JX),B(JX,X)) \\ &= \tilde{K}(X) - 2|B(X,X)|^2 \end{aligned}$$

Recall that a submanifold is called totally geodesic if its second fundamental form vanishes. Since B is symmetric, this is equivalent to saying that $B(X,X) = 0$ for all vectors X tangent to the submanifold. The equation above has the following important consequence:

Proposition 5.6. *A complex submanifold M in a Kahler manifold of constant holomorphic sectional curvature is totally geodesic if and only if it also has constant holomorphic sectional curvature in the induced metric, with the same curvature constant as the ambient space.*

Proof. This is immediate since:

$$K(X) - \tilde{K}(X) = |B(X,X)|^2$$

so $K = \tilde{K}$ if and only if $B = 0$. □

We can use this to completely classify the totally geodesic submanifolds of Kahler manifolds of constant holomorphic curvature.

Proposition 5.7. *A totally geodesic Kahler submanifold in one of the standard models for a simply connected Kahler manifold of constant holomorphic sectional curvature must be one of the following:*

- *An affine subspace in $\mathbb{C}^n$ with its flat Kahler metric.*
- *An affine subspace in $\mathbb{C}P^n$ with its Fubini-Study metric; more precisely, the projection of an affine subspace of $\mathbb{C}^{n+1} - \{0\}$ to $\mathbb{C}P^n$.*
- *The image of a linear subspace of $\mathbb{D}^n$ with its hyperbolic metric which passes through the point $z = 0$ under an isometry of the ambient hyperbolic space.*

Proof. We start by recalling a fact from Riemannian geometry: given a Riemannian manifold M and a subspace of some tangent space, say $V \subset T_P M$,

there is at most one totally geodesic submanifold of M passing through p whose tangent space is V. Briefly, this is because such a submanifold, if it exists, must coincide with $\exp_p V$.

Now we prove the proposition case by case, though the general structure of the argument is the same regardless of the ambient space.

Case 1. *Flat metric.*

Up to an isometry of the ambient space we may assume that the submanifold passes through the point $z = 0$, and that the tangent plane at this point is given by $z_{k+1} = \cdots = z_n = 0$. Further, due to the uniqueness of totally geodesic submanifolds we alluded to above, it suffices for us to construct a single totally geodesic submanifold with this tangent plane. Easy enough, let:

$$M = \{z \in \mathbb{C}^n : z_{k+1} = \cdots = z_n = 0\}$$

The restriction of the metric on the ambient space to M is easily seen to be another flat metric on M, so it also has constant holomorphic curvature zero. Owing to the previous proposition, this guarantees that it is totally geodesic.

The other two cases are technically a little more complicated, but the idea is exactly the same.

Case 2. *Fubini-Study metric.*

Again, let's use the transitive property of the isometry group to arrange it so that our submanifold passes through the point $[1, 0, \ldots, 0]$ and shares a tangent plane with the affine subspace:

$$M = \{[z_0, \ldots, z_n] \in \mathbb{C}P^n : z_{k+1} = \cdots = z_n = 0\}$$

It then suffices for us to show that M is in fact totally geodesic. First, note that this affine subspace is in fact an embedded copy of $\mathbb{C}P^k$, with the embedding function given in homogeneous coordinates by:

$$[z_0, \ldots, z_k] \mapsto [z_0, \ldots, z_k, 0, \ldots, 0]$$

Now, as we saw in a previous section, a Kahler potential for the Fubini-Study metric on $\mathbb{C}P^n$ in the standard coordinate system centred at $[1, 0, \ldots, 0]$ is:

$$f(z) = \log(1 + |z|^2) = \log(1 + z_0\bar{z}_0 + \cdots + z_n\bar{z}_n)$$

By a previous lemma, the restriction of this to M is a Kahler potential for the induced metric on M. But:

$$f|_M(z_0, \ldots, z_k) = \log(1 + z_0\bar{z}_0 + \cdots + z_k\bar{z}_k)$$

which is the Kahler potential for the standard Fubini-Study metric on $\mathbb{C}P^k$. Since M is a Kahler manifold of constant holomorphic sectional curvature whose curvature constant is the same as the ambient space, we conclude that M is totally geodesic.

Case 3. *Hyperbolic Metric*

The proof is the same as for the Fubini-Study metric, but instead uses the potential

$$f(z) = -\log(1 - |z|^2)$$

for the hyperbolic metric.

Since we have covered all three possible cases, the proposition has been proved. □

6. Examples of Kahler-Einstein Submanifolds of Complex Projective Space

In this section we give some examples of Einstein submanifolds of the complex projective space (equipped with the Fubini-Study metric).

6.1. **Totally Geodesic Subspaces.** Recall that in a previous section we were able to completely classify the totally geodesic subspaces in a Kahler manifold of constant holomorphic curvature. In the case of a Fubini study space these were (projections of) intersections of linear subspaces with the ambient space. Of course, these subspaces are also complex projective spaces, and they inherit a Fubini study metric of the same holomorphic curvature as the ambient space. In particular, they are Einstein submanifolds.

6.2. **Embedded Fubini-Study spaces.** Now we move on to a more general class of embedded Fubini-Study spaces, which include the totally geodesic subspaces as a special case. We begin with an example to illustrate the general case.

Example 6.3. We show how to embedd the complex projective space $\mathbb{C}P^2$ into the Fubini-Study space $\mathbb{C}P^5$ so that the induced metric is also Fubini-Study, but so that the image does not lie in any totally geodesic subspace.

Let $[z_0, z_1, z_2]$ be homogeneous coordinates on $\mathbb{C}P^2$, and $[w_0, w_1, w_2, w_3, w_4, w_5]$ homogeneous coordinates on $\mathbb{C}P^5$. Our embedding will be given by the mapping:

$$\phi([z_0, z_1, z_2]) = [z_0^2, \sqrt{2}z_0z_1, \sqrt{2}z_0z_2, z_1^2, \sqrt{2}z_1z_2, z_2^2]$$

which is inspired by the algebraic identity:

$$(z_0 + z_1 + z_2)^2 = z_0^2 + 2z_0z_1 + 2z_0z_2 + z_1^2 + 2z_1z_2 + z_2^2$$

Note that since each of the terms in the embedding is homogeneous of degree 2, we have indeed defined a mapping $\mathbb{C}P^2 \to \mathbb{C}P^5$.

Now consider the inhomogeneous coordinate systems $[1, z_1, z_2]$ and $[1, w_1, w_2, w_3, w_4]$, and notice that the mapping we defined above sends the region covered by the first coordinate system into that covered by the second. The differential of ϕ in these coordinate systems is now easy to compute and is given by:

$$d\phi = \begin{pmatrix} \sqrt{2} & 0 \\ 0 & \sqrt{2} \\ 2z_1 & 0 \\ \sqrt{2}z_2 & \sqrt{2}z_1 \\ 0 & 2z_2 \end{pmatrix}$$

It is easy to see that this matrix has full rank, and so our mapping is an immersion. To see further that it is an embedding we need to check that it is globally one to one. This is simple enough. If $\phi([z_0, z_1, z_2]) = \phi([x_1, x_2, x_3])$ then there is a complex number λ so that:

$$z_0^2 = \lambda^2 x_0^2 \text{ and } z_1^2 = \lambda^2 x_1^2 \text{ and } z_2^2 = \lambda^2 x_2^2$$

Upon taking square roots, we get:

$$z_0 = \pm\lambda x_0 \text{ and } z_1 = \pm\lambda x_1 \text{ and } z_2 = \pm\lambda x_2$$

Now, if the signs in three of these equalities agree, then we get $[z_0, z_1, z_2] = [\lambda x_1, \lambda x_2, \lambda x_3]$, as we wanted. So suppose to the contrary that a pair of signs disagrees, say:

$$z_0 = \lambda x_0 \text{ and } z_1 = -\lambda x_1$$

Then:

$$\sqrt{2}z_0z_1 = -\lambda^2\sqrt{2}x_0x_1$$

But this contradicts the relation $\sqrt{2}z_0z_1 = \lambda^2\sqrt{2}x_0x_1$, which is a consequence of our assumption. Therefore all the signs must agree, and the mapping is globally one to one.

Now we show that the image of the mapping does not lie in any totally geodesic subspace. If this were so then there would be a linear relation:

$$az_0^2 + b\sqrt{2}z_0z_1 + c\sqrt{2}z_0z_2 + dz_1^2 + e\sqrt{z}_1z_2 + fz_2^2) = 0$$

which holds for all values of (z_0, z_1, z_2). Successively setting each of the variables equal to one while letting the two remaining be zero gets us that:

$$a = 0 \text{ and } d = 0 \text{ and } f = 0$$

Similarly, if we successively let two of the variables take the value one, while letting the other be zero, them we get that:

$$\lambda(1,1,0)b = 0 \text{ and } \lambda(1,0,1)c = 0 \text{ and } \lambda(0,1,1)e = 0$$

Therefore, the only linear relation possible is the trivial one, and we conclude that the image of ϕ does not lie in a totally geodesic subspace.

Finally, we verify that the induced metric is Fubini-Study, and explicitly calculate the holomorphic curvature. We continue to use the coordinate systems above. Recall that a Kahler potential for the Fubini-Study metric on the ambient space is:

$$f(w_1, w_2, w_3, w_4, w_5) = \frac{2}{c}\log(1 + |w_1|^2 + |w_2|^2 + |w_3|^2 + |w_4|^2 + |w_5|^2)$$

The potential for the induced metric on $\mathbb{C}P^2$ is the restriction of the potential for the ambient space to the image. Substituting in, we get that the induced potential is:

$$\begin{aligned} f \circ \varphi(z_1, z_2) &= \frac{2}{c}\log(1 + |z_0^2|^2 + 2|z_0 z_1|^2 + 2|z_0 z_2|^2 + |z_1^2|^2 + 2|z_1 z_2|^2 + |z_2^2|^2) \\ &= \frac{2}{c}\log((1 + |z_1|^2 + |z_2|^2)^2) \\ &= \frac{4}{c}\log(1 + |z_1|^2 + |z_2|^2) \end{aligned}$$

This is the potential for the Fubini-Study metric on $\mathbb{C}P^2$ with constant holomorphic curvature $\frac{c}{2}$.

Now we turn to the general case.

To begin, we consider the algebraic identity:

$$(z_0 + z_1 + \cdots z_m)^k = \sum_I c_I z^I$$

where the I's are multi-indices. We include the following combinatorial result for completeness:

Lemma 6.4. *The number of terms in the sum above is* $\begin{pmatrix} m+k \\ k \end{pmatrix}$

Proof. Each term in the sum above corresponds to a monomial of the shape $z_0^{i_1} z_1^{i_1} \cdots z_m^{i_m}$, where each of the i's is a non-negative integer, and $i_0 + i_1 + \cdots + i_m = k$. Think of forming such a monomial by first writing down a list of k stars, for instance, say $k = 5$, then we would start with: $* * * * *$ To communicate a monomial we then group these stars into $m + 1$ collections. We do this by separating the stars with bars, for instance, when $m = 2$ we get the following correspondences:

$$*|**|** \mapsto z_0 z_1^2 z_2^2$$

$$** \,||\, *** \mapsto z_0^2 z_2^3$$

$$|\, ***** \,| \mapsto z_1^5$$

and so on (we hope the reader can see the correspondence from these examples). Now to count the number of possible star and bar diagrams we simply observe that we have $m+n$ spaces we want to fill with k stars and m bars. Choosing the k places to put the stars automatically determines where the bars go, because they must fill up the remaining spaces. Since there are $n+k$ spaces, and we are placing k stars in them, we get the count that were after. □

For notational convenience we will use the abbreviation $C(m+k,k) = \begin{pmatrix} m+k \\ k \end{pmatrix}$ below.

Now we define an embedding $\varphi : \mathbb{C}P^m \to \mathbb{C}P^{C(m+k,k)}$ by:

$$[z_0, z_1, \ldots, z_m] \mapsto [\sqrt{c_I} z^I]_I$$

For example, in the case $m = 2$ and $k = 2$ we get the mapping from our example above. Furthermore, in the case $m = 1$, we get the following embedding of $\mathbb{C}P^1$ into $\mathbb{C}P^k$:

$$[z_0, z_1] \mapsto \left[\sqrt{\begin{pmatrix} n \\ 0 \end{pmatrix}} z_0^k, \sqrt{\begin{pmatrix} n \\ 1 \end{pmatrix}} z_0^{k-1} z_1, \cdots, \sqrt{\begin{pmatrix} n \\ n \end{pmatrix}} z_1^k \right]$$

which was mentioned in the introduction.

We want to determine what metric is induced on $\mathbb{C}P^m$ by this embedding. Working in the coordinate system $[1, w_1, \ldots, w_{C(k,m)}]$ we recall that a potential for the Fubini-Study metric of constant holomorphic curvature c on the ambient space is:

$$f(w_1, \ldots, w_{C(k,m)}) = \frac{2}{c} \log(1 + |w_1|^2 + \cdots + |w_{C(k,m)}|^2)$$

The map φ, when expressed in this coordinate system for the ambient space and the corresponding coordinate system for $\mathbb{C}P^m$, is given by:

$$\varphi[1, z_1, \ldots, z_m] = \left[\sqrt{c_I} z^I|_{z_0=1}\right]_I$$

So a Kahler potential for the induced metric on $\mathbb{C}P^m$ is:

$$\begin{aligned} f \circ \varphi[1, z_1, \ldots, z_m] &= \frac{2}{c} \log \left(\sum_I |\sqrt{c_I} z^I|^2_{z_0=1} \right) \\ &= \frac{2}{c} \log \left(\sum_I c_I |z_I|^2_{z_0=1} \right) \\ &= \frac{2}{c} \log(\ (1 + |z_1|^2 + \cdots + |z_m|^2)^k\ ) \\ &= \frac{2k}{c} \log(1 + |z_1|^2 + \cdots + |z_m|^2) \end{aligned}$$

which is the potential for the fubini-study metric on $\mathbb{C}P^m$ of constant holomorphic sectional curvature c, scaled by a factor of k. Hence the induced metric has constant holomorphic sectional curvature $\frac{c}{k}$, and is consequently a Kahler-Einstein submanifold with Einstein constant $\lambda = \frac{(m+1)c}{2k}$.

6.5. **Products of Fubini-Study Spaces.** We now want to expand the discussion in the previous section to embedded products of Fubini-Study spaces. Again, we start with a simple example to illustrate.

Example 6.6. We show how to embed $\mathbb{C}P^1 \times \mathbb{C}P^1$ into $\mathbb{C}P^3$ so that the induced geometry is a product of Fubini-Study metrics. Again, this embedding will be full in the sense that the image does not lie in any proper totally geodesic subspace.

Let $[x_0, x_1]$ and $[y_0, y_1]$ be homogeneous coordinates on the two $\mathbb{C}P^1$ factors, and let $[w_0, w_1, w_2, w_3]$ be homogeneous coordinates on $\mathbb{C}P^3$. Our embedding is given by the mapping:

$$\varphi([x_0, x_1], [y_0, y_1]) = [x_0 y_0, x_0 y_1, x_1 y_0, x_1 y_1]$$

It is easy to see that this mapping is well defined.

We now want to show that φ is an immersion. To do so, we consider the inhomogeneous coordinate systems $[1, x_1]$, $[1, y_1]$ and $[1, w_1, w_2, w_3]$. Note that the region covered in $\mathbb{C}P^1 \times \mathbb{C}P^1$ by the product of the first two coordinate systems is mapped by φ into the region covered by the third in $\mathbb{C}P^3$. Therefore we can compute the differential of φ in these coordinates, and the result is:

$$d\varphi = \begin{pmatrix} 0 & 1 \\ 1 & 0 \\ y_1 & x_1 \end{pmatrix}$$

This matrix clearly has full rank, so φ is an immersion.

Now we check that φ is globally one to one, and is hence an embedding. Suppose that $\varphi([x_0, x_1], [y_0, y_1]) = \varphi([x'_0, x'_1], [y'_0, y'_1])$. Then there is a non-zero complex number λ so that:

$$x_0y_0 = \lambda x'_0y'_0 \text{ and } x_0y_1 = \lambda x'_0y'_1 \text{ and } x_1y_0 = \lambda x'_1y'_0 \text{ and } x_1y_1 = \lambda x'_1y'_1$$

Dividing the first of these equations by the second, and the first by the third, (or the other way around if necessary) gives us:

$$\frac{y_0}{y_1} = \frac{y'_0}{y'_1} \text{ and } \frac{x_0}{x_1} = \frac{x'_0}{x'_1}$$

So there are non-zero complex numbers a and b such that:

$$y_0 = ay'_0 \text{ and } y_1 = ay'_1$$

and:

$$x_0 = bx'_0 \text{ and } x_1 = bx'_1$$

So we get $[x_0, x_1] = [x'_0, x'_1]$ and $[y_0, y_1] = [y'_0, y'_1]$, which is what we wanted to show.

Now let's show that the image of φ does not lie in any proper totally geodesic subspace. Let:

$$ax_0y_0 + bx_0y_1 + cx_1y_0 + dx_1y_0 = 0$$

be a linear relation that holds for all x_0, x_1, y_0 and y_1. Setting pairs of variables equal to one while letting the other pair be zero immediately allows us to conclude that $a = b = c = d = 0$. So the only linear relation is the trivial one, and consequently the image of φ does not lie in a proper totally geodesic subspace.

We now want to determine the metric induced on $\mathbb{C}P^1 \times \mathbb{C}P^1$ by this embedding. Again, we work in the coordinate systems we already used above, and use the Kahler potential of the ambient space, which we remind the reader is:

$$f(w_1, w_2, w_3) = \frac{2}{c} \log(1 + |w_1|^2 + |w_2|^2 + |w_3|^2)$$

Therefore, a potential for the induced metric on $\mathbb{C}P^1 \times \mathbb{C}P^1$ is:

$$\begin{aligned} f \circ \varphi(x_1, y_1) &= \frac{2}{c} \log(1 + |x_1|^2 + |y_1|^2 + |x_1y_1|^2) \\ &= \frac{2}{c} \log((1 + |x_1|^2)(1 + |y_1|^2)) \\ &= \frac{2}{c} \log(1 + |x_1|^2) + \frac{2}{c} \log(1 + |y_1|^2) \end{aligned}$$

Which the reader will recognise as a potential for the product of two Fubini-Study spaces of constant holomorphic curvature c. Note that since the curvatures of each factor agree, this is an Einstein submanifold.

Now we will discuss how to embed products in general. Before we begin in earnest, let's first discuss when a product of Fubini-Study spaces is a Kahler-Einstein manifold:

Lemma 6.7. *A metric product $\mathbb{C}P^{n_1}(c_1) \times \mathbb{C}P^{n_2}(c_2) \times \cdots \times \mathbb{C}P^{n_k}(c_k)$ of Fubini-Study spaces is Kahler-Einstein if and only if $(n_1+1)c_1 = (n_2+1)c_2 = \cdots = (n_k+1)c_k$.*

Proof. Recall that a Fubini-Study space $\mathbb{C}P^n(c)$ is Kahler-Einstein with constant $\frac{(n+1)c}{2}$, and that a product of Einstein manifolds is itself Einstein if and only if the Einstein constants of the individual factors agree. The lemma follows immediately from these two facts. □

We will only consider the case of a product of two projective spaces in this exposition, but it will be clear to the reader how to generalize our results to the case of more factors. Consider the algebraic identity:

$$(x_0 + x_1 + \cdots + x_n)^r (y_0 + y_1 + \cdots + y_m)^s = \sum_{IJ} c_{IJ} x^I y^J$$

where the I's and the J's are multi-indicies. Note that there are $\begin{pmatrix} n+r \\ r \end{pmatrix}\begin{pmatrix} m+s \\ s \end{pmatrix}$ terms in the sum on the right, which we abbreviate $C(n+r,r)C(m+s,s)$ for the same typographical reasons as before.

Now consider the product manifold $\mathbb{C}P^n \times \mathbb{C}P^m$. Similarly to before, we define a mapping:

$$\varphi : \mathbb{C}P^n \times \mathbb{C}P^m \to \mathbb{C}P^{C(n+r,r)C(m+s,s)-1}$$

whose expression in the obvious homogeneous coordinates is:

$$\varphi([x_0, x_1, \ldots, x_n], [y_0, y_1, \ldots, y_m]) = [\sqrt{c_{IJ}} x^I y^J]_{IJ}$$

Notice that the previous example concerned the case $n = m = r = s = 1$.

We want to determine what metric is induced on the product by this embedding. To do so we, as usual, restrict the Kahler potential f for the metric

on the ambient space to our submanifold. We express the result in inhomogeneous coordinates (as before). The result is:

$$\begin{aligned} f \circ \varphi([1, x_1, \ldots, x_n], [1, y_1, \ldots, y_m]) &= \frac{2}{c} \log \left(\sum_{IJ} |\sqrt{c_{IJ}} x^I y^J|^2_{x_0=y_0=1} \right) \\ &= \frac{c}{2} \log \left(\sum_{IJ} c_{IJ} |x^I y^J|^2_{x_0=y_0=1} \right) \\ &= \frac{c}{2} \log \left((1 + |x_1|^2 + \cdots + |x_n|^2)^r (1 + |y_1|^2 + \cdots + |y_m|^2)^s \right) \\ &= \frac{cr}{2} \log(1 + |x_1|^2 + \cdots + |x_n|^2) + \frac{cs}{2} \log(1 + |y_1|^2 + \cdots + |y_m|^2) \end{aligned}$$

Which we recognise as the potential for a product of Fubini-Study metrics, of holomorphic curvature rc on the first factor, and holomorphic curvature sc on the second factor. By the previous lemma, this is an Einstein metric if and only if:

$$(n+1)rc = (m+1)sc \Leftrightarrow (n+1)r = (m+1)s$$

The Einstein constant of the metric in this case is the same as the Einstein constant of the metric on either of the factors, which is:

$$\lambda = \frac{(n+1)c}{2r} = \frac{(m+1)c}{2s}$$

6.8. **The Hypersphere.** Consider the variety Q^{n-1} in $\mathbb{C}P^n$ defined by the homogeneous polynomial equation:

$$z_0^2 + z_1^2 + \cdots + z_n^2 = 0$$

which is called the hypersphere. It follows easily from the implicit function theorem that this is a codimension one, holomorphically embedded submanifold in $\mathbb{C}P^n$. We will show in this section that it is Kahler-Einstein.

Let $Z_0 : Q^{n-1} \to \mathbb{C}^{n+1} - \{0\}$ be a local section of the projection map $\pi : \mathbb{C}^{n+1} \to \mathbb{C}P^n$ over Q^{n-1}, and let $\{e_1, \ldots, e_{n-1}\}$ be a local unitary frame field for TQ^{n-1} with the same domain of definition as Z_0. Recall from a previous section that there is a unitary frame $\{Z_1, \ldots, Z_{n-1}\}$ in the unitary complement of Z_0 corresponding to $\{e_1, \ldots, e_{n-1}\}$. We can extend this frame to a basis in the following way:

Lemma 6.9. *The vectors $\{Z_0, Z_1, \ldots, Z_{n-1}, \bar{Z}_0\}$ form a unitary frame for $\mathbb{C}^{n+1}$.*

Proof. Since $\{Z_1, Z_2, \ldots, Z_{n-1}\}$ lie in the unitary complement of Z_0, we need to verify that the equations $\bar{Z}_0 \cdot Z_0$ and $Z_i \cdot \bar{Z}_0$ hold for $i = 1, \ldots, n-1$, where $\cdot$ denotes the standard Hermitian inner product in $\mathbb{C}^{n+1}$.

For the first equation, note that the components of the vector

$$Z_0 = (z_{00}, z_{01}, \ldots, z_{0n+1})$$

satisfy the defining equation of Q^{n-1}, and hence:

$$Z_0 \cdot \bar{Z}_0 = z_{00}^2 + \cdots + z_{0n+1}^2 = 0$$

To check the other equations, recall that Z_i is an isometric lift of the tangent vector e_i. Let $e_i(t)$ be a curve in Q^{n-1} so that $e_i'(0) = e_i$, and let $Z_i(t)$ be a lift of this curve satisfying $Z_i'(0) = Z_i$. Then the components of $Z_i(t)$ satisfy the defining equation of Q^{n-1} for every t, that is, $Z_i(t) \cdot \bar{Z}_0 = 0$ for every t. Differentiating this equation gives:

$$0 = Z_i'(t) \cdot \bar{Z}_0$$

for every t. Evaluating this at $t = 0$ gives the result that we want. □

We are almost ready to show that Q^{n-1} is an Einstein submanifold, but we need a little more preparation. First, differentiating the equation $Z_j \cdot \bar{Z}_0 = 0$ gives us:

$$0 = d(Z_j \cdot \bar{Z}_0) = dZ_j \cdot \bar{Z}_0 + Z_j \cdot d\bar{Z}_0$$

so:

$$dZ_j \cdot \bar{Z}_0 = -Z_j \cdot \bar{Z}_0 = -dZ_0 \cdot \bar{Z}_j$$

Now since $\{Z_1, \ldots, Z_{n-1}\}$ corresponds isometrically to the unitary frame field $\{e_1, \ldots, e_{n-1}\}$, it follows that the unitary frame $\{\bar{Z}_1, \ldots, \bar{Z}_{n-1}\}$ corresponds to $\{\bar{e}_1, \ldots, \bar{e}_{n-1}\}$. The calculation above shows that:

$$\phi_j^n = -dZ_0 \cdot \bar{Z}_j = -\bar{e}^j$$

Corresponding to the unitary frame $\{Z_1, \ldots, Z_{n-1}, \bar{Z}_0\}$ for the unitary complement of Z_0, there is a unitary frame $\{e_1, \ldots, e_{n-1}, e_n\}$ for $T\mathbb{C}P^n|_{Q^{n-1}}$, let θ be the connection form for the Fubini-Study metric in this frame field. Then the $(n-1) \times (n-1)$ submatrix in the upper left hand corner of θ is the connection form matrix for the induced metric on Q^{n-1}. Let Ω be the corresponding curvature form for Q^{n-1}. In the subsequent calculation we consider all differential forms as restricted to Q^{n-1}, note in this context that we have:

$$\phi_0^n = e^n = 0$$

Now we can calculate. Below the greek index α always runs over the range $1, \ldots, n$ and k runs over the range $1, \ldots, n-1$:

$$\begin{aligned}
\text{tr } \Omega = \Omega_j^j &= d\theta_j^j + \theta_k^j \wedge \theta_j^k \\
&= d(\phi_j^j - \delta_j^j \phi_0^0) + (\phi_k^j - \delta_k^j \phi_0^0) \wedge (\phi_j^k - \delta_j^k \phi_0^0) \\
&= -\phi_\alpha^j \wedge \phi_j^\alpha + \delta_j^j \phi_\alpha^0 \wedge \phi_0^\alpha + \phi_k^j \wedge \phi_j^k - \delta_j^k \phi_k^j \wedge \phi_0^0 \\
&\quad - \delta_k^j \phi_0^0 \wedge \phi_j^k + \delta_k^j \delta_j^k \phi_0^0 \wedge \phi_0^0 \\
&= -\phi_n^j \wedge \phi_j^n - \phi_k^j \wedge \phi_j^k + (n-1)\phi_n^0 \wedge \phi_0^n + (n-1)\phi_k^0 \wedge \phi_0^k \\
&\quad + \phi_k^j \wedge \phi_j^k - \phi_j^j \wedge \phi_0^0 - \phi_0^0 \wedge \phi_j^j \\
&= -\phi_n^j \wedge \phi_j^n + (n-1)\phi_k^0 \wedge \phi_0^k \\
&= -\phi_j^n \wedge \bar{\phi}_j^n + (n-1)\phi_0^k \wedge \bar{\phi}_0^k \\
&= -\bar{e}^j \wedge e^j + (n-1)e^k \wedge \bar{e}^k \\
&= ne^k \wedge \bar{e}^k \\
&= -2ni\frac{i}{2}e^k \wedge \bar{e}^k \\
&= -2i\omega
\end{aligned}$$

So we see that Q^{n-1} is Einstein of constant 2. Since our ambient space in this calculation had constant holomorphic curvature 4, we see that in general the hypersphere is a codimension 1 Einstein submanifold of $\mathbb{C}P(c)$, with Einstein constant $\frac{nc}{2}$.

7. Calabi's Theorem

In this section we give a purely differential geometric proof of Calabi's theorem on one dimensional complex submanifolds of a complex space form.

First we need some setup. Fix a one dimensional holomorphically immersed manifold M in the space form M_c, and denote with angle brackets $\langle \quad , \quad \rangle$ the Hermitian metric on M_c, which restricts to the Hermitian metric on M.

Let $\{e_0, e_1, e_2, \ldots, e_n\}$ be a local unitary frame field for the bundle TM_c^+, chosen so that $\{e_1, \ldots e_n\}$ is a unitary frame for the normal bundle NM^+, and $\{e_0\}$ is a unit vector in the tangent bundle TM^+. Let $\{e^0, e^1, \ldots, e^n\}$ be the dual unitary frame field. We study the connection forms Θ and θ for M_c and M with respect to these frames. Recall that the connection form for M_c is a matrix of one forms satisfying:

$$\nabla e_\alpha = \Theta_\alpha^\beta e_\beta$$

where ∇ is the connection on M_c. The connection form θ (which in this context is just a single one form, not a matrix of one forms) is defined analogously using the connection on M.

Remark 7.1. In the subsequent calculations we make the following conventions on all indices: the lower case Greek index α will always have the range $0, 1, 2, \ldots, n$, and lower case Roman indices i, j, k will always have the range $1, 2, \ldots, n$. The reader should also note that repeated indices in a single term are automatically summed over their ranges.

Furthermore, every differential form in the calculations below is always considered as a form on M by pulling back through the immersion $M \to M_c$. In this context we have, for instance, that $e^i = 0$ for all i, and $\Theta^0_0 = \theta$.

Finally, we remind the reader that since we are using a unitary frame field, the curvature forms satisfy the conjugation relations $\bar{\theta} = -\theta$ and $\bar{\Theta}^i_j = -\Theta^j_i$.

To begin our work, we differentiate the expression $e^i = 0$ and use the torsion-free property of the Kahler connection. The result is:

$$0 = de^i = -\Theta^i_\alpha \wedge e^\alpha = -\Theta^i_0 \wedge e^0$$

It follows from the above equation that Θ^i_0 is a $(1,0)$-form on M, and therefore there are some locally defined functions b^i_0 satisfying:

$$\Theta^i_0 = b^i_0 e^0 \tag{1}$$

Lemma 7.2. *The immersion $M \to M_c$ is totally geodesic if and only if $b^i_0 = 0$ for all i.*

Proof. Let $B : TM^+ \times TM^+ \to NM^+$ be the second fundamental form of the immersion. We have:

$$\begin{aligned}\langle B(e_0, e_0), e_i\rangle &= \left\langle \tilde{\nabla}_{e_0} e_0, e_i \right\rangle = \langle \Theta^\alpha_0(e_0) e_\alpha, e_i\rangle \\ &= \Theta^i_0(e_0) = b^i_0\end{aligned}$$

So B vanishes exactly when $b^i_0 = 0$ for every i. □

It follows from the proof above that the second fundamental form can be expressed as:

$$B(e_0, e_0) = b^i_0 e_i$$

Now let ω and Ω be the curvature forms of M and M_c respectively. The fundamental equations for ω and Ω are:

$$\omega = d\theta + \theta \wedge \theta = d\theta$$

$$\Omega = d\Theta + \Theta \wedge \Theta$$

Recalling that $\Omega_0^0 = \theta$, and using (1), we get:

$$\omega = d\Theta_0^0 + \Theta_0^0 \wedge \Theta_0^0 = \Omega_0^0 - \Theta_i^0 \wedge \Theta_0^i = \Omega_0^0 - b_0^i \bar{b}_0^i e^0 \wedge \bar{e}^0$$

Now, by assumption, M_c has constant holomorphic sectional curvature c and M is Einstein of constant λ, which in this case (dimension 1) means that M has constant holomorphic sectional curvature λ. Therefore we have:

$$\omega = \frac{\lambda}{2} e^0 \wedge \bar{e}^0$$
$$\Omega_0^0 = \frac{c}{2} e^0 \wedge \bar{e}^0$$

Substituting these into the above equation we get:

$$\frac{\lambda}{2} e^0 \wedge \bar{e}^0 = \frac{c}{2} e^0 \wedge \bar{e}^0 - b_0^i \bar{b}_0^i e^0 \wedge \bar{e}^0$$

Hence, since $e^0 \wedge \bar{e}^0$ spans the space of $(1,1)$ forms on M, it follows that:

$$b_0^i \bar{b}_0^i = \frac{1}{2}(c - \lambda) \tag{2}$$

Corollary 7.3. *M is totally geodesic if and only if $c = \lambda$.*

Proof. This follows immediately from equation 2 and the previous lemma. □

To continue, we begin by differentiating (1):

$$d\Theta_0^i = db_0^i \wedge e^0 + b_0^i de^0$$

Using the torsion free property of the metric again we have:

$$de^0 = -\theta \wedge e^0$$

Applying the fundamental equations, and the formula for the curvature in a holomorphic space form, we get:

$$\begin{aligned} d\Theta_0^i &= \Omega_0^i - \Theta_\alpha^i \wedge \Theta_0^\alpha \\ &= \frac{c}{4} e^i \wedge \bar{e}^0 - \Theta_0^i \wedge \theta - \Theta_j^i \wedge \Theta_0^j \\ &= b_0^i \theta \wedge e^0 - b_0^j \Theta_j^i \wedge e^0 \end{aligned}$$

Which, upon substituting into the above, gives us:

$$0 = (db_0^i - 2b_0^i \theta + b_0^j \Theta_j^i) \wedge e^0$$

It follows from the above equation that $d_0^i - 2b_0^i\theta + b_0^j\Theta_j^i$ is a $(1,0)$ form, and hence there must be some functions b_1^i so that:

$$b_1^i e^0 = db_0^i - 2b_0^i \theta + b_0^j \Theta_j^i \tag{3}$$

Lemma 7.4. *The following orthogonality relations hold:*

(1) $b_1^i \bar{b}_0^i = 0$

(2) $b_1^i \bar{b}_1^i = \frac{3}{8}(c-\lambda)(c-2\lambda)$

Proof. We begin by differentiating (2):

$$0 = \bar{b}_0^i db_0^i + b_0^i d\bar{b}_0^i$$

Now (3), along with the skew-Hermitian property for the connection form, can be used to simplify the expression above:

$$\bar{b}_0^i db_0^i = \bar{b}_0^i b_1^i e^0 + 2\bar{b}_0^i b_0^i \theta - \bar{b}_0^i b_0^j \Theta_j^i$$
$$b_0^i d\bar{b}_0^i = b_0^i \bar{b}_1^i \bar{e}^0 - 2\bar{b}_0^i b_0^i + b_0^i \bar{b}_0^j \Theta_i^j$$

Upon substituting this into the first equation,and switching the summation indices to cancel some terms, we get:

$$0 = \bar{b}_0^i b_1^i e^0 + b_0^i \bar{b}_1^i \bar{e}^0$$

So, since e^0 and $\bar{e}^0$ are linearly independent, we conclude that:

$$b_1^i \bar{b}_0^i = 0 \tag{4}$$

Which proves part (a) of the lemma.

To prove (b), we differentiate the equation $\bar{b}_0^i b_1^i e^0 = 0$, which is a consequence of (4) above. This gives us:

$$0 = b_1^i d\bar{b}_0^i \wedge e^0 + \bar{b}_0^i d(b_1^i e^0)$$

For the first of these terms, we use (3) to substitute in for $d\bar{b}_0^i$, then simplify with (4) where appropriate:

$$b_1^i d\bar{b}_0^i \wedge e^0 = b_1^i \bar{b}_1^i \bar{e}^0 \wedge e^0 + b_1^i \bar{b}_0^i \Theta_i^j \wedge e^0$$

For the second of the terms, we first differentiate equation (3) and substitute in the result:

$$\bar{b}_0^i d(b_1^i e^0) = -2\bar{b}_0^i db_0^i \wedge \theta - 2\bar{b}_0^i b_0^i d\theta + \bar{b}_0^i db_0^j \wedge \Theta_j^i + \bar{b}_0^i b_0^j d\Theta_j^i$$

Of the four terms above, we simplify the first and third by using (3) to substitute in for the occurrences of db_0^i, then apply the orthogonality relations already derived. Equation (2) and the Einstein equation $2d\theta = \lambda e^0 \wedge \bar{e}^0$ are used to simplify the second term. The results are:

$$-2\bar{b}_0^i db_0^i \wedge \theta = 2\bar{b}_0^i b_0^j \Theta_j^i \wedge \theta$$
$$-2\bar{b}_0^i b_0^i d\theta = -\frac{1}{2}\lambda(c-\lambda) e^0 \wedge \bar{e}^0$$
$$\bar{b}_0^i db_0^j \wedge \Theta_j^i = \bar{b}_0^i b_1^j e^0 \wedge \Theta_j^i + 2\bar{b}_0^i b_0^j \theta \wedge \Theta_j^i - \bar{b}_0^i b_0^k \Theta_k^j \wedge \Theta_j^i$$

For the last term, we recall our formula for the curvature form in a space of constant holomorphic curvature:

$$\begin{aligned} d\Theta_j^i &= \Omega_j^i - \Theta_\alpha^i \wedge \Theta_i^\alpha \\ &= \frac{c}{4}\delta_j^i e^0 \wedge \bar{e}^0 + b_0^i \bar{b}_0^j e^0 \wedge \bar{e}^0 - \Theta_k^i \wedge \Theta_j^k \end{aligned}$$

Substituting this expression in, and using the various orthogonality relations we have already obtained, we get:

$$\bar{b}_0^i b_0^j d\Theta_j^i = \frac{1}{8}c(c-\lambda)e^0 \wedge \bar{e}^0 + \frac{1}{4}(c-\lambda)^2 e^0 \wedge \bar{e}^0 - \bar{b}_0^i b_0^j \Theta_k^i \wedge \Theta_j^k$$

Now we substitute these results into the first equation of the paragraph, and observe that all the terms involving a θ or a Θ cancel. We are left with:

$$0 = b_1^i \bar{b}_1^i \bar{e}^0 \wedge e^0 - \frac{1}{2}\lambda(c-\lambda)e^0 \wedge \bar{e}^0 + \frac{1}{8}c(c-\lambda)e^0 \wedge \bar{e}^0 + \frac{1}{4}(c-\lambda)^2 e^0 \wedge \bar{e}^0$$

A short calculation yields:

$$\begin{aligned} &\frac{1}{8}c(c-\lambda)e^0 \wedge \bar{e}^0 + \frac{1}{4}(c-\lambda)^2 e^0 \wedge \bar{e}^0 - \frac{1}{2}\lambda(c-\lambda)e^0 \wedge \bar{e}^0 \\ &\qquad = \frac{3}{8}(c-\lambda)(c-2\lambda)e^0 \wedge \bar{e}^0 \end{aligned}$$

and therefore:

$$b_1^i \bar{b}_1^i = \frac{3}{8}(c-\lambda)(c-2\lambda) \tag{5}$$

which completes the proof of (b), and hence of the lemma. □

At this point we would like to start an inductive process based on the procedure we used to arrive at the previous lemma. Unfortunately, to do this honestly, one more step must be worked out separately. We spare the readers the details of this, and state the results as another lemma, as the argument differs only slightly from the actual induction step we will outline below.

Lemma 7.5. *There are locally defined functions b_2^i which satisfy the relations:*

$$b_2^i e^0 = db_1^i - 3b_1^i\theta + b_1^j \Theta_j^i + \frac{3}{4}(c-2\lambda)b_0^i \bar{e}^0$$

These functions satisfy the orthogonality relations:

(1) $b_2^i \bar{b}_1^i = b_2^i \bar{b}_0^i = 0$
(2) $b_2^i \bar{b}_2^i = \frac{3}{8}(c-\lambda)(c-2\lambda)(c-3\lambda)$

Proof. As stated, the proof is omitted. □

Now we have assembled enough facts to start an inductive procedure. We prove the following lemma:

Lemma 7.6. *Set $b^i_{-1} = 0$. For each $r = 1, 2, \dots$ there are locally defined functions b^i_r satisfying:*

$$b^i_r e^0 = db^i_{r-1} - (r+1)b^i_{r-1}\theta + b^j_{r-1}\Theta^i_j + \frac{r+1}{4}(c - r\lambda)b^i_{r-2}\bar{e}^0 \tag{6}$$

For $0 \leq s \leq r$ these functions satisfy the orthogonality relations:

$$b^i_r \bar{b}^i_s = \delta^r_s \frac{(r+2)!}{4^{r+1}}(c-\lambda)(c-2\lambda)\cdots(c-(r+1)\lambda) \tag{7}$$

Proof. The reader may easily check that we have already verified formula (6) for $r = 1, 2$, and formula (7) for $r, s = 0, 1, 2$. So we assume, towards induction, that we have verified the formula for the cases $1, 2, \dots, r$, and work on proving them for $r + 1$.

We begin by differentiating (6):

$$\begin{aligned} db^i_r \wedge e^0 + b^i_r de^0 = &- (r+1)db^i_{r-1} \wedge \theta - (r+1)b^i_{r-1}d\theta \\ &+ db^j_{r-1} \wedge \Theta^i_j + b^j_{r-1}d\Theta^i_j \\ &+ \frac{r+1}{4}(c - r\lambda)db^i_{r-2}\bar{e}^0 + \frac{r+1}{4}(c-r\lambda)b^i_{r-2}d\bar{e}^0 \end{aligned}$$

For the term on the left hand side, we use the torsion free metric property to get:

$$b^i_r \wedge de^0 = -b^i_r \theta \wedge e^0$$

To simplify the terms on the right hand side, we use the inductive assumptions and the procedures outlined in the exposition above. The results are:

$$\begin{aligned} -(r+1)db^i_{r-1} \wedge \theta = &- (r+1)b^i_r e^0 \wedge \theta + (r+1)b^j_{r-1}\Theta^i_j \wedge \theta \\ &+ \frac{(r+1)^2}{4}(c - r\lambda)b^i_{r-2}\bar{e}^0 \wedge \theta \end{aligned}$$

$$-(r+1)b^i_{r-1}d\theta = -\frac{r+1}{2}\lambda b^i_{r-1}e^0 \wedge \bar{e}^0$$

$$\begin{aligned} db^j_{r-1} \wedge \Theta^i_j = &\, b^j_r e^0 \wedge \Theta^i_j + (r+1)b^j_{r-1}\theta \wedge \Theta^i_j \\ &- b^k_{r-1}\Theta^j_k \wedge \Theta^i_j - \frac{r+1}{4}(c-r\lambda)b^j_{r-2}\bar{e}^0 \wedge \Theta^i_j \end{aligned}$$

$$b^j_{r-1}d\Theta^i_j = \frac{c}{4}b^i_{r-1}e^0 \wedge \bar{e}^0 - b^j_{r-1}\Theta^i_k \wedge \Theta^k_j$$

$$\begin{aligned} \frac{r+1}{4}(c-r\lambda)db^i_{r-2} \wedge \bar{b}^0 = &\frac{r+1}{4}(c-r\lambda)b^i_{r-1}e^0 \wedge \bar{e}^0 \\ &+ \frac{(r+1)r}{4}(c-r\lambda)b^i_{r-2}\theta \wedge \bar{e}^0 \\ &- \frac{r+1}{4}(c-r\lambda)b^j_{r-2}\Theta^i_j \wedge \bar{e}^0 \end{aligned}$$

$$\frac{r+1}{4}(c-r\lambda)b^i_{r-2}d\bar{e}^0 = \frac{r+1}{4}(c-r\lambda)b^i_{r-2}\theta\wedge\bar{e}^0$$

Substituting these results in, we see that three pairs of terms cancel, and we are left, upon gathering like terms, with:

$$\begin{aligned} db^i_r\wedge e^0 = (r+2)b^i_r\theta\wedge e^0 - b^j_r\Theta^i_j\wedge e^0 \\ -\left(\frac{c}{4}-\frac{r+1}{2}\lambda+\frac{r+1}{4}(c-r\lambda)\right)b^i_{r-1}\bar{e}^0\wedge e^0 \\ +\left(\frac{(r+1)r}{4}-\frac{(r+1)^2}{4}+\frac{r+1}{4}\right)(c-r\lambda)b^i_{r-2}\theta\wedge\bar{e}^0 \end{aligned}$$

A short calculation reveals that:

$$\frac{c}{4}-\frac{r+1}{2}\lambda+\frac{r+1}{4}(c-r\lambda) = \frac{r+2}{4}(c-(r+1)\lambda)$$

and:

$$\frac{(r+1)r}{4}-\frac{(r+1)^2}{4}+\frac{r+1}{4} = 0$$

Therefore:

$$0 = \left(db^i_r-(r+2)b^i_r\theta+b^j_r\Theta^i_j+\frac{r+2}{4}(c-(r+1)\lambda)b^i_{r-1}\bar{e}^0\right)\wedge e^0$$

As before, we observe that the coefficient of e^0 above must be a (1,0) form, and hence there are some locally defined functions b^i_{r+1} so that:

$$b^i_{r+1}e^0 = db^i_r-(r+2)b^i_r\theta+b^j_r\Theta^i_j+\frac{r+2}{4}(c-(r+1)\lambda)b^i_{r-1}\bar{e}^0$$

Which completes the inductive verification of formula (6).

Remark 7.7. We included lemma 3 separately from the inductive procedure to be justified in our use of formula (6) to simplify the term involving db^i_{r-2}.

We now go on to verify the orthogonality relations. To simplify our notation, we write:

$$K_r = \frac{(r+2)!}{4^{r+1}}(c-\lambda)(c-2\lambda)\cdots(c-(r+1)\lambda)$$

Observe that these numbers satisfy the recursive relation:

$$K_r = \frac{r+2}{4}(c-(r+1)\lambda)K_{r-1}$$

Suppose now that $0\leq s\leq r$. We differentiate the equation $b^i_r\bar{b}^i_s = \delta^r_s K_r$, which is a consequence of (7). The result is:

$$\bar{b}^i_s db^i_r + b^i_r d\bar{b}^i_s = 0$$

Using equations (6) and (7) to simplify yields the results:

$$\bar{b}^i_s db^i_r = \bar{b}^i_s b^i_{r+1} e^0 - (r+2)\delta^r_s K_r \theta - \bar{b}^i_s b^j_r \Theta^i_j - \frac{r+2}{4}\delta^s_{r-1} K_{r-1}(c-(r+1)\lambda)\bar{e}^0$$

$$b^i_r d\bar{b}^i_s = b^i_r \bar{b}^i_{s+1}\bar{e}^0 + (s+2)\delta^s_r K_r \theta + b^i_r \bar{b}^j_s \Theta^j_i - \frac{r+2}{4}\delta^r_{s-1} K_r (c-(r+1)\lambda)e^0$$

We examine these two equations on a case by case basis.

Case 4. $s \neq r-1, r, r+1$

The above equations yield that:

$$\begin{aligned} 0 &= \bar{b}^i_s b^i_{r+1} e^0 - \bar{b}^i_s b^j_r \Theta^i_j + b^i_r \bar{b}^j_s \Theta^j_i \\ &= \bar{b}^i_s b^i_{r+1} e^0 \end{aligned}$$

Which proves (7) for $s = 1, 2, \ldots, r-2$.

Case 5. $s = r-1$

We get in this case that:

$$\begin{aligned} 0 &= \bar{b}^i_{r-1} b^i_{r+1} e^0 - b^j_r \bar{b}^i_{r-1}\Theta^i_j - \frac{r+2}{4}(c-(r+1)\lambda)K_{r-1}\bar{e}^0 + K_r \bar{e}^0 + b^i_r \bar{b}^j_{r-1}\Theta^j_i \\ &= \bar{b}^i_{r-1} b^i_{r+1} e^0 \end{aligned}$$

Which proves (7) for $s = r-1$

Case 6. $s = r$

We have:

$$\begin{aligned} 0 = b^i_{r+1}\bar{b}^i_r e^0 &- (r+2)K_r\theta - \bar{b}^i_r b^j_r \Theta^i_j \\ &+ b^i_r \bar{b}^i_{r+1}\bar{e}^0 + (r+2)K_r\theta + b^i_r \bar{b}^j_r \Theta^j_i \end{aligned}$$

Which, upon cancelling terms becomes:

$$0 = b^i_{r+1}\bar{b}^i_r e^0 + b^i_r \bar{b}^i_{r+1}\bar{e}^0$$

Which proves (7) for $s = r$.

At this point we have completed our argument, except for verifying the single relation:

$$b^i_{r+1}\bar{b}^i_{r+1} = \frac{(r+3)!}{4^{r+2}}(c-\lambda)\cdots(c-(r+2)\lambda)$$

To get this, we differentiate the equation:

$$\bar{b}^i_r b^i_{r+1} e^0 = 0$$

Which becomes:

$$0 = b^i_{r+1} d\bar{b}^i_r \wedge e^0 + \bar{b}^i_r d(b^i_{r+1}e^0)$$

To simplify the first term, we use equation (6) and the orthogonality relations we have already proved to get:

$$b^i_{r+1}d\bar{b}^i_r \wedge e^0 = b^i_{r+1}\bar{b}^i_{r+1}\bar{e}^0 \wedge e^0 + b^i_{r+1}\bar{b}^j_r\Theta^j_i \wedge e^0$$

For the second term, we differentiate equation (6) (using $r+1$ instead of r) to get:

$$\begin{aligned}\bar{b}^i_r d(b^i_{r+1}e^0) = &-(r+2)\bar{b}^i_r db^i_r \wedge \theta - (r+2)\bar{b}^i_r b^i_r d\theta + \bar{b}^i_r db^j_r \wedge \Theta^i_j + \bar{b}^i_r b^j_r d\Theta^i_j \\ &+ \frac{r+2}{4}(c-(r+1)\lambda)\bar{b}^i_r db^i_{r-1} \wedge \bar{e}^0 + \frac{r+1}{4}(c-(r+1)\lambda)\bar{b}^i_r b^i_{r+1}d\bar{e}^0\end{aligned}$$

and now simplify these terms using the usual techniques:

$$-(r+2)\bar{b}^i_r db^i_r \wedge \theta = (r+2)\bar{b}^i_r b^j_r \Theta^i_j \wedge \theta$$

$$-(r+2)\bar{b}^i_r b^i_r d\theta = -\frac{(r+2)^2(r+1)!}{2\cdot 4^{r+1}}\lambda(c-\lambda)\cdots(c-(r+1)\lambda)e^0 \wedge \bar{e}^0$$

$$\begin{aligned}\bar{b}^i_r db^j_r \wedge \Theta^i_j = &\bar{b}^i_r b^j_{r+1}e^0 \wedge \Theta^i_j + (r+2)\bar{b}^i_r b^j_r \theta \wedge \Theta^i_j - \bar{b}^i_r b^k_r \Theta^j_k \wedge \Theta^i_j \\ &- \frac{r+2}{4}(c-(r+1)\lambda)\bar{b}^i_r b^j_{r-1}\bar{e}^0 \wedge \Theta^i_j\end{aligned}$$

$$\bar{b}^i_r b^j_r d\Theta^i_j = \frac{c(r+2)!}{4^{r+2}}(c-\lambda)\cdots(c-(r+1)\lambda)e^0 \wedge \bar{e}^0 - \bar{b}^i_r b^j_r \Theta^i_k \wedge \Theta^k_j$$

$$\begin{aligned}\frac{r+2}{4}(c-(r+1)\lambda)\bar{b}^i_r db^i_{r-1} \wedge \bar{e}^0 = &\\ = \frac{(r+2)^2(r+1)!}{4^{r+2}}&(c-\lambda)\cdots(c-(r+1)\lambda)^2 e^0 \wedge \bar{e}^0 \\ - \frac{r+2}{4}&(c-(r+1)\lambda)\bar{b}^i_r b^j_{r-1}\Theta^i_j \wedge \bar{e}^0\end{aligned}$$

$$\frac{r+1}{4}(c-(r+1)\lambda)\bar{b}^i_r b^i_{r+1}d\bar{e}^0 = 0$$

Substituting these results in above, cancelling, and gathering the remaining like terms, we get:

$$\begin{aligned}&b^i_{r+1}\bar{b}^i_{r+1}e^0 \wedge \bar{e}^0 = \\ &\quad\frac{(r+2)!}{4^{r+1}}(c-\lambda)\cdots(c-(r+1)\lambda)\left(\frac{c}{4} + \frac{r+2}{4}(c-(r+1)\lambda) - \frac{r+2}{2}\lambda\right)e^0 \wedge \bar{e}^0\end{aligned}$$

A quick calculation shows that:

$$\frac{c}{4} + \frac{r+2}{4}(c-(r+1)\lambda) - \frac{r+2}{2}\lambda = \frac{r+3}{4}(c-(r+2)\lambda)$$

so we get:

$$b^i_{r+1}\bar{b}^i_{r+1}e^0 \wedge \bar{e}^0 = \frac{(r+3)!}{4^{r+2}}(c-\lambda)\cdots(c-(r+2)\lambda)e^0 \wedge \bar{e}^0$$

which completes the proof of (7), and hence, of the lemma. □

With this result, we can immediately give a proof of Calabi's theorem.

Theorem 7.8. *Suppose M is a complex curve. Then, if $c \leq 0$, M is totally geodesic, and if $c > 0$, the Einstein constant λ is one of the numbers $c, \frac{c}{2}, \frac{c}{3}, \ldots, \frac{c}{n+1}$, where $n+1 = \dim M_c$.*

Proof. For a fixed point on M we consider the vectors $b_r = (b_r^1, \ldots, b_r^n)$ for $r = 0, 1, 2, \ldots, n$. The lemma above tells us that b_r is orthogonal to b_s for $r \neq s$, and that:

$$|b_r|^2 = \frac{(r+2)!}{4^{r+1}}(c-\lambda)(c-2\lambda)\cdots(c-(r+1)\lambda)$$

Since these vectors reside in an n dimensional vector space, and there are $n+1$ of them, they cannot all be nonzero. Therefore:

$$0 = \frac{(r+2)!}{4^{r+1}}(c-\lambda)(c-2\lambda)\cdots(c-(r+1)\lambda)$$

for some $0 \leq r \leq n$. It follows immediately that λ must be one of the numbers $c, \frac{c}{2}, \frac{c}{3}, \ldots, \frac{c}{n+1}$. Furthermore, if $c < 0$, then the Gauss equation of the immersion implies that $\lambda \leq c$, and in this case the only possibility is $c = \lambda$. As we saw in lemma 2, this means that M must be totally geodesic in M_c. □

8. Chern's Theorem

We now turn to a more general situation. Let M be any complex manifold, holomorphically immersed in the space form M_c so that the induced Kahler metric on M is Einstein, with Einstein constant λ. Suppose the dimension of M is n, and the dimension of M_c is $n+m$. Let $\{e_1, \ldots, e_{n+m}\}$ be a local unitary frame field for TM_c^+, chosen so that, when restricted to M, $\{e_1, \ldots, e_n\}$ is a local unitary frame field for TM^+, and $\{e_{n+1}, \ldots, e_{n+m}\}$ is a local unitary frame field for NM^+. Let $\{e^1, \ldots, e^{n+m}\}$ be the dual unitary frame field. Again, we study the connection forms Θ and θ for M_c and M.

Remark 8.1. We make use of index conventions similar to those in the previous section. Lower case Greek indices α, β, γ will have the range $1, 2, \ldots, n+m$, lower case Roman indices i, j, k, l, r, s, t will have the range $1, 2, \ldots, n$, and the lower case Roman indices a, b, c will have the range $n+1, n+2, \ldots, n+m$. Note that repeated indices in a single term are automatically summed over their ranges, though we will occasionally drop this convention for the sake of clarity when convenient.

Furthermore, every differential form in the subsequent calculation is always restricted to a form on M. In this context we have $e^a = 0$ for all a, and $\Theta_j^i = \theta_j^i$.

Finally, since we are using a unitary frame field, the curvature forms satisfy the skew-hermitian properties $\bar{\Theta}^{\alpha}_{\beta} = -\Theta^{\beta}_{\alpha}$, and $\bar{\theta}^{i}_{j} = -\theta^{j}_{i}$.

We begin by differentiating the equation $e^a = 0$ and using the torsion free property of the metric connection. The result is:

$$0 = de^a = -\Theta^a_\alpha \wedge e^\alpha = -\Theta^a_i \wedge e^i$$

It follows that Θ^a_i is a $(1,0)$ form on M, so there are some locally defined functions A^a_{ij} so that:

$$\Theta^a_i = A^a_{ij} e^j \tag{8}$$

Lemma 8.2. *The second fundamental form of the immersion is locally expressed as:*

$$B(e_i, e_j) = A^a_{ij} e_a \tag{9}$$

Consequently, the immersion is totally geodesic if and only if $A^a_{ij} = 0$ for all a, i and j.

Proof. We simply calculate:

$$\langle B(e_i, e_j), e_a \rangle = \left\langle \tilde{\nabla}_{e_i} e_j, e_a \right\rangle = \langle \Theta^\alpha_i(e_j) e_\alpha, e_a \rangle = \Theta^a_i(e_j) = A^a_{ij}$$

Recalling that $B(e_i, \bar{e}_j) = 0$, and $B(\bar{e}_i, \bar{e}_j) = \overline{B(e_i, e_j)}$, we see that the lemma immediately follows. □

Since the second fundamental form is symmetric, an immediate consequence of the lemma above is that the symmetry condition $A^a_{ij} = A^a_{ji}$ holds for all i, j and a.

Now we consider the curvature forms ω and Ω for M and M_c respectively, which satisfy the usual fundamental equations. In particular:

$$\omega^i_i = d\theta^i_i + \theta^i_k \wedge \theta^k_i$$

and:

$$\Omega^i_i = d\Theta^i_i + \Theta^i_\alpha \wedge \Theta^\alpha_i = d\theta^i_i + \Theta^i_\alpha \wedge \Theta^\alpha_i$$

Hence:

$$\omega^i_i = \Omega^i_i - \Theta^i_a \wedge \Theta^a_i = \Omega^i_i - A^a_{ik} \bar{A}^a_{ij} e^k \wedge \bar{e}^j$$

Now, since the metric on M is Einstein, we have:

$$\omega^i_i = \frac{\lambda}{2} e^i \wedge \bar{e}^i$$

And since M_c has constant holomorphic curvature c, we have:

$$\Omega^i_i = \frac{(n+1)c}{4} e^i \wedge \bar{e}^i$$

Substituting these expressions in above yields:

$$\frac{\lambda}{2}e^i \wedge \bar{e}^i = \frac{(n+1)c}{4}e^i \wedge \bar{e}^i - A^a_{ik}\bar{A}^a_{ij}e^k \wedge \bar{e}^j$$

So:

$$A^a_{ik}\bar{A}^a_{ij}e^k \wedge \bar{e}^j = \left(\frac{(n+1)c}{4} - \frac{\lambda}{2}\right)e^i \wedge \bar{e}^i$$

Therefore, since the forms $e^k \wedge \bar{e}^j$ form a basis for the space of $(1,1)$ forms on M, we conclude that:

$$A^a_{ij}\bar{A}^a_{ik} = \delta^j_k\left(\frac{(n+1)c}{4} - \frac{\lambda}{2}\right) \tag{10}$$

Corollary 8.3. *The Einstein constant satisfies $\lambda \leq \frac{(n+1)c}{2}$, with equality if and only if M is totally geodesic. Furthermore, if $\lambda < \frac{(n+1)c}{2}$, then the linear map $TM^+ \to Hom(TM^+, NM^+)$ defined by:*

$$X \mapsto (Y \mapsto B(X,Y))$$

is an injection.

Proof. Using equation (10) with $j = k$ we get:

$$0 \leq A^a_{ij}\bar{A}^a_{ij} = n\left(\frac{(n+1)c}{4} - \frac{\lambda}{2}\right)$$

So it must be the case that $\lambda \leq \frac{(n+1)c}{2}$. Equality holds only when $A^a_{ij} = 0$ for all a, i, j, which by the previous lemma, is equivalent to M being totally geodesic.

Now, assume that $\lambda < \frac{(n+1)c}{2}$, and let $X = x_i e_i$ be a non-zero vector in TM^+. We need to see that the linear map $Y \mapsto B(X,Y)$ is non-zero, so we calculate:

$$\begin{aligned}\langle B(X,e_i), B(X,e_i)\rangle &= x_j\bar{x}_k \langle B(e_i,e_j), B(e_i,e_k)\rangle = x_j\bar{x}_k\delta^a_b A^a_{ij}\bar{A}^b_{ik} \\ &= x_j\bar{x}_k A^a_{ij}\bar{A}^a_{jk} = x_j\bar{x}_k\delta^j_k\left(\frac{(n+1)c}{4} - \frac{\lambda}{2}\right) \\ &= x_j\bar{x}_j\left(\frac{(n+1)c}{4} - \frac{\lambda}{2}\right)\end{aligned}$$

Which is positive when $\lambda < \frac{(n+1)c}{2}$. Therefore, there must be some j so that $B(X,e_j) \neq 0$, which completes the proof of the corollary. □

To continue, we differentiate equation (8):

$$d\Theta^a_i = dA^a_{ik} \wedge e^k + A^a_{ij}de^j$$

The fundamental equation gives:

$$d\Theta_i^a = \Omega_i^a - \Theta_\alpha^a \wedge \Theta_i^\alpha$$

And using the curvature formula for space forms, we get:

$$\Omega_i^a = \frac{c}{4} e^a \wedge \bar{e}^i = 0$$

So:

$$d\Theta_i^a = -\Theta_\alpha^a \wedge \Theta_i^\alpha = -\Theta_j^a \wedge \Theta_i^j - \Theta_b^a \wedge \Theta_i^b = A_{jk}^a \theta_i^j \wedge e^k - A_{ik}^b \Theta_b^a \wedge e^k$$

On the other hand, the torsion free metric property gives:

$$A_{ij}^a de^j = -A_{ij}^a \theta_k^j \wedge e^k$$

Substituting all these ingredients in now yields:

$$0 = \left(A_{jk}^a \theta_i^j - A_{ik}^b \Theta_b^a - dA_{ik}^a + A_{ij}^a \theta_k^j \right) \wedge e^k$$

Therefore, the form appearing as a coefficient of e^k must be a $(1,0)$ form, and so there are some locally defined functions B_{ijl}^a that satisfy:

$$B_{ijl}^a e^l = dA_{ij}^a + A_{ij}^b \Theta_b^a - A_{kj}^a \theta_i^k - A_{ik}^a \theta_j^k \tag{11}$$

Lemma 8.4. *The first covariant derivative of the second fundamental form has the local expression:*

$$\nabla B(e_i, e_j, e_l) = B_{ijl}^a e_a$$

Consequently, the second fundamental form is parallel if and only if $B_{ijl}^a = 0$ for all a, i, j and l.

Proof. We use the definition of the covariant derivative to calculate:

$$\begin{aligned}
\nabla B(e_i, e_j, e_l) &= (\nabla_{e_l} B)(e_i, e_j) \\
&= \nabla_{e_l}^\perp (B(e_i, e_j)) - B(\nabla_{e_l} e_i, e_j) - B(e_i, \nabla_{e_j} e_j) \\
&= \nabla_{e_l}^\perp \left(A_{ij}^a e_a \right) - B(\theta_i^k(e_l) e_k, e_j) - B(e_i, \theta_j^k(e_j) e_k) \\
&= \left(dA_{ij}^a + A_{ij}^b \Theta_b^a - A_{kj}^a \theta_i^k - A_{ik}^a \theta_j^k \right) (e_l) e_a \\
&= B_{ijl}^a e_a
\end{aligned}$$

which completes the proof. □

Corollary 8.5. *The numbers B_{ijl}^i are invariant with regards to all permutations of the lower three indicies. That is, $B_{ijl}^a = B_{jil}^a = B_{ilj}^a$.*

Proof. The relation $B_{ijl}^a = B_{jil}^a$ is an easy consequence of the symmetry of A_{ij}^a and equation (11).

Verifying the other symmetry relation is considerably more subtle. By the previous lemma, it suffices to check the tensor equation:

$$\nabla B(e_i, e_j, e_l) - \nabla B(e_i, e_l, e_j) = 0$$

Fix a general point p in M, and for the moment assume (for convenience) that the frame $\{e_1, e_2, \ldots, e_{n+m}\}$ is parallel at p. The following calculations are always evaluated at the point p. We have:

$$\begin{aligned}\nabla B(e_i, e_j, e_l) - \nabla B(e_i, e_l, e_j) &= \nabla^{\perp}_{e_l}(B(e_i, e_j)) - \nabla^{\perp}_{e_j}(B(e_i, e_l)) \\ &= dA^a_{ij}(e_l)e_a - dA^a_{il}(e_j)e_a\end{aligned}$$

Now since:

$$A^a_{ij} = \Theta^a_i(e_j) = e^a(\nabla_{e_j} e_i) = \left\langle \nabla_{e_j} e_i, e_a \right\rangle$$

compatibility of the connection with the metric yields:

$$dA^a_{ij}(e_l) = e_l \left\langle \nabla_{e_j} e_i, e_a \right\rangle = \left\langle \nabla_{e_l} \nabla_{e_j} e_i, e_a \right\rangle$$

where we have used that $\nabla_{e_l} e_a = 0$ at p. Therefore:

$$\begin{aligned}dA^a_{ij}(e_l) - dA^a_{il}(e_j) &= \left\langle \nabla_{e_l} \nabla_{e_j} e_i, e_a \right\rangle - \left\langle \nabla_{e_j} \nabla_{e_l} e_i, e_a \right\rangle \\ &= \langle R(e_l, e_j)e_i, e_a \rangle \\ &= 0\end{aligned}$$

since e_l and e_j lie in TM^+. □

We now want to find an orthogonality relation between the A^a_{ij} and the B^a_{ijl}. To begin, we differentiate equation (10):

$$0 = \bar{A}^a_{ik} dA^a_{ij} + A^a_{ij} d\bar{A}^a_{ik}$$

To simplify these two terms we use equation (11), the results are:

$$\bar{A}^a_{ik} dA^a_{ij} = \bar{A}^a_{ik} B^a_{ijl} e^l + \bar{A}^a_{ik} A^a_{il} \theta^l_j + \bar{A}^a_{ik} A^a_{lj} \theta^l_i - \bar{A}^a_{ik} A^b_{ij} \Theta^a_b$$

and:

$$A^a_{ij} d\bar{A}^a_{ik} = A^a_{ij} \bar{B}^a_{ikl} \bar{e}^l - A^a_{ij} \bar{A}^a_{lk} \theta^i_l - A^a_{ij} \bar{A}^a_{il} \theta^k_l + A^a_{ij} \bar{A}^b_{ik} \Theta^b_a$$

Substituting in these two formula, we see, upon re-indexing, that two pairs of terms cancel, and we are left with:

$$0 = \bar{A}^a_{ik} B^a_{ijl} e^l + A^a_{ij} \bar{B}^a_{ikl} \bar{e}^l + \bar{A}^a_{ik} A^a_{il} \theta^l_j - A^a_{ij} \bar{A}^a_{il} \theta^k_l$$

To deal with the last two terms here, we call upon formula (10), which yields:

$$\begin{aligned}\bar{A}^a_{ik} A^a_{il} \theta^l_j - A^a_{ij} \bar{A}^a_{il} \theta^k_l &= \delta^k_l \left(\frac{(n+1)c}{4} - \frac{\lambda}{2} \right) \theta^l_j - \delta^j_l \left(\frac{(n+1)c}{4} - \frac{\lambda}{2} \right) \theta^k_l \\ &= \left(\frac{(n+1)c}{4} - \frac{\lambda}{2} \right) \left(\theta^k_j - \theta^k_j \right) \\ &= 0\end{aligned}$$

Therefore:

$$0 = \bar{A}^a_{ik} B^a_{ijl} e^l + A^a_{ij} \bar{B}^a_{ikl} \bar{e}^l$$

and we conclude the orthogonality relation:

$$\bar{A}^a_{ik} B^a_{ijl} = 0 \tag{12}$$

For all k, j and l.

For the next corollary, we denote by $TM^+ \odot TM^+$ the space of symmetric 2-tensors on TM^+.

Corollary 8.6. *The image of the linear map $TM^+ \odot TM^+ \to Hom(TM^+, NM^+)$ defined by:*

$$X \odot Y \mapsto (Z \mapsto (\nabla B)(X, Y, Z))$$

is orthogonal to the image of the map $TM^+ \to Hom(TM^+, NM^+)$.

Proof. Let $X = x_j e_j$ and $Y = y_{rs} e_r \odot e_s$. Then we have:

$$\begin{aligned} \langle B(X, e_i), (\nabla_{e_i} B)(Y) \rangle &= x_j \bar{y}_{rs} \langle B(e_i, e_j), (\nabla_{e_i} B)(e_r, e_s) \rangle \\ &= x_j \bar{y}_{rs} \delta^a_b A^a_{ij} \bar{B}^b_{irs} = x_j \bar{y}_{rs} A^a_{ij} \bar{B}^a_{irs} \\ &= 0 \end{aligned}$$

Which completes the proof. □

We now want to get an expression for $B^a_{ijl} \bar{B}^a_{ijr}$. To do so, we begin by differentiating the equation $\bar{A}^a_{ik} B^a_{ijl} e^l = 0$. The result is:

$$0 = B^a_{ijl} d\bar{A}^a_{ik} \wedge e^l + \bar{A}^a_{ik} d(B^a_{ijl} e^l)$$

Using equation (11) to substitute into the first term, and making use of the orthogonality relation (12), we get:

$$B^a_{ijl} d\bar{A}^a_{ik} \wedge e^l = B^a_{ijl} \bar{B}^a_{ikr} \bar{e}^r \wedge e^l + \bar{A}^b_{ik} B^a_{ikl} \Theta^b_a \wedge e^l - \bar{A}^a_{rk} B^a_{ijl} \theta^i_r \wedge e^l$$

For the second term, we differentiate equation (11) to get:

$$\begin{aligned} \bar{A}^a_{ik} d(B^a_{ijl} e^l) = &\bar{A}^a_{ik} dA^b_{ij} \wedge \Theta^a_b + \bar{A}^a_{ik} A^b_{ij} d\Theta^a_b - \bar{A}^a_{ik} dA^a_{rj} \wedge \theta^r_i \\ &- \bar{A}^a_{ik} A^a_{rj} d\theta^r_i - \bar{A}^a_{ik} dA^a_{ir} \wedge \theta^r_j - \bar{A}^a_{ik} A^a_{ir} d\theta^r_j \end{aligned}$$

We simplify these terms one by one, using always the orthogonality relations (11), (12), and the following remark:

Remark 8.7. We take a look at the $d\Theta$ and $d\theta$ terms. For instance, to calculate $d\Theta^a_b$, we apply the fundamental equation, and the formula for the

curvature in a space form to get:

$$\begin{aligned} d\Theta^a_b &= \Omega^a_b - \Theta^a_\alpha \wedge \Theta^\alpha_b \\ &= \frac{c}{4}(\delta^a_b e^l \wedge \bar{e}^l + e^a \wedge \bar{e}^b) - \Theta^a_l \wedge \Theta^l_b - \Theta^a_c \wedge \Theta^c_b \\ &= \frac{c}{4}\delta^a_b e^l \wedge \bar{e}^l + A^a_{lr}\bar{A}^b_{ls} e^r \wedge \bar{e}^s - \Theta^a_c \wedge \Theta^c_b \end{aligned}$$

Similar analysis yields:

$$d\theta^r_i = \frac{c}{4}\delta^r_i e^l \wedge \bar{e}^l + \frac{c}{4} e^r \wedge \bar{e}^i - \theta^r_l \wedge \theta^l_i + \bar{A}^b_{rl} A^b_{is} \bar{e}^l \wedge e^s$$

and:

$$d\theta^r_j = \frac{c}{4}\delta^r_j e^l \wedge \bar{e}^l + \frac{c}{4} e^r \wedge \bar{e}^j - \theta^r_l \wedge \theta^l_j + \bar{A}^b_{rl} A^b_{js} \bar{e}^l \wedge e^s$$

Now, carefully simplifying each of the six terms yields:

$$\begin{aligned} \bar{A}^a_{ik} dA^b_{ij} \wedge \Theta^a_b =& B^b_{ijl}\bar{A}^a_{ik} e^l \wedge \Theta^a_b - \bar{A}^a_{ik} A^c_{ij} \Theta^b_c \wedge \Theta^a_b \\ &+ \bar{A}^a_{ik} A^b_{rj} \theta^r_i \wedge \Theta^a_b + \bar{A}^a_{ik} A^b_{ir} \theta^r_j \wedge \Theta^a_b \end{aligned}$$

$$\bar{A}^a_{ik} A^b_{ij} d\Theta^a_b = \frac{c}{4}\bar{A}^a_{ik} A^a_{ij} e^l \wedge \bar{e}^l + \bar{A}^a_{ik}\bar{A}^b_{ls} A^a_{lr} A^b_{ij} e^r \wedge \bar{e}^s - \bar{A}^a_{ik} A^b_{ij} \Theta^a_c \wedge \Theta^c_b$$

$$\begin{aligned} -\bar{A}^a_{ik} dA^a_{rj} \wedge \theta^r_i =& -\bar{A}^a_{ik} B^a_{rjl} e^l \wedge \theta^r_i + \bar{A}^a_{ik} A^b_{rj} \Theta^a_b \wedge \theta^r_i \\ &- \bar{A}^a_{ik} A^a_{lj} \theta^l_r \wedge \theta^r_i - \bar{A}^a_{ik} A^a_{rl} \theta^l_j \wedge \theta^r_i \end{aligned}$$

$$\begin{aligned} -\bar{A}^a_{ik} A^a_{rj} d\theta^r_i =& -\frac{c}{4}\bar{A}^a_{ik} A^a_{ij} e^l \wedge \bar{e}^l - \frac{c}{4}\bar{A}^a_{ik} A^a_{rj} e^r \wedge \bar{e}^i \\ &+ \bar{A}^a_{ik} A^a_{rj} \theta^r_l \wedge \theta^l_i - \bar{A}^a_{ik}\bar{A}^b_{rl} A^a_{rj} A^b_{is} \bar{e}^l \wedge e^s \end{aligned}$$

$$-\bar{A}^a_{ik} dA^a_{ir} \wedge \theta^r_j = \bar{A}^a_{ik} A^b_{ir} \Theta^a_b \wedge \theta^r_j - \bar{A}^a_{ik} A^a_{lr} \theta^l_i \wedge \theta^r_j - \bar{A}^a_{ik} A^a_{il} \theta^l_r \wedge \theta^r_j$$

$$\begin{aligned} -\bar{A}^a_{ik} A^a_{ir} d\theta^r_j =& -\frac{c}{4}\bar{A}^a_{ik} A^a_{ij} e^l \wedge \bar{e}^l - \frac{c}{4}\bar{A}^a_{ik} A^a_{ir} e^r \wedge \bar{e}^j \\ &+ \bar{A}^a_{ik} A^a_{ir} \theta^r_l \wedge \theta^l_j - \bar{A}^a_{ik}\bar{A}^b_{rl} A^a_{ir} A^b_{js} \bar{e}^l \wedge e^s \end{aligned}$$

Substituting in all these results, we see that all the terms containing either a θ or a Θ cancel, and we are left with:

$$\begin{aligned} 0 =& B^a_{ijl}\bar{B}^a_{ikr}\bar{e}^e \wedge e^l - \frac{c}{4}\bar{A}^a_{ik} A^a_{ij} e^l \wedge \bar{e}^l - \frac{c}{4}\bar{A}^a_{ik} A^a_{rj} e^r \wedge \bar{e}^i - \frac{c}{4}\bar{A}^a_{ik} A^a_{ir} e^r \wedge \bar{e}^j \\ &+ \bar{A}^a_{ik}\bar{A}^b_{ls} A^a_{lr} A^b_{ij} e^r \wedge \bar{e}^s + \bar{A}^a_{ik}\bar{A}^b_{rl} A^a_{ij} A^b_{is} e^s \wedge \bar{e}^l + \bar{A}^a_{ik}\bar{A}^b_{rl} A^a_{ir} A^b_{js} e^s \wedge \bar{e}^l \end{aligned}$$

We can simplify three of these terms further by applying formula (10):

$$\begin{aligned}-\frac{c}{4}\bar{A}^a_{ik}A^a_{ij}e^l\wedge\bar{e}^l &= -\frac{c}{4}\delta^j_k\left(\frac{(n+1)c}{4}-\frac{\lambda}{2}\right)e^l\wedge\bar{e}^l\\ &= -\frac{c}{4}\delta^j_k\delta^r_s\left(\frac{(n+1)c}{4}-\frac{\lambda}{2}\right)e^s\wedge\bar{e}^r\end{aligned}$$

$$\begin{aligned}-\frac{c}{4}\bar{A}^a_{ik}A^a_{ir}e^r\wedge\bar{e}^j &= -\frac{c}{4}\delta^k_r\left(\frac{(n+1)c}{4}-\frac{\lambda}{2}\right)e^r\wedge\bar{e}^j\\ &= -\frac{c}{4}\delta^k_r\delta^j_s\left(\frac{(n+1)c}{4}-\frac{\lambda}{2}\right)e^r\wedge\bar{e}^s\\ &= -\frac{c}{4}\delta^k_s\delta^j_r\left(\frac{(n+1)c}{4}-\frac{\lambda}{2}\right)e^s\wedge\bar{e}^r\end{aligned}$$

$$\begin{aligned}\bar{A}^a_{ik}\bar{A}^b_{rl}A^a_{ir}A^b_{js}e^s\wedge\bar{e}^l &= \delta^k_r\left(\frac{(n+1)c}{4}-\frac{\lambda}{2}\right)\bar{A}^b_{rl}\bar{A}^b_{js}e^s\wedge\bar{e}^l\\ &= \left(\frac{(n+1)c}{4}-\frac{\lambda}{2}\right)\bar{A}^a_{rk}\bar{A}^a_{js}e^s\wedge\bar{e}^r\end{aligned}$$

We get, upon gathering like terms and changing around some of the summation indices:

$$\begin{aligned}B^a_{ijs}\bar{B}^a_{ikr}e^s\wedge\bar{e}^r =& -\frac{c}{4}\left(\frac{(n+1)c}{4}-\frac{\lambda}{2}\right)\left(\delta^j_k\delta^r_s+\delta^k_s\delta^j_r\right)e^s\wedge\bar{e}^r\\ &+\left(\frac{nc}{4}-\frac{\lambda}{2}\right)\bar{A}^a_{rk}A^a_{sj}e^s\wedge\bar{e}^r+\bar{A}^a_{ik}\bar{A}^b_{lr}A^a_{ls}A^b_{ij}e^s\wedge\bar{e}^r\\ &+\bar{A}^a_{ik}\bar{A}^b_{lr}A^a_{lj}A^b_{is}e^s\wedge\bar{e}^r\end{aligned}$$

Since the forms $e^s\wedge\bar{e}^r$ form a basis for the $(1,1)$ forms on M, we can isolate the coefficients in the equation above to finally get:

$$\begin{aligned}B^a_{ijs}\bar{B}^a_{ikr} =& -\frac{c}{4}\left(\frac{(n+1)c}{4}-\frac{\lambda}{2}\right)\left(\delta^j_k\delta^r_s+\delta^k_s\delta^j_r\right)+\left(\frac{nc}{4}-\frac{\lambda}{2}\right)\bar{A}^a_{rk}A^a_{sj}\\ &+\bar{A}^a_{ik}\bar{A}^b_{lr}A^a_{ls}A^b_{ij}+\bar{A}^a_{ik}\bar{A}^b_{lr}A^a_{lj}A^b_{is}\end{aligned}\tag{13}$$

Corollary 8.8. *If $\lambda<\frac{(n+1)c}{2}$ then, in fact, $\lambda\le\frac{nc}{2}$, with equality holding if and only if the second fundamental form B is parallel.*

Proof. For convenience, write $K=\left(\frac{(n+1)c}{4}-\frac{\lambda}{2}\right)$. Set $k=j$ and $r=s$ in formula (13) to get:

$$\begin{aligned}B^a_{ijs}\bar{B}^a_{ijs} =& -\frac{c}{4}K(\delta^j_j\delta^r_r+\delta^r_j\delta^r_j)+\left(K-\frac{c}{4}\right)\bar{A}^a_{sj}A^a_{sj}\\ &+\bar{A}^a_{ij}\bar{A}^b_{sr}A^a_{sr}A^b_{ij}+\bar{A}^a_{ij}\bar{A}^b_{sr}A^a_{sj}A^b_{ir}\end{aligned}$$

It is easy to see that:

$$-\frac{c}{4}K(\delta^j_j\delta^r_r + \delta^r_j\delta^r_j) = -\frac{c}{4}Kn(n+1)$$

The second and fourth terms can be computed using formula (10):

$$\left(K - \frac{c}{4}\right)\bar{A}^a_{sj}A^a_{sj} = nK\left(K - \frac{c}{4}\right)$$

$$\bar{A}^a_{ij}\bar{A}^b_{sr}A^a_{sj}A^b_{ir} = \delta^i_s\delta^i_s K^2 = nK^2$$

The third term we can merely estimate, using the Cauchy-Schwarz inequality. For sake of clarity, in the estimate we temporarily drop our summation convention. We have:

$$\begin{aligned}
\sum_{abijsr} \bar{A}^a_{ij}\bar{A}^b_{sr}A^a_{sr}A^b_{ij} &= \sum_{ijsr}\left(\sum_a \bar{A}^a_{ij}A^a_{sr}\right)\left(\sum_b \bar{A}^b_{sr}A^b_{ij}\right) = \sum_{ijsr}\left|\sum_a \bar{A}^a_{ij}A^a_{sr}\right|^2 \\
&\leq \sum_{ijrs}\left|\sum_a \bar{A}^a_{ij}\right|^2\left|\sum_b A^b_{sr}\right|^2 \\
&= \sum_{ijrs}\left(\sum_a \bar{A}^a_{ij}A^a_{ij}\right)\left(\sum_b A^b_{sr}\bar{A}^b_{sr}\right) \\
&= n^2K^2
\end{aligned}$$

Where we have used (10) in the last equality. Putting this all together, and doing a little calculation, we get:

$$\begin{aligned}
B^a_{ijs}\bar{B}^a_{ijs} &\leq -\frac{c}{4}Kn(n+1) + nK\left(K - \frac{c}{4}\right) + nK^2 + n^2K^2 \\
&= n(n+2)\left(\frac{(n+1)c}{4} - \frac{\lambda}{2}\right)\left(\frac{nc}{4} - \frac{\lambda}{2}\right)
\end{aligned}$$

Now $B^a_{ijs}\bar{B}^a_{ijs}$ and $\left(\frac{(n+1)c}{4} - \frac{\lambda}{2}\right)$ are both non-negative, so it follows that $\left(\frac{nc}{4} - \frac{\lambda}{2}\right)$ must be as well. That is, $\lambda \leq \frac{nc}{2}$. Equality occurs if and only if $B^a_{ijs} = 0$ for every a, i, j, s, which by lemma 6, means that the second fundamental form B is parallel. □

With the formula for $B^a_{ijk}\bar{B}^a_{irs}$ in hand we want to study the map $TM^+ \odot TM^+ \to \mathrm{Hom}(TM^+, NM^+)$ in more detail. So let $X = x_{jk}e_j \odot e_k$ be a non-zero vector in $TM^+ \odot TM^+$. Then:

$$\begin{aligned}
\langle \nabla B(X, e_i), \nabla B(X, e_i)\rangle &= x_{jk}\bar{x}_{rs}\langle \nabla B(e_j, e_k, e_i), \nabla B(e_r, e_s, e_i)\rangle \\
&= x_{jk}\bar{x}_{rs}B^a_{ijk}\bar{B}^a_{irs}
\end{aligned}$$

Using (13) we get that this is equal to:

$$-\frac{c}{4}\left(\frac{(n+1)c}{4}-\frac{\lambda}{2}\right)\left(\delta_r^j\delta_k^s+\delta_r^k\delta_j^s\right)x_{jk}\bar{x}_{rs}+\left(\frac{nc}{4}-\frac{\lambda}{2}\right)A_{rs}^b\bar{A}_{jk}^b x_{jk}\bar{x}_{rs}$$
$$+\bar{A}_{mr}^b\bar{A}_{ls}^c A_{lk}^b A_{mj}^c x_{jk}\bar{x}_{rs}+\bar{A}_{mr}^b\bar{A}_{ls}^c A_{lj}^b A_{mk}^c x_{jk}\bar{x}_{rs}$$

Looking at the first two of these terms, and recalling that $x_{ij}=x_{ji}$, we see that:

$$-\frac{c}{4}\left(\frac{(n+1)c}{4}-\frac{\lambda}{2}\right)\left(\delta_r^j\delta_k^s+\delta_r^k\delta_j^s\right)x_{jk}\bar{x}_{rs}=-\frac{c}{2}\left(\frac{(n+1)c}{4}-\frac{\lambda}{2}\right)x_{jk}\bar{x}_{jk}$$
$$=-\frac{c}{2}\left(\frac{(n+1)c}{4}-\frac{\lambda}{2}\right)\sum_{jk}|x_{jk}|^2$$

For the second term we observe that:

$$\left(\frac{nc}{4}-\frac{\lambda}{2}\right)A_{rs}^b\bar{A}_{jk}^b x_{jk}\bar{x}_{rs}=\left(\frac{nc}{4}-\frac{\lambda}{2}\right)\sum_b\left|A_{jk}^b x_{jk}\right|^2$$

For the final two terms, we calculate:

$$\begin{aligned}
&\bar{A}_{mr}^b\bar{A}_{ls}^c A_{lk}^b A_{mj}^c x_{jk}\bar{x}_{rs}+\bar{A}_{mr}^b\bar{A}_{ls}^c A_{lj}^b A_{mk}^c x_{jk}\bar{x}_{rs}\\
&\quad=\bar{A}_{mr}^b\bar{A}_{ls}^c A_{lk}^b A_{mj}^c x_{jk}\bar{x}_{rs}+\bar{A}_{mr}^c\bar{A}_{ls}^b A_{lj}^c A_{mk}^b x_{jk}\bar{x}_{rs}\\
&\quad=\sum_{lmbc}\left(\sum_{rs}(\bar{A}_{mr}^b\bar{A}_{ls}^c-\bar{A}_{mr}^c\bar{A}_{ls}^b)\bar{x}_{rs}\right)\left(\sum_{jk}(A_{lk}^b A_{mj}^c-A_{lj}^c A_{mk}^b)x_{jk}\right)\\
&\qquad+\bar{A}_{mr}^b\bar{A}_{ls}^c A_{lj}^c A_{mk}^b x_{jk}\bar{x}_{rs}+\bar{A}_{mr}^c\bar{A}_{ls}^b A_{lk}^b A_{mj}^c x_{jk}\bar{x}_{rs}\\
&\quad=-\sum_{lmbc}\left|\sum_{jk}\left(A_{lk}^b A_{mj}^c-A_{lj}^c A_{mk}^b\right)x_{jk}\right|^2+2\delta_k^r\delta_s^j\left(\frac{(n+1)c}{4}-\frac{\lambda}{2}\right)^2 x_{jk}\bar{x}_{rs}\\
&\quad=-\sum_{lmbc}\left|\sum_{jk}\left(A_{lk}^b A_{mj}^c-A_{lj}^c A_{mk}^b\right)x_{jk}\right|^2+2\left(\frac{(n+1)c}{4}-\frac{\lambda}{2}\right)^2\sum_{jk}|x_{jk}|^2
\end{aligned}$$

Therefore, we have:

$$\begin{aligned}\langle \nabla B(X,e_i), \nabla B(X,e_i)\rangle &= -\sum_{lmbc}\left|\sum_{jk}\left(A^b_{lk}A^c_{mj} - A^c_{lj}A^b_{mk}\right)x_{jk}\right|^2 \\ &\quad + \left(\frac{nc}{4} - \frac{\lambda}{2}\right)\sum_b \left|A^b_{jk}x_{jk}\right|^2 \\ &\quad + \left(\frac{(n+1)c}{4} - \frac{\lambda}{2}\right)\left(2\left(\frac{(n+1)c}{4} - \frac{\lambda}{2}\right) - \frac{c}{2}\right)\sum_{jk}|x_{jk}|^2 \\ &= -\sum_{lmbc}\left|\sum_{jk}\left(A^b_{lk}A^c_{mj} - A^c_{lj}A^b_{mk}\right)x_{jk}\right|^2 \\ &\quad + \left(\frac{nc}{4} - \frac{\lambda}{2}\right)\sum_b \left|A^b_{jk}x_{jk}\right|^2 \\ &\quad + 2\left(\frac{(n+1)c}{4} - \frac{\lambda}{2}\right)\left(\frac{nc}{4} - \frac{\lambda}{2}\right)\sum_{jk}|x_{jk}|^2\end{aligned}$$

We now have all the pieces in hand that we need to prove Chern's theorem.

Theorem 8.9. *Suppose that $m = 1$, then the Einstein constant λ is either $\frac{(n+1)c}{2}$ or $\frac{nc}{2}$. In the first case, the immersion is totally geodesic, and in the second, the second fundamental form B is parallel.*

Proof. When $m = 1$ the first term in the previous formula is easily seen to vanish, so:

$$\begin{aligned}\langle \nabla B(X,e_i), \nabla B(X,e_i)\rangle &= \left(\frac{nc}{4} - \frac{\lambda}{2}\right)\sum_b \left|A^b_{jk}x_{jk}\right|^2 \\ &\quad + \left(\frac{(n+1)c}{4} - \frac{\lambda}{2}\right)\left(2\left(\frac{(n+1)c}{4} - \frac{\lambda}{2}\right) - \frac{c}{2}\right)\sum_{jk}|x_{jk}|^2\end{aligned}$$

If $\lambda \neq \frac{(n+1)c}{2}, \frac{nc}{2}$, then, by a previous corollary, it must be the case that $\lambda < \frac{nc}{2}$, hence the right hand side is positive. Therefore:

$$\langle \nabla B(X,e_i), \nabla B(X,e_i)\rangle > 0$$

and hence the map $TM^+ \odot TM^+ \to \mathrm{Hom}(TM^+, NM^+)$ is injective. We know already that the map $TM^+ \to \mathrm{Hom}(TM^+, NM^+)$ is injective, and that the images of these two maps are orthogonal. Since the dimension of TM^+ is n, the dimension of $TM^+ \odot TM^+$ is $\frac{n(n-1)}{2}$. The dimension of

$\mathrm{Hom}(TM^+, NM^+)$ is $n \times 1 = n$, so we must conclude that:

$$n + \frac{n(n-1)}{2} < n$$

which is a contradiction unless $n = 1$, a case we have already studied in the previous chapter. □

References

[C1] E. Calabi, *Isometric imbedding of complex manifolds*, Ann. of Math. 58 (1953), 1-23.

[C2] E. Calabi, *Metric Riemann surfaces*, Contributions to the Theory of Riemann Surfaces, Ann. of Math. Studies, Princeton, 1953.

[CE] J. Cheeger, D. Ebin, *Comparison Theorems in Riemannian Geometry*, AMS Chelsea Publishing, 2008.

[Ch] S. S. Chern, *Einstein hypersurfaces in a Kählerian manifold of constant holomorphic curvature*, J. Differential Geometry. 1 (1967), 21-31.

[dC] M. P. doCarmo *Riemannian Geometry*, Birkhauser. 1992.

[S] B. Smyth, *Differential geometry of complex hypersurfaces*, Ann. of Math. 85 (1967), 246-266.

[T] K. Tsukada, *Einstein Kähler submanifolds with codimension 2 in a complex space form*, Math Ann. 274 (1986), 503-516.

[Um1] M. Umehara, *Einstein Kähler submanifolds of a complex linear or hyperbolic space*, Tôhoku Math. Journ. 39 (1987), 385-389.

[Um2] M. Umehara, *Diastases and real analytic functions on complex manifolds*, J. Math. Soc. Japan Vol. 40, No. 3 (1988), 519-539.