

# Mathematical Induction

Madiba Hudson-Quansah

March 2024

# CONTENTS

<b>CHAPTER 1</b>	<b>MATHEMATICAL INDUCTION</b>	<b>PAGE 2</b>
1.1	Introduction	2
1.2	Why Mathematical Induction is Valid	3
1.3	Examples of Proofs by Mathematical Induction	4
	Proving Summation Formulae — 4	
	1.3.1.0.1 Sums of Geometric Progressions . . . . .	6
1.4	Exercises	7
<b>CHAPTER 2</b>	<b>EXERCISES</b>	<b>PAGE 12</b>

# Chapter 1

## Mathematical Induction

### 1.1 Introduction

#### Definition 1.1.1: Mathematical Induction

Mathematical induction can be used to prove statements that assert  $P(n)$  is true for all positive integers  $n$ , where  $P(n)$  is a propositional function. A proof by mathematical induction has two parts, a **basis step**, where we show that  $P(1)$  is true and an **inductive step**, where we show that for all positive integers  $k$  if  $P(k)$  is true then  $P(k+1)$  is also true.

Mathematical Induction can be expressed as the rule of inference  $P(1) \wedge (P(k) \rightarrow P(k+1)) \rightarrow \forall n P(n)$ , where the domain is the set of positive integers.

#### Example 1.1.1

##### Question 1

Show that

$$P(n) : 1 + 2 + 3 + \dots + n = \frac{n(n+1)}{2}$$

**Solution:** We first start with the basis step,  $P(1)$ , which is

$$\begin{aligned} P(1) : 1 &= \frac{1(1+1)}{2} \\ 1 &= 1 \end{aligned}$$

$\therefore P(1)$  is true.

Next is the inductive step, where we need to show that  $P(k) \rightarrow P(k+1)$

We assume

$$P(k) : 1 + 2 + 3 + \dots + k = \frac{k(k+1)}{2}$$

and prove

$$P(k+1) : 1 + 2 + 3 + \dots + (k+1) = \frac{(k+1)(k+2)}{2}$$

To prove this we need to relate these two predicates. And because  $P(k+1)$  can be expressed as  $P(k) + (k+1)$  we can write

$$P(k+1) : 1 + 2 + 3 + \dots + (k+1) = \frac{k(k+1)}{2} + (k+1)$$

So we simplify

$$\begin{aligned}\frac{k(k+1)}{2} + (k+1) &= \frac{k(k+1) + 2(k+1)}{2} \\ &= \frac{(k+1)(k+2)}{2}\end{aligned}$$

### Example 1.1.2

#### Question 2

Show that

$$P(n) : 1 + 3 + 5 + \dots + (2n - 1) = n^2 \quad \forall n \in \mathbb{Z}$$

**Solution:** The basis step is  $P(1)$

$$P(1) : 1 = 1^2$$

Assume

$$P(k) : 1 + 3 + 5 + \dots + (2k - 1) = k^2$$

and show that

$$P(k+1) : 1 + 3 + 5 + \dots + (2(k+1) - 1) = (k+1)^2$$

$P(k+1)$  can be expressed as  $P(k) + (2k+1)$

$$\begin{aligned}P(k+1) : 1 + 3 + 5 + \dots + (2k - 1) + (2k + 1) \\ &= k^2 + 2k + 1 \\ &= (k+1)^2\end{aligned}$$

$\therefore P(k+1)$  is true

Conclusion: Having completed the basis and induction steps we can conclude that  $1 + 3 + 5 + \dots + (2n - 1) = n^2$

## 1.2 Why Mathematical Induction is Valid

### Definition 1.2.1: Well Ordering Property of Positive Integers

This axiom states that every non-empty subset of positive integers has a least element

The validity of mathematical induction as a proof technique comes from the well ordering property, as an axiom of positive integers. So suppose we know that  $P(1)$  is true for all positive integers  $n$  and that the proposition  $P(k) \rightarrow P(k+1)$  is true for all positive integers  $k$ . To show that  $P(n)$  must be true for all positive integers  $n$  we can use a proof by contradiction.

We first assume that there is at least positive integer for which  $P(n)$  is false, i.e.  $\exists x \in \mathbb{Z}^+ \neg P(x)$

Then the set of positive integers for which  $P(n)$  is false, let this be  $S$ , is non-empty. Thus by the well ordering property,  $S$  has a least element, which we will denote as  $m$ .

We know that  $m$  cannot be 1, because  $P(1)$  is true, and since  $m$  is a positive integer it must be greater than one ergo  $m - 1$  is also a positive integer.

Again due to well ordering property we know the least value in  $S$  is  $m$  therefore  $m - 1$  is not in set  $S$ , as a result  $P(m - 1)$  is true.

Because the statements  $P(m - 1) \rightarrow P(m)$  is also true, it must be the case that  $P(m)$  is true, which is a contradiction of the statement that  $P(m)$  must be false.

Hence  $P(n)$  must be true for every positive integer.

## 1.3 Examples of Proofs by Mathematical Induction

### 1.3.1 Proving Summation Formulae

#### Question 3

Conjecture a formula for the sum of the first  $n$  positive odd integers. Then prove your conjecture using mathematical induction.

**Solution:** The sums of the first  $n$  positive odd integers for  $n = 1, 2, 3, 4, 5$  are

$$\begin{aligned} 1 &= 1 \\ 1 + 3 &= 4 \\ 1 + 3 + 5 &= 9 \\ 1 + 3 + 5 + 7 &= 16 \\ 1 + 3 + 5 + 7 + 9 &= 25 \end{aligned}$$

From these values it is reasonable to conjecture that the sum of the first  $n$  odd integers is  $n^2$ , i.e.

$$1 + 3 + 5 + \dots + (2n - 1) = n^2$$

Let  $P(n)$  be  $1 + 3 + 5 + \dots + (2n - 1) = n^2$

Basis Step:

$$P(1) : 1 = 1^2$$

Inductive Step:

To complete this step we need to prove that  $\forall k \in \mathbb{Z}^+ (P(k) \rightarrow P(k + 1))$

Assume  $\exists k \in \mathbb{Z}^+ P(k) = T$ , then

$$P(k) : 1 + 3 + 5 + \dots + (2k - 1) = k^2$$

$$P(k+1) : 1 + 3 + 5 + \dots + (2k+1) = (k+1)^2$$

$$P(k+1) : 1 + 3 + 5 + \dots + (2k-1) + (2k+1) = (k+1)^2$$

$$P(k+1) : P(k) + (2k+1)$$

$$1 + 3 + 5 + \dots + (2k+1) = k^2 + 2k + 1$$

$$= (k+1)(k+1)$$

$$= (k+1)^2$$

$\therefore P(k+1)$  is  $T$

Hence we can conclude that  $P(n)$  is true for all positive integers  $n$ .

#### Question 4

Use mathematical induction to show that

$$1 + 2 + 2^2 + \dots + 2^n = 2^{n+1} - 1$$

For all non-negative integers  $n$

**Solution:**

Let  $P(n)$  be  $1 + 2 + 2^2 + \dots + 2^n = 2^{n+1} - 1$

Basis Step:

$$P(0) : 2^0 = 2^{0+1} - 1$$

$$1 = 2 - 1$$

$$1 = 1$$

Induction Step

To complete this step I must prove  $P(k) \rightarrow P(k+1)$  where  $k$  is any non-negative integer.

Assume  $P(k)$  is true for some non-negative integer  $k$ , then

$$P(k) : 1 + 2 + 2^2 + \dots + 2^k = 2^{k+1} - 1$$

Then  $P(k+1)$  is

$$P(k+1) : 1 + 2 + 2^2 + \dots + 2^k + 2^{k+1} = 2^{k+2} - 1$$

$P(k+1)$  can also be expressed as  $P(k+1) = P(k) + 2^{k+1}$

$$\begin{aligned}
P(k+1) : 1 + 2 + 2^2 + \dots + 2^{k+1} &= 2^{k+1} - 1 + 2^{k+1} \\
&= 2^k \times 2^1 + 2^k \times 2^1 - 1 \\
&= 2^k (2^1 + 2^1) - 1 \\
&= 2^k (2^2) - 1 \\
&= 2^{k+2} - 1
\end{aligned}$$

$\therefore P(k+1)$  is  $T$

Hence we can conclude that  $P(n)$  is true for all non-negative integers  $n$ .

### 1.3.1.0.1 Sums of Geometric Progressions

#### Question 5

Use mathematical induction to prove this formula for the sum of a finite number of terms of a geometric progression with initial term  $a$  and a common ratio  $r$

$$\sum_{j=0}^n ar^j = a + ar + ar^2 + \dots + ar^n = \frac{ar^{n+1} - a}{r - 1} \quad \text{where } r \neq 1$$

where  $n$  is a non-negative integer.

**Solution:** Let  $P(n)$  be "the sum of the first  $n + 1$  terms of a geometric progression in this formula is correct".

#### Basis Step

$$\begin{aligned}
P(0) : \sum_{j=0}^0 ar^j &= \frac{ar^{0+1} - a}{r - 1} \\
a &= \frac{ar - a}{r - 1} \\
a &= \frac{a(r - 1)}{r - 1} \\
a &= a
\end{aligned}$$

#### Induction Step

To complete this step I must prove  $P(k) \rightarrow P(k+1)$  where  $k$  is any non-negative integer.

Assume  $P(k)$  for some non-negative integer  $k$ , then

$$P(k) : \sum_{j=0}^k ar^j = a + ar + ar^2 + \dots + ar^k = \frac{ar^{k+1} - a}{r - 1}$$

Then  $P(k+1)$  is

$$P(k+1) : \sum_{j=0}^{k+1} ar^j = a + ar + ar^2 + \dots + ar^{k+1} = \frac{ar^{k+2} - a}{r - 1}$$

And can be expressed as  $P(k+1) = P(k) + ar^{k+1}$ , therefore:

$$\begin{aligned}
 P(k+1) : \sum_{j=0}^{k+1} ar^j &= a + ar + ar^2 + \dots + ar^k + ar^{k+1} = \frac{ar^{k+1} - a}{r-1} + ar^{k+1} \\
 &= \frac{ar^{k+1} - a}{r-1} + \frac{ar^{k+1}}{1} \\
 &= \frac{ar^{k+1} - a + (ar^{k+1})(r-1)}{r-1} \\
 &= \frac{ar^{k+1} + ar^{k+2} - ar^{k+1} - a}{r-1} \\
 &= \frac{ar^{k+2} - a}{r-1}
 \end{aligned}$$

$\therefore P(k+1)$  is  $T$

Hence we can conclude that  $P(n)$  is true for all non-negative integers  $n$

## 1.4 Exercises

### Question 6

Prove that

$$3 + 3 \times 5 + 3 \times 5^2 + \dots + 3 \times 5^n = \frac{3(5^{n+1} - 1)}{4}$$

whenever  $n$  is non-negative integer

**Solution:**

Let  $P(n)$  be  $3 + 3 \times 5 + 3 \times 5^2 + \dots + 3 \times 5^n = \frac{3(5^{n+1}-1)}{4}$

Basis Step

$$\begin{aligned}
 P(0) : 3 &= \frac{3(5^1 - 1)}{4} \\
 3 &= \frac{12}{4} \\
 3 &= 3
 \end{aligned}$$

Induction Step

To complete this step I must prove  $P(k) \rightarrow P(k+1)$  for any non-negative integer  $k$

Assume  $P(k)$  is  $T$  for some non-negative integer  $k$ , then

$$P(k) : 3 + 3 \times 5 + 3 \times 5^2 + \dots + 3 \times 5^k = \frac{3(5^{k+1} - 1)}{4}$$

Then  $P(k+1)$  is



$$P(k+1) : 3 + 3 \times 5 + 3 \times 5^2 + \dots + 3 \times 5^{k+1} = \frac{3(5^{k+2} - 1)}{4}$$

And can be expressed as  $P(k+1) = P(k) + 3 \times 5^{k+1}$ , therefore:

$$\begin{aligned} P(k+1) : 3 + 3 \times 5 + 3 \times 5^2 + \dots + 3 \times 5^k + 3 \times 5^{k+1} &= \frac{3(5^{k+1} - 1)}{4} + (3 \times 5^{k+1}) \\ &= \frac{3(5^{k+1} - 1) + 4(3 \times 5^{k+1})}{4} \\ &= \frac{3 \times 5^{k+1} - 3 + 12 \times 5^{k+1}}{4} \\ &= \frac{3(5^{k+1} + 4 \times 5^{k+1} - 1)}{4} \end{aligned}$$

Let  $x = 5^{k+1}$

$$\begin{aligned} &= \frac{3(x + 4x - 1)}{4} \\ &= \frac{3(5x - 1)}{4} \\ &= \frac{3(5^1 \times 5^{k+1}) - 1}{4} \\ &= \frac{3(5^{k+2} - 1)}{4} \end{aligned}$$

$\therefore P(k+1)$  is  $T$

Hence we can conclude that  $P(n)$  is true for all non-negative integers  $n$

#### Question 7

Prove that

$$1 \times 1! + 2 \times 2! + \dots + n \times n! = (n+1)! - 1$$

Whenever  $n$  is a positive integer.

**Solution:**

Let  $P(n)$  be  $1 \times 1! + 2 \times 2! + \dots + n \times n! = (n+1)! - 1$

Basis Step

$$\begin{aligned} P(1) : 1 \times 1! &= (1+1)! - 1 \\ 1 &= 2 - 1 \end{aligned}$$

Induction Step

To conclude this step I need to prove  $P(k) \rightarrow P(k+1)$  for any positive integer  $k$

Assume  $P(k)$  is  $T$  for some positive integer  $k$ , then

$$P(k) : 1 \times 1! + 2 \times 2! + \dots + k \times k! = (k+1)! - 1$$

Then  $P(k+1)$  is:

$$P(k+1) : 1 \times 1! + 2 \times 2! + \dots + (k+1) \times (k+1)! = (k+2)! - 1$$

And can be expressed as  $P(k+1) : P(k) + (k+1) \times (k+1)!$ , therefore

$$\begin{aligned} P(k+1) : 1 \times 1! + 2 \times 2! + \dots + k \times k! + (k+1) \times (k+1)! &= ((k+1)! - 1) + ((k+1) \times (k+1)!) \\ &= (k+1)! + (k+1) \times (k+1)! - 1 \\ &= (k+1)! (k+2) - 1 \\ &= (k+2)! - 1 \end{aligned}$$

### Question 8

1. Find a formula for

$$\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots + \frac{1}{2^n}$$

by examining the values of this expression for small values of  $n$

2. Prove the formula you conjectured in part 1.

**Solution:**

- 1.

$$\begin{aligned} S_n &= \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots + \frac{1}{2^n} \\ S_1 &= \frac{1}{2} \\ S_2 &= \frac{3}{4} \\ S_3 &= \frac{7}{8} \\ S_4 &= \frac{15}{16} \\ S_n &= \frac{2^n - 1}{2^n} \end{aligned}$$

2. **Proof:** Let  $P(n)$  be  $\frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots + \frac{1}{2^n} = \frac{2^n - 1}{2^n}$

Basis Step

$$\begin{aligned} P(1) : \frac{1}{2} &= \frac{2^1 - 1}{2^1} \\ \frac{1}{2} &= \frac{1}{2} \end{aligned}$$

### Induction Step

To complete this step I need to prove  $P(k) \rightarrow P(k+1)$  for any positive integer  $k$ .

Assume  $P(k)$  is  $T$  for some positive integer  $k$ . Then:

$$P(k) : \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots + \frac{1}{2^k} = \frac{2^k - 1}{2^k}$$

Then  $P(k+1)$  is:

$$P(k+1) : \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots + \frac{1}{2^{k+1}} = \frac{2^{k+1} - 1}{2^{k+1}}$$

And can be expressed as  $P(k+1) : P(k) + \frac{1}{2^{k+1}}$ , therefore

$$\begin{aligned} P(k+1) : \frac{1}{2} + \frac{1}{4} + \frac{1}{8} + \dots + \frac{1}{2^{k+1}} &= \frac{2^k - 1}{2^k} + \frac{1}{2^{k+1}} \\ &= \frac{(2^k - 1)(2^{k+1}) + 2^k}{(2^k)(2^{k+1})} \end{aligned}$$



### Question 9

Show that 3 divides  $n^3 + 2n$ , whenever  $n \in \mathbb{Z}^+$

**Proof:** This has the same meaning as:

" $n^3 + 2n$  is divisible by 3"

" $n^3 + 2n$  is a multiple of 3"

Let  $P(n)$  be  $n^3 + 2n$  is a multiple of 3

### Basis Step

$$P(1) : 1^3 + 2 \times 1 \text{ is a multiple of 3}$$

True because 3 is a multiple of three.

### Induction Step

Assume  $P(k)$  is  $T$  then

$$"k^3 + 2k \text{ is multiple of 3}" - \exists m \in \mathbb{Z}, k^3 + 2k = 3m$$

And show  $P(k+1)$  is

$$"(k+1)^3 + 2(k+1) \text{ is a multiple of 3}"$$

We have

$$\begin{aligned}(k+1)^3 + 2(k+1) &= k^3 + 3k^2 + 3k + 1 + 2k + 2 \\&= (k^3 + 2k) + 3k^2 + 3k + 3 \\&= 3m + 3k^2 + 3k + 3 \\&= 3(m + k^2 + k + 1) \\ \text{Let } z &= m + k^2 + k + 1 \\&= 3z\end{aligned}$$

Since  $z$  is made up of positive integers  $z$  is a positive integer

$\therefore P(k+1)$

Hence we can conclude that  $P(n)$  is true for all positive integers.

⊙

## Chapter 2

## Exercises

### Question 10

Use mathematical induction to prove that 43 divides  $6^{n+1} + 7^{2n-1}$  for every positive integer  $n$

**Proof:** The given statements can be written as " $6^{n+1} + 7^{2n-1}$  is a multiple of 43".  
Let  $P(n)$  be  $6^{n+1} + 7^{2n-1}$  is a multiple of 43.

#### Basis Step

$$\begin{aligned} P(1) : 6^{1+1} + 7^{2 \times 1 - 1} &\text{ is a multiple of 43} \\ &: 43 \text{ is a multiple of 43} \end{aligned}$$

$P(1)$  is true as 43 is a multiple of 43.

#### Induction Step

Assume  $P(k)$  is  $T$ , then:

$$\text{"}6^{k+1} + 7^{2k-1}\text{" is a multiple of 43, means } \exists m \in \mathbb{Z}, 6^{k+1} + 7^{2k-1} = 43m$$

I must now show that  $P(k+1)$ : " $6^{k+2} + 7^{2(k+1)-1}$ " is a multiple of 43.

$$\begin{aligned}
6^{k+2} + 7^{2(k+1)-1} &= 6^{k+2} + 7^{2k+1} \\
&= 6^{k+1+1} + 7^{2k+2-1} \\
&= 6^{k+1} \times 6^1 + 7^{2k-1} \times 7^2 \\
&= 6^{k+1} \times 6^1 + 7^{2k-1} \times 49 \\
&= 6^{k+1} \times 6^1 + 7^{2k-1} \times 6 + 7^{2k-1} \times 43 \\
&= 6 \left( 6^{k+1} + 7^{2k-1} \right) + 7^{2k-1} \times 43 \\
&= 6 (43m) + 7^{2k-1} \times 43 \\
&= 43 \left( 6m + 7^{2k-1} \right)
\end{aligned}$$

$$\begin{aligned}
\text{Let } u &= 6m + 7^{2k-1} \\
&= 43u
\end{aligned}$$

Since  $u$  is made up of integers  $u$  is an integer

$\therefore P(k+1)$  is  $T$ .

Hence I can conclude  $P(n)$  is true for all positive integers



### Question 11

Prove by induction that  $\sum_{j=0}^n \left(-\frac{1}{2}\right)^j = \frac{2^{n+1} + (-1)^n}{3 \times 2^n}$ , whenever  $n$  is a non-negative integer

**Proof:** Let  $P(n)$  be:

$$1 + \left(-\frac{1}{2}\right)^1 + \left(-\frac{1}{2}\right)^2 + \dots + \left(-\frac{1}{2}\right)^n = \frac{2^{n+1} + (-1)^n}{3 \times 2^n}$$

Basis Step

$$\begin{aligned}
P(0) : 1 &= \frac{2+1}{3 \times 1} \\
1 &= 1
\end{aligned}$$

$\therefore P(0)$  is  $T$

Induction Step

To complete this step I must prove  $P(k) \rightarrow P(k+1)$  for every non non-negative integer  $k$

Assume  $P(k)$  is true for some post integer  $k$ , then:

$$P(k) : 1 + \left(-\frac{1}{2}\right)^1 + \left(-\frac{1}{2}\right)^2 + \dots + \left(-\frac{1}{2}\right)^k = \frac{2^{k+1} + (-1)^k}{3 \times 2^k}$$

Then  $P(k+1)$  is:

$$P(k+1) : 1 + \left(-\frac{1}{2}\right)^1 + \left(-\frac{1}{2}\right)^2 + \dots + \left(-\frac{1}{2}\right)^{k+1} = \frac{2^{k+2} + (-1)^{k+1}}{3 \times 2^{k+1}}$$

And can be expressed as  $P(k+1) : P(k) + \left(-\frac{1}{2}\right)^{k+1}$ , therefore:

$$\begin{aligned}
 P(k+1) : 1 + \left(-\frac{1}{2}\right) + \left(-\frac{1}{2}\right)^2 + \dots + \left(-\frac{1}{2}\right)^k + \left(-\frac{1}{2}\right)^{k+1} &= \frac{2^{k+1} + (-1)^k}{3 \times 2^k} + \left(-\frac{1}{2}\right)^{k+1} \\
 &= \frac{2^{2k+2} + (-1)^k \times 2^{k+1} - 3 \times 2^k \times 1^{k+1}}{3 \times 2^{2k+1}} \\
 &= \frac{2^k (2^{k+2} + (-1)^k \times 2 - 3)}{2^k \times 3 \times 2^{k+1}} \\
 &= \frac{2^{k+2} + (-1)^k \times 2 - 3}{3 \times 2^{k+1}}
 \end{aligned}$$

☺