

Limits and Derivatives

Case 1

Changing input values, output values remain the same \therefore The change in y with respect to changes x values is 0.

This function is called a constant function

$$f(x) = k, \text{ for all input } x$$

Case 2

Changing input values result in changing output values/ different output values.

$$f(x) = \text{not a constant function}$$

Case 3

Changing input values result in a constant change in output values.

$$\Delta f(x) = \text{constant}$$

This function is called a linear function.

Case 4

Changing input values result in a non constant change in output values

$$\Delta f(x) = \text{not constant}$$

Questions

1. Find the derivative of $f(x) = x^2$ at $x = 1$

$$\begin{aligned}\frac{d}{dx}f(x)_{x \rightarrow 1^-} &= 2 \\ \frac{d}{dx}f(x)_{x \rightarrow 1^+} &= 2 \\ \frac{d}{dx}x^2 &= 2\end{aligned}$$

2. Investigate the existence or otherwise the derivatives if the following functions:

$$f(x) = \begin{cases} x^2, & \text{if } x \leq 2 \\ 1 + 2x & \text{if } x > 2 \end{cases} \text{ at } x = 2$$
$$g(x) = |x| = \begin{cases} -x, & \text{if } x < 0 \\ x, & \text{if } x \geq 0 \end{cases} \text{ at } x = 0$$

Q1.

$$\frac{d}{dx} f(x)_{x \rightarrow 2^-} = 4$$

$$\frac{d}{dx} f(x)_{x \rightarrow 2^+} = \text{is not constant}$$

Q2.

$$\frac{d}{dx} g(x)_{x \rightarrow 0^-} = 1$$

$$\frac{d}{dx} g(x)_{x \rightarrow 0^+} = 1$$

$$= \frac{d}{dx} g(x) = 1$$

Derivatives

- Slope or Gradient of a non-linear function
- Derivative = $\frac{dy}{dx}$ = Instantaneous rate of change of a function
- Slope of a line = $\frac{\Delta y}{\Delta x}$ = Average rate of change
- Leverage average rate of change to obtain the instantaneous rate of change = **First Principle**

$$\frac{dy}{dx} = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

- The derivative is a function at a point and a number at another point

Secant

| A line going through two points on a curve

Questions

Q1. Find the derivative of $f(x) = x^2$ using first principle

$$\frac{df}{dx} = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

$$\frac{(x+h)^2 - x^2}{h}$$

$$\frac{2hx + h^2}{h}$$

$$2x + (0)$$

$$\frac{df}{dx} = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = 2x$$

Q2. Find the derivative of $f(x) = \frac{1}{x}$ using first principle

$$\begin{aligned}
\frac{df}{dx} &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
\frac{df}{dx} &= \lim_{h \rightarrow 0} \left(\frac{1}{(x+h)} - \frac{1}{x} \right) \div h \\
&\quad \frac{-h}{x^2 + hx} \div h \\
&\quad \frac{-h}{x^2 + hx} \times \frac{1}{h} \\
&\quad - \frac{1}{x^2 + hx} \\
&\quad - \frac{1}{x^2 + 0x} \\
\frac{df}{dx} &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = -\frac{1}{x^2}
\end{aligned}$$

Q3. Find the derivative of $f(x) = \sqrt{x}$

$$\begin{aligned}
\frac{df}{dx} &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} \\
\frac{df}{dx} &= \lim_{h \rightarrow 0} = \frac{\sqrt{x+h} - \sqrt{x}}{h} \times \frac{\sqrt{x+h} + \sqrt{x}}{\sqrt{x+h} + \sqrt{x}} \\
\frac{df}{dx} &= \lim_{h \rightarrow 0} = \frac{(\sqrt{x+h} - \sqrt{x}) - (\sqrt{x+h} + \sqrt{x})}{h(\sqrt{x+h} + \sqrt{x})} \\
\frac{df}{dx} &= \lim_{h \rightarrow 0} = \frac{x+h-x}{h(\sqrt{x+h} + \sqrt{x})} \\
\frac{df}{dx} &= \lim_{h \rightarrow 0} = \frac{h}{h(\sqrt{x+h} + \sqrt{x})} \\
\frac{df}{dx} &= \lim_{h \rightarrow 0} = \frac{1}{\sqrt{x+h} + \sqrt{x}} \\
&\quad \frac{1}{\sqrt{x+0} + \sqrt{x}} \\
&\quad \frac{1}{2\sqrt{x}} \\
\frac{df}{dx} &= \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h} = \frac{1}{2\sqrt{x}}
\end{aligned}$$

Techniques of Differentiation

Power Rule

$y = x^n$, n is a real number

$$\frac{dy}{dx} = nx^{n-1}$$

Questions

$$y = x^2 \tag{1}$$

$$y = \frac{1}{x^2} \tag{2}$$

$$y = \frac{1}{\sqrt{x}} + x^3 - 1 \tag{3}$$

$$y = x^{-8} + 3x^2 \tag{4}$$

Q1.

$$\begin{aligned}y &= x^2, \quad n = 2 \\ \frac{dy}{dx} &= 2x^{2-1} \\ &= 2x^1 \\ &= 2x\end{aligned}$$

Q2.

$$\begin{aligned}y &= \frac{1}{x^2} \\ y &= x^{-2}, \quad n = -2 \\ y' &= -2x\end{aligned}$$

Q3.

$$\begin{aligned}y &= \sqrt{x} \\ y &= x^{\frac{1}{2}} \\ y' &= \frac{x^{-\frac{1}{2}}}{2} \\ y' &= \frac{1}{2\sqrt{x}}\end{aligned}$$

Q4.

$$\begin{aligned}y &= \frac{1}{\sqrt{x}} + x^3 - 1 \\ y' &= x^{-\frac{1}{2}} + x^3 - 1 \\ y' &= -\frac{1}{2}x^{-\frac{3}{2}} + 3x^2\end{aligned}$$

Qi.

$$\begin{aligned}y &= -x^{-8} + 3x^2 \\ y' &= -8x^{-9} + 6x\end{aligned}$$

Chain Rule

Chain Function

┆ A composite function i.e. $f[g]$

$$\begin{aligned}y &= (2x + 1)^2 \\ y' &= 4(2x + 1)\end{aligned}$$

$$\begin{aligned}y &= (3x^2 + 2x)^5 \\ y &= (5)(6x + 2)(3x^2 + 2x)^4 \\ y &= (30x + 10)(3x^2 + 2x)\end{aligned}$$

Questions

$$y = 15x^9 - 3x^{12} + 5x - 46$$

$$y = 2t^6 + 7t^{-6}$$

$$y = 8x^3 - \frac{1}{3x^5} + x - 23$$

$$y = \sqrt{x} + 9\sqrt[3]{x^4} - \frac{2}{\sqrt[5]{x^2}}$$

$$y = \sqrt[3]{x^2}(2x - x^2)$$

$$y = \frac{2t^5 + t^2 - 5}{t^2}$$

$$y = 2x^3 + \frac{300}{x^3} + 4$$

$$\text{Q1. } y = 15x^9 - 3x^{12} + 5x - 46$$

$$y' = 185x^8 - 36x^{11} + 5$$

$$\text{Q2. } y = 2t^6 + 7t^{-6}$$

$$y' = 12t^5 = 42t^{-7}$$

$$\text{Q3. } y = 8x^3 - \frac{1}{3x^5} + x - 23$$

$$y = 8x^3 - \frac{1}{3x^5} + x - 23$$

$$y = 8x^3 - \frac{1}{3}x^{-5} + x - 23$$

$$y' = 24x^2 + \frac{5}{3}x^{-6} + 1$$

$$\text{Q4. } y = \sqrt{x} + 9\sqrt[3]{x^4} - \frac{2}{\sqrt[5]{x^2}}$$

$$y = \sqrt{x} + 9\sqrt[3]{7} - \frac{2}{\sqrt[5]{x^2}}$$

$$y = x^{\frac{1}{2}} + 9(x^7)^{\frac{1}{3}} - 2(x^2)^{-\frac{1}{5}}$$

$$y = x^{\frac{1}{2}} + 9x^{\frac{7}{3}} - 2x^{-\frac{2}{5}}$$

$$y' = \frac{1}{2}x^{-\frac{1}{2}} + 21x^{\frac{4}{3}} + \frac{4}{5}x^{-\frac{7}{5}}$$

$$y' = \frac{1}{2\sqrt{x}} + 21x^{\frac{4}{3}} + \frac{4}{5}x^{-\frac{7}{5}}$$

$$\text{Q5. } y = \sqrt[3]{x^2}(2x - x^2)$$

$$y = (x^2)^{\frac{1}{3}}(2x - x^2)$$

$$y = x^{\frac{2}{3}}(2x - x^2)$$

$$y = 2x^{\frac{5}{3}} - x^{\frac{2}{3}}$$

$$y' = \frac{10}{3}x^{\frac{2}{3}} - \frac{8}{3}x^{\frac{5}{3}}$$

$$\text{Q6. } y = \frac{2t^5 + t^2 - 5}{t^2}$$

$$y = 2t^3 + 1 - \frac{5}{t^2}$$

$$y = 2t^3 + 1 - 5t^{-2}$$

$$y' = 6t^2 + 10t^{-3}$$

$$\text{Q7. } y = 2x^3 + \frac{300}{x^3} + 4$$

$$y = 2x^3 + 300x^{-3} + 4$$

$$y' = 6x^2 - 900x^{-4}$$

Product Rule

Given $y = u \times v$

$$\frac{dy}{dx} = u' \times v + v' \times u$$

Q1. $y = (x^2 + 1)(x^3 - x)$

$$\begin{aligned}y &= (x^2 + 1)(x^3 - x) \\y' &= (2x)(x^3 - x) + (x^2 + 1)(3x^2 - 1) \\y' &= 2x^4 - 2x^2 + 3x^4 + 2x^2 - 1 \\y' &= 5x^4 - 1\end{aligned}$$

Q2. $y = (6x^3 - x)(10 - 20x)$

$$\begin{aligned}y' &= (18x^2 - 1)(10 - 20x) + (-20)(6x^3 - x) \\y' &= 180x^2 - 10 - 360x^3 + 20x - 120x^3 + 20x \\y' &= -480x^3 + 180x^2 + 40x - 10\end{aligned}$$

Questions

$$\begin{aligned}y &= (4t^2 - t)(t^3 - 8t^2 + 12) \\y &= (1 + \sqrt{x^3})(x^{-3} - 2\sqrt[3]{x}) \\y &= (4 - t^2)(1 + 5t^2) \\y &= (x - \frac{2}{x^2})(7 - 2x^3) \\y &= (3 - x)(1 - 2x + x^2)\end{aligned}$$

Q1. $y = (4t^2 - t)(t^3 - 8t^2 + 12)$

$$\begin{aligned}y' &= (8t - 1)(t^3 - 8t^2 + 12) + (3t^2 - 16t)(4t^2 - t) \\y' &= 8t^4 - 64t^3 + 96t - t^3 + 8t^2 - 12 + 12t^4 - 3t^3 - 64t^3 + 16t^2 \\y' &= 20t^4 - 132t^3 + 24t^2 + 96t - 12\end{aligned}$$

Q2. $y = (1 + \sqrt{x^3})(x^{-3} - 2\sqrt[3]{x})$

$$\begin{aligned}y &= (1 + (x^3)^{\frac{1}{2}})(x^{-3} - 2\sqrt[3]{x}) \\y &= (1 + x^{\frac{1}{2}})(x^{-3} - 2(x^{\frac{1}{3}})) \\y' &= (\frac{3}{2}x^{\frac{1}{2}})(x^{-3} - 2x^{\frac{1}{3}}) + (-3x^4 - \frac{2}{3}x^{-\frac{2}{3}} - \frac{2}{3}x^{\frac{5}{6}}) \\y' &= \frac{3}{2}x^{-\frac{5}{2}} - 3x^{\frac{5}{6}} - 3x^{-4} - 3x^{-\frac{5}{2}} - \frac{2}{3}x^{-\frac{2}{3}} - \frac{2}{3}x^{\frac{5}{6}} \\y' &= -\frac{11}{3}x^{\frac{5}{6}} - \frac{3}{2}x^{-\frac{5}{2}} - \frac{2}{3}x^{-\frac{2}{3}} - 3x^{-4}\end{aligned}$$

Q3. $y = (4 - t^2)(1 + 5t^2)$

$$\begin{aligned}y' &= (-2t)(1 + 5t^2) + (10t)(4 - t^2) \\y' &= -2t - 10t^3 + 40t - 10t^3 \\y' &= 20t^3 + 38t\end{aligned}$$

Q4. $y = (x - \frac{2}{x^2})(7 - 2x^3)$

$$\begin{aligned}
 y &= (x - 2x^{-2})(7 - 2x^3) \\
 y' &= (1 + 4x^{-3})(7 - 2x^3) + (-6x^2)(x - 2x^{-2}) \\
 y' &= 7 - 2x^3 + 28x^{-3} - 8x^0 - 6x^3 + 12x^0 \\
 y' &= 7 - 2x^3 + 28x^{-3} - 8 - 6x^3 + 12 \\
 y' &= -8x^3 + \frac{28}{x^3} + 11
 \end{aligned}$$

Q5. $y = (3 - x)(1 - 2x + x^2)$

$$\begin{aligned}
 y' &= (-1)(1 - 2x + x^2) + (2x - 2)(3 - x) \\
 y' &= -1 + 2x - x^2 + 6x - 2x^2 - 6 + 2x \\
 y' &= -7 + 10x - 3x^2
 \end{aligned}$$

Quotient Rule

Where $y = \frac{f(x)}{g(x)}$

$$\frac{dy}{dx} = \frac{g(x) \times f'(x) - f(x) \times g'(x)}{(g(x))^2}$$

Where $y = \frac{u}{v}$

$$y' = \frac{v \times u' - u \times v'}{v^2}$$

Questions