

Duality theory in linear optimization and its extensions—formally verified

Abstract: Farkas established that a system of linear inequalities has a solution if and only if we cannot obtain a contradiction by taking a linear combination of the inequalities. We state and formally prove several Farkas-like theorems in Lean 4. Furthermore, we consider a linearly ordered field extended with two special elements denoted by \perp and \top where \perp is below every element and \top is above every element. We define $\perp + a = \perp = a + \perp$ of all a and we define $\top + b = \top = b + \top$ for all $b \neq \perp$. For multiplication, we define $\perp \cdot c = \perp = c \cdot \perp$ for every $c \geq 0$ but $\top \cdot d = \top = d \cdot \top$ only for $d > 0$ because $\top \cdot 0 = 0 = 0 \cdot \top$. We extend certain Farkas-like theorems to a setting where coefficients are from an extended linearly ordered field.

1 Introduction

Rouché, Capelli, Kronecker, Fonténe, Frobenius (TODO citations) established that a system of linear equalities has a solution if and only if we cannot obtain a contradiction by taking a linear combination of the equalities. We state their theorem as follows.

Theorem (equalityFrobenius): Let I and J be finite types. Let F be a linearly ordered field. Let A be a matrix of type $(I \times J) \rightarrow F$. Let b be a vector of type $I \rightarrow F$. Exactly one of the following exists:

- vector $x : J \rightarrow F$ such that $A \cdot x = b$
- vector $y : I \rightarrow F$ such that $A^T \cdot y = 0$ and $b \cdot y \neq 0$

Geometric interpretation of equalityFrobenius is straightforward. The column vectors of A generate a hyperplane in the $|I|$ -dimensional Euclidean space that goes contains the origin. The point b either lies in this hyperplane (in this case, the entries of x give coefficients which, when applied to the column vectors of A , give a vector from the origin to the point b), or there exists a line through the origin that is orthogonal to all the column vectors of A (i.e., orthogonal to the entire hyperplane) such that b projected onto this line falls outside of the origin (in this case, y gives a direction of this line), i.e., to a different point from where all column vectors of A get projected.

This theorem can be given in much more general settings. In our paper, however, this is the only version we provide. This staple of linear algebra is not our main focus but a byproduct of the other theorems we prove. In particular, we obtain equalityFrobenius as an immediate corollary of the following theorem.

Theorem (equalityFrobenius_lt): Let I and J be finite types. Let F be a linearly ordered field. Let A be a matrix of type $(I \times J) \rightarrow F$. Let b be a vector of type $I \rightarrow F$. Exactly one of the following exists:

- vector $x : J \rightarrow F$ such that $A \cdot x = b$
- vector $y : I \rightarrow F$ such that $A^T \cdot y = 0$ and $b \cdot y < 0$

The way we state the theorem exemplifies certain patterns that permeate through our work. Our results are phrased as “there are two systems of (in)equalities; exactly one of them has a solution”. This goes hand-in-hand with our decision to focus mostly on “symmetric” Farkas-like theorems. Note that it must be impossible to satisfy the second system by $y = 0$. Had it been allowed, we would have said nothing about the first system as it would have to lead to a contradiction every time. The constraint $b \cdot y < 0$ disqualifies the zero solution in most of our theorems. Intuitively, it should make sense that one of the systems is always “strict” (to easily see why, consider I and J singletons), which also means that our two systems will never be “fully symmetric”.

Farkas (TODO citation) gave a similar characterization for systems of linear equalities with nonnegative variables. We state his theorem as follows.

Theorem (equalityFarkas): Let I and J be finite types. Let F be a linearly ordered field. Let A be a matrix of type $(I \times J) \rightarrow F$. Let b be a vector of type $I \rightarrow F$. Exactly one of the following exists:

- nonnegative vector $x : J \rightarrow F$ such that $A \cdot x = b$
- vector $y : I \rightarrow F$ such that $A^T \cdot y \geq 0$ and $b \cdot y < 0$

Geometric interpretation of equalityFarkas is easy. The column vectors of A generate a cone in the $|I|$ -dimensional Euclidean space from the origin towards some infinity. The point b either lies inside this cone (in this case, the entries of x give nonnegative coefficients which, when applied to the column vectors of A , give a vector from the origin to the point b), or there exists a hyperplane that contains the origin and that strictly separates b from given cone (in this case, y gives a normal vector of this hyperplane).

We prove equalityFrobenius_lt by applying equalityFarkas to the matrix $(A \mid -A)$. However, equalityFarkas will be proved later, from a more general theorem.

Minkowski (TODO citation) similarly established that a system of linear inequalities has a nonnegative solution if and only if we cannot obtain a contradiction by taking a nonnegative linear combination of the inequalities. For reasons that will be apparent later, we give two versions of this theorem.

Theorem (inequalityFarkas): Let I and J be finite types. Let F be a linearly ordered field. Let A be a matrix of type $(I \times J) \rightarrow F$. Let b be a vector of type $I \rightarrow F$. Exactly one of the following exists:

- nonnegative vector $x : J \rightarrow F$ such that $A \cdot x \leq b$
- nonnegative vector $y : I \rightarrow F$ such that $A^T \cdot y \geq 0$ and $b \cdot y < 0$

Geometric interpretation of inequalityFarkas is a bit harder. The column vectors of A generate a cone in the $|I|$ -dimensional Euclidean space from the origin to some infinity. The point b determines an orthogonal cone that starts in b and goes to negative infinity in the direction of all coordinate axes. Either these two cones intersect (in this case, the entries of x give nonnegative coefficients which, when applied to the column vectors of A , give a vector from the origin to a point in the intersection), or there exists a hyperplane that contains the origin and that strictly separates b from the cone generated by A but does not cut through the positive orthant, i.e., the origin is the only nonnegative point contained in the hyperplane (in this case, y gives a normal vector of this hyperplane).

Theorem (inequalityFarkas_neg): Let I and J be finite types. Let F be a linearly ordered field. Let A be a matrix of type $(I \times J) \rightarrow F$. Let b be a vector of type $I \rightarrow F$. Exactly one of the following exists:

- nonnegative vector $x : J \rightarrow F$ such that $A \cdot x \leq b$
- nonnegative vector $y : I \rightarrow F$ such that $(-A^T) \cdot y \leq 0$ and $b \cdot y < 0$

Obviously, inequalityFarkas_neg is an immediate corollary of inequalityFarkas. We prove inequalityFarkas by applying equalityFarkas to the matrix $(1 \mid A)$ where 1 is the identity matrix of type $(I \times I) \rightarrow F$.

The next theorem generalizes equalityFarkas to structures where multiplication does not have to be commutative. Furthermore, it supports infinitely many equations.

Theorem (coordinateFarkas): Let I be any type. Let J be a finite type. Let R be a linearly ordered division ring. Let A be an R -linear map from from $(I \rightarrow R)$ to $(J \rightarrow R)$. Let b be an R -linear map from from $(I \rightarrow R)$ to R . Exactly one of the following exists:

- nonnegative vector $x : J \rightarrow R$ such that, for all $w : I \rightarrow R$, we have $\sum_{j:J} (A \ w)_j \bullet x_j = b \ w$
- vector $y : I \rightarrow R$ such that $A \ y \geq 0$ and $b \ y < 0$

In the next generalization, we replace the partially ordered module $I \rightarrow R$ by a general R -module W .

Theorem (scalarFarkas): Let J be a finite type. Let R be a linearly ordered division ring. Let W be an R -module. Let A be an R -linear map from from W to $(J \rightarrow R)$. Let b be an R -linear map from from W to R . Exactly one of the following exists:

- nonnegative vector $x : J \rightarrow R$ such that, for all $w : W$, we have $\sum_{j:J} (A \ w)_j \bullet x_j = b \ w$
- vector $y : W$ such that $A \ y \geq 0$ and $b \ y < 0$

In the most general theorem, stated below, we replace certain occurrences of R by a linearly ordered R -module V whose order respects order on R . This result origins from TODO.

Theorem (fintypeFarkasBartl): Let J be a finite type. Let R be a linearly ordered division ring. Let W be an R -module. Let V be a linearly ordered R -module¹. Let A be an R -linear map from W to $(J \rightarrow R)$. Let b be an R -linear map from W to V . Exactly one of the following exists:

- nonnegative bivector $x : J \rightarrow V$ such that, for all $w : W$, we have $\sum_{j:J} (A \ w)_j \bullet x_j = b \ w$
- vector $y : W$ such that $A \ y \geq 0$ and $b \ y < 0$

In the last branch, $A \ y \geq 0$ uses the partial order² on $(J \rightarrow R)$ whereäs $b \ y < 0$ uses the linear order³ on V . Note that fintypeFarkasBartl subsumes scalarFarkas (as well as the other versions based on equality), since R can be viewed as a linearly ordered module over itself. We prove fintypeFarkasBartl in Section TODO, which is where the heavy lifting comes.

Until now, we have talked about known results. What follows is a new extension of the theory.

Definition: Let F be a linearly ordered field. We define an **extended** linearly ordered field F_∞ as $F \cup \{\perp, \top\}$ with the following properties. Let p and q be numbers from F . We have $\perp < p < \top$. We define addition and scalar action on F_∞ as follows:

+		\perp	q	\top
\perp		\perp	\perp	\perp
p		\perp	$p+q$	\top
\top		\perp	\top	\top

\bullet		\perp	q	\top
0		\perp	0	0
$p > 0$		\perp	$p \cdot q$	\top

¹We furthermore require monotonicity of scalar multiplication by nonnegative elements on the left. This assumption will be implicit in later occurrences.

²The order on $(J \rightarrow R)$ is always the coordinate-wise application of R 's linear order.

³In case V has finite dimension, you can choose an arbitrary direction and project vectors from V onto it, or you can order elements of V lexicographically.

Informally speaking, \top represents the positive infinity, \perp represents the negative infinity, and we say that \perp is “stronger” than \top in all arithmetic operations. The surprising parts are $\perp + \top = \perp$ and $0 \bullet \perp = \perp$. Because of that, F_∞ is not a field. However, F_∞ is still a densely linearly ordered abelian monoid with characteristic zero.

Theorem (extendedFarkas): Let I and J be finite types. Let F be a linearly ordered field. Let A be a matrix of type $(I \times J) \rightarrow F_\infty$. Let b be a vector of type $I \rightarrow F_\infty$. Assume that A does not have \perp and \top in the same row. Assume that A does not have \perp and \top in the same column. Assume that A does not have \top in any row where b has \top . Assume that A does not have \perp in any row where b has \perp . Exactly one of the following exists:

- nonnegative vector $x : J \rightarrow F$ such that $A \cdot x \leq b$
- nonnegative vector $y : I \rightarrow F$ such that $(-A^T) \cdot y \leq 0$ and $b \cdot y < 0$

Note that extendedFarkas looks pretty much like equalityFarkas_neg and, in certain sense, generalizes it. Indeed, in Section TODO, we prove extendedFarkas using equalityFarkas_neg and some additional machinery.

Definition: Let I and J be finite types. Let R be a partially ordered semiring. Let A be a matrix of type $(I \times J) \rightarrow R$. Let b be a vector of type $I \rightarrow R$. Let c be a vector of type $J \rightarrow R$. We say that $P = (A, b, c)$ is a **linear program** over R . We say that its **constraints** are indexed by I . We say that its **variables** are indexed by J . We say that a nonnegative vector $x : J \rightarrow R$ is a **solution** to P if and only if $A \cdot x \leq b$. We say that P **reaches** an objective value r if and only if there exists x such that x is a solution to P and $c \cdot x = r$.

TODO define feasibility and dualize.

TODO state weakDuality and strongDuality.

2 Formalization

2.1 We start with a review of algebraic typeclasses that our project depends on

Additive semigroup is a structure on any type with addition where the addition is associative:

```
class AddSemigroup (G : Type u) extends Add G where
  protected add_assoc : ∀ a b c : G, (a + b) + c = a + (b + c)
```

Additive monoid is an additive semigroup with the zero element, thanks to which we can define a scalar multiplication by the natural numbers:

```
class AddMonoid (M : Type u) extends AddSemigroup M, AddZeroClass M where
  protected nsmul : ℕ → M → M
  protected nsmul_zero : ∀ x, nsmul 0 x = 0 := by intros; rfl
  protected nsmul_succ : ∀ (n : ℕ) (x), nsmul (n + 1) x = nsmul n x + x := by intros; rfl
```

Subtractive monoid is an additive monoid that adds two more operations (unary and binary minus) that satisfy some basic properties:

```
class SubNegMonoid (G : Type u) extends AddMonoid G, Neg G, Sub G where
  protected sub := SubNegMonoid.sub'
  protected sub_eq_add_neg : ∀ a b : G, a - b = a + -b := by intros; rfl
  protected zsmul : ℤ → G → G
  protected zsmul_zero' : ∀ a : G, zsmul 0 a = 0 := by intros; rfl
  protected zsmul_succ' (n : ℕ) (a : G) :
    zsmul (Int.ofNat n.succ) a = zsmul (Int.ofNat n) a + a := by intros; rfl
  protected zsmul_neg' (n : ℕ) (a : G) : zsmul (Int.negSucc n) a = -zsmul n.succ a := by intros; rfl
```

Additive group is a subtractive monoid in which the unary minus acts as an inverse with respect to addition:

```
class AddGroup (A : Type u) extends SubNegMonoid A where
  protected add_left_neg : ∀ a : A, -a + a = 0
```

Abelian group is defined as an additive group that is a commutative additive monoid at the same time (TODO AddCommMonoid):

```
class AddCommGroup (G : Type u) extends AddGroup G, AddCommMonoid G
```

Ring is defined as a semiring that is an abelian group at the same time and has 1 that behaves well (TODO Semiring):

```
class Ring (R : Type u) extends Semiring R, AddCommGroup R, AddGroupWithOne R
```

Division ring is a ring with a lot of extra requirements (TODOs DivInvMonoid, Nontrivial, NNRatCast, RatCast):

```
class DivisionRing (α : Type*) extends Ring α, DivInvMonoid α, Nontrivial α, NNRatCast α, RatCast α where
  protected mul_inv_cancel : ∀ (a : α), a ≠ 0 → a * a⁻¹ = 1
  protected inv_zero : (0 : α)⁻¹ = 0
  protected nnratCast := NNRat.castRec
```

We define a linearly ordered division ring as a division ring that is a linearly ordered ring at the same time (TODO all about order):

```
class LinearOrderedDivisionRing (R : Type*) extends LinearOrderedRing R, DivisionRing R
```

Linearly ordered field is defined as a linearly ordered commutative ring that is a field at the same time (TODO Field):

```
class LinearOrderedField (α : Type*) extends LinearOrderedCommRing α, Field α
```

Note that LinearOrderedDivisionRing is not a part of the algebraic hierarchy provided by Mathlib, hence LinearOrderedField does not inherit LinearOrderedDivisionRing, thus we provide a custom instance that converts LinearOrderedField to LinearOrderedDivisionRing:

```
instance LinearOrderedField.toLinearOrderedDivisionRing {F : Type*} [instF : LinearOrderedField F] :  
  LinearOrderedDivisionRing F := { instF with }
```

This instance is needed for the step from coordinateFarkas to equalityFarkas.

2.2 Vectors and stuff

We distinguish two types of vectors; implicit vectors and explicit vectors. Implicit vectors are members of a vector space; they don't have any internal structure. Explicit vectors are functions from coordinates to values. The set of coordinates needn't be ordered. Matrices live next to explicit vectors. They are also functions; they take a row index and a column index and they output a value at the given spot. Neither the row indices nor the column vertices are required to form an ordered set. That's why multiplication between matrices and vectors is defined only in structures where addition forms a commutative semigroup. Consider the following example:

$$\begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \end{pmatrix} \cdot \begin{pmatrix} 7 \\ 8 \\ 9 \end{pmatrix} = \begin{pmatrix} ? \\ - \end{pmatrix}$$

We don't know whether the value at the question mark is equal to $(1 \cdot 7 + 2 \cdot 8) + 3 \cdot 9$ or to $(2 \cdot 8 + 1 \cdot 7) + 3 \cdot 9$ or to any other ordering of summands. This is why commutativity of addition is necessary for the definition to be valid. On the other hand, we don't assume any property of multiplication in the definition of multiplication between matrices and vectors; they don't even have to be of the same type; we only require the elements of the vector to have an action on the elements of the matrix (this is not a typo – normally, we would want matrices to have an action on vectors – not in our work).

TODO formal definitions.

We start from bottom up!

Theorem equalityFrobenius is stated as follows:

```
theorem equalityFrobenius (A : Matrix I J F) (b : I → F) :  
  (∃ x : J → F, A *v x = b) ≠ (∃ y : I → F, AT *v y = 0 ∧ b ·v y ≠ 0)
```

Theorem equalityFrobenius_lt is stated as follows:

```
theorem equalityFrobenius_lt (A : Matrix I J F) (b : I → F) :  
  (∃ x : J → F, A *v x = b) ≠ (∃ y : I → F, AT *v y = 0 ∧ b ·v y < 0)
```

Theorem equalityFarkas is stated as follows:

```
theorem equalityFarkas (A : Matrix I J F) (b : I → F) :  
  (∃ x : J → F, 0 ≤ x ∧ A *v x = b) ≠ (∃ y : I → F, 0 ≤ AT *v y ∧ b ·v y < 0)
```

Theorem inequalityFarkas is stated as follows:

```
theorem inequalityFarkas [DecidableEq I] (A : Matrix I J F) (b : I → F) :  
  (∃ x : J → F, 0 ≤ x ∧ A *v x ≤ b) ≠ (∃ y : I → F, 0 ≤ y ∧ 0 ≤ AT *v y ∧ b ·v y < 0)
```

Theorem inequalityFarkas_neg is stated as follows:

```
theorem inequalityFarkas_neg [DecidableEq I] (A : Matrix I J F) (b : I → F) :  
  (∃ x : J → F, 0 ≤ x ∧ A *v x ≤ b) ≠ (∃ y : I → F, 0 ≤ y ∧ -AT *v y ≤ 0 ∧ b ·v y < 0)
```

Theorem coordinateFarkas is stated as follows:

```
theorem coordinateFarkas {I J : Type*} [Fintype J] [LinearOrderedDivisionRing R]  
  (A : (I → R) →l [R] J → R) (b : (I → R) →l [R] R) :  
  (∃ x : J → R, 0 ≤ x ∧ ∀ w : I → R, ∑ j : J, A w j • x j = b w) ≠ (∃ y : I → R, 0 ≤ A y ∧ b y < 0)
```

Theorem scalarFarkas is stated as follows:

```
theorem scalarFarkas {J : Type*} [Fintype J] [LinearOrderedDivisionRing R] [AddCommGroup W] [Module R W]  
  (A : W →l [R] J → R) (b : W →l [R] R) :  
  (∃ x : J → R, 0 ≤ x ∧ ∀ w : W, ∑ j : J, A w j • x j = b w) ≠ (∃ y : W, 0 ≤ A y ∧ b y < 0)
```

Theorem fintypeFarkasBartl is stated as follows:

```
theorem fintypeFarkasBartl {J : Type*} [Fintype J] [LinearOrderedDivisionRing R]  
  [LinearOrderedAddCommGroup V] [Module R V] [PosSMulMono R V] [AddCommGroup W] [Module R W]  
  (A : W →l [R] J → R) (b : W →l [R] V) :  
  (∃ x : J → V, 0 ≤ x ∧ ∀ w : W, ∑ j : J, A w j • x j = b w) ≠ (∃ y : W, 0 ≤ A y ∧ b y < 0)
```

TODO.

Also TODO somehow politely say that the Mathlib's API for block matrices is a mess and needs an overhaul.

3 Proving the Farkas-Bartl theorem

We prove `finFarkasBartl` and, in the end, we obtain `fintypeFarkasBartl` as corollary.

Theorem (`finFarkasBartl`): Let n be a natural number. Let R be a linearly ordered division ring. Let W be an R -module. Let V be a linearly ordered R -module. Let A be an R -linear map from W to $([n] \rightarrow R)$. Let b be an R -linear map from W to V . Exactly one of the following exists:

- nonnegative bivector $x : [n] \rightarrow V$ such that, for all $w : W$, we have $\sum_{j:[n]} (A\ w)_j \bullet x_j = b\ w$
- vector $y : W$ such that $A\ y \geq 0$ and $b\ y < 0$

The only difference is that `finFarkasBartl` uses $[n] = \{0, \dots, n-1\}$ instead of an arbitrary (unordered) finite type J .

Proof idea: We first prove that both cannot exist at the same time. Assume we have x and y of said properties. We plug y for w and obtain $\sum_{j:[n]} (A\ y)_j \bullet x_j = b\ y$. On the left-hand side, we have a sum of nonnegative vectors, which contradicts $b\ y < 0$.

We prove “at least one exists” by induction on n . If $n = 0$ then $A\ y \geq 0$ is a tautology. We consider b . Either b maps everything to the zero vector, which allows x to be the empty bivector, or some w gets mapped to a nonzero vector, where we choose y to be either w or $(-w)$. Since V is linearly ordered, one of them satisfies $b\ y < 0$. The induction step will be proved for all A and all b , not only for given A and b .

Lemma (`industepFarkasBartl`): Let m be a natural number. Let R be a linearly ordered division ring. Let W be an R -module. Let V be a linearly ordered R -module. Assume (induction hypothesis) that for all R -linear maps $A_0 : W \rightarrow ([m+1] \rightarrow R)$ and $b_0 : W \rightarrow V$, the formula “ $\forall y_0 : W, A_0\ y_0 \geq 0 \implies b_0\ y_0 \geq 0$ ” implies existence of a nonnegative bivector $x_0 : [m] \rightarrow V$ such that, for all $w_0 : W$, $\sum_{i:[m]} (A_0\ w_0)_i \bullet (x_0)_i = b_0\ w_0$. Let A be an R -linear map from W to $([m+1] \rightarrow R)$. Let b be an R -linear map from W to V . Assume that, for all $y : W$, $A\ y \geq 0$ implies $b\ y \geq 0$. We claim there exists a nonnegative bivector $x : [m+1] \rightarrow V$ such that, for all $w : W$, we have $\sum_{i:[m+1]} (A\ w)_i \bullet x_i = b\ w$.

Proof idea: Let $A_{<m}$ mean $A \upharpoonright [m]$. We distinguish two cases. If, for all $y : W$, the inequality $A_{<m}\ y \geq 0$ implies $b\ y \geq 0$, then plug $A_{<m}$ for A_0 , obtain x_0 , and construct a bivector x such that $x_m = 0$ and otherwise x copies x_0 . We easily check that x is nonnegative and that $\sum_{i:[m+1]} (A\ w)_i \bullet x_i = b\ w$ holds.

In the second case, we have y' such that $A_{<m}\ y' \geq 0$ holds but $b\ y' < 0$ also holds. We realize that $(A\ y')_m < 0$. We now declare $y := (A\ y')_m \bullet y'$ and observe $(A\ y)_m = 1$. We establish the following facts (proofs are omitted):

- for all $w : W$, we have $A\ (w - ((A\ w)_m \bullet y)) = 0$
- for all $w : W$, the inequality $A_{<m}\ (w - ((A\ w)_m \bullet y)) \geq 0$ implies $b\ (w - ((A\ w)_m \bullet y)) \geq 0$
- for all $w : W$, the inequality $A_{<m}\ w - A_{<m}\ ((A\ w)_m \bullet y) \geq 0$ implies $b\ w - b\ ((A\ w)_m \bullet y) \geq 0$
- for all $w : W$, the inequality $(A_{<m} - (z \mapsto (A\ z)_m \bullet (A_{<m}\ y)))\ w \geq 0$ implies $(b - (z \mapsto (A\ z)_m \bullet (b\ y)))\ w \geq 0$

We observe that $A_0 := A_{<m} - (z \mapsto (A\ z)_m \bullet (A_{<m}\ y))$ and $b_0 := b - (z \mapsto (A\ z)_m \bullet (b\ y))$ are R -linear maps. Thanks to the last fact, we can apply induction hypothesis to A_0 and b_0 . We obtain a nonnegative bivector x' such that, for all $w_0 : W$, $\sum_{i:[m]} (A_0\ w_0)_i \bullet x'_i = b_0\ w_0$. It remains to construct a nonnegative bivector $x : [m+1] \rightarrow V$ such that, for all $w : W$, we have $\sum_{i:[m+1]} (A\ w)_i \bullet x_i = b\ w$. We choose $x_m = b\ y - \sum_{i:[m]} (A_{<m}\ y)_i \bullet x'_i$ and otherwise x copies x' . We check that our x has the required properties. Qed.

We complete the proof of `finFarkasBartl` by applying `industepFarkasBartl` to $A_{\leq n}$ and b . Finally, we obtain `fintypeFarkasBartl` from `finFarkasBartl` using some boring mechanisms regarding equivalence between finite types.

4 Extended Farkas theorem

Theorem (`extendedFarkas`): Let I and J be finite types. Let F be a linearly ordered field. Let A be a matrix of type $(I \times J) \rightarrow F_\infty$. Let b be a vector of type $I \rightarrow F_\infty$. Assume that A does not have \perp and \top in the same row. Assume that A does not have \perp and \top in the same column. Assume that A does not have \top in any row where b has \top . Assume that A does not have \perp in any row where b has \perp . Exactly one of the following exists:

- nonnegative vector $x : J \rightarrow F$ such that $A \cdot x \leq b$
- nonnegative vector $y : I \rightarrow F$ such that $(-A^T) \cdot y \leq 0$ and $b \cdot y < 0$

(restated)

4.1 Proof idea

We need to do the following steps in given order:

1. Delete all rows of both A and b where A has \perp or b has \top (they are tautologies).
2. Delete all columns of A that contain \top (they force respective variables to be zero).
3. If b contains \perp , then $A \cdot x \leq b$ cannot be satisfied, but $y = 0$ satisfies $(-A^T) \cdot y \leq 0$ and $b \cdot y < 0$. Stop here.
4. Assume there is no \perp in b . Use `inequalityFarkas_neg`. In either case, extend x or y with zeros on all deleted positions.

4.2 Counterexamples

If A has \perp and \top in the same row, it may happen that both x and y exist:

$$A = \begin{pmatrix} \perp & \top \\ 0 & -1 \end{pmatrix} \quad b = \begin{pmatrix} 0 \\ -1 \end{pmatrix} \quad x = \begin{pmatrix} 1 \\ 1 \end{pmatrix} \quad y = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

If A has \perp and \top in the same column, it may happen that both x and y exist:

$$A = \begin{pmatrix} \perp \\ \top \end{pmatrix} \quad b = \begin{pmatrix} -1 \\ 0 \end{pmatrix} \quad x = \begin{pmatrix} 0 \end{pmatrix} \quad y = \begin{pmatrix} 1 \\ 1 \end{pmatrix}$$

If A has \top in a row where b has \top , it may happen that both x and y exist:

$$A = \begin{pmatrix} \top \\ -1 \end{pmatrix} \quad b = \begin{pmatrix} \top \\ -1 \end{pmatrix} \quad x = \begin{pmatrix} 1 \end{pmatrix} \quad y = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$$

If A has \perp in a row where b has \perp , it may happen that both x and y exist:

$$A = \begin{pmatrix} \perp \end{pmatrix} \quad b = \begin{pmatrix} \perp \end{pmatrix} \quad x = \begin{pmatrix} 1 \end{pmatrix} \quad y = \begin{pmatrix} 0 \end{pmatrix}$$

5 Proving the (extended) strong LP duality

We note that it is intentional that both PRIM and DUAL are defined as minimization problems.
TODO.

6 Related work

TODO presents an overcomplicated proof in Isabelle by analyzing the Simplex algorithm that already had been formally verified. It took them 30 pages to get to the basic Farkas; no generalization was provided.

In Lean, it would be possible to prove Farkas for reals using the Hahn-Banach separation theorem. However, we do not yet know that the set of feasible solutions is closed.

TODO.

7 Conclusion

We formally verified several Farkas-like theorems in Lean 4. We extended the existing theory to a new setting where some coefficient can carry infinite values. We realized that the abstract work with modules over linearly ordered division rings and linear maps between them was fairly easy to carry on in Lean 4 thanks to the library Mathlib that is perfectly suited for such tasks. In contrast, manipulation with matrices got tiresome whenever we needed a not-fully-standard operation. It turns out Lean 4 cannot automate case analyses unless they take place in the “outer layers” of formulas. Summation over subtypes and summation of conditional expression made us develop a lot of ad-hoc machinery which we would have preferred to be handled by existing tactics. Another area where Lean 4 is not yet helpful is the search for counterexamples. Despite these difficulties, we find Lean 4 to be an extremely valuable tool for elegant expressions of mathematical formulas and for proving them formally.