

**Theorem 1.** *Recursively-enumerable languages are closed under the Kleene star.*

*Proof.* Let  $\mathbf{L} \subseteq \mathbf{T}^*$  be a language generated by grammar  $\mathbf{G} = (\mathbf{T}, \mathbf{N}, S, \mathbf{P})$ . We construct a grammar  $\mathbf{G}^* = (\mathbf{T}, \mathbf{N}^*, Z, \mathbf{P}^*)$  to generate the language  $\mathbf{L}^*$ .

$$\mathbf{N}^* = \mathbf{N} \cup \{Z, \#, R\}$$

We expanded the set of nonterminals with a new starting symbol, a delimiter, and a symbol that denotes final rewriting.

$$\mathbf{P}^* = \mathbf{P} \cup \{Z \rightarrow ZS\#, Z \rightarrow R\#, R\# \rightarrow R, R\# \rightarrow \epsilon\} \cup \{Rt \rightarrow tR \mid t \in \mathbf{T}\}$$

To be continued!

**Lemma 2.** *If  $\alpha \in (\mathbf{T} \cup \mathbf{N})^*$  can be derived by  $\mathbf{G}^*$ , then one of the following holds.*

- $(\exists x_1, x_2, \dots, x_m \in (\mathbf{T} \cup \mathbf{N})^*)$   
 $((\forall i \in [m])(S \rightarrow^* x_i)) \wedge (\alpha = Zx_1\#x_2\# \dots x_m\#)$
- $(\exists x_1, x_2, \dots, x_m \in (\mathbf{T} \cup \mathbf{N})^*)$   
 $((\forall i \in [m])(S \rightarrow^* x_i)) \wedge (\alpha = R\#x_1\#x_2\# \dots x_m\#)$
- $(\exists w_1, w_2, \dots, w_n \in \mathbf{L})(\exists \beta \in \mathbf{T}^*)(\exists \gamma, x_1, x_2, \dots, x_m \in (\mathbf{T} \cup \mathbf{N})^*)$   
 $((S \rightarrow^* \beta\gamma) \wedge ((\forall i \in [m])(S \rightarrow^* x_i)) \wedge$   
 $(\alpha = w_1w_2 \dots w_n \beta R\gamma \#x_1\#x_2\# \dots x_m\#))$
- $\alpha \in \mathbf{L}^*$
- $(\exists \sigma \in (\mathbf{T} \cup \mathbf{N})^*)(\alpha = \sigma R)$   
*(This case happens if the rule  $R\# \rightarrow R$  is used in the final position (where  $R\# \rightarrow \epsilon$  should be used instead).  $R$  in the final position prevents the derivation from terminating.)*
- $(\exists \omega \in (\mathbf{T} \cup \mathbf{N} \cup \{\#\})^*)(\alpha = \omega\#)$   
*(This case happens if the rule  $R\# \rightarrow \epsilon$  is used too early (that is, anywhere but the final  $\#$  position). The derivation will never terminate in this case.)*

*Proof.* Induction on  $\mathbf{G}^*$  derivation steps. Base case  $\alpha = Z$  holds by  $m = 0$ . Now assume  $Z \rightarrow^* \alpha \rightarrow \alpha'$  and proceed by case analysis.

- $(\exists x_1, x_2, \dots, x_m \in (\mathbf{T} \cup \mathbf{N})^*)$   
 $((\forall i \in [m])(S \rightarrow^* x_i)) \wedge (\alpha = Zx_1\#x_2\#\dots x_m\#)$ 
  - If  $\alpha \rightarrow \alpha'$  used a rule from  $\mathbf{P}$ , it could be applied only in some  $x_i$ . Hence  $S \rightarrow^* x_i \rightarrow x'_i$  so the same condition holds after replacing  $x_i$  by  $x'_i$ .
  - If  $\alpha \rightarrow \alpha'$  used the rule  $Z \rightarrow ZS\#$ , it was applied at the beginning of  $\alpha$ . Therefore, we put  $m' := m + 1$ ,  $x'_1 := S$ , and increase all indices by one, that is,  $x'_2 := x_1$ ,  $x'_3 := x_2$ ,  $\dots$ ,  $x'_{m'} := x_m$ . The same condition holds.
  - If  $\alpha \rightarrow \alpha'$  used the rule  $Z \rightarrow R\#$ , we keep all variables the same and the second condition holds.
  - The rules  $R\# \rightarrow R$ ,  $R\# \rightarrow \epsilon$ , and  $Rt \rightarrow tR$  are not applicable ( $\alpha$  does not contain  $R$ ).
- $(\exists x_1, x_2, \dots, x_m \in (\mathbf{T} \cup \mathbf{N})^*)$   
 $((\forall i \in [m])(S \rightarrow^* x_i)) \wedge (\alpha = R\#x_1\#x_2\#\dots x_m\#)$ 
  - If  $\alpha \rightarrow \alpha'$  used a rule from  $\mathbf{P}$ , it could be applied only in some  $x_i$ . Hence  $S \rightarrow^* x_i \rightarrow x'_i$  so the same condition holds after replacing  $x_i$  by  $x'_i$ .
  - The rules  $Z \rightarrow ZS\#$  and  $Z \rightarrow R\#$  are not applicable ( $\alpha$  does not contain  $Z$ ).
  - If  $\alpha \rightarrow \alpha'$  used the rule  $R\# \rightarrow R$ , it was applied at the beginning of  $\alpha$ . If  $m = 0$ , the fifth condition holds (dead end). Otherwise, we put  $m' := m - 1 \geq 0$ ,  $\gamma := x_1$ , and decrease all indices by one, that is,  $x'_1 := x_2$ ,  $x'_2 := x_3$ ,  $\dots$ ,  $x'_{m'} := x_m$ . As there is nothing before the nonterminal  $R$ , we put  $n := 0$  and  $\beta := \epsilon$ . Now, the third condition holds.
  - If  $\alpha \rightarrow \alpha'$  used the rule  $R\# \rightarrow \epsilon$  then; if  $m = 0$ , we obtain the empty word (which belongs to  $\mathbf{L}^*$ , satisfying the fourth condition); if  $m > 0$ , the last condition holds (because  $\#$  stayed at the end of  $\alpha'$ , at the same time  $R$  disappeared, and  $Z$  did not appear).
  - The rule  $Rt \rightarrow tR$  is not applicable (the only  $R$  in  $\alpha$  is immediately followed by  $\#$ ).

- $(\exists w_1, w_2, \dots, w_n \in \mathbf{L})(\exists \beta \in \mathbf{T}^*)(\exists \gamma, x_1, x_2, \dots, x_m \in (\mathbf{T} \cup \mathbf{N})^*)$   
 $((S \rightarrow^* \beta\gamma) \wedge ((\forall i \in [m])(S \rightarrow^* x_i)) \wedge$   
 $(\alpha = w_1 w_2 \dots w_n \beta R \gamma \# x_1 \# x_2 \# \dots x_m \#))$ 
  - If  $\alpha \rightarrow \alpha'$  used a rule from  $\mathbf{P}$ , it could be applied in  $\gamma$  or in some  $x_i$ .  
In the first case,  $\gamma \rightarrow \gamma'$  implies  $\beta\gamma \rightarrow \beta\gamma'$ , hence  $S \rightarrow^* \beta\gamma \rightarrow \beta'\gamma'$ .  
In the remaining cases, we observe  $S \rightarrow^* x_i \rightarrow x'_i$  as we did at the beginning of our case analysis. As a result, the same condition still holds.
  - The rules  $Z \rightarrow ZS\#$  and  $Z \rightarrow R\#$  are not applicable ( $\alpha$  does not contain  $Z$ ).
  - If  $\alpha \rightarrow \alpha'$  used the rule  $R\# \rightarrow R$ ,  $\gamma$  must have been empty. If  $m = 0$ , the fifth condition holds (dead end). Otherwise, we put  $n' := n + 1$ ,  $w_{n'} := \beta$ ,  $\beta' := \epsilon$ ,  $\gamma' := x_1$ ,  $m' := m - 1$ , and decrease indices of  $x_i$  by one, that is,  $x'_1 := x_2$ ,  $x'_2 := x_3$ ,  $\dots$ ,  $x'_{m'} := x_m$ . Since  $w_{n'} = \beta = \beta\gamma \in \mathbf{T}^*$  and  $S \rightarrow^* \beta\gamma$ , we have  $w_{n'} \in \mathbf{L}$ . The same condition holds.
  - If  $\alpha \rightarrow \alpha'$  used the rule  $R\# \rightarrow \epsilon$ ,  $\gamma$  must have been empty. If  $m = 0$ , we get  $\alpha = w_1 w_2 \dots w_n \beta$  and  $\beta \in \mathbf{L}$ , hence the fourth condition  $\alpha' \in \mathbf{L}^*$  gets satisfied. If  $m > 0$ , the last condition now holds (because  $\#$  stayed at the end of  $\alpha'$ , at the same time  $R$  disappeared, and  $Z$  did not appear).
  - If  $\alpha \rightarrow \alpha'$  used a rule of the form  $Rt \rightarrow tR$  ( $t \in \mathbf{T}$ ), we have  $\delta \in (\mathbf{T} \cup \mathbf{N})^*$  such that  $\gamma = t\delta$ . We put  $\beta' := \beta t$  and  $\gamma' := \delta$ . As  $\beta\gamma = \beta t\delta = \beta'\gamma'$ , the same condition holds.
- $\alpha \in \mathbf{L}^*$ 
  - No rule is applicable ( $\alpha$  contains only terminals). The step  $\alpha \rightarrow \alpha'$  did not happen.
- $(\exists \sigma \in (\mathbf{T} \cup \mathbf{N})^*)(\alpha = \sigma R)$ 
  - No matter which rule was applied, it happened within  $\sigma$ . No rule could match the final  $R$ . The same condition holds for  $\alpha' = \sigma' R$ .

- $(\exists \omega \in (\mathbf{T} \cup \mathbf{N} \cup \{\#\})^*)(\alpha = \omega\#)$ 
  - If  $\alpha \rightarrow \alpha'$  used a rule from  $\mathbf{P}$ , the same condition still holds because the nonterminal  $\#$  is not on the left side of any rule from  $\mathbf{P}$  and neither  $Z$  nor  $R$  is on the right side of any rule from  $\mathbf{P}$ .
  - The rules  $Z \rightarrow ZS\#$ ,  $Z \rightarrow R\#$ ,  $R\# \rightarrow R$ ,  $R\# \rightarrow \epsilon$ , and  $Rt \rightarrow tR$  are not applicable ( $\alpha$  contains neither  $Z$  nor  $R$ ).

□

**Lemma 3.** *Let  $w_1, w_2, \dots, w_n \in \mathbf{L}$ . Then  $\mathbf{G}^*$  can derive  $Zw_1\#w_2\#\dots w_n\#$ .*

*Proof.* Induction on  $n$ . The base case holds because  $Z \rightarrow^* Z$  in zero steps.

Now assume  $Z \rightarrow^* Zw_1\#w_2\#\dots w_n\#$  and  $S \rightarrow^* w_{n+1}$ . We start with the rule  $Z \rightarrow ZS\#$ . We observe  $ZS\# \rightarrow^* Zw_1\#w_2\#\dots w_n\#S\#$  and  $Zw_1\#w_2\#\dots w_n\#S\# \rightarrow^* Zw_1\#w_2\#\dots w_n\#w_{n+1}\#$ . From transitivity, we obtain  $Z \rightarrow^* Zw_1\#w_2\#\dots w_n\#w_{n+1}\#$ . □

*Proof of Theorem 1.* We need to show that the language of  $\mathbf{G}^*$  equals  $\mathbf{L}^*$ .

For “ $\subseteq \mathbf{L}^*$ ”, use Lemma 2 and observe that, if  $\mathbf{G}^*$  generates  $\alpha \in \mathbf{T}^*$ , then  $\alpha \in \mathbf{L}^*$  because all the remaining options require  $\alpha$  to contain a nonterminal.

For “ $\supseteq \mathbf{L}^*$ ”, use Lemma 3. If  $w \in \mathbf{L}^*$ , there are  $w_1, w_2, \dots, w_n \in \mathbf{L}$  such that  $w_1w_2\dots w_n = w$ . We see  $Zw_1\#w_2\#\dots w_n\# \rightarrow R\#w_1\#w_2\#\dots w_n\#$ . Since all  $w_i$  are made of terminals only, by repeated application of  $R\# \rightarrow R$  and  $Rt \rightarrow tR$  (for all  $t \in \mathbf{T}$ ) we get  $R\#w_1\#w_2\#\dots w_n\# \rightarrow^* w_1w_2\dots w_nR\#$ . Finally, we get  $w_1w_2\dots w_nR\# \rightarrow^* w_1w_2\dots w_n = w$  by a single application of  $R\# \rightarrow \epsilon$ . We conclude that  $\mathbf{G}^*$  generates  $w$ . □