

# Magma equations

Matthew Bolan, Jose Brox, Mario Carneiro, Martin Dvořák,  
Andrés Goens, Harald Husum, Zoltan Kocsis, Alex Meiburg,  
Pietro Monticone, David Renshaw, Cody Roux,  
Jérémy Scanvic, Shreyas Srinivas, Anand Rao Tadipatri,  
Terence Tao, Vlad Tsyrklevich, Daniel Weber, Fan Zheng,  
et al.

2025-01-28

## Is there an identity between the commutative identity and the constant identity?

[Ask Question](#)

Asked 1 year, 6 months ago   Modified 2 months ago   Viewed 3k times



**26**



I asked this on Math Stack Exchange, but it didn't get a single answer. So, I am now asking it here. Let our signature be that of a single binary operation  $+$ . I define the constant identity to be  $x + y = z + w$ . The commutative identity is, of course, the well-known identity  $x + y = y + x$ . I wonder, is there an identity strictly between those two, meaning, is there a single identity  $E$  such that the constant identity implies  $E$ , but not conversely, and  $E$  implies the commutative identity, but not conversely?

# Mathoverflow answer — before the project started

Yes:

$$(x + x) + y = y + x$$

The constant identity implies this because both sides are  $+$ es. This does not imply the constant identity because it is true about any set with an operation that is commutative, associative, and idempotent (meaning  $x + x = x$  for all  $x$ ), the smallest nontrivial example is  $\{x, y\}$  where  $x + x = x + y = y + x = x$  and  $y + y = y$ .

This implies commutativity.  $(x + x) + (x + x) = (x + x) + x = x + x$ , so  $x + x$  is idempotent.  $(x + x) + y = ((x + x) + (x + x)) + y = y + (x + x)$ , so  $x + x$  commutes with everything.  $(x + x) + (y + y) = (y + y) + x = x + y$ , and  $(x + x) + (y + y) = (y + y) + (x + x)$  because  $x + x$  commutes with everything, so  $x + y = y + x$ , so  $+$  is commutative.

This is not implied by commutativity, because (for example) addition of natural numbers is commutative but does not satisfy this identity.


Share Cite Improve this answer Follow

answered Jul 16, 2023 at 23:50



paste bee

# Mathoverflow answer — after the project started



The commutative law (43) is the red box on top; the constant law (46) is in the blue box on the bottom

## A pilot project in universal algebra to explore new ways to collaborate and use machine assistance?

---

25 September, 2024 in [math.RA](#), [polymath](#) | Tags: [Artificial Intelligence](#), [Equational Theory Project](#), [machine assisted proof](#), [universal algebra](#) | by [Terence Tao](#)

Traditionally, mathematics research projects are conducted by a small number (typically one to five) of expert mathematicians, each of which are familiar enough with all aspects of the project that they can verify each other's contributions. It has been challenging to organize mathematical projects at larger scales, and particularly those that involve contributions from the general public, due to the need to verify all of the contributions; a single error in one component of a mathematical argument could invalidate the entire project. Furthermore, the sophistication of a typical math project is such that it would not be realistic to expect a member of the public, with say an undergraduate level of mathematics education, to contribute in a meaningful way to many such projects.

# Announcement

**Proposition 1** Equation4 implies Equation7.

*Proof:* Suppose that  $G$  obeys Equation4, thus

$$(x \circ x) \circ y = y \circ x \quad (1)$$

for all  $x, y \in G$ . Specializing to  $y = x \circ x$ , we conclude

$$(x \circ x) \circ (x \circ x) = (x \circ x) \circ x$$

and hence by another application of (1) we see that  $x \circ x$  is idempotent:

$$(x \circ x) \circ (x \circ x) = x \circ x. \quad (2)$$

Now, replacing  $x$  by  $x \circ x$  in (1) and then using (2), we see that

$$(x \circ x) \circ y = y \circ (x \circ x),$$

so in particular  $x \circ x$  commutes with  $y \circ y$ :

$$(x \circ x) \circ (y \circ y) = (y \circ y) \circ (x \circ x). \quad (3)$$

Also, from two applications (1) one has

$$(x \circ x) \circ (y \circ y) = (y \circ y) \circ x = x \circ y.$$

Thus (3) simplifies to  $x \circ y = y \circ x$ , which is Equation7.  $\square$

# Announcement



```
1 import Mathlib.Tactic
2
3 universe u
4
5 class Magma (α : Type u) where
6   op : α → α → α
7
8 infixl:65 " ∘ " => Magma.op
9
10 def Equation4 (G: Type*) [Magma G] := ∀ x y : G, (x ∘ x) ∘ y = y ∘ x
11
12 def Equation7 (G: Type*) [Magma G] := ∀ x y : G, x ∘ y = y ∘ x
13
14 theorem Equation4_implies_Equation7 (G: Type*) [Magma G] (h: Equation4 G) : Equation7 G := by
15   have idem (x : G) : (x ∘ x) ∘ (x ∘ x) = (x ∘ x) := by
16     rw [h x (x ∘ x), h x x]
17   have comm (x y : G) : (x ∘ x) ∘ y = y ∘ (x ∘ x) := by
18     rw [←idem x, h (x ∘ x) y, idem x]
19   have op_idem (x y : G) : (x ∘ x) ∘ (y ∘ y) = x ∘ y := by
20     rw [h x (y ∘ y), h y x]
21   intro x y
22   rw [← op_idem x y, comm x (y ∘ y), op_idem y x]
```

# Easy implication

Prove that

$$\forall x, \forall y, \quad x = x \diamond y$$

implies

$$\forall x, \forall y, \forall z, \forall w, \quad x \diamond y = (x \diamond z) \diamond (w \diamond y)$$

.



# Easy implication

Prove that

$$\forall x, \forall y, \quad x = x \diamond y$$

implies

$$\forall x, \forall y, \forall z, \forall w, \quad x \diamond y = (x \diamond z) \diamond (w \diamond y)$$

.

Solution:

$$x \diamond y = x = x \diamond z = (x \diamond z) \diamond (w \diamond y)$$

# Easy counterexample

Show that

*todo*

does not imply

*todo*

.

# Easy counterexample

Show that

*todo*

does not imply

*todo*

.

Solution:

*todo*

# Why equations?

Howbeit, for easie alteration of equations. I will propounde a few crâples, because the extraction of their rootes, maie the more aptly bee wroughte. And to avoid the tedious repetition of these wordes: is equall to: I will sette as I doe often in worke use, a paire of paraleles, or twenowe lines of one lengthe, thus:  $=====$ , because noe. 2. thynges, can be more equalle. And now marke these numbers.

1.  $14.ze. \text{---} | \text{---} .15.9 \text{=====} 71.9.$

2.  $20.ze. \text{-----} .18.9 \text{=====} 102.9.$

"I will sette as I doe often in worke use, a paire of paraleles, or twin lines of one lengthe, thus:  $=$ , because noe 2 thynges can be more equalle."

Robert Recorde's mathematical book The Whetstone of Witte (1557)

# Why magmas?



# Scope of the project

Order-0 equations:

$$x = x \quad (\text{E1})$$

$$x = y \quad (\text{E2})$$

Order-1 equations e.g.:

$$x = y \diamond z \quad (\text{E7})$$

Order-2 equations e.g.:

$$x \diamond y = y \diamond x \quad (\text{E43})$$

Order-3 equations e.g.:

$$x = (y \diamond x) \diamond (x \diamond z) \quad (\text{E168})$$

Order-4 equations e.g.:

$$(x \diamond y) \diamond z = (u \diamond v) \diamond w \quad (\text{E4694})$$

# Lean and manual vs automated contributions


# Implication graph completion

- ▶ If  $P \implies Q$  and  $Q \implies R$ , then  $P \implies R$ .
- ▶ If  $P \implies Q$  and  $P \not\Rightarrow R$ , then  $Q \not\Rightarrow R$ .
- ▶ If  $Q \implies R$  and  $P \not\Rightarrow R$ , then  $P \not\Rightarrow Q$ .



## Constructivism?

**Intuitionists be like:**



# Dashboard

The implication graph is **99.99986%** complete.

An implication is considered *explicitly true* or *explicitly false* if we have a proof of the corresponding proposition formalised in Lean. It is *implicitly true* or *implicitly false* if the proposition can be derived by taking the reflexive transitive closure of explicitly proven implications.

Our current counts of implications in each of those categories are:

explicitly true	implicitly true	explicitly false	implicitly false	no proof
10,657	8,167,622	586,915	13,268,412	30

The *no proof* column above represents work that we still need to do. Among the *no proof* implications, we have the following conjecture counts:

explicitly true	implicitly true	explicitly false	implicitly false	no conjecture
0	0	9	21	0

The implication graph is **100.00000%** complete if we include conjectures.

# Equation explorer

# Difficult implication

Prove that

$$\forall x, \forall y, \forall z, \quad x = y \diamond ((x \diamond x) \diamond (z \diamond z))$$

implies

$$\forall x_1, \forall y_1, \quad x_1 = y_1$$

.

# Difficult implication

Prove that

$$\forall x, \forall y, \forall z, \quad x = y \diamond ((x \diamond x) \diamond (z \diamond z))$$

implies

$$\forall x_1, \forall y_1, \quad x_1 = y_1$$

.

Solution (found by egg):

let  $x_2 := x_1 \diamond x_1$

let  $x_4 := x_2 \diamond x_2$

let  $x_8 := x_4 \diamond x_4$

let  $y_2 := y_1 \diamond y_1$

let  $y_4 := y_2 \diamond y_2$

let  $y_8 := y_4 \diamond y_4$

# Difficult implication

ass :  $\forall x, \forall y, \forall z, \quad x = y \diamond ((x \diamond x) \diamond (z \diamond z))$

Solution (found by egg):

$x_1 = y_1 \diamond x_4$	ass $x_1 \ y_1 \ x_1$
$= y_1 \diamond (y_1 \diamond (x_8 \diamond x_8))$	ass $x_4 \ y_1 \ x_4$
$= y_1 \diamond (y_1 \diamond (x_1 \diamond x_1))$	ass $x_1 \ x_4 \ x_1$
$= y_1 \diamond (y_8 \diamond (x_1 \diamond x_1))$	ass $y_1 \ y_4 \ y_1$
$= y_4$	ass $y_4 \ y_1 \ x_1$
$= y_1 \diamond (y_8 \diamond (y_1 \diamond y_1))$	ass $y_4 \ y_1 \ y_1$
$= y_1 \diamond (y_1 \diamond (y_1 \diamond y_1))$	ass $y_1 \ y_4 \ y_1$
$= y_1 \diamond (y_1 \diamond (y_8 \diamond y_8))$	ass $y_1 \ y_4 \ y_1$
$= y_1 \diamond y_4$	ass $y_4 \ y_1 \ y_4$
$= y_1$	ass $y_1 \ y_1 \ y_1$

# Metatheorems

# Counterexample constructions

- ▶ Finite magmas
- ▶ Linear models
- ▶ Translation-invariant models
- ▶ Twisting semigroups
- ▶ Greedy constructions
- ▶ Ad-hoc modifications
- ▶ Combinations of the above



# Greedy constructions

## Preliminaries

We want to build a magma operation  $\diamond: M \times M \rightarrow M$  that obeys one equation  $E$  but not another  $E'$ .

We one can first consider *partial magma operations*  $\diamond: \Omega \rightarrow M$  defined on some  $\Omega \subseteq M \times M$ .

We say that a partial operation  $\diamond': \Omega' \rightarrow M$  *extends*  $\diamond: \Omega \rightarrow M$  iff:

- ▶  $\Omega \subseteq \Omega'$
- ▶  $(x, y) \in \Omega \implies x \diamond y = x \diamond' y$

Given a sequence  $\diamond_n: \Omega_n \rightarrow M$  of partial operations, each of which is an extension of the previous, we can define the *direct limit*

$\diamond_\infty: \bigcup_n \Omega_n \rightarrow M$  to be the partial operation defined by  $x \diamond_\infty y := x \diamond_n y$  whenever  $(x, y) \in \Omega_n$ .

# Greedy constructions

## Abstract greedy algorithm

Let  $E$  and  $E'$  be equations. Let  $\Gamma$  be a first-order theory regarding a partial magma operations  $\diamond: \Omega \rightarrow M$  on a carrier  $M$ . Assume the following:

- ▶ (Seed) There exists a finitely supported partial magma operation  $\diamond_0: \Omega_0 \rightarrow M$  satisfying  $\Gamma$  that contradicts  $E'$ , in the sense that there is some assignment of variables in  $E'$  in  $M$  such that both sides of  $E'$  are defined using  $\diamond_0$  but not equal to each other.
- ▶ (Soundness) If  $\diamond_n: \Omega_n \rightarrow M$  is a sequence of partial magma operations obeying  $\Gamma$  with each  $\diamond_{n+1}$  an extension of  $\diamond_n$ , and the direct limit  $\diamond_\infty$  is total, then this limit obeys  $E$ .
- ▶ (Greedy extension) If  $\diamond: \Omega \rightarrow M$  is a finitely supported partial magma operation obeying  $\Gamma$ , and  $a, b \in M$ , then there exists a finitely supported extension  $\diamond': \Omega' \rightarrow M'$  of  $\diamond$  to a possibly larger carrier  $M'$  such that  $a \diamond' b$  is defined.

Then  $E \not\models E'$ .

# Greedy constructions

## Trial and error

1. Start with a minimal rule set  $\Gamma$  that has just enough axioms to imply the soundness property for the given hypothesis  $E$ .
2. Attempt to establish the greedy extension property for this rule set by setting  $a \diamond' b$  equal to a new element  $c \notin M$ , and then defining additional values of  $\diamond'$  as necessary to recover the axioms of  $\Gamma'$ .
3. If this can be done in all cases, then locate a seed  $\diamond_0$  refuting the given target  $E'$ , stop.
4. If there is an obstruction (often due to a collision in which a given operation  $x \diamond' y$  is required to equal two different values), add one or more rules to  $\Gamma$  to avoid this obstruction, and return to Step 2.

# Greedy constructions

Example (E73 does not imply E4380)

Show that the equation

$$x = y \diamond (y \diamond (x \diamond y))$$

does not imply the equation

$$x \diamond (x \diamond x) = (x \diamond x) \diamond x$$

.

# Greedy constructions

Example (E73 does not imply E4380)

Show that the equation

$$x = y \diamond (y \diamond (x \diamond y))$$

does not imply the equation

$$x \diamond (x \diamond x) = (x \diamond x) \diamond x$$

.

►  $y \diamond (x \diamond y) = d \implies y \diamond d = x$

# Greedy constructions

Example (E73 does not imply E4380)

Show that the equation

$$x = y \diamond (y \diamond (x \diamond y))$$

does not imply the equation

$$x \diamond (x \diamond x) = (x \diamond x) \diamond x$$

.

▶  $y \diamond (x \diamond y) = d \implies y \diamond d = x$

▶  $x \diamond y = z \diamond y \implies x = z$

# Greedy constructions

Example (E73 does not imply E4380)

Show that the equation

$$x = y \diamond (y \diamond (x \diamond y))$$

does not imply the equation

$$x \diamond (x \diamond x) = (x \diamond x) \diamond x$$

.

- ▶  $y \diamond (x \diamond y) = d \implies y \diamond d = x$
- ▶  $x \diamond y = z \diamond y \implies x = z$
- ▶  $x \diamond y \neq y$

# Finite magmas

## Finite graph

Some implications are true specifically only for finite magmas.

The finite implication graph is **99.99999%** complete.

explicitly true	implicitly true	explicitly false	implicitly false	no proof
10,750	8,168,349	586,220	13,268,315	2

The finite implication graph is **99.99999%** complete if we include conjectures.

explicitly true	implicitly true	explicitly false	implicitly false	no conjecture
0	0	0	0	2



# About the project

The Atlantic

## We're Entering Uncharted Territory for Math

Terence Tao, the world's greatest living mathematician, has a vision for AI.

By Matteo Wong



# About the project

Substack

## On Math Platform



MICHAEL BUCKO

Okt. 05, 2024



Teilen

I was thinking a combination of *mathematics as a collaborative game (simulations)* and *engineering mathematics*. A platform with the map of mathematics as well as the pull request capability — that could take an idea (or a group of ideas and insights), that kind of seed, modeled as an e-graph (or something similar), given the mathlib-like math knowledge base context, to something that gets formalized, and is fully reusable. With enough compute, one could get agents to work on that thing in the background — it'd become a small island (rather than only a point) in that special embedding space.