

1 Inference rules

1.1 Presuppositivity

MLTT rule:

$$\frac{\Gamma, x:A \vdash B(x) \text{ type}}{\Gamma \vdash A \text{ type}}$$

MLTT rule:

$$\frac{\Gamma \vdash A \equiv B \text{ type}}{\Gamma \vdash A \text{ type}}$$

MLTT rule:

$$\frac{\Gamma \vdash A \equiv B \text{ type}}{\Gamma \vdash B \text{ type}}$$

MLTT rule:

$$\frac{\Gamma \vdash a:A}{\Gamma \vdash A \text{ type}}$$

MLTT rule:

$$\frac{\Gamma \vdash a \equiv b:A}{\Gamma \vdash a:A}$$

MLTT rule:

$$\frac{\Gamma \vdash a \equiv b:A}{\Gamma \vdash b:A}$$

1.2 Judgemental equality is an equivalence relation

MLTT rule:

$$\frac{\Gamma \vdash A \text{ type}}{\Gamma \vdash A \equiv A \text{ type}}$$

MLTT rule:

$$\frac{\Gamma \vdash A \equiv B \text{ type}}{\Gamma \vdash B \equiv A \text{ type}}$$

MLTT rule:

$$\frac{\Gamma \vdash A \equiv B \text{ type} \quad \Gamma \vdash B \equiv C \text{ type}}{\Gamma \vdash A \equiv C \text{ type}}$$

MLTT rule:

$$\frac{\Gamma \vdash a : A}{\Gamma \vdash a \equiv a : A}$$

MLTT rule:

$$\frac{\Gamma \vdash a \equiv b : A}{\Gamma \vdash b \equiv a : A}$$

MLTT rule:

$$\frac{\Gamma \vdash a \equiv b : A \quad \Gamma \vdash b \equiv c : A}{\Gamma \vdash a \equiv c : A}$$

1.3 Variable conversion

MLTT rule:

$$\frac{\Gamma \vdash A \equiv A' \text{ type} \quad \Gamma, x:A, \Delta \vdash B(x) \text{ type}}{\Gamma, x:A', \Delta \vdash B(x) \text{ type}}$$

MLTT rule:

$$\frac{\Gamma \vdash A \equiv A' \text{ type} \quad \Gamma, x:A, \Delta \vdash \mathcal{J}}{\Gamma, x:A', \Delta \vdash \mathcal{J}}$$

Derived rule:

$$\frac{\Gamma \vdash A \equiv A' \text{ type} \quad \Gamma \vdash a : A}{\Gamma \vdash a : A'}$$

1.4 Substitution

Metasyntactic notation (used in many places):

$$t[a/x] := t \text{ with all occurrences of } x \text{ replaced by } a$$

MLTT rule:

$$\frac{\Gamma \vdash a : A \quad \Gamma, x:A, \Delta \vdash \mathcal{J}}{\Gamma, \Delta[a/x] \vdash \mathcal{J}[a/x]} S$$

MLTT rule:

$$\frac{\Gamma \vdash a \equiv a' : A \quad \Gamma, x:A, \Delta \vdash B \text{ type}}{\Gamma, \Delta[a/x] \vdash B[a/x] \equiv B[a'/x] \text{ type}}$$

MLTT rule:

$$\frac{\Gamma \vdash a \equiv a' : A \quad \Gamma, x:A, \Delta \vdash b : B}{\Gamma, \Delta[a/x] \vdash b[a/x] \equiv b[a'/x] : B[a/x]}$$

1.5 Weakening

MLTT rule:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma, \Delta \vdash \mathcal{J}}{\Gamma, x:A, \Delta \vdash \mathcal{J}} W$$

MLTT rule:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash B \text{ type}}{\Gamma, x:A \vdash B \text{ type}} W$$

1.6 Generic elements

MLTT rule:

$$\frac{\Gamma \vdash A \text{ type}}{\Gamma, x:A \vdash x:A} \delta$$

1.7 Manipulating variables

Derived rule:

$$\frac{\Gamma, x:A, \Delta \vdash \mathcal{J}}{\Gamma, x':A, \Delta[x'/x] \vdash \mathcal{J}[x'/x]} x'/x$$

Derived rule:

$$\frac{\Gamma \vdash B \text{ type} \quad \Gamma, x:A, y:B, \Delta \vdash \mathcal{J}}{\Gamma, y:B, x:A, \Delta \vdash \mathcal{J}}$$

2 Π types

2.1 Dependent functions

2.1.1 Π formation

MLTT rule:

$$\frac{\Gamma, x:A \vdash B(x) \text{ type}}{\Gamma \vdash \Pi_{(x:A)} B(x) \text{ type}} \Pi$$

MLTT rule:

$$\frac{\Gamma \vdash A \equiv A' \text{ type} \quad \Gamma, x:A \vdash B(x) \equiv B'(x) \text{ type}}{\Gamma \vdash \Pi_{(x:A)} B(x) \equiv \Pi_{(x:A')} B'(x) \text{ type}} \Pi\text{-eq}$$

2.1.2 Π introduction

MLTT rule:

$$\frac{\Gamma, x:A \vdash b(x) : B(x)}{\Gamma \vdash \lambda x.b(x) : \Pi_{(x:A)} B(x)} \lambda$$

MLTT rule:

$$\frac{\Gamma, x:A \vdash b(x) \equiv b'(x) : B(x)}{\Gamma \vdash \lambda x.b(x) \equiv \lambda x.b'(x) : \Pi_{(x:A)} B(x)} \lambda\text{-eq}$$

2.1.3 Π elimination

MLTT rule:

$$\frac{\Gamma \vdash f : \Pi_{(x:A)} B(x)}{\Gamma, x:A \vdash f(x) : B(x)} \text{ev}$$

MLTT rule:

$$\frac{\Gamma \vdash f \equiv f' : \Pi_{(x:A)} B(x)}{\Gamma, x:A \vdash f(x) \equiv f'(x) : B(x)} \text{ev-eq}$$

2.1.4 Π computation

MLTT rule:

$$\frac{\Gamma, x:A \vdash b(x) : B(x)}{\Gamma, x:A \vdash (\lambda y.b(y))(x) \equiv b(x) : B(x)} \beta$$

MLTT rule:

$$\frac{\Gamma \vdash f : \Pi_{(x:A)} B(x)}{\Gamma \vdash \lambda x.f(x) \equiv f : \Pi_{(x:A)} B(x)} \eta$$

2.2 Ordinary functions

2.2.1 Basic

MLTT rule:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash B \text{ type}}{\Gamma \vdash A \rightarrow B \text{ type}} \rightarrow$$

MLTT rule:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash B \text{ type}}{\Gamma \vdash A \rightarrow B \equiv \Pi_{(x:A)} B \text{ type}}$$

Derived rule:

$$\frac{\Gamma \vdash B \text{ type} \quad \Gamma, x:A \vdash b(x) : B}{\Gamma \vdash \lambda x.b(x) : A \rightarrow B} \lambda$$

Derived rule:

$$\frac{\Gamma \vdash f : A \rightarrow B}{\Gamma, x:A \vdash f(x) : B} \text{ev}$$

Derived rule:

$$\frac{\Gamma \vdash B \text{ type} \quad \Gamma, x:A \vdash b(x) : B}{\Gamma, x:A \vdash (\lambda y.b(y))(x) \equiv b(x) : B} \beta$$

Derived rule:

$$\frac{\Gamma \vdash f : A \rightarrow B}{\Gamma \vdash \lambda x.f(x) \equiv f : A \rightarrow B} \eta$$

The symbol \rightarrow is right-associative:

$$X \rightarrow Y \rightarrow Z \quad := \quad X \rightarrow (Y \rightarrow Z)$$

2.2.2 Identity function

MLTT rule:

$$\frac{\Gamma \vdash A \text{ type}}{\Gamma \vdash \text{id}_A : A \rightarrow A}$$

MLTT rule:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash B \text{ type}}{\Gamma \vdash \text{id}_A \equiv \lambda x.x : A \rightarrow A}$$

2.2.3 Function composition

MLTT rule:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash B \text{ type} \quad \Gamma \vdash C \text{ type}}{\Gamma \vdash \text{comp} : (B \rightarrow C) \rightarrow (A \rightarrow B) \rightarrow (A \rightarrow C)}$$

MLTT rule:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash B \text{ type} \quad \Gamma \vdash C \text{ type}}{\Gamma \vdash \text{comp} \equiv \lambda g.\lambda f.\lambda x.g(f(x)) : (B \rightarrow C) \rightarrow (A \rightarrow B) \rightarrow (A \rightarrow C)}$$

Convention:

$$g \circ f \quad := \quad \text{comp}(g, f) \equiv \lambda x. g(f(x))$$

Derived rule:

$$\frac{\Gamma \vdash f : A \rightarrow B \quad \Gamma \vdash g : B \rightarrow C \quad \Gamma \vdash h : C \rightarrow D}{\Gamma \vdash (h \circ g) \circ f \equiv h \circ (g \circ f) : A \rightarrow D}$$

3 The natural numbers

3.1 Basic

MLTT rule:

$$\frac{}{\vdash \mathbb{N} \text{ type}} \text{N-form}$$

MLTT rule:

$$\frac{}{\vdash 0_{\mathbb{N}} : \mathbb{N}}$$

MLTT rule:

$$\frac{}{\vdash \text{succ}_{\mathbb{N}} : \mathbb{N} \rightarrow \mathbb{N}}$$

3.2 Induction principle

Convention (used in many places):

$$f(x, y) \quad := \quad f(x)(y) \equiv (f(x))(y)$$

Metasyntactic shortcut (only for this document):

$$\textcircled{\mathbb{P}} \quad := \quad \Gamma, n : \mathbb{N} \vdash P(n) \text{ type}$$

MLTT rule:

$$\frac{\textcircled{\mathbb{P}} \quad \Gamma \vdash p_0 : P(0_{\mathbb{N}}) \quad \Gamma \vdash p_S : \Pi_{(n:\mathbb{N})} (P(n) \rightarrow P(\text{succ}_{\mathbb{N}}(n)))}{\Gamma \vdash \text{ind}_{\mathbb{N}}(p_0, p_S) : \Pi_{(n:\mathbb{N})} P(n)} \text{N-ind}$$

MLTT rule:

$$\frac{\textcircled{\mathbb{P}} \quad \Gamma \vdash p_0 : P(0_{\mathbb{N}}) \quad \Gamma \vdash p_S : \Pi_{(n:\mathbb{N})} (P(n) \rightarrow P(\text{succ}_{\mathbb{N}}(n)))}{\Gamma \vdash \text{ind}_{\mathbb{N}}(p_0, p_S, 0_{\mathbb{N}}) \equiv p_0 : P(0_{\mathbb{N}})}$$

MLTT rule:

$$\frac{\textcircled{\text{P}} \quad \Gamma \vdash p_0 : P(0_{\mathbb{N}}) \quad \Gamma \vdash p_S : \Pi_{(n:\mathbb{N})} (P(n) \rightarrow P(\text{succ}_{\mathbb{N}}(n)))}{\Gamma, n : \mathbb{N} \vdash \text{ind}_{\mathbb{N}}(p_0, p_S, \text{succ}_{\mathbb{N}}(n)) \equiv p_S(n, \text{ind}_{\mathbb{N}}(p_0, p_S, n)) : P(\text{succ}_{\mathbb{N}}(n))}$$

3.3 Addition

MLTT rule:

$$\overline{m : \mathbb{N} \vdash \text{add-zero}_{\mathbb{N}}(m) : \mathbb{N}}$$

MLTT rule:

$$\overline{m : \mathbb{N} \vdash \text{add-zero}_{\mathbb{N}}(m) \equiv m : \mathbb{N}}$$

MLTT rule:

$$\overline{m : \mathbb{N} \vdash \text{add-succ}_{\mathbb{N}}(m) : \mathbb{N} \rightarrow \mathbb{N} \rightarrow \mathbb{N}}$$

MLTT rule:

$$\overline{m : \mathbb{N} \vdash \text{add-succ}_{\mathbb{N}}(m) \equiv \lambda n. \text{succ}_{\mathbb{N}} : \mathbb{N} \rightarrow \mathbb{N} \rightarrow \mathbb{N}}$$

MLTT rule:

$$\overline{m : \mathbb{N} \vdash \text{add}_{\mathbb{N}}(m) : \mathbb{N} \rightarrow \mathbb{N}}$$

MLTT rule:

$$\overline{m : \mathbb{N} \vdash \text{add}_{\mathbb{N}}(m) \equiv \text{ind}_{\mathbb{N}}(\text{add-zero}_{\mathbb{N}}(m), \text{add-succ}_{\mathbb{N}}(m)) : \mathbb{N} \rightarrow \mathbb{N}}$$

Derived rule (sugared):

$$\frac{\Gamma \vdash m : \mathbb{N}}{\Gamma \vdash m + 0_{\mathbb{N}} \equiv m : \mathbb{N}}$$

Derived rule (sugared):

$$\frac{\Gamma \vdash m : \mathbb{N} \quad \Gamma \vdash n : \mathbb{N}}{\Gamma \vdash m + \text{succ}_{\mathbb{N}}(n) \equiv \text{succ}_{\mathbb{N}}(m + n) : \mathbb{N}}$$

4 More inductive types

4.1 The unit type

MLTT rule:

$$\overline{\vdash \mathbf{1} \text{ type}}$$

MLTT rule:

$$\overline{\vdash \star : \mathbf{1}}$$

MLTT rule:

$$\frac{\Gamma, x:\mathbf{1} \vdash P(x) \text{ type}}{\Gamma \vdash \text{ind}_\mathbf{1} : P(\star) \rightarrow \Pi_{(x:\mathbf{1})} P(x)}$$

MLTT rule:

$$\frac{\Gamma, x:\mathbf{1} \vdash P(x) \text{ type} \quad \Gamma \vdash p : P(\star)}{\Gamma \vdash \text{ind}_\mathbf{1}(p, \star) \equiv p : P(\star)}$$

Derived rule:

$$\frac{\Gamma \vdash A \text{ type}}{\Gamma \vdash \text{ind}_\mathbf{1} : A \rightarrow (\mathbf{1} \rightarrow A)}$$

Derived rule:

$$\frac{\Gamma \vdash A \text{ type}}{\Gamma \vdash \text{ind}_\mathbf{1} \equiv \lambda a. \lambda x. a : A \rightarrow (\mathbf{1} \rightarrow A)}$$

4.2 The empty type

MLTT rule:

$$\overline{\vdash \emptyset \text{ type}}$$

MLTT rule:

$$\frac{\Gamma, x:\emptyset \vdash P(x) \text{ type}}{\Gamma \vdash \text{ind}_\emptyset : \Pi_{(x:\emptyset)} P(x)}$$

MLTT rule:

$$\frac{\Gamma \vdash A \text{ type}}{\Gamma \vdash \text{ex-falso} : \emptyset \rightarrow A}$$

MLTT rule:

$$\frac{\Gamma \vdash A \text{ type}}{\Gamma \vdash \text{ex-falso} \equiv \text{ind}_\emptyset : \emptyset \rightarrow A}$$

4.3 Negation

MLTT rule:

$$\frac{\Gamma \vdash A \text{ type}}{\Gamma \vdash \neg A \text{ type}}$$

MLTT rule:

$$\frac{\Gamma \vdash A \text{ type}}{\Gamma \vdash \neg A \equiv (A \rightarrow \emptyset) \text{ type}}$$

4.4 Coproducts

MLTT rule:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash B \text{ type}}{\Gamma \vdash A+B \text{ type}}$$

MLTT rule:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash B \text{ type}}{\Gamma \vdash \text{inl} : A \rightarrow A+B}$$

MLTT rule:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash B \text{ type}}{\Gamma \vdash \text{inr} : B \rightarrow A+B}$$

MLTT rule:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash B \text{ type} \quad \Gamma, z : A+B \vdash P(z) \text{ type}}{\Gamma \vdash \text{ind}_+ : (\Pi_{(x:A)} P(\text{inl}(x))) \rightarrow (\Pi_{(y:B)} P(\text{inr}(y))) \rightarrow (\Pi_{(z:A+B)} P(z))}$$

MLTT rule:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash B \text{ type} \quad \Gamma, z : A+B \vdash P(z) \text{ type}}{\Gamma, f : \Pi_{(x:A)} P(\text{inl}(x)), g : \Pi_{(y:B)} P(\text{inr}(y)), a : A \vdash \text{ind}_+(f, g, \text{inl}(a)) \equiv f(a) : P(\text{inl}(a))}$$

MLTT rule:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash B \text{ type} \quad \Gamma, z : A+B \vdash P(z) \text{ type}}{\Gamma, f : \Pi_{(x:A)} P(\text{inl}(x)), g : \Pi_{(y:B)} P(\text{inr}(y)), b : B \vdash \text{ind}_+(f, g, \text{inr}(b)) \equiv g(b) : P(\text{inr}(b))}$$

Derived rule:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash B \text{ type} \quad \Gamma \vdash X \text{ type}}{\Gamma \vdash \text{ind}_+ : (A \rightarrow X) \rightarrow (B \rightarrow X) \rightarrow (A+B \rightarrow X)}$$

Derived rule:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash A' \text{ type} \quad \Gamma \vdash B \text{ type} \quad \Gamma \vdash B' \text{ type}}{\Gamma, f : A \rightarrow A', g : B \rightarrow B' \vdash f + g : A + B \rightarrow A' + B'}$$

Derived rule:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash A' \text{ type} \quad \Gamma \vdash B \text{ type} \quad \Gamma \vdash B' \text{ type}}{\Gamma, f : A \rightarrow A', g : B \rightarrow B', x : A \vdash (f + g)(\text{inl}(x)) \equiv \text{inl}(f(x)) : A' + B'}$$

Derived rule:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash A' \text{ type} \quad \Gamma \vdash B \text{ type} \quad \Gamma \vdash B' \text{ type}}{\Gamma, f : A \rightarrow A', g : B \rightarrow B', y : B \vdash (f + g)(\text{inr}(y)) \equiv \text{inr}(g(y)) : A' + B'}$$

4.5 Σ types

4.5.1 Dependent pairs

MLTT rule:

$$\frac{\Gamma, x : A \vdash B(x) \text{ type}}{\Gamma \vdash \Sigma_{(x:A)} B(x) \text{ type}}$$

MLTT rule:

$$\frac{\Gamma, x : A \vdash B(x) \text{ type}}{\Gamma \vdash \text{pr}_1 : (\Sigma_{(x:A)} B(x)) \rightarrow A}$$

MLTT rule:

$$\frac{\Gamma, x : A \vdash B(x) \text{ type}}{\Gamma, x : A, y : B(x) \vdash \text{pr}_1((x, y)) \equiv x : A}$$

MLTT rule:

$$\frac{\Gamma, x : A \vdash B(x) \text{ type}}{\Gamma \vdash \text{pr}_2 : \Pi_{(p : \Sigma_{(x:A)} B(x))} B(\text{pr}_1(p))}$$

MLTT rule:

$$\frac{\Gamma, x : A \vdash B(x) \text{ type}}{\Gamma, x : A, y : B(x) \vdash \text{pr}_2((x, y)) \equiv y : B(x)}$$

Disclaimer: Many rules about dependent pairs are missing (hopefully, none of them is needed for the homework).

4.5.2 Ordinary pairs

MLTT rule:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash B \text{ type}}{\Gamma \vdash A \times B \text{ type}}$$

MLTT rule:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash B \text{ type}}{\Gamma \vdash A \times B \equiv \Sigma_{(x:A)} B \text{ type}}$$

MLTT rule:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash B \text{ type} \quad \Gamma, x:A, y:B \vdash P(x, y) \text{ type}}{\Gamma \vdash \text{ind}_\times : (\Pi_{(x:A)} \Pi_{(y:B)} P(x, y)) \rightarrow (\Pi_{(z:A \times B)} P(z))}$$

MLTT rule:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash B \text{ type} \quad \Gamma, x:A, y:B \vdash P(x, y) \text{ type}}{\Gamma, g: \Pi_{(x:A)} \Pi_{(y:B)} P(x, y), x:A, y:B \vdash \text{ind}_\times(g, (x, y)) \equiv g(x, y) : P(x, y)}$$

Derived rule:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash B \text{ type}}{\Gamma \vdash \text{pr}_1 : A \times B \rightarrow A}$$

Derived rule:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash B \text{ type}}{\Gamma, x:A, y:B \vdash \text{pr}_1((x, y)) \equiv x : A}$$

Derived rule:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash B \text{ type}}{\Gamma \vdash \text{pr}_2 : A \times B \rightarrow B}$$

Derived rule:

$$\frac{\Gamma \vdash A \text{ type} \quad \Gamma \vdash B \text{ type}}{\Gamma, x:A, y:B \vdash \text{pr}_2((x, y)) \equiv y : B}$$

5 Propositional equality

MLTT rule:

$$\frac{\Gamma \vdash a:A \quad \Gamma \vdash b:A}{\Gamma \vdash a =_A b \text{ type}}$$

MLTT rule:

$$\frac{\Gamma \vdash a:A}{\Gamma \vdash \text{refl}_a : a =_A a}$$

MLTT rule:

$$\frac{\Gamma, x:A, y:A, p : x =_A y \vdash P(x, y, p) \text{ type}}{\Gamma \vdash J : (\Pi_{(x:A)} P(x, x, \text{refl}_x)) \rightarrow (\Pi_{(a:A)} \Pi_{(b:A)} \Pi_{(q:a=_A b)} P(a, b, q))}$$

MLTT rule:

$$\frac{\Gamma, x:A, y:A, p : x =_A y \vdash P(x, y, p) \text{ type} \quad \Gamma \vdash d : \Pi_{(x:A)} P(x, x, \text{refl}_x) \quad \Gamma \vdash a:A}{\Gamma \vdash J(d, a, a, \text{refl}_a) \equiv d(a) : P(a, a, \text{refl}_a)}$$

Derived rule:

$$\frac{\Gamma \vdash p : a =_A b \quad \Gamma \vdash a \equiv a' : A \quad \Gamma \vdash b:B}{\Gamma \vdash p : a' =_A b}$$

Derived rule:

$$\frac{\Gamma \vdash a \equiv b : A}{\Gamma \vdash p : a =_A b}$$